11-3-2007

Understanding Barriers for Adherence to Follow-Up Care for Abnormal Pap Tests

Katherine S. Eggleston
University of Texas Health Science Center at Houston

Ann L. Coker
University of Texas Health Science Center at Houston, ann.coker@uky.edu

Irene Prabhu Das
South Carolina Department of Health & Environmental Control, prabhudasi@mail.nih.gov

Suzanne T. Cordray
University of South Carolina - Columbia

Kathryn J. Luchok
University of South Carolina, luchok@mailbox.sc.edu

Follow this and additional works at: https://uknowledge.uky.edu/crvaw_facpub

Part of the [Obstetrics and Gynecology Commons](#), [Oncology Commons](#), and the [Public Health Commons](#)

[Right click to open a feedback form in a new tab to let us know how this document benefits you.](#)

Repository Citation

Eggleston, Katherine S.; Coker, Ann L.; Prabhu Das, Irene; Cordray, Suzanne T.; and Luchok, Kathryn J., "Understanding Barriers for Adherence to Follow-Up Care for Abnormal Pap Tests" (2007). *CRVAW Faculty Journal Articles*. 110.
https://uknowledge.uky.edu/crvaw_facpub/110

This Article is brought to you for free and open access by the Center for Research on Violence Against Women at UKnowledge. It has been accepted for inclusion in CRVAW Faculty Journal Articles by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Understanding Barriers for Adherence to Follow-Up Care for Abnormal Pap Tests

Digital Object Identifier (DOI)

<http://dx.doi.org/10.1089/jwh.2006.0161>

Notes/Citation Information

Dr. Ann Coker had not been a faculty member of the University of Kentucky at the time of publication.

This is a copy of an article published in the *Journal of Women's Health* © 2007 Mary Ann Liebert, Inc.; *Journal of Women's Health* is available online at: <http://online.liebertpub.com/loi/jwh>

Understanding Barriers for Adherence to Follow-Up Care for Abnormal Pap Tests

KATHERINE S. EGGLESTON, M.S.P.H.,¹ ANN L. COKER, Ph.D.,¹
IRENE PRABHU DAS, M.P.H., Ph.D.,² SUZANNE T. CORDRAY, M.P.H.,³
and KATHRYN J. LUCHOK, Ph.D.⁴

ABSTRACT

Objective: Approximately 4000 women annually will die from preventable and treatable cervical cancer. Failure to adhere to follow-up recommendations after an abnormal Pap test can lead to development of cervical cancer. This paper summarizes the body of literature on adherence to follow-up after an abnormal Pap test in order to facilitate development of interventions to decrease morbidity and mortality due to cervical cancer.

Methods: We conducted a comprehensive search of published literature addressing risk factors for adherence or interventions to improve adherence following an abnormal Pap test as the outcome. We included peer-reviewed original research conducted in the United States from 1990 to 2005.

Results: Fourteen analytical and twelve experimental studies that met our criteria were reviewed. Lesion severity and health beliefs were consistently associated with adherence rates. Communication interventions, including telephone reminders, counseling, and educational sessions, increased follow-up compliance across intervention studies. Inconsistent evidence for associations among race, income, and age were found.

Conclusions: Further research is needed to reinforce current studies addressing health beliefs and social support. Interventions that focus on the interplay among psychological, educational, and communication barriers are necessary. These interventions should be adapted and applied across various racial/ethnic and socioeconomic groups to reach all women with a high-risk profile for invasive cervical cancer.

¹University of Texas Health Science Center, School of Public Health, Houston, Texas.

²South Carolina Department of Health & Environmental Control, Division of Cancer Prevention & Control, Columbia, South Carolina.

³University of South Carolina, Norman J. Arnold School of Public Health, Columbia, South Carolina.

⁴Southern Institute of Children and Families, Columbia, South Carolina.

This research was funded through a cooperative agreement with the Centers for Disease Control and Prevention entitled Network for Understanding Lack of Pap Follow-up, U48/CCU409664-10.

INTRODUCTION

WHEN CERVICAL CANCER IS DETECTED EARLY, the likelihood of survival is almost 100% with appropriate follow-up and treatment.¹ Despite the preventable and treatable nature of this cancer, the American Cancer Society (ACS) estimates that almost 4000 women will have died from cervical cancer in 2006 in the United States.² A key part of this discrepancy is failure to obtain follow-up care after an abnormal Pap test. The Pap test, which screens for precursor lesions to cervical cancer, is one of the most used cancer screening tools currently available; countrywide estimates from the Behavioral Risk Factor Surveillance System (BRFSS) in 2004 report that 78%–90% of women aged >18 years had a Pap test within the preceding 3 years.³

Follow-up diagnostic examinations and, if necessary, treatment are central to the effectiveness of the Pap test in the prevention of invasive cervical cancer.⁴ In a recent study of long-term members of a comprehensive medical care program, 13% of invasive cervical carcinomas were attributable to failure to follow up with abnormal Pap test results.⁵ With the majority of women adhering to Pap test guidelines, more research is needed to determine why women who receive an abnormal result are not complying with necessary follow-up and treatment schedules.

A large body of literature suggests multiple factors are associated with adherence to follow-up recommendations, including factors associated with the patient, both demographic and psychosocial in nature, and with the healthcare system.^{6–9}

The purpose of this systematic review is to summarize reports from recent analytical or experimental research that addressed factors associated with the lack of adequate and timely follow-up care for an abnormal Pap test (nonadherence). The future aim is to facilitate both patient and clinic-based interventions to improve adherence to follow-up recommendations and, thus, reduce needless deaths due to cervical cancer. This is the first review to include both analytical and intervention studies addressing adherence to abnormal Pap test follow-up recommendations and to address risk factors for lack of follow-up by individual, psychosocial, and attributes of the healthcare system.

MATERIALS AND METHODS

A systematic approach to all literature was used to identify original research addressing follow-up of an abnormal Pap test. Our review included academic peer-reviewed sources from the following databases: Medline, Pubmed, Science Direct, Medline-Ovid, Med Science, and Ebsco-Medline. The search terms used were: abnormal Pap, colposcopy, Pap test, screening, gynecologist, or cervical cancer, adherence, follow-up, barriers. Reference lists from papers identified were also reviewed for inclusion of additional papers.

To develop a comprehensive review, we included papers published from 1990 to 2005 regardless of whether they had been included in past reviews. The outcome of interest for all papers reviewed was adherence to recommended follow-up care for an abnormal Pap test. In addition, all papers included in the review had to address risk factors for adherence or involve interventions to improve adherence. A total of 73 abstracts were originally identified. Table 1 outlines the criteria for inclusion in this systematic review and the numbers excluded based on these criteria. We required that each study be evidence based, in that the research involved following women either prospectively or retrospectively through follow-up care after an abnormal Pap test (11 excluded). Studies conducted outside the United States were excluded because different systems of medical care are not comparable to that of the United States ($n = 12$). Papers not published in English were also excluded ($n = 7$). Finally, qualitative research ($n = 4$), recent reviews ($n = 5$), and literature older than 1990 ($n = 8$) were also excluded from this review.

TABLE 1. SELECTION OF PAPERS INCLUDED IN SYSTEMATIC REVIEW

<i>Inclusion criteria</i>	<i>Number excluded</i>
Total studies originally identified	–73
Evidenced based	–11
Conducted in the United States	–12
English language	–7
Quantitative results	–4
Original research	–5
Published since 1990	–8
Total studies reviewed	26

Defining abnormal Pap tests

For the purposes of this review, an abnormal Pap test includes any result that requires additional diagnostic or follow-up procedures. Therefore, the following Pap test results were considered abnormal: insufficient Pap test, infection, atypia, atypical squamous cells of undetermined significance (ASCUS), low-grade squamous intraepithelial lesions (LSIL), high-grade squamous intraepithelial lesion (HSIL), and atypical glandular cells (AGC).

Defining adherence

In this review, definitions of adherence varied across studies, but all were evaluated based on follow-up after an abnormal Pap test result within a time period specified by each healthcare office or governing organization.

RESULTS

Twenty-six papers were identified that met our criteria.^{10–35} All studies were quantitative, defined here as a study analyzing data using statistics that reported a *p* value for correlations and predictors. Table 2 summarizes the key design elements of the 26 quantitative papers that addressed adherence for abnormal Pap test follow-up. The studies are arranged alphabetically by the first author's last name and are presented in two groups by the study design (analytical and experimental studies). Fourteen papers used an analytical study design and addressed risk factors for adherence.^{10–14,18–20,23,27–29,31,34} Twelve studies used an experimental or quasi-experimental design: 9 were randomized trials^{16,17,24–26,30,32,33,35} and 3 were intervention studies based on nonrandomized designs.^{15,21,22} Table 2 provides the study setting and sample, the number of subjects, the data source and list of independent variables assessed, and the data source and operational definition of the outcome (adherence to follow-up recommendations for an abnormal Pap test).

Study sites included public health and community clinics,^{11,14,18,19,27–29} hospitals,^{13,31} academic clinics,^{12,23,34} the National Breast and Cervical Cancer Early Detection Program (NBCCEDP),¹⁰ and a large consortium of laboratories.²⁰ The sample sizes ranged from 119²³ to 16,132 women.²⁰

With the exception of four studies that relied on self-report through telephone interviews and questionnaires,^{15,27,30,31} all used medical records to obtain data on independent variables. All studies addressed demographic factors as independent variables correlated with adherence; among these were age, race/ethnicity, insurance, marital status, education, income, place of residence, primary language spoken, tobacco use, and cervical lesion severity.

The reviewed research used medical records or self-report to determine adherence and defined adherence as the receipt of recommended follow-up care. However, studies differed in the definition of adherence. Most studies used a restricted time frame^{10–12,15,16,18,20,21,23,26,28–30,34,35} that ranged from receipt of follow-up care within 4–6 weeks³⁵ to receipt of follow up care within 18 months.¹⁶ Other studies measured adherence by the number of follow-up appointments kept,¹⁴ ever completing the recommended procedure,^{19,22,24,25,27,31} or adherence within the time frame set by the physician.^{32,33}

Table 3 lists the measures of association, a summary of the study findings, and adherence rates. Adherence rates for the analytical studies ranged from 27%²⁸ to 90%,¹⁹ whereas those of the experimental studies ranged from 40%²¹ to 93%¹⁵ for those receiving the intervention. Nine of twelve intervention studies^{15–17,24–26,30,32,35} reported higher adherence among those participating in the intervention. The findings are summarized by patient, healthcare system, and social/environmental characteristics.

Patient characteristics

Age. Of the 14 studies^{10,11,14,16–18,20,23–27,31,34} that addressed age and adherence, 7 found that younger women were less likely to receive follow-up care.^{16,17,20,25–27,34} Specifically, McKee et al.²⁷ reported that teenagers and women older than 30 were less likely to receive follow-up care. Additionally, a study conducted by Fox et al.¹⁸ reported that increased age was a significant risk factor for noncompliance.

Race/ethnicity. Of the 10 analytical^{10–12,14,18,23,27,28,31,34} and 7 experimental^{16,17,24–26,30,32} studies to address race/ethnicity and adherence,^{7,10,11,17,18,25,26,32} found that African American women were less likely to schedule follow-up visits, to keep appointments,

TABLE 2. DESCRIPTIONS OF ORIGINAL RESEARCH ADDRESSING ADHERENCE TO FOLLOW-UP CARE FOR ABNORMAL PAP TEST (N = 26)

<i>Author, date</i>	<i>Setting/sample</i>	<i>Study design</i>	<i>Sample size</i>	<i>Independent variables source and list</i>	<i>Outcome source and definition</i>
Analytical studies addressing risk factors for adherence					
Bernard et al., 2005 ¹⁰	Nationwide, women with 2 abnormal Pap tests within 2 years enrolled in the NBCCEDPa, 1991–2000	Retrospective cohort	10,004	Source: Medical record Variables: Age, race, lesion severity	Source: Medical record Outcome: Followed up within the 2-year observation period
Cardin et al., 2001 ¹¹	Houston, TX Health Department, women with abnormal Pap test, 1996–2000	Retrospective cohort	1,216, 24% HSIL, 76% LSIL	Source: Medical record Variables: Age, race, lesion severity.	Source: Medical record Outcome: Completed referral evaluation/currently evaluated/treated within 9 months of abnormal Pap test Source: Medical record Outcome: Followed up by 1992
Carey and Gjerdingen, 1993 ¹²	St. Paul, University of Minnesota Family Practice Clinic, women with an abnormal Pap test; 1989–1992	Retrospective cohort	190	Source: Medical record Variables: Race.	Source: Medical record Outcome: Followed up by 1992
Crane, 1996 ¹³	Los Angeles, CA, two county hospitals, women with abnormal Pap test, 1986–1988	Cohort	498	Source: Medical record Variables: Social support, knowledge.	Source: Medical record Outcome: Followed up within 4 months of abnormal Pap test
Eger and Peipert, 1996 ¹⁴	Providence, RI, Women and Infants Colpocopy Clinic, women referred for colposcopy, 1992–1993	Retrospective case-control	179	Source: Medical record Census Variables: Age, race, insurance, lesion severity, marital status, tobacco use, gravidity, parity, income (ZIP code)	Source: Medical record Outcome: Number of appointments kept or missed
Fox et al., 1997 ¹⁸	CA breast and cervical cancer control program, women with abnormal Pap test, 1994	Retrospective cohort	1,738	Source: Medical record Variables: Age, race, lesion severity, rural residence.	Source: Medical record Outcome: Followed up at least 60 days after abnormal Pap test
Hartz and Fenaughty, 2001 ¹⁹	Anchorage, AK reproductive health clinic, low-income women with CIN1, 1995–1999	Cohort	219	Source: Medical record Variables: Management choice (cryotherapy vs. serial Pap), income.	Source: Medical record Outcome: Completion of follow-up
Jones and Novis, 2000 ²⁰	American College of Pathologists Q-Probes program, women with abnormal cytology, 1998	Retrospective cohort	16,132; 306 laboratories; 60 cases per laboratory	Source: Medical record Variables: Age, race, lesion severity, pregnancy status, laboratory characteristics	Source: Medical record Outcomes: Followed up within 1 year of abnormal Pap test
Lavin et al., 1997 ²³	Boston, MA departments of pediatric and obstetrics and gynecology, adolescents with abnormal Pap tests, 1994–1995	Retrospective cohort	119	Source: Medical record Variables: Age, race, insurance, lesion severity, previous referral	Source: Medical record, telephone Outcome: Completion of follow-up

McKee et al., 1999 ²⁷	Bronx, NY, Montefiore Family Health Center, women with an abnormal Pap test; 1993–1994	Telephone survey	279	Source: Questionnaire Variables: Age, race, insurance, lesion severity, education, communication, psychological, provider, logistics Source: Medical record Variables: Race, insurance, lesion severity, pregnancy status, provider training, colposcopy on site, discussion of care plan Source: Medical record Variables: Lesion severity, insurance, language, proximity to clinic	Source: Telephone survey, medical record Outcome: Completion of colposcopy
McKee et al., 2001 ²⁸	7 urban community health centers, women with abnormal Pap test, 1996	Retrospective cohort	387		Source: Medical record Outcome: Percent of women receiving colposcopy or repeat Pap tests every 6 months until 3 are normal
Melnikow et al., 1999 ²⁹	CA Family Planning Clinics, women with abnormal Pap tests referred for colposcopy, 1994–1996	Retrospective cohort	243		Source: Medical record Outcome: Followed up within 8 months of abnormal Pap test
Nelson et al., 2002 ³¹	Los Angeles, CA, Kaiser Permanente Medical Center, women with abnormal Pap test, 1998–1999	Cohort	1,049	Source: Mail/phone survey Variables: Age, race, education, income, health beliefs, cancer knowledge	Source: Medical record Outcome: Delay in first scheduled clinic visit for follow-up or multiple contacts needed to schedule follow-up appointment
Peterson, et al., 2003 ³⁴	Boston, MA urban academic clinics, women with abnormal Pap test, 1999–2000	Retrospective cohort	423	Source: Medical record Variables: Age, race, lesion severity, insurance, primary language, provider type, source of care	Source: Medical record/pathology Outcome: Followed-up within 4 months of abnormal Pap test for HSIL and 7 months for LSIL
Experimental studies addressing risk factors for adherence Ell et al., 2002 ¹⁵	Los Angeles, CA, Screening Adherence Follow-up Program (SAFe), women with LSIL/HSIL, 1998–2000	Intervention	196	Source: Phone questionnaire Variables: Access to care, knowledge, health beliefs, depression, anxiety. Intervention: Stepped care with mental health assessment (case management addressing reminders, knowledge, psychosocial factors, and system barriers)	Source: Medical record report Outcome: Completed clinic appointments during 1-year study period (partial, full, or nonadherent)
Engelstad et al., 2001 ¹⁶	Oakland, CA, Alameda County Medical Center ER, women 18–74 with an abnormal Pap test, October 1993–June 1995	Randomized trial	108	Source: Medical record Variables: Age, race, insurance status, lesion severity Intervention: Case management (reminder protocol) and universal colposcopy vs. traditional care.	Source: Medical record Outcome: Followed up within 6 months of abnormal Pap test and completed diagnostic resolution in 18 months

(continued)

TABLE 2. DESCRIPTIONS OF ORIGINAL RESEARCH ADDRESSING ADHERENCE TO FOLLOW-UP CARE FOR ABNORMAL PAP TEST (N = 26) (CONT'D)

<i>Author, date</i>	<i>Setting/sample</i>	<i>Study design</i>	<i>Sample size</i>	<i>Independent variables source and list</i>	<i>Outcome source and definition</i>
Engelstad et al., 2005 ¹⁷	Oakland, CA, Alameda County Medical Center ER, women 18–74 with abnormal Pap test, September 1999–August 2001	Randomized trial status	348	Source: Medical record Variables: Age, race, insurance status, lesion severity, language, obstetrical Intervention: Community health advisor (addressing reminders, knowledge, psychosocial factors) Source: Medical record and data system Intervention: Tracking follow-up, transportation and financial incentives	Source: Medical record Outcome: Initial follow-up within 6 months of abnormal Pap test
Kaplan et al., 2000 ²¹	Los Angeles, CA, hospitals and health care centers, women with abnormal Pap test, 1989–1990	Intervention	4,488	Source: Medical record Intervention: Tracking follow-up, transportation and financial incentives	Source: Medical record/data system Outcome: Followed up within 9 months of abnormal Pap test; categorized as complete, partial, or no follow-up
Lacey et al., 1993 ²²	Chicago, IL community-based health clinic, women with abnormal Pap test, 1989–1990	Intervention	101	Source: Medical record Intervention: Referral to public hospital vs. private physician	Source: Telephone call to private physician OR computerized tracking system Outcome: Completed scheduled follow-up appointment
Lerman et al., 1992 ²⁴	Philadelphia, PA, Temple University Hospital, low-income minority women who missed a colposcopy	Randomized trial	90	Source: Medical record Variables: Age, race, marital status, education Intervention: Telephone counseling (addressing knowledge, psychosocial factors, cost, transportation)	Source: Medical records, validated by telephone interview Outcome: Compliance with rescheduled colposcopy
Marcus et al., 1992 ²⁵	Los Angeles, CA, 12 primary healthcare clinics, women with abnormal Pap test, 1984–1986	Randomized trial	2,044	Source: Medical record Variables: Age, race, insurance, lesion severity, marital status, education Interventions (1) letter (2) tape on Pap tests (3), transportation incentives	Source: Medical record, telephone interview, mail questionnaire Outcome: At least one return visit vs. no return visit
Marcus et al., 1998 ²⁶	Los Angeles, CA, County Department of Health Services, women with abnormal Pap test	Randomized trial	1,453	Source: Medical record Variables: Age, race, lesion severity, live-in relationship, insurance, hospital Interventions: (1) reminder+ pamphlet (2) voucher	Source: Medical record and telephone interview Outcome: Followed-up within 4– 6 months after abnormal Pap test

Miller et al., 1997 ³⁰	Philadelphia, PA, Temple University Hospital, women with abnormal Pap test, 1992– 1995	Randomized trial	828	Source: Telephone interview Variables: Race, marital status, education. Intervention: Telephone counseling and reminder protocol (addressing knowledge, psychosocial factors)	Source: Sign-in list at clinic Outcome: Followed up within 6 months after abnormal Pap test
Paskett et al., 1990 ³³	Seattle, WA, Women's Care Center, women with abnormal Pap tests, 1987–1988	Randomized trial	161	Source: Medical record Intervention: Pamphlet	Source: Medical record Outcome: Obtained repeat Pap by date specified by physician
Paskett et al., 1995 ³²	6 clinics: 2 family planning, 2 dysplasia, 2 family practice	Randomized trial	541	Source: Medical record Variables: Race, lesion severity, number of children, smoking status Intervention: Motivational brochures	Source: Medical record Outcome: Followed-up within 1 week of date specified by physician
Takacs et al., 2004 ³⁵	Miami-Dade County, FL colposcopy clinic, women with abnormal Pap test, June 2002– December 2002	Randomized trial	60	Source: Medical record Intervention: Video or traditional colposcopy	Source: Medical record Outcome: Followed-up within 4– 6 weeks after abnormal Pap test

^aHSIL, high-grade squamous intraepithelial lesion; LSIL, low-grade squamous intraepithelial lesion; STI, sexually transmitted infection; NBCCEDP, National Breast and Cervical Cancer Early Detection Program.

TABLE 3. SUMMARY OF RESULTS OF ORIGINAL RESEARCH ADDRESSING ADHERENCE TO RECOMMENDATIONS FOR FOLLOW-UP CARE FOR ABNORMAL PAP TEST (N = 26)

<i>Author, date</i>	<i>Measure of association and confidence interval or p value</i>	<i>Summary of findings</i>	<i>Adherence rate</i>
Analytical studies addressing risk factors for adherence			
Bernard et al., 2005 ¹⁰	Adherence After 2 abnormal Pap tests; race: $p = 0.01$ ASCUS-ASCUS: ^a 23%, no follow-up ASCUS-LSIL: 12.4%, no follow-up	African Americans had a higher percentage of no-follow up compared with other racial groups Those with ASCUS on second Pap test result were less likely to follow up than those with LSIL	72.3%
Cardin et al., 2001 ¹¹	Appointment scheduling African American vs. other race: OR = 0.47, 0.33–0.67	African American women with ASCUS/LSIL were less likely to be compliant than women of other races Women with LSIL were less likely to be compliant than women with HSIL	62%
Carey and Gjerdingen, 1993 ¹²	Adherence African American vs. other race: OR = 0.55, 0.41–0.75 Follow-up rate difference Asian vs. whites: $p < 0.001$; Asian vs. African American: $p < 0.05$	Southeast Asian women were less likely than white and African American women to comply with follow-up	86.3% diagnostic 79.5%: treatment
Crane, 1996 ¹³	Adherence By support: any social support: OR = 3.12, $p < 0.001$ By knowledge: Informational support, OR = 1.88, $p = 0.03$ Knowledge of Pap test: OR = 1.92, $p = 0.003$ By race/ethnicity All: Emotional support, $p = 0.01$ Latinas: Tangible support, $p < 0.05$ African Americans: Emotional support, $p < 0.05$	Women who received informational, emotional, and/or tangible support were more likely to be compliant Women who had less Pap test knowledge were less likely to adhere to recommendations Latinas who received tangible support and African Americans who received emotional support were more likely to comply	
Eger and Peipert, 1996 ¹⁴	Adherence Lesion grade: OR = 0.34, $p = 0.01$	Women who were noncompliant were less likely to have high-grade histology than women who were compliant Age, race, insurance, marital status, tobacco use, gravidity, parity, income were not associated with adherence	77%
Fox et al., 1997 ¹⁸	Nonadherence Urban vs. rural, OR = 1.73, $p < 0.006$ African American vs. white, OR = 3.80, $p < 0.001$ Asian vs. white, OR = 1.94, $p < 0.009$ Hispanic vs. white, OR = 1.84, $p < 0.0001$ SIL vs. negative, OR = 9.43, $p > 0.001$ Age 65+ vs. >30, OR = 3.03, $p > 0.001$	Increased age, nonwhite race/ethnicity, urban setting, and severity of Pap test result are significant predictors of not completing follow-up	35.6%
Hartz and Fenaughty, 2001 ¹⁹	Adherence First follow-up visit Surveillance vs. cryotherapy, $p = 0.01$ High income vs. low income, $p = 0.008$	Patients who chose surveillance only were more likely to adhere to initial recommendations but not for long-term follow-up Women who had highest mean income had highest completion of follow-up rate	90%: 1st visit after colposcopy 37%: complete

Jones and Noves, 2000 ²⁰	Adherence HSIL/CIN vs. LSIL/GIL, $p = 0.001$ Pregnant < nonpregnant, $p < 0.01$	Women with HSIL/CIN were more likely to receive follow-up than women with LSIL or GIL Younger and pregnant patients were less likely to receive follow-up	85%: 12 months 77%: 6 months 65%: 3 months 61.7%
Lavin et al., 1997 ²³	Adherence Adolescent clinic visit vs. no visit, 79% vs. 45%, $p = 0.007$	Visit to an adolescent clinic predicted adherence No differences in race, insurance status, age, Pap test results, recommendations for previous colposcopy or Pap tests for compliers vs. noncompliers	
McKee et al., 1999 ²⁷	Adherence rate Age, $p = 0.02$; Pap test severity, $p = 0.06$ Knowledge of results, $p = 0.001$ Recall receiving letter, $p = 0.001$ Doctor understands my needs, $p = 0.03$	Teenagers were less likely to receive follow-up care, as well as women older than 30 Lack of effective communication was associated with low rates of return for colposcopy Pap test severity, ethnicity, insurance, and education were not significant predictors of adherence No psychological, logistical, or provider characteristics were associated with barriers to colposcopy Severity of lesion, availability of colposcopy on site, discussion of plan and site predicted adherence Ethnicity, insurance, provider training, and pregnancy were not associated with adherence	75%
McKee et al., 2001 ²⁸	Adherence ASCUS vs. atypia, OR = 1.56, $p < 0.05$ Colposcopy available vs. not, OR = 0.39, $p < 0.01$ Discussed plan vs. not, OR = 3.44, $p < 0.01$	Women with LSIL/HSIL were less likely to adhere to colposcopy compared with repeat Pap tests Women with ASCUS were more likely to adhere to colposcopy compared with repeat Pap tests Women with ASCUS were less likely to adhere than all others to any appointment	27%: complete 28%: moderate
Melnikow et al., 1999 ²⁹	Adherence Colposcopy vs. repeat Pap test LSIL/HSIL, HR = 0.37 (0.14–0.99) ASCUS, HR = 2.67 (1.22–5.86) Adherence to any appointment LSIL/HSIL vs. ASCUS, HR = 3.59 (1.40–9.25) No insurance, HR = 0.43 (0.20–0.93) Delay in care Need a Pap test only with abnormal bleeding (yes/no), OR = 1.1 (1.1–15.5) Fatalism score/point increase, OR = 1.1 (1.0–1.2)	Women with LSIL/HSIL were less likely to adhere to colposcopy compared with repeat Pap tests Women with ASCUS were more likely to adhere to colposcopy compared with repeat Pap tests Women with ASCUS were less likely to adhere than all others to any appointment Women with no insurance were less likely to follow up Proximity to clinic and language did not predict adherence When fatalism and health beliefs were included in model, race/ethnicity was not related to delay in care Women who delayed care had higher mean fatalism score Women who believe a Pap test is needed only with abnormal bleeding were more likely to delay care Income, education, and race were not associated with delays in care	56%
Nelson et al., 2002 ³¹	Inadequate follow-up Adjusted OR (95% CI) Age 18–29 vs. $\geq 50 = 2.7$ (1.1–6.4) Medicaid vs. HMO = 1.9 (1.0–3.5) Uninsured vs. HMO = 1.6 (0.9–2.9) Adjusted HR, ≤ 30 years = 0.6 (0.4–0.9) Uninsured = 0.8 (0.6–1.0) Medicaid = 0.8 (0.6–1.0)	Young age and either having Medicaid or being uninsured was a significant predictor of inadequate follow-up Neither race/ethnicity, specialty of provider, language, nor lesion severity were significant predictors of inadequate follow-up	87%: timely 13%: delayed
Peterson et al., 2003 ³⁴	Adherence Intervention LSIL-dichotomous: OR = 3.14 (1.5–7.7)	Women enrolled in program had significantly better adherence rates than nonenrollees Barriers to care cited included fear of finding cancer, costs,	62%
Experimental studies addressing risk factors for adherence Ell et al., 2002 ¹⁵			83%: overall 93%: HSIL (continued)

TABLE 3. SUMMARY OF RESULTS OF ORIGINAL RESEARCH ADDRESSING ADHERENCE TO RECOMMENDATIONS FOR FOLLOW-UP CARE FOR ABNORMAL PAP TEST (N = 26) (CONT'D)

<i>Author, date</i>	<i>Measure of association and confidence interval or p value</i>	<i>Summary of findings</i>	<i>Adherence rate</i>
Engelstad et al., 2001 ¹⁶	Incremental adherence: 41% full, 42% partial, 17% nonadherent HSIL-dichotomous: OR = 6.5 (2.5–16.9) Incremental adherence: 61% full, 42% partial, 17% nonadherent Adherence Intervention: OR = 3.98 (1.63–9.74) Age: OR = 1.08 (1.02, 1.13) Race: Asian, OR = 0.16 (0.03–0.85) No insurance: 2.78 (1.00–7.71) Adherence and resolution Intervention: OR = 6.53 (2.39–17.84) Race: Asian, OR = 0.06 (0.01–0.61) HG/LGSIL vs. ASC/AGUS: OR = 4.24 (1.21–14.81)	worries about examination treatment, lack of understanding of follow-up examination, and transportation or child care problems The proportion of women who received initial follow-up or diagnostic resolution was greater in the interventional arm Older women, those without insurance, and non-Asian women were more likely to follow up in 6 months Non-Asian women and all women with less severe lesions (ASCUS/AGUS) were more likely to follow up in 6 months and have diagnostic resolution in 18 months	50%: intervention 19%: control
Engelstad et al., 2005 ¹⁷	Adherence Overall intervention: OR = 3.33, $p = 0.001$ Age: OR = 1.04, $p = 0.001$ Race: white; OR = 3.3, $p = 0.001$; Asian: OR = 0.3, $p = 0.03$	Adherence to follow-up was greater in the intervention arm overall and by all lesion types Older women and white women were more likely to follow-up than African Americans; Asians were less likely to follow-up than African Americans Language, lesion severity, insurance, Hispanic race, and obstetrical status were not associated with adherence No significant difference in follow-up among control and intervention groups by location of treatment: hospitals, CHC, or PHC	60.7%: intervention 32.4%: control
Kaplan et al., 2000 ²¹	Adherence: Baseline vs. all intervention years Hospital, OR = 2.74 (0.75–10.00) CHC, OR = 2.28 (0.47–11.13) PHC, OR = 1.61 (0.17–14.90)		40%: intervention 35.6%: control
Lacey et al., 1993 ²²	Adherence Rate of adherence: public hospital vs. private physician, $p > 0.05$	No difference in follow-up rates occurred between women who were referred to public hospital and those who chose to follow-up with their private physician	69%: overall 70%: public hospital 57%: private 43%: control 67%: intervention
Lerman et al., 1992 ²⁴	Adherence Intervention: OR = 2.6 $p < 0.003$	No SES factors were associated with adherence; telephone intervention improved adherence for colposcopy Half of all nonadherer's reported the reason for missing appointment was not knowing purpose of test African American, Hispanic, younger patients; patients with less than HS education, those with no insurance, using a county facility, or those with a less severe Pap test result were less likely to return for follow-up care Transportation incentives had significant impact on county patients, those with more severe Pap tests, and noninsured patients	
Marcus et al., 1992 ²⁵	Adherence Race/ethnicity: $p < 0.02$; age: $p < 0.001$; education: $p < 0.01$; insurance: $p < 0.01$; health care: $p < 0.001$; severity of Pap test: $p < 0.001$ County patient: incentive: OR = 1.51 (1.06–2.15)		Ranged 56%–87% based on clinic site 71%: overall

Marcus et al., 1998 ²⁶	Noncounty patients: personalized follow-up with slide-tape program, OR = 4.54 (1.12–18.40) Severe Pap tests: incentive: OR = 1.71 (1.10–2.67) Less severe Pap test: personalized follow-up with slide-tape program, OR = 5.16 (1.47–18.04) No insurance: incentive OR = 1.77 (1.19–2.65) Risk of returning for follow-up care Follow-up intervention: OR = 1.56 (1.12–2.17) Voucher intervention: OR = 01.50 (1.09–2.05) Age: OR = 1.03 (1.02–1.05) Live-in relationship (yes/no): OR = 1.54 (1.22–1.95) African American vs. white: OR = 0.49 (0.27–0.90) HSIL vs. LSIL: OR = 1.64 (1.24–2.17) Adherence Standard care vs. telephone confirmation: OR = 0.47 (0.32–0.73) Telephone counseling vs. telephone confirmation: OR = 1.50 (1.04–2.17) Forgets appointments: OR = 0.31 (0.19–0.51) Scheduling conflicts: OR = 0.45 (0.25–0.72)	Personalized follow-up with slide-tape program had significant impact on noncounty patients and those with less severe Pap test results	70%
		Interventions had significant and independent effect on follow-up rates African American women were less likely to return for follow-up care than white women Women in a live-in relationship vs. no live-in relationship and those with HSIL vs. LSIL were more likely to return for follow-up care Younger women were less likely than older women to return for follow-up care Insurance and Hispanic race were not predictor of adherence Women with standard care were less likely to adhere than those who had telephone confirmation; those with telephone confirmation were less likely to adhere than those with telephone counseling Forgetting appointments and conflicts were both independently associated with nonadherence No socioeconomic variables were associated with adherence	
Miller et al., 1997 ³⁰			60%
Paskett et al., 1990 ³³	Adherence Compliance rate: control = 51%, intervention = 64%, OR = 1.71 (0.91–3.20) $p = 0.10$ (two tailed)	Intervention pamphlets increased compliance in 13% of women who would have been noncompliers, but difference was not statistically significant at the $p = 0.05$ level	51.3%: control 64.2%: intervention
Paskett et al., 1995 ³²	Adherence Intervention: OR = 2.6, 1.2–5.6 Dysplasia vs. atypia: OR = 4.2, 1.4–12.3 White vs. African American: OR = 3.0 (1.1–7.9) Nonsmoker vs. smoker: OR = 2.5 (1.3–5.7) Nulliparous: OR = 2.9 (1.1–8.2)	Significant predictors of adherence were being in intervention clinic, having dysplasia, and being white, a nonsmoker, and nulliparous	33.8%–69.3%: atypia 33.3%–87.5%:
Takacs et al., 2004 ³⁵	Adherence Rate of follow-up: video vs. resular colposcopy = 80% vs. 50%, $p = 0.014$, adjusted OR = 4.9, 95% CI = 1.2–19.4	Patients undergoing video colposcopy were almost 5 times more likely to return for follow-up than patients undergoing regular colposcopy	50%: control 80%: intervention

^aAGUS, atypical glandular cells of undetermined significance; ASCUS, atypical squamous cells of undetermined significance; CHC, community health clinics; CIN, cervical intraepithelial neoplasia; GIL, glandular epithelial lesion; HMO, health maintenance organization; HSIL, high-grade squamous intraepithelial lesion; LSIL, low-grade squamous intraepithelial lesion; OR, odds ratio; PHC, provider health clinics; SES, socioeconomic status.

or to receive follow-up care. Specifically, Cardin et al.¹¹ reported that African American women with ASCUS/LSIL were less likely to be compliant than African American women with HSIL. Four studies^{12,16–18} reported that Asian women were less likely than women of other race/ethnicities (whites or African Americans) to comply with follow-up recommendations. Finally, two studies found that Hispanics were significantly less likely to adhere to follow-up recommendations than white¹⁸ and African American women.²⁵ However, the only study to address fatalism and health beliefs,³¹ reported that race/ethnicity was no longer associated with adherence after adjusting for these factors.

Primary language. All three studies that investigated a primary language other than English as a barrier to follow-up care^{17,29,34} concluded that this demographic was not associated with adherence.

Place of residence. Two studies examined place of residence by assessing urban vs. rural residence¹⁸ and proximity to clinic.²⁹ Proximity to clinic was not a predictor of adherence,²⁹ but urban setting was a significant predictor of nonadherence.¹⁸

Education. Five studies investigated educational attainment, measured by receipt of a high school diploma,^{24,25,27,30,31} as a predictor of adherence to follow-up care. Only one study found that those with less than a high school education were less likely to return for follow-up care.²⁵

Pregnancy status. Three studies addressed pregnancy status and adherence^{17,20,28}; only and Novis²⁰ reported that pregnant women were less likely to receive timely follow-up. Gravidity and parity were not associated with adherence in one study,¹⁴ although being nulliparous was positively associated with being adherent in another.³²

Tobacco use. Two studies that assessed tobacco use as a predictor of adherence to follow-up found conflicting results. Eger and Peipert¹⁴ found that tobacco use was not associated with the number of follow-up appointments kept, whereas, Paskett et al.³² found that nonsmokers were more likely than smokers to follow up within 1 week of the date specified by the physician.

Income/insurance/cost of follow-up. Thirteen studies addressed income,^{14,19,31} insurance,^{14,16,17,23,25–29,34}

or cost as barriers to receiving follow-up care.^{21,25,26} Four found that those with higher income¹⁹ or private insurance^{25,29,34} were more likely to adhere to recommended follow-up. Conversely, Engelstad et al.¹⁶ found that women with no insurance were more likely to have a follow-up visit in 6 months than those with insurance. The remaining eight studies found no association between adherence and income^{14,21,31} or insurance.^{14,17,23,26–28} Of three interventions^{21,25,26} to address economic strain, however, two reported that transportation incentives²⁵ and economic vouchers had a significant impact on adherence.²⁶ Furthermore, cost, transportation, and child care problems were among the most frequent barriers reported by others.^{15,24,30}

Knowledge of Pap test. Several studies addressed the influence of Pap test knowledge on adherence to follow-up.^{13,15,17,24–26,30–33} Both Crane¹³ and Nelson et al.³¹ found that women who did not know the purpose of a Pap test were less likely to adhere to recommendations than those who correctly identified the purpose. Two studies that investigated barriers to care reported that lack of understanding of the purpose of a follow-up examination was a reason for nonadherence.^{15,24} Studies that used educational brochures^{26,32,33} or a tape on Pap tests²⁵ all reported a significant increase in compliance. Telephone counseling interventions, including education on abnormal Pap tests and the importance of follow-up, also predicted adherence.^{15,17,24,30} Finally, Lavin et al.²³ found that a visit to an adolescent clinic positively predicted adherence; the authors speculated that during this visit, more information on the purpose and importance of follow-up care may have been given.

Lesion severity. Of the 15 studies that addressed lesion severity and adherence,^{10,11,14,16–18,20,23,25–29,32,34} 11^{10,11,14,16–18,20,25,26,28,29,32} found that women with less severe lesions were less likely to adhere to follow-up recommendations. Melnikow et al.²⁹ reported that, overall, women with LSIL/HSIL were more likely to adhere to any appointment compared with women with ASCUS; however, women with LSIL/HSIL were less likely to adhere to colposcopy compared with repeat Pap tests, whereas women with ASCUS were more likely to adhere to colposcopy compared with repeat Pap tests. When following women after two abnormal Pap tests, Bernard et al.¹⁰ stated that women whose results were less severe on the second assessment were least likely to receive the recommended colposcopy.

Psychosocial factors

Psychological barriers. When addressing health beliefs and adherence, Nelson et al.³¹ found that fatalism and health beliefs (specifically, that Pap tests were needed only when one had abnormal bleeding) were associated with poor adherence. Another study addressed psychological factors, including fear of cancer, embarrassment of pelvic examinations, and belief in early detection and follow-up. Researchers reported no significant differences in adherence among women who differed in responses on these items.²⁷ Four intervention studies used a case management approach to address psychosocial factors, including attitudes, coping skills, anxiety, and fear resulting from an abnormal Pap test result.^{15,17,24,30} Psychological barriers to care reported by women in each study included fear of finding cancer, worries about examination/treatment, and fertility concerns. All interventions addressing these concerns resulted in significant differences in adherence; women in the intervention group in each respective study were more than twice as likely to follow up as women in the control group.^{15,17,24,30}

Social support. Six studies addressed social support and marital status or live-in relationship as a predictor of adherence to follow-up after an abnormal Pap test.^{13,14,24–26,30} Two found that women with a live-in relationship²⁶ or those with any type of social support¹³ were more likely to follow up within 4–6 months than those without a live-in relationship or any kind of social support. Crane¹³ found that the type of support most likely to affect adherence differed by ethnic group. African American women were more likely to adhere when emotional support was available, whereas tangible support significantly influenced adherence among Latinas.¹³

Healthcare system characteristics

A number of studies assessed the relationship of adherence to existing clinician and site characteristics, including appointment reminder protocols.^{15–17,19–22,26–28,30,34,35}

Patient involvement/communication. Clinician involvement of patients in decisions, referrals, and management options were assessed in two studies.^{19,28} Hartz and Fenaughty¹⁹ reported that patients who chose surveillance had improved initial adherence but not long-term adherence. McKee et al.²⁸ reported that discussion of follow-

up options and plans at a prior visit increased adherence rates. Communication between clinicians and patients was also noted as a factor influencing adherence to follow-up. Lack of effective communication was associated with lower rates of adherence in one study,²⁷ as both “knowledge of results” and “getting the doctor to understand my needs” were significantly associated with adherence in univariate analysis.

Healthcare facility/clinician specialty. Site attributes were also assessed as an indicator of adherence to follow-up in some studies. McKee et al.²⁸ reported that clinics with colposcopy on-site had higher adherence rates. Takacs et al.³⁵ reported that patients randomized to the video colposcopy were five times more likely to return for follow-up care relative to those receiving traditional colposcopy. Lacey et al.²² reported no difference in follow-up rates between women who were referred to the public hospital and those who chose to follow up with their private physician. Furthermore, three studies^{27,28,34} found that clinician specialty was not associated with improved adherence. The only study to address laboratory characteristics²⁰ reported no association between adherence and the type of follow-up system used by clinics, laboratory reporting summaries, or the proficiency of cytologists in reading slides.

Reminder protocols. Several experimental studies included protocols in which the healthcare facility called or sent a reminder letter approximately 1 week before the woman’s appointment for follow-up care.^{15–17,21,26,30} Five^{15–17,26,30} of six studies reported a positive association between the reminder protocol and adherence to follow-up care; three of the studies were coupled with telephone counseling or an educational pamphlet.^{15,17,26} Miller et al.³⁰ reported that the telephone reminder system was effective at increasing adherence rates, but not as effective as additional telephone counseling. The only study to report no change in adherence used a letter instead of a telephone reminder system.²¹

Strengths and limitations of reviewed research

Table 4 outlines the strength and quality of studies reviewed and provides a corresponding score based on the following categories: power/sample size, use of a theoretical model, study design, response rate, outcome validation, and measurement of lesion severity. Studies that used a

TABLE 4. QUALITY OF STUDIES REVIEWED FOR BARRIERS TO ADHERENCE TO FOLLOW-UP CARE FOR ABNORMAL PAP TESTS

Author, year	Power/ sample size 0 = <100 1 = 100–399 2 = 400–699 3 = 700–999 4 = 1000+	Theoretical model 0 = No 1 = Yes	Study design 0 = Case series 1 = CC/CS 2 = Cohort 3 = Intervention 4 = RCT	Response rate 0 = None, or >50% 1 = 50%–70% 2 = 70%+	Outcome validation 0 = Self- report 1 = Medical record 2 = Validation	Measurement of lesion severity 0 = All lesion levels combined 1 = Separate analysis by lesion severity	Total score (Points/total)
Analytical studies							
Bernard et al., 2005 ¹⁰	4	0	2	NA ^a	1	1	8/12 = 0.67 ^b
Cardin et al., 2001 ¹¹	4	0	2	NA ^a	1	1	8/12 = 0.67 ^b
Carey and Gjerdingen Crane, 1996 ¹³	1	0	2	NA ^a	1	0	4/12 = 0.33
Eger and Peipert, 1996 ¹⁴	2	1	2	1	2	0	8/14 = 0.57
Fox et al., 1997 ¹⁸	1	0	1	NA ^a	1	0	3/12 = 0.25
Hartz and Fenaughty, 2001 ¹⁹	4	0	2	NA ^a	1	1	8/12 = 0.67 ^b
Jones and Hovis, 2000 ²⁰	1	0	2	1	1	1	6/14 = 0.42
Lavin et al., 1997 ²³	4	0	2	NA ^a	1	1	8/12 = 0.67 ^b
McKee et al., 1999 ²⁷	1	0	2	NA ^a	1	0	4/12 = 0.33
McKee et al., 2001 ²⁸	1	1	1	1	2	0	6/14 = 0.42
Melnikow et al., 1999 ²⁹	1	0	2	NA ^a	1	1	5/12 = 0.42
Nelson et al., 2002 ³¹	4	1	2	NA ^a	1	1	5/12 = 0.42
Peterson et al., 2003 ³⁴	2	0	2	2	1	0	10/14 = 0.71 ^b
Experimental studies							
Ell et al., 2002 ¹⁵	1	1	3	NA ^a	1	0	5/12 = 0.42
Engelstad et al., 2001 ¹⁶	1	0	4	0	2	1	8/14 = 0.57
Kaplan et al., 2000 ²¹	4	1	3	2	1	1	9/14 = 0.64
Lacey et al., 1993 ²²	1	1	3	2	1	1	11/14 = 0.78 ^b
Lerman et al., 1992 ²⁴	0	0	4	2	1	0	8/14 = 0.57
Marcus et al., 1998 ²⁶	4	1	4	2	1	0	7/14 = 0.50
Marcus et al., 1998 ²⁶	4	0	4	2	1	1	13/14 = 0.92 ^b
Miller et al., 1997 ³⁰	3	1	4	1	2	1	12/14 = 0.86 ^b
Paskett et al., 1990 ³³	1	1	4	2	1	0	11/14 = 0.78 ^b
Paskett et al., 1995 ³²	2	1	4	2	1	0	9/14 = 0.64
Takacs et al., 2004 ³⁵	0	0	4	2	1	1	11/14 = 0.78 ^b
						0	7/14 = 0.50

^aRetrospective study; no applicable response rate (total score out of 12 points).^bHigh-quality study (score $\geq$ 0.67 or top 33% of studies).

retrospective cohort design were not considered when assessing response rates.^{10-12,14,18,20,23,28,29,34} Therefore, the total score was out of 14 points for those with a response rate and 12 points for those without a response rate. Studies that scored within the top third of the sample were determined to be of high quality (score ≥ 0.67) (Table 4).^{10,11,17,18,20,21,25,26,30-32}

Sample size indicates study power and the ability to correctly rule out the possibility that a type 2 error occurred (not having sufficient power to detect a true association). Study power is, however, a function of sample size, the prevalence of the independent variable, and the prevalence of the outcome in the study population. Among the reviewed research, nine had good study power.^{10,11,18,20,21,25,26,30,31}

The randomized trial design was a strength for nine studies.^{16,17,24-26,30,32,33,35} Eleven studies used a theoretical model to guide their research, which was considered a strength.^{13,15,17,21,22,25,27,30-33} Only one study reported a low response rate (35%),¹⁵ which may indicate questionable validity of the study results.

Misclassification of both the independent and outcome variables can be a concern for observational studies and may lead to information bias. Studies that relied exclusively on medical records for information on patient characteristics^{10-12,14,16,18-25,28-35} may have misclassified these variables if they are not routinely available in medical records. Further, women who received follow-up care at other clinics may have been misidentified as nonadherent if self-reported follow-up was not available. Four studies obtained adherence information through self-report in addition to medical record abstraction; these studies both validated their outcomes using another source and were able to obtain complete documentation from other sites.^{13,15,26,27} Another study¹⁷ excluded women who reported follow-up at another clinic to avoid this bias.

Because lesion severity may directly impact the importance of and timeliness of adherence, we included assessment by lesion severity as a component of quality. Fourteen of 26 studies either conducted separate analysis by lesion severity or included only one type of lesion in their research.^{10,11,15-21,25,26,28,29,32}

As previously noted, having a range of adherence definitions makes comparisons across the studies challenging. In general, those studies with shorter time intervals for defining adherence may

be more prone to outcome misclassification. However, different types of noncompliance exist (e.g., delay in seeking care, nonparticipation in care, and cancellation of appointments), so all may be considered an appropriate measure.

Summary of reviewed research

Table 5 provides a summary of factors addressed by the literature included in this systematic review. Table 5 provides a breakdown of the associations by patient factors, psychosocial factors, and healthcare system factors. The strength of association corresponds to the range of point estimates (odds ratio [OR]/hazards ratio [HR]) for studies reporting a statistically significant association (Weak, OR/HR: 1.0-1.49 or 0.68-1.0; moderate, OR/HR: 1.5-3.0 or 0.33-0.67; strong, OR/HR: 3.0-10.0 or 0.10-0.32). Furthermore, Table 5 indicates the number of high-quality studies that contributed to the results.

DISCUSSION

Over the past 15 years, many factors have been hypothesized to affect adherence to follow-up procedures following an abnormal Pap test. In this discussion, we focus on modifiable risk factors for adherence at the patient, psychosocial, and healthcare system levels and address future directions for research.

Patient factors

Among those modifiable factors associated with the patient, the majority of studies included in this review agreed that women with less severe lesions were less likely to be adherent to follow-up care.^{10,11,14,16,18,20,25,26,28,29,32} Reasons for this association may include a belief of both women and clinicians that follow-up is less important for less severe Pap test results. Although this may in general be true, all women recommended for follow-up should receive follow-up care in a timely manner, as some less severe lesions may become more severe without follow-up. Potential interventions may include stressing the importance of follow-up regardless of lesion severity during patient-provider meetings and within educational materials addressing abnormal Pap tests.

The Pap test can detect lesions that will never progress to cervical cancer; therefore, women

TABLE 5. SUMMARY OF FACTORS ADDRESSING BARRIERS TO ADHERENCE TO FOLLOW-UP CARE

<i>Barriers</i>	<i>Number of studies that addressed factor</i>	<i>Number of studies that reported an association with factor</i>	<i>Strength of association^a</i>	<i>Number of high-quality studies^b</i>
Patient factors				
Age	14 (10, 11, 14, 16–18, 20, 23–27, 31, 34)	7 (16, 17, 20, 25–27, 34)	Weak–Moderate	8 (10, 11, 17, 18, 20, 25, 26, 30)
Race	16 (10, 11, 14, 16–18, 23–28, 30–32)	7 (10, 11, 17, 18, 25, 26, 32)	Moderate–strong	9 (10, 11, 17, 18, 25, 26, 30, 31, 32)
African American	7 (12, 16–18, 27, 31, 34)	4 (12, 16, 17, 18)	Moderate–strong	3 (17, 18, 31)
Asian	12 (16–18, 23–28, 30, 31, 34)	2 (18, 25)	Moderate	6 (17, 18, 25, 26, 30, 31)
Hispanic	3 (17, 29, 34)	No evidence	—	—
Language	2 (18, 29)	1 (18)	Moderate	1 (18)
Residence	5 (24, 25, 27, 30, 31)	1 (25)	No point estimate*	3 (25, 30, 31)
Education	3 (17, 20, 28)	1 (20)	No point estimate*	2 (17, 20)
Pregnancy	2 (14, 32)	1 (32)	Moderate	1 (32)
Gravity/parity	2 (14, 32)	1 (32)	Moderate	1 (32)
Tobacco	10 (14, 16, 17, 23, 25, 29, 34)	3 (25, 29, 34)	Moderate	2 (17, 25, 26)
Insurance	3 (14, 19, 31)	1 (19)	No point estimate*	1 (31)
Income/cost	3 (21, 25, 26)	2 (25, 26)	Moderate	3 (21, 25, 26)
interventions	2 (13, 31)	2 (13, 31)	Weak–moderate	1 (31)
Knowledge	7 (15, 17, 24–26, 30, 32, 33)	7 (15, 17, 24–26, 30, 32, 33)	Moderate–strong	5 (17, 25, 26, 30, 32)
interventions	15 (10, 11, 14, 16–18, 20, 23, 25–29, 32, 34)	11 (10, 11, 14, 16, 18, 20, 25, 26, 28, 29, 32)	Moderate–strong	8 (10, 11, 17, 18, 20, 25, 26, 32)
Lesion severity				

Psychosocial factors			
Psychological	2 (27, 31)	1 (31)	Weak
interventions	4 (15, 17, 24, 30)	4 (15, 17, 24, 30)	Moderate-strong
Social support/marital status	6 (13, 14, 24-26, 30)	2 (13, 26)	Moderate-strong
Healthcare system factors			
Communication/management choice	3 (19, 27, 28)	3 (19, 27, 28)	Strong
Reminder protocols	6 (15-17, 21, 26, 30)	5 (15-17, 26, 30)	Moderate-strong
interventions	3 (27, 28, 34)	No evidence	—
Clinician specialty	1 (22)	No evidence	—
interventions	3 (20, 28, 35)	2 (28)	Moderate-strong
Facility characteristics			
			1 (20)

^aStrength of association is defined by the range of point estimates (OR/HR) for statistically significant studies: weak, OR/HR: 1.0–1.49 or 0.68–1.0; moderate, OR/HR: 1.5–3.0 or 0.33–0.67; strong, OR/HR: 3.0–10.0 or 0.10–0.32.

^bQuality of study defined by: power/sample size, use of theoretical model, study design, response rate, outcome misclassification, and separation of lesion severity in analysis (Table 4).

* $p \leq 0.001$; no measure of strength or point estimate was reported in the study.

who have an abnormal Pap test result of ASCUS are now routinely tested for human papillomavirus (HPV). Because most cervical cancer cases result from persistent HPV infection, proper education and counseling of the potential impact of a positive HPV result may motivate women to adhere to follow-up care in a timely manner.

Many women have noted lack of time, money, or insurance as barriers to receiving timely follow-up care.^{8,15} Although the majority of studies found no association between economic factors and adherence,^{14,17,21,23,26–28,31} transportation incentives and economic vouchers did significantly increase adherence.^{25,26} Currently, the Breast and Cervical Cancer Treatment Act provides medical assistance through Medicaid to eligible women who were screened for and found to have breast or cervical cancer, including precancerous conditions. However, women who are ineligible for this service still may feel they cannot afford follow-up care; these women may benefit from targeted interventions to address economic barriers to follow-up care.

Misunderstanding of the purpose of a Pap test and the need for follow-up after an abnormal result were frequent barriers to care.^{13,15,24,31} All interventions that addressed knowledge of the Pap test and cervical cancer significantly increased adherence rates.^{15,17,24–26,30,32,33} The current knowledge gap can be addressed on a personal level by nurses, community health workers, and patient navigators tailoring informational needs to each woman. Furthermore, a need clearly exists for a comprehensive, easy to understand, and culturally acceptable brochure to aid in the education of women who have an abnormal Pap test.

Psychosocial factors

Social factors, such as the ability to cope effectively with an abnormal result,³⁰ as well as other attitudinal beliefs that may be influenced by social support³⁶ or culture may be important and potentially modifiable risk factors for nonadherence. To illustrate, greater fatalism was associated with lower adherence rates in two studies reviewed;^{15,31} further, fatalism appears to be a mediator in the association between race and adherence and, unlike race, is potentially modifiable.³¹ The literature supports that the emotional reaction (e.g., fear, anxiety, and depression) to the news of an abnormal Pap test result may reduce

a woman's ability to return for follow-up care in a timely manner.^{8,15,17,24,30,31,37–39} A small literature suggests that stressful events occurring in women's lives around the time of the abnormal Pap test may serve as competing life priorities, thereby reducing the likelihood of obtaining follow-up care.^{40–42} Effective coping strategies to reduce psychological distress from an abnormal Pap result have been noted.^{13,43} In order to reduce clinic no-show rates, we need a better understanding of the interactions among anxiety, distress, effective coping, and support as they affect adherence.

Healthcare system characteristics

Although patients are ultimately responsible for following the recommendations of their clinicians, clear patient-provider or laboratory communications may have a positive impact on adherence. Enhanced communication between the patient and the provider in experimental studies included in this review resulted in increased adherence rates.^{16,17,24,25,30,32,33} Interventions aimed at reducing this communication barrier included telephone reminder systems, counseling/educational sessions, and instructive and culturally relevant pamphlets. The success of these interventions is consistent with studies reviewed by Yabroff et al.⁴ and Abercrombie.⁷ These interventions may differentially affect women across a range of socioeconomic levels, and were tailored to meet women's individual needs.

Little research has addressed individual characteristics of the provider in influencing the effectiveness of communication. Future directions for research could include elements of effective patient-provider communication. These may include observational studies with permissible videotaped interactions. The type of healthcare provider may vary (e.g., physician, nurse, health education counselor), but the ability of the provider to simply and effectively communicate medical information as well as personal concern for that patient would be an important contribution.

CONCLUSIONS

Inconsistent evidence for risk factors and barriers to receipt of follow-up care as well as extreme differences in reported adherence rates

across the 26 scholarly papers reviewed indicate that the influence of demographic, individual, and healthcare system level factors on adherence to abnormal Pap tests is undoubtedly complex. Women differ in their knowledge of cervical cancer screening, their attitudes toward preventive follow-up care, and their overall access to this care, including financial and transportation barriers. Similarly, healthcare providers differ in their attitudes and communication skills. The healthcare system may determine follow-up availability on site and extended clinic hours.

Future directions

A range of barriers for adherence to follow-up after an abnormal Pap test have been identified in the existing literature. Therefore, the most cost-effective strategy to overcome diverse barriers among at-risk populations will likely include tailored interventions through case management or patient navigation. Additionally, although many modalities of patient education are used (e.g., letter, pamphlets, videotape/DVD, phone calls, e-mails, websites), little research has addressed the content of the material in terms of the intended audience, the cultural competence for a range of audiences, or the accuracy in light of new HPV testing methods, screening, and follow-up options.

Successful interventions, if incorporated into everyday practice, may increase the short-term cost of healthcare services. A recent study by Wagner et al.⁴⁴ however, suggests that tailored counseling interventions may be more cost-effective than usual care among high-risk populations.⁴⁴ By reducing barriers to follow-up care for abnormal Pap tests, the financial cost of late-stage treatment for cervical cancer will decrease. Further, decreases in the adverse physical, psychological, and emotional effects of cervical cancer morbidity and mortality would be significant.

REFERENCES

- Schiffman MH, Brinton LA, Devesa SS, Fraumeni J, Joseph F. Cervical cancer. In: Schottenfeld D, Fraumeni J, Joseph F, eds. *Cancer epidemiology and prevention*. New York: Oxford University Press, 1996.
- Cancer facts and figures 2006. Atlanta: American Cancer Society, 2006.
- Hughes E, McCracken M, Roberts H, et al. Surveillance for certain health behaviors among states and selected local areas—Behavioral Risk Factor Surveillance System, United States, 2004. *MMWR Surveill Summ* 2006;55:1.
- Yabroff KR, Washington KS, Leader A, Neilson E, Mandelblatt JS. Is the promise of cancer-screening programs being compromised? Quality of follow-up care after abnormal screening results. *Med Care Res Rev* 2003;60:294.
- Leyden WA, Manos MM, Geiger AM, et al. Cervical cancer in women with comprehensive health care access: Attributable factors in the screening process. *J Natl Cancer Inst* 2005;97:675.
- Bastani R, Yabroff KR, Myers RE, Glenn B. Interventions to improve follow-up of abnormal findings in cancer screening. *Cancer* 2004;101:1188.
- Abercrombie PD. Improving adherence to abnormal Pap smear follow-up. *J Obstet Gynecol Neonat Nurs* 2001;30:80.
- Khanna N, Phillips MD. Adherence to care plan in women with abnormal Papanicolaou smears: A review of barriers and interventions. *J Am Board Fam Pract* 2001;14:123.
- McKee D. Improving the follow-up of patients with abnormal Papanicolaou smear results. *Arch Fam Med* 1997;6:574.
- Benard VB, Lawson HW, Ehemann CR, Anderson C, Helsel W. Adherence to guidelines for follow-up of low-grade cytologic abnormalities among medically underserved women. *Obstet Gynecol* 2005;105:1323.
- Cardin VA, Grimes RM, Jiang ZD, Pomeroy N, Harrell L, Cano P. Low-income minority women at risk for cervical cancer: A process to improve adherence to follow-up recommendations. *Public Health Rep* 2001;116:608.
- Carey P, Gjerdingen DK. Follow-up of abnormal Papanicolaou smears among women of different races. *J Fam Pract* 1993;37:583.
- Crane LA. Social support and adherence behavior among women with abnormal Pap smears. *J Cancer Educ* 1996;11:164.
- Eger RR, Peipert JF. Risk factors for noncompliance in a colposcopy clinic. *J Reprod Med* 1996;41:671.
- Ell K, Vourlekis B, Muderbach L, et al. Abnormal cervical screen follow-up among low-income Latinas: Project safe. *J Womens Health Gend Based Med* 2002;11:639.
- Engelstad LP, Stewart SL, Nguyen BH, et al. Abnormal Pap smear follow-up in a high-risk population. *Cancer Epidemiol Biomarkers Prev* 2001;10:1015.
- Engelstad LP, Stewart SL, Ostero-Sabogal R, Leung MS, Davis PI, Pasick RJ. The effectiveness of a community outreach intervention to improve follow-up among underserved women at highest risk for cervical cancer. *Prev Med* 2005;41:741.
- Fox P, Amsberger P, Zhang X. An examination of differential follow-up rates in cervical cancer screening. *J Community Health* 1997;22:199.
- Hartz LE, Fenaughty AM. Management choice and adherence to follow-up after colposcopy in women with cervical intraepithelial neoplasia 1. *Obstet Gynecol* 2001;98:674.

20. Jones BA, Novis DA. Follow-up of abnormal gynecologic cytology. *Arch Pathol Lab Med* 2000;124:665.
21. Kaplan CP, Bastani R, Belin TR, Marcus A, Nasseri K, Hu MY. Improving follow-up after an abnormal Pap smear: Results from a quasi-experimental intervention study. *J Womens Health Gend Based Med* 2000;9:779.
22. Lacey L, Whitfield J, DeWhite W, et al. Referral adherence in an inner city breast and cervical cancer screening program. *Cancer* 1993;72:950.
23. Lavin C, Goodman E, Perlman S, Kelly LS, Emans SJ. Follow-up of abnormal papanicolaou smears in a hospital-based adolescent clinic. *J Pediatr Adolesc Gynecol* 1997;10:141.
24. Lerman C, Hanjani P, Caputo C, et al. Telephone counseling improves adherence to colposcopy among lower-income minority women. *J Clin Oncol* 1992;10:330.
25. Marcus AC, Crane LA, Kaplan CP, et al. Improving adherence to screening follow-up among women with abnormal Pap smears: Results from a large clinic-based trial of three intervention strategies. *Med Care* 1992;30:216.
26. Marcus AC, Kaplan CP, Crane LA, et al. Reducing loss-to-follow-up among women with abnormal Pap smears. Results from a randomized trial testing an intensive follow-up protocol and economic incentives. *Med Care* 1998;36:397.
27. McKee MD, Lurio J, Marantz P, Burton W, Mulvihill M. Barriers to follow-up of abnormal Papanicolaou smears in an urban community health center. *Arch Fam Med* 1999;8:129.
28. McKee MD, Schechter C, Burton W, Mulvihill M. Predictors of follow-up of atypical and ASCUS Pap test results in a high-risk population. *J Fam Pract* 2001;50:609.
29. Melnikow J, Chan BK, Stewart GK. Do follow-up recommendations for abnormal Papanicolaou smears influence patient adherence? *Arch Fam Med* 1999;8:510.
30. Miller SM, Siejak KK, Schroeder CM, Lerman C, Hernandez E, Helm CW. Enhancing adherence following abnormal Pap smears among low-income minority women: A preventive telephone counseling strategy. *J Natl Cancer Inst* 1997;89:703.
31. Nelson K, Geiger AM, Mangione CM. Effect of health beliefs on delays in care for abnormal cervical cytology in a multiethnic population. *J Gen Intern Med* 2002;17:709.
32. Paskett ED, Phillips KC, Miller ME. Improving compliance among women with abnormal Papanicolaou smears. *Obstet Gynecol* 1995;86:353.
33. Paskett ED, White E, Carter WB, Chu J. Improving follow-up after an abnormal Pap smear: A randomized controlled trial. *Prev Med* 1990;19:630.
34. Peterson NB, Han J, Freund KM. Inadequate follow-up for abnormal Pap smears in an urban population. *J Natl Med Assoc* 2003;95:825.
35. Takacs P, Chakhtoura N, DeSantis T. Video colposcopy improves adherence to follow-up compared to regular colposcopy: A randomized trial. *Arch Gynecol Obstet* 2004;270:182.
36. Trimble EL, Harlan LC, Clegg LX. Untreated cervical cancer in the United States. *Gynecol Oncol* 2005;96:271.
37. Bell S, Porter M, Kitchener H, Fraser C, Fisher P, Mann E. Psychological response to cervical screening. *Prev Med* 1995;24:610.
38. Fylan F. Screening for cervical cancer: A review of women's attitudes, knowledge, and behavior. *Br J Gen Pract* 1998;48:1509.
39. Lerman C, Miller SM, Scarborough R, Hanjani P, Nolte S, Smith D. Adverse psychologic consequences of cytologic cervical screening. *Am J Obstet Gynecol* 1991;165:658.
40. Breitkopf CR, Catero J, Jaccard J, Berenson AB. Psychological and sociocultural perspectives on follow-up of abnormal Papanicolaou results. *Obstet Gynecol* 2004;104:1347.
41. Coker AL, Bond SM, Brandt HM. Reasons women discontinue paid follow-up care for pre-invasive neoplasia. *Am J Epidemiol* 2002;155(11)Suppl:S3.
42. Lovejoy NC, Roche N, McLean D. Life stress and risk of precancerous cervical lesions: A pretest directed by the life stress model. *Oncol Nurs Forum* 1997;24:63.
43. Lauver DR, Baggot A, Kruse K. Women's experiences in coping with abnormal Papanicolaou results and follow-up colposcopy. *J Obstet Gynecol Neonat Nurs* 1999;28:283.
44. Wagner TH, Engelstad LP, McPhee SJ, Pasick RJ. The cost-effectiveness of an outreach intervention for women with abnormal pap smears: VA Palo Alto and Stanford University, 2005.

Address reprint requests to:

Katherine S. Eggleston, M.S.P.H.

University of Texas Health Science Center

School of Public Health

1200 Herman Pressler Drive

Houston, TX 77025

E-mail: Katherine.S.Eggleston@uth.tmc.edu

This article has been cited by:

1. Joy Melnikow, Shalini Kulasingam, Christina Slee, L. Jay Helms, Miriam Kuppermann, Stephen Birch, Colleen E. McGahan, Andrew Coldman, Benjamin K. S. Chan, George F. Sawaya. 2010. Surveillance After Treatment for Cervical Intraepithelial Neoplasia. *Obstetrics & Gynecology* **116**:5, 1158-1170. [[CrossRef](#)]
2. Sanja Percac-Lima, Leslie S. Aldrich, Gloria B. Gamba, Adriana M. Bearse, Steven J. Atlas. 2010. Barriers to Follow-Up of an Abnormal Pap Smear in Latina Women Referred for Colposcopy. *Journal of General Internal Medicine* **25**:11, 1198-1204. [[CrossRef](#)]
3. Monique A. Tello , Mollie Jenckes , Jennifer Gaver , Jean R. Anderson , Richard D. Moore , Geetanjali Chander . 2010. Barriers to Recommended Gynecologic Care in an Urban United States HIV ClinicBarriers to Recommended Gynecologic Care in an Urban United States HIV Clinic. *Journal of Women's Health* **19**:8, 1511-1518. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
4. J. Zapka, S. H. Taplin, R. Anhang Price, C. Cranos, R. Yabroff. 2010. Factors in Quality Care--The Case of Follow-Up to Abnormal Cancer Screening Tests--Problems in the Steps and Interfaces of Care. *JNCI Monographs* **2010**:40, 58-71. [[CrossRef](#)]
5. Tzy-Mey Kuo, Vicki B. Benard, Nancy D. Berkman, Carolyn K. Martin, Lisa C. Richardson. 2010. Timing of Colposcopy After Cervical Cytologic Abnormalities. *Obstetrics & Gynecology* **115**:3, 629-636. [[CrossRef](#)]
6. Tracy A. Battaglia, M. Christina Santana, Sharon Bak, Manjusha Gokhale, Timothy L. Lash, Arlene S. Ash, Richard Kalish, Stephen Tringale, James O. Taylor, Karen M. Freund. 2010. Predictors of timely follow-up after abnormal cancer screening among women seeking care at urban community health centers. *Cancer* **116**:4, 913-921. [[CrossRef](#)]
7. Jessica Warman. 2010. Cervical Cancer Screening in Young Women: Saving Lives With Prevention and Detection. *Oncology Nursing Forum* **37**:1, 33-38. [[CrossRef](#)]
8. Su-Ying Fang, Bih-Ching Shu. 2009. Adherence Characteristics After Abnormal Screening Results Between Mammogram and Papanicolaou Test Groups. *Cancer Nursing* **32**:6, 437-445. [[CrossRef](#)]
9. Meg Watson, Mona Saraiya, Vicki Benard, Steven S. Coughlin, Lisa Flowers, Vilma Cokkinides, Molly Schwenn, Youjie Huang, Anna Giuliano. 2008. Burden of cervical cancer in the United States, 1998-2003. *Cancer* **113**:S10, 2855-2864. [[CrossRef](#)]
10. Vicki B. Benard, William Howe, Mona Saraiya, William Helsel, Herschel W. Lawson. 2008. Assessment of Follow-Up for Low-Grade Cytological Abnormalities in the National Breast and Cervical Cancer Early Detection Program, 2000-2005. *Journal of Lower Genital Tract Disease* **12**:4, 300-306. [[CrossRef](#)]
11. Carlos E. Pineda, J. Michael Berry, Naomi Jay, Joel M. Palefsky, Mark L. Welton. 2008. High-Resolution Anoscopy Targeted Surgical Destruction of Anal High-Grade Squamous Intraepithelial Lesions: A Ten-Year Experience. *Diseases of the Colon & Rectum* **51**:6, 829-837. [[CrossRef](#)]
12. Ann Scheck McAlearney, Katherine W. Reeves, Stephanie L. Dickinson, Kimberly M. Kelly, Cathy Tatum, Mira L. Katz, Electra D. Paskett. 2008. Racial differences in colorectal cancer screening practices and knowledge within a low-income population. *Cancer* **112**:2, 391-398. [[CrossRef](#)]