7-2010

White Matter Diffusion Alterations in Normal Women at Risk of Alzheimer's Disease

Charles D. Smith

University of Kentucky, csmith@mri.uky.edu

Himachandra Chebrolu

University of Kentucky

Anders H. Andersen

University of Kentucky, anders@mri.uky.edu

David A. Powell

University of Kentucky

Mark A. Lovell

University of Kentucky, malove2@pop.uky.edu

See next page for additional authors

Follow this and additional works at: https://uknowledge.uky.edu/neurology_facpub

Part of the [Chemistry Commons](#), [Medical Neurobiology Commons](#), and the [Neurology Commons](#)

Right click to open a feedback form in a new tab to let us know how this document benefits you.

Repository Citation

Smith, Charles D.; Chebrolu, Himachandra; Andersen, Anders H.; Powell, David A.; Lovell, Mark A.; Xiong, Shuling; and Gold, Brian T., "White Matter Diffusion Alterations in Normal Women at Risk of Alzheimer's Disease" (2010). *Neurology Faculty Publications*. 1.

https://uknowledge.uky.edu/neurology_facpub/1

This Article is brought to you for free and open access by the Neurology at UKnowledge. It has been accepted for inclusion in Neurology Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

White Matter Diffusion Alterations in Normal Women at Risk of Alzheimer's Disease

Digital Object Identifier (DOI)

<http://dx.doi.org/10.1016/j.neurobiolaging.2008.08.006>

Notes/Citation Information

Published in *Neurobiology of Aging*, v. 31, issue 7, p. 1122–1131.

© 2008 Elsevier Inc. All rights reserved.

This manuscript version is made available under the CC-BY-NC-ND 4.0 license

<http://creativecommons.org/licenses/by-nc-nd/4.0/>

Authors

Charles D. Smith, Himachandra Chebrolu, Anders H. Andersen, David A. Powell, Mark A. Lovell, Shuling Xiong, and Brian T. Gold

© 2008 Elsevier Inc. All rights reserved.

**This manuscript version is made available under
the CC-BY-NC-ND 4.0 license**

<http://creativecommons.org/licenses/by-nc-nd/4.0/>

Published in final edited form as:

Neurobiol Aging. 2010 July ; 31(7): 1122–1131. doi:10.1016/j.neurobiolaging.2008.08.006.

White matter diffusion alterations in normal women at risk of Alzheimer's disease

Charles D. Smith^{a,b,c,*}, Himachandra Chebrolu^b, Anders H. Andersen^{b,c}, David A. Powell^b, Mark A. Lovell^d, Shuling Xiong^d, and Brian T. Gold^c

^aDepartment of Neurology, University of Kentucky, Lexington, KY 40536, United States

^bMagnetic Resonance Imaging and Spectroscopy Center, University of Kentucky, Lexington, KY 40536, United States

^cDepartment of Anatomy and Neurobiology, University of Kentucky, Lexington, KY 40536, United States

^dDepartment of Chemistry, University of Kentucky, Lexington, KY 40536, United States

Abstract

Increased white matter mean diffusivity and decreased fractional anisotropy (FA) has been observed in subjects diagnosed with mild cognitive impairment (MCI) and Alzheimer's disease (AD). We sought to determine whether similar alterations of white matter occur in normal individuals at risk of AD. Diffusion tensor images were acquired in 42 cognitively normal right-handed women with both a family history of dementia and at least one apolipoprotein E4 allele. These were compared with images from 23 normal women without either AD risk factor. Group analyses were performed using tract-based spatial statistics. Reduced FA was observed in the fronto-occipital and inferior temporal fasciculi (particularly posteriorly), the splenium of the corpus callosum, subcallosal white matter and the cingulum bundle. These findings demonstrate that specific white matter pathways are altered in normal women at increased risk of AD years before the expected onset of cognitive symptoms.

Keywords

Diffusion tensor; Fractional anisotropy; Magnetic resonance imaging; Alzheimer's disease; Apolipoprotein E

1. Introduction

There is now strong evidence that significant Alzheimer's neuropathology is present years before any symptoms of mild cognitive impairment (MCI) or Alzheimer's disease (AD) appear (Braak and Braak, 1997; Bennett et al., 2006). The underlying theory of this study is that the first appearance of cognitive symptoms due to AD pathology, e.g., amnesic MCI, represents

© 2008 Elsevier Inc. All rights reserved

*Corresponding author at: Magnetic Resonance Imaging and Spectroscopy Center, Room 62, University of Kentucky Medical Center, 800 Rose Street, Lexington, KY 40536-0098, United States. Tel.: +1 859 323 1113; fax: +1 859 323 1068. *E-mail address:* csmith@mri.uky.edu (C.D. Smith)..

Conflict of interest None.

Appendix A. Supplementary data Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.neurobiolaging.2008.08.006.

early failure of brain compensatory mechanisms and cerebral reserve, a critical point in the further evolution of neuronal injury produced by that pathology (Smith, 2007). We assume that AD neuropathology progresses with time. But even if a treatment could theoretically slow or arrest MCI/AD, other processes of aging and vascular injury could operate on this substrate of critical reserve and result in further cognitive decline (Schneider et al., 2007). Therefore, identifying the brain at risk *before* MCI is important to the goal of preventing the critical point from ever being reached, by using AD disease-modifying treatments, combined with treatments aimed at the comorbid pathologies and enhancement of brain reserve, before symptoms appear.

Several methodologies have been applied, or have potential to be applied, to asymptomatic persons to assess the brain at AD risk, including cerebrospinal fluid protein patterning, genetic analysis, resting glucose positron emission tomography, and functional and structural magnetic resonance imaging. Studies of white matter integrity using magnetic resonance diffusion tensor imaging have demonstrated alterations in the frontal, temporal and parietal lobes in subjects with MCI and AD (Medina et al., 2006; Xie et al., 2006; Fellgiebel et al., 2007; Fjorbank et al., 2007; Huang et al., 2007; Stahl et al., 2007; Sydykova et al., 2007; Teipel et al., 2007; Zhang et al., 2007; Ding et al., 2008; Rose et al., 2008; Yasmin et al., 2008). There is pathological evidence that disruption of white matter in Alzheimer's disease may be an early event (Englund et al., 1988). In line with this data, there is important recent evidence suggesting that normal persons at increased risk for late-onset AD by virtue of an apolipoprotein-E4 (APOE4) allele show reduced fractional anisotropy (FA) in certain brain regions compared to low-risk groups (Nierenberg et al., 2005; Persson et al., 2006a,b). However, these studies have either explored only a few regions with ROI analyses, or used automated techniques known to have limitations related to registration and smoothing (Bookstein, 2001; Smith et al., 2006). In the present study we applied rigorously validated methods for comparing group fractional anisotropy maps across the brain's white matter (Smith et al., 2006) between well-characterized normal subjects who vary in their risk of AD. The main advantages of these methods are improved registration of white matter, the lack of a need to apply an arbitrary smoothing kernel, and a common reference framework (mean FA skeleton). In addition, because the method is voxel-based, arbitrary definitions of region of interest are not required.

2. Methods

2.1. Subjects

All subjects are participants in a University of Kentucky longitudinal imaging study of normal persons who vary in their risk of AD based on family history of dementia and APOE allele status (Smith et al., 1999, 2002). Inclusion criteria include generally good health without memory complaints, stable medication regimen, right handed female between the age of 40 and 90 years, vision correctable to at least 20/50, and knowledge of family history (FH). Family history was considered positive if one or more first-degree relatives had late-onset progressive dementia. Only right-handed females were recruited because they are part of a larger multimodal imaging study that includes functional magnetic resonance imaging during language-based tasks. Language task-based fMRI is known to demonstrate gender and handedness activation differences.

Exclusions include current medical evaluation or treatment for acute or worsening complaints (e.g., unstable asthma), stroke or other cerebral injury, psychoactive medications except stable doses of serotonin reuptake inhibitors and absence of active depression, significant head injury (operationally loss of consciousness greater than 5 min), significant psychiatric history, claustrophobia, metallic implants in the head or neck, pacemakers, major surgery within 3 months, or death of one or both parents before age 65.

Participants who are selected for scanning based on risk undergo standardized psychometric testing which includes the Mini Mental State Examination (MMSE), Trails A (TA) and B (TB), FAS Letter Fluency (LF), Wechsler Adult Intelligence Scale (WAIS) Block Design (Block Des) and Vocabulary (Vocab), 15-item Boston Naming Test (BNT), Wechsler Memory Scale (WMS) Logical Memory I (LM Imm) and II (LM Del), Visual Reproduction immediate and delayed (VR and VR Del), Digit Span (DS), Spatial Span (SS), Selective Reminding Test Immediate and Delayed (SRT & SRT Del), and Beck depression inventory (Beck).

Abbreviations are given for reference to Table 1B. These tests are given to exclude persons with cognitive impairment and to document equivalent normal cognitive test performance between risk groups. All subjects had an APOE allele determination. APOE alleles were determined using polymerase chain reaction (PCR) amplification followed by enzymatic digestion and separation of DNA fragments by agarose gel electrophoresis. Informed consent was obtained from each enrollee under an approved University of Kentucky medical institutional review board protocol.

Forty-four high risk (both FH dementia and one or more APOE4 alleles) and 27 low risk (neither FH nor APOE4) participants underwent diffusion tensor magnetic resonance imaging (DT-MRI) between the years 2006 and 2007. One of each risk group were excluded because of poor memory performance on cognitive testing (1.5 S.D. below the mean on Logical Memory II), and an additional one of the high risk and three of the low-risk group were not analyzed because of poor quality MRI scans. None of the included scans demonstrated small or large-vessel infarcts. Consequently there were 42 in the high-risk and 23 in the low-risk group available for DT analysis. Demographic information and testing results are summarized in Table 1. Psychometric scores that sample general mental ability, processing speed, and delayed verbal and nonverbal memory are provided in the table.

Demographic and psychometric scores were compared between risk groups. First age and education were entered as response variables in a standard least-squares analysis with risk group as the independent variable (Table 1A). Age and education were chosen as the most likely to variables to affect psychometric test scores. Because all subjects were right-handed females, gender and handedness were not factors. A standard least-squares model was then developed for the psychometric test scores, with risk group, age, education and risk by age interaction as the independent variables *versus* the following independent variables: MMSE, Logical Memory immediate and delayed, Block Design, Visual Reproduction immediate and delayed, Boston naming, Vocabulary, Selective Reminding Test immediate and delayed, Trails A and B, Letter Fluency, Spatial Span, Digit Symbol, and Beck inventory score (Table 1B).

To test for potential differences between risk groups related to vascular factors or estrogen replacement that might affect white matter, Chi-Square analysis using Pearson's criterion without correction for multiple comparisons was performed on the following categorical variables (Present/Not Present): current smoking, past smoking, current estrogen replacement, past estrogen replacement, hypercholesterolemia, hypertension and diabetes. There were three subjects with stable diabetes in the study, one in the low-and two in the high-AD risk group. None of the *p*-values in these comparisons was less than $p = 0.16$, demonstrating that the frequencies of these factors were the same between risk groups. More detailed data is presented in a supplementary table on the journal web site (<http://www.elsevier.com>).

2.2. MRI protocol

Scanning was performed on a Siemens Trio 3T instrument. Whole-brain (40-slice) diffusion tensor images were acquired with 12 encoding directions with 3 averages each (B_0 images were averaged 6 times; TR = 14600 ms, TE = 96 ms; resolution 1.8 mm × 1.8 mm × 3.0 mm) using a fluid attenuated inversion recovery (FLAIR) double-refocused spin echo sequence.

2.3. Image processing and statistics

Voxelwise statistical analysis of the FA data was carried out using TBSS (Tract-Based Spatial Statistics (Smith et al., 2006)), part of the publicly available image processing software FSL 4.0 (<http://www.fmrib.ox.ac.uk/fsl/tbss/index.html>). TBSS was used because its algorithms for alignment of FA images from multiple subjects into a common space have been extensively tested and validated (Smith et al., 2006). In addition, the technique does not require spatial smoothing of images, which can produce different results based upon the kernel size (Jones et al., 2005). All of the processing modules referred to below by their acronym are available for download at the FSL web site. First, FA images were created by fitting a tensor model to the raw diffusion data using FSL's Diffusion Toolbox, which consists of a series of tools for low-level diffusion parameter reconstruction and probabilistic tractography.

In order to minimize the extent of warping required for registration to a common target, a subset of 25 subjects was used to select a target image for nonlinear registration. Subjects forming both the whole set and the subset were scanned over the same time interval. The subset was selected in two steps. First, the brain volumes of all subjects were calculated and rank ordered. Second, age-volume means within the interquartile range were arranged in chronological order, and the 25 subjects closest in age to the median were selected. From the 25 subjects, the optimum target image was found.

The subset subjects' FA images were aligned with every other subset subjects' FA images using the nonlinear registration toolkit IRTK (Rueckert et al., 1999) (<http://www.doc.ic.ac.uk/~dr/software>). The common target FA image for the nonlinear registration was chosen as the FA image for which the least amount of total nonlinear warping across all subset subjects was necessary for registration to the same space. The least amount of warping was determined by registration of each subject to every other subject, and summarization of each warp field by its mean displacement. The resulting target FA image was then affine registered to MNI space. A single transformation matrix for each of the 65 subjects, combining the nonlinear registration of the subject's FA image to the target FA image and the affine registration of the target FA image to MNI space, was then applied. The individual FA images were resampled at $1\text{ mm} \times 1\text{ mm} \times 1\text{ mm}$ isotropic voxels using sinc interpolation.

Next, a mean FA image from all of the subjects was created and thinned with a FA threshold of 0.2 to create a mean FA skeleton which represents the centers of all tracts common to the group. Each subject's aligned FA data was then projected onto this skeleton and the resulting data analyzed using voxelwise cross-subject statistics. A false discovery rate (FDR) threshold scaled to an alpha of $p = 0.05$ was determined via random simulations (using 5000 permutations) to correct for multiple comparisons (implemented as in: <http://www.fmrib.ox.ac.uk/fsl/randomise/fdr.html>). The FDR was then applied to all voxel-based group comparisons. Centered age was used as a covariate in the analysis, because the risk groups showed an overall difference in age (see Section 3).

3. Results

Demographic and testing characteristics of the study groups are given in Table 1. Because no correction was made for the 16 multiple comparisons in Table 1B, we chose to conservatively interpret p between 0.01 and 0.05 as indicating marginal significance, and p less than 0.01 as significant. Low-AD risk subjects were older than high-risk subjects due to the convenience nature of the sample. We speculate that normal working-age high-risk younger subjects may be more motivated to volunteer their time for this study than similarly aged low-risk volunteers because of personal experience with Alzheimer's disease in family members. In previous studies using APOE allele status for risk stratification, the average age was approximately 67

years (Nierenberg et al., 2005; Persson et al., 2006a,b), compared to approximately 58 years for high-risk subjects in the present study. The time to expected development of MCI or AD is therefore about 9 years greater in the present study. Education level was not different between risk groups. The frequencies of past and present hypertension, hypercholesterolemia, smoking and estrogen replacement were similar between risk groups (Supplemental Table).

Correlation with age was in the expected positive direction for Trails A and B, and in the negative direction for Block Design, Visual Reproduction Immediate and Delayed, Selective Reminding Test Delayed, and Spatial Span. Education effects were observed only marginally on the Beck inventory; the absence of these effects overall is attributed to the similar mean and narrow range of education in the study subjects (Table 1B). The age $\times$ risk interaction was marginally significant for LM immediate interaction, scores decreased with increasing age in the low-risk group but not for the high-risk group, whereas VR immediate scores decreased with increasing age in the high-risk group but not for the low-risk group.

Trails A was the only psychometric test of the 16 performed meeting our threshold p -value of <0.01 for risk group difference. Trails A is a visuomotor task combining visuospatial and motor speed components. High-risk subjects were slower on this test than low risk. High-risk subjects were also slower on Trails B, but not significantly. The difference between Trails B and A is thought to produce a variable with reduced dependence on motor speed. This B minus A difference when compared between groups was not significant, suggesting that lower performance on Trails A in high-AD risk subjects was due to decreased motor speed and not visuospatial factors (high risk 39.9 ± 3.7 S.E.M., low risk 35.7 ± 5.2 S.E.M. seconds; $p = 0.5$). As expected, age was positively correlated with B minus A ($p = 0.02$). Spatial span was lower in high-risk subjects but not within our cutoff for group difference (Table 1B). In contrast, there were no differences between risk groups in general mental ability (MMSE, Vocabulary, BNT) or in memory measures (LM delayed, or in the more sensitive Selective Reminding Test delayed).

The Beck Inventory score was slightly higher in high-risk subjects but well below the cutoff of 13 for mild depression (mean 7.3 in high risk and 3.2 in low risk; $p = 0.05$). The lower Block Design score in high-risk subjects, although only marginally significant, appeared interesting at first because of the association with visuospatial intelligence. However, in a direct comparison of Block Design scaled score between risk groups adjusted for age and education, the difference was not significant (high risk 11.4 ± 3.704 S.E.M., low risk 12.5 ± 0.5 S.E.M.; $p = 0.08$), tempering the potential significance of this finding.

Figs. 1-3 illustrate statistical maps rendered on the common FA target image in Montreal Neurologic Institute (MNI) spatial coordinates, using the contrast: FA high-AD risk less than FA low risk. The reverse contrast yielded only a few scattered voxels. Maps are referenced to a standard human white matter atlas (Mori et al., 2005). The maps demonstrate lower FA in the high-AD risk subjects in: (a) the inferior longitudinal fasciculus of the temporal lobe posteriorly (particularly inferior fronto-occipital fasciculus bilaterally) and cingulum bundle posteriorly on the left (Fig. 1), (b) splenium of the corpus callosum, white matter of the parahippocampal gyrus (left), and right mid cingulum (Fig. 2), and (c) inferior fronto-occipital fasciculus/uncinate fasciculus anteriorly (Fig. 3). Decreased white matter FA in these specific regions is seen in the younger high-AD risk group of subjects compared to the older low-AD risk subjects, the reverse of the increased FA expected with aging; therefore the difference cannot be attributed to aging. A listing of the center locations of clusters with ≥ 50 contiguous voxels (at least 0.05 cm^3 volume) is shown in Table 2. These listings provide a concise summary of relatively spatially coherent FA decreases from results in the figures, which were corrected for multiple comparisons at the significance level but did not impose a cluster threshold. Cluster thresholding refers to comparisons based on spatial extent of FA differences.

4. Discussion

The most important finding was a clearly delineated and highly selective decrease in FA in the inferior temporal lobe white matter bilaterally (inferior longitudinal fasciculus/inferior fronto-occipital fasciculus). This decrease has a strong linear spatial character suggesting involvement of a specific group of white matter fibers bridging the amygdala/hippocampal head region anteriorly and the ventral visual association areas posteriorly (Fig. 1).

We also found significant FA decreases in the splenium of the corpus callosum, white matter of the cingulum, and fronto-occipital fasciculus anteriorly. Downward curvature of the involved fronto-occipital fasciculus shown in Fig. 3, particularly well seen on the left, is characteristic of this pathway, which parallels the uncinate fasciculus in part of its course. In general our findings were more robust in the temporal and frontal regions than in parietal lobes, and more spatially coherent in the ventral visual association regions, medial temporal lobe, cingulum and frontal lobe than in the parietal lobe or occipital lobe.

Previous studies using DTI in aging and MCI/AD have used several different acquisition and analysis strategies. Studies of normal aging have shown consistently decreased FA and related measures with age, particularly in the frontal white matter with relative sparing of the temporal, parietal and occipital regions (Pfefferbaum and Sullivan, 2003; Furutani et al., 2005; Charlton et al., 2006; Persson et al., 2006a,b; Sullivan et al., 2006; Grieve et al., 2007; Kochunov et al., 2007; Madden et al., 2007; Taylor et al., 2007; Abe et al., 2008; Hsu et al., 2008; Hugenschmidt et al., 2008). Increases of FA with age have not been reliably observed in any region.

In contrast, only a few studies have reported selective findings in the frontal lobe in MCI or AD (Choi et al., 2005; Sydykova et al., 2007). The remainder have demonstrated more consistent overlapping findings of decreased FA in the cingulum bundle (Fellgiebel et al., 2007; Teipel et al., 2007; Zhang et al., 2007), temporal lobe generally (Xie et al., 2006; Huang et al., 2007; Stahl et al., 2007), parietal lobe and posterior cingulate (Medina et al., 2006; Firbank et al., 2007; Huang et al., 2007; Ding et al., 2008; Rose et al., 2008), medial temporal white matter (Fellgiebel et al., 2007; Huang et al., 2007; Yasmin et al., 2008), inferior temporal lobe, superior and inferior fronto-occipital fasciculi (Xie et al., 2006; Fellgiebel et al., 2007), and splenium of the corpus callosum (Stahl et al., 2007). Findings in MCI tend to overlap with AD, particularly in posterior areas. These studies have differed in that some selected a restricted target region of a priori interest, e.g., the medial temporal white matter or cingulate bundles (Zhang et al., 2007; Ding et al., 2008; Yasmin et al., 2008), others picked several general regions of interest within lobar white matter (Fellgiebel et al., 2004, 2007; Choi et al., 2005; Stahl et al., 2007), whereas still others used a voxel-based technique (Medina et al., 2006; Rose et al., 2006, 2008; Xie et al., 2006; Firbank et al., 2007; Huang et al., 2007; Sydykova et al., 2007; Teipel et al., 2007). For comparison the analysis technique we used in this study is voxel-based, and therefore unbiased with respect to a specific hypothesis regarding where regional FA differences might be found.

It is important to make the distinction between studies described above that examine subjects with symptoms of a clinically diagnosed cognitive disorder, e.g., MCI and AD, and studies of normal persons without symptoms or deficits on psychometric testing who may later develop these conditions. There are many fewer investigations like the present one that examine the latter issue. Brain FA alterations that are associated with the presence of a diagnosed clinical disorder due to an underlying pathology may be assumed to be different in degree, or perhaps kind, from alterations that predict later development of those symptoms. But it is currently unknown whether such findings would predict MCI or AD. This is implied but not demonstrated in studies of at-risk normal persons. The first step in the validation of the current

findings is to follow these currently (c. 2008) normal study subjects longitudinally to show whether the observed FA alterations are predictive of later clinically diagnosed MCI or AD.

The voxel-based studies have not been entirely consistent in localization of FA decreases in AD, e.g., some find frontal and internal capsule alterations, but the overall pattern across studies suggests alterations in the white matter of the parahippocampal gyrus, inferior longitudinal fasciculus, splenium of the corpus callosum, parietal white matter, and cingulum bundle (Medina et al., 2006; Rose et al., 2006, 2008; Xie et al., 2006; Firbank et al., 2007; Huang et al., 2007; Sydykova et al., 2007; Teipel et al., 2007).

Of particular relevance, a recent study reported reduced FA in the fornix and left frontal WM in a young group of mildly impaired (CDR = 0.5) subjects (mean age = 35 years old) at risk for early onset-AD by virtue of fully penetrant genetic alterations (Ringman et al., 2007). Only two studies have explored WM integrity differences in cognitively normal subjects at risk for the more common late-onset form of AD due to possession of an APOE4 allele. Using ROI or SPM techniques, respectively, these studies observed reduced FA in the left parahippocampal WM (Nierenberg et al., 2005), or the splenium of the corpus callosum, and left posterior hippocampus (Persson et al., 2006a,b). Our results show some overlap with findings in these studies, specifically, the findings of decreased FA in the splenium of the corpus callosum and inferomedial temporal lobes. The location of the present FA decreases in presymptomatic carriers of AD overlap with previous FA reductions observed in AD in the inferior fronto-occipital fasciculi (Xie et al., 2006; Fellgiebel et al., 2007), and cingulum bundle (Fellgiebel et al., 2007; Teipel et al., 2007; Zhang et al., 2007). The term presymptomatic in this context is a group characterization and should not be interpreted to mean that each individual subject should be considered presymptomatic. These findings therefore suggest that AD-related FA declines are already present in groups of individuals at high risk for AD, even when they are thinking, behaving and functioning normally.

The strengths of our study are the confirmation of cognitive normality and test performance equivalence between risk groups using comprehensive psychometric testing of all participants, rigorous exclusion of confounding neurologic disease, reduction of FA variance due to gender and cerebral language representation by exclusive recruitment of right handed women (Hsu et al., 2008), exclusion of potential confounds due to smoking, hypertension, hypercholesterolemia and estrogen replacement, a consistent imaging protocol across all subjects using a 3T MRI instrument, and use of an advanced, validated and regionally unbiased DTI methodology.

Several cautions should be kept in mind in interpreting our results. Despite equivalent cognitive performance between risk groups, the low-risk subjects were older than the high-AD risk. We used age as a regressor in all our analyses, but statistical adjustments are not perfect. Because FA is known to decrease - not increase - with age, a finding of an FA decrease in the younger high-risk group is the reverse of expectation on an age basis, so we believe the difference in age strengthens rather than weakens our findings. Nonetheless, the difference in age between groups should be noted.

Another more important caveat is in interpretation of results. We can only infer that the alterations we have observed are related to the risk of AD. The study participants were selected to differ in no respect other than AD risk, but it is possible the groups differed in some other unknown but important way for which we could not adjust or control. Furthermore, we do not know that presence of these white matter alterations is due to underlying AD pathology, a basic hypothesis of the study, because we have no independent verification of its degree or presence, e.g., by autopsy or brain biopsy.

The issue of the functional significance of the FA declines in presymptomatic groups also needs further investigation. This is a tricky issue because presymptomatic high-risk subjects display cognitive function that is, by definition, comparable to that of low-risk subjects. However, there is accumulating evidence that FA is of functional significance in young healthy subjects (Madden et al., 2004; Tuch et al., 2005; Gold et al., 2007), and that age-related FA declines contribute to some cognitive performance declines (Madden et al., 2004; Gold et al., 2008). The view that FA declines are functionally relevant is also in-line with our previous finding of decreased ventral temporal activation in asymptomatic high-AD risk women in an fMRI naming task (Smith et al., 1999). The fMRI under-recruitments in this previous study of subjects were in regions adjacent to the postero-inferior FA alterations observed in the present study. Although both studies were performed using similar subjects selected by the same criteria, the specific individuals studied were different.

Finally, like other studies of presymptomatic participants, we cannot yet claim a role for altered FA in predicting AD because the participants have not been followed long enough to observe even the earliest symptoms for validation. Longitudinal follow up of the participants described in this study is in place at our center and should yield relevant results in the future.

In summary this report represents further progress toward the goal of identifying the brain at risk of AD using DTI. We have identified restricted regions of the inferomedial temporal lobes bilaterally, and other regions, that demonstrate decreased FA in normal high-AD risk subjects, that may in future serve as a biomarker for the presence of asymptomatic AD pathology. What we have observed here are specific regions of decreased FA in high-AD risk subjects. The next step in validating this work is to follow subjects longitudinally to show whether the observed changes are truly predictive of MCI or AD, and with what accuracy. Therefore, this method is promising but needs further study before it could be used for that purpose in trials of preventive agents in AD.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Acknowledgements

This study was supported by NINDS Grant R01 NS-36660. We thank Agnes Bogner, R.T., Kim Wilson, B.A., LuAnn Hamon, B.A., Dorothy Ross, B.A., and Barbara Martin, B.A. for their invaluable assistance in recruiting, scanning and testing the participants, and for treating them professionally and so well.

References

- Abe O, Yamasue H, Aoki S, Suga M, Yamada H, Kasai K, Masutani Y, Kato N, Kato N, Ohtomo K. Aging in the CNS: comparison of gray/white matter volume and diffusion tensor data. *Neurobiol. Aging* 2008;29:102–116. [PubMed: 17023094]
- Bennett DA, Schneider JA, Arvanitakis Z, Kelly JF, Aggarwal NT, Shah RC, Wilson RS. Neuropathology of older persons without cognitive impairment from two community-based studies. *Neurology* 2006;66:1837–1844. [PubMed: 16801647]
- Bookstein FL. “Voxel-based morphometry” should not be used with imperfectly registered images. *Neuroimage* 2001;14:1454–1462. [PubMed: 11707101]
- Braak H, Braak E. Frequency of stages of Alzheimer-related lesions of different age categories. *Neurobiol. Aging* 1997;18:351–357. [PubMed: 9330961]
- Charlton RA, Barrick TR, McIntyre DJ, Shen Y, O'Sullivan M, Howe FA, Clark CA, Morris RG, Markus HS. White matter damage on diffusion tensor imaging correlates with age-related cognitive decline. *Neurology* 2006;66:217–222. [PubMed: 16434657]

- Choi SJ, Lim KO, Monteiro I, Reisberg B. Diffusion tensor imaging of frontal white matter microstructure in early Alzheimer's disease: a preliminary study. *J. Geriatr. Psychiatry Neurol* 2005;18:12–19. [PubMed: 15681623]
- Ding B, Chen KM, Ling HW, Zhang H, Chai WM, Li X, Wang T. Diffusion tensor imaging correlates with proton magnetic resonance spectroscopy in posterior cingulate region of patients with Alzheimer's disease. *Dement. Geriatr. Cogn. Disord* 2008;25:218–225. [PubMed: 18219201]
- Englund E, Brun A, Alling C. White matter changes in dementia of Alzheimer's type. Biochemical and neuropathological correlates. *Brain* 1988;111(Part 6):1425–1439. [PubMed: 3208064]
- Fellgiebel A, Schermuly I, Gerhard A, Keller I, Albrecht J, Weibrich C, Muller MJ, Stoeter P. Functional relevant loss of long association fibre tracts integrity in early Alzheimer's disease. *Neuropsychologia*. 2007
- Fellgiebel A, Wille P, Muller MJ, Winterer G, Scheurich A, Vucurevic G, Schmidt LG, Stoeter P. Ultrastructural hippocampal and white matter alterations in mild cognitive impairment: a diffusion tensor imaging study. *Dement. Geriatr. Cogn. Disord* 2004;18:101–108. [PubMed: 15087585]
- Firbank MJ, Blamire AM, Krishnan MS, Teodorczuk A, English P, Gholkar A, Harrison R, O'Brien JT. Atrophy is associated with posterior cingulate white matter disruption in dementia with Lewy bodies and Alzheimer's disease. *Neuroimage* 2007;36:1–7. [PubMed: 17412610]
- Furutani K, Harada M, Minato M, Morita N, Nishitani H. Regional changes of fractional anisotropy with normal aging using statistical parametric mapping (SPM). *J. Med. Invest* 2005;52:186–190. [PubMed: 16167537]
- Gold BT, Powell DK, Xuan L, Jiang Y, Hardy PA. Speed of lexical decision correlates with diffusion anisotropy in left parietal and frontal white matter: evidence from diffusion tensor imaging. *Neuropsychologia* 2007;45:2439–2446. [PubMed: 17509627]
- Gold BT, Powell DK, Xuan L, Jicha GA, Smith CD. Agerelated slowing of task switching is associated with decreased integrity of frontoparietal white matter. *Neurobiol. Aging*. 2008 [Epub ahead of print].
- Grieve SM, Williams LM, Paul RH, Clark CR, Gordon E. Cognitive aging, executive function, and fractional anisotropy: a diffusion tensor MR imaging study. *Am. J. Neuroradiol* 2007;28:226–235. [PubMed: 17296985]
- Hsu JL, Leemans A, Bai CH, Lee CH, Tsai YF, Chiu HC, Chen WH. Gender differences and age-related white matter changes of the human brain: a diffusion tensor imaging study. *Neuroimage* 2008;39:566–577. [PubMed: 17951075]
- Huang J, Friedland RP, Auchus AP. Diffusion tensor imaging of normal-appearing white matter in mild cognitive impairment and early Alzheimer disease: preliminary evidence of axonal degeneration in the temporal lobe. *Am. J. Neuroradiol* 2007;28:1943–1948. [PubMed: 17905894]
- Hugenschmidt CE, Peiffer AM, Kraft RA, Casanova R, Deibler AR, Burdette JH, Maldjian JA, Laurienti PJ. Relating imaging indices of white matter integrity and volume in healthy older adults. *Cereb Cortex* 2008;18:433–442. [PubMed: 17575289]
- Jones DK, Symms MR, Cercignani M, Howard RJ. The effect of filter size on VBM analyses of DT-MRI data. *Neuroimage* 2005;26:546–554. [PubMed: 15907311]
- Kochunov P, Thompson PM, Lancaster JL, Bartzokis G, Smith S, Coyle T, Royall DR, Laird A, Fox PT. Relationship between white matter fractional anisotropy and other indices of cerebral health in normal aging: tract-based spatial statistics study of aging. *Neuroimage* 2007;35:478–487. [PubMed: 17292629]
- Madden DJ, Spaniol J, Whiting WL, Bucur B, Provenzale JM, Cabeza R, White LE, Huettel SA. Adult age differences in the functional neuroanatomy of visual attention: a combined fMRI and DTI study. *Neurobiol. Aging* 2007;28:459–476. [PubMed: 16500004]
- Madden DJ, Whiting WL, Huettel SA, White LE, MacFall JR, Provenzale JM. Diffusion tensor imaging of adult age differences in cerebral white matter: relation to response time. *Neuroimage* 2004;21:1174–1181. [PubMed: 15006684]
- Medina D, DeToledo-Morrell L, Urresta F, Gabrieli JD, Moseley M, Fleischman D, Bennett DA, Leurgans S, Turner DA, Stebbins GT. White matter changes in mild cognitive impairment and AD: a diffusion tensor imaging study. *Neurobiol. Aging* 2006;27:663–672. [PubMed: 16005548]
- Mori, S.; Wakana, S.; Nagae-Poetscher, LM.; van Zijl, PCM. *MRI Atlas of Human White Matter*. Elsevier; Amsterdam: 2005.

- Nierenberg J, Pomara N, Hoptman MJ, Sittis JJ, Ardekani BA, Lim KO. Abnormal white matter integrity in healthy apolipoprotein E epsilon4 carriers. *Neuroreport* 2005;16:1369–1372. [PubMed: 16056141]
- Persson J, Lind J, Larsson A, Ingvar M, Cruts M, Van Broeckhoven C, Adolfsson R, Nilsson LG, Nyberg L. Altered brain white matter integrity in healthy carriers of the APOE epsilon4 allele: a risk for AD? *Neurology* 2006a;66:1029–1033. [PubMed: 16606914]
- Persson J, Nyberg L, Lind J, Larsson A, Nilsson LG, Ingvar M, Buckner RL. Structure-function correlates of cognitive decline in aging. *Cereb Cortex* 2006b;16:907–915. [PubMed: 16162855]
- Pfefferbaum A, Sullivan EV. Increased brain white matter diffusivity in normal adult aging: relationship to anisotropy and partial voluming. *Magn. Reson. Med* 2003;49:953–961. [PubMed: 12704779]
- Ringman JM, O'Neill J, Geschwind D, Medina L, Apostolova LG, Rodriguez Y, Schaffer B, Varpetian A, Tseng B, Ortiz F, Fitten J, Cummings JL, Bartzokis G. Diffusion tensor imaging in preclinical and presymptomatic carriers of familial Alzheimer's disease mutations. *Brain* 2007;130:1767–1776. [PubMed: 17522104]
- Rose SE, Janke AL, Chalk JB. Gray and white matter changes in Alzheimer's disease: a diffusion tensor imaging study. *J. Magn. Reson. Imaging* 2008;27:20–26. [PubMed: 18050329]
- Rose SE, McMahon KL, Janke AL, O'Dowd B, de Zubicaray G, Strudwick MW, Chalk JB. Diffusion indices on magnetic resonance imaging and neuropsychological performance in amnesic mild cognitive impairment. *J. Neurol. Neurosurg. Psychiatry* 2006;77:1122–1128. [PubMed: 16754694]
- Rueckert D, Sonoda LI, Hayes C, Hill DL, Leach MO, Hawkes DJ. Nonrigid registration using free-form deformations: application to breast MR images. *IEEE Trans. Med. Imaging* 1999;18:712–721. [PubMed: 10534053]
- Schneider JA, Arvanitakis Z, Bang W, Bennett DA. Mixed brain pathologies account for most dementia cases in community-dwelling older persons. *Neurology* 2007;69:2197–2204. [PubMed: 17568013]
- Smith CD. Mild cognitive impairment is too late: the case for presymptomatic detection and treatment of Alzheimer's disease. *Cogn. Sci* 2007;3:127–177.
- Smith CD, Andersen AH, Kryscio RJ, Schmitt FA, Kindy MS, Blonder LX, Avison MJ. Altered brain activation in cognitively intact individuals at high risk for Alzheimer's disease. *Neurology* 1999;53:1391–1396. [PubMed: 10534240]
- Smith CD, Andersen AH, Kryscio RJ, Schmitt FA, Kindy MS, Blonder LX, Avison MJ. Women at risk for AD show increased parietal activation during a fluency task. *Neurology* 2002;58:1197–1202. [PubMed: 11971086]
- Smith SM, Jenkinson M, Johansen-Berg H, Rueckert D, Nichols TE, Mackay CE, Watkins KE, Ciccarelli O, Cader MZ, Matthews PM, Behrens TE. Tract-based spatial statistics: voxelwise analysis of multi-subject diffusion data. *Neuroimage* 2006;31:1487–1505. [PubMed: 16624579]
- Stahl R, Dietrich O, Teipel SJ, Hampel H, Reiser MF, Schoenberg SO. White matter damage in Alzheimer disease and mild cognitive impairment: assessment with diffusion-tensor MR imaging and parallel imaging techniques. *Radiology* 2007;243:483–492. [PubMed: 17456872]
- Sullivan EV, Adalsteinsson E, Pfefferbaum A. Selective age-related degradation of anterior callosal fiber bundles quantified in vivo with fiber tracking. *Cereb Cortex* 2006;16:1030–1039. [PubMed: 16207932]
- Sydykova D, Stahl R, Dietrich O, Ewers M, Reiser MF, Schoenberg SO, Moller HJ, Hampel H, Teipel SJ. Fiber connections between the cerebral cortex and the corpus callosum in Alzheimer's disease: a diffusion tensor imaging and voxel-based morphometry study. *Cereb Cortex* 2007;17:2276–2282. [PubMed: 17164468]
- Talairach, J.; Tournoux, P. Co-Planar Stereotaxic Atlas of the Human Brain. Thieme Medical Publishers; New York: 1988.
- Taylor WD, Bae JN, MacFall JR, Payne ME, Provenzale JM, Steffens DC, Krishnan KR. Widespread effects of hyperintense lesions on cerebral white matter structure. *Am. J. Roentgenol* 2007;188:1695–1704. [PubMed: 17515396]
- Teipel SJ, Stahl R, Dietrich O, Schoenberg SO, Perneczky R, Bokde AL, Reiser MF, Moller HJ, Hampel H. Multivariate network analysis of fiber tract integrity in Alzheimer's disease. *Neuroimage* 2007;34:985–995. [PubMed: 17166745]

- Tuch DS, Salat DH, Wisco JJ, Zaleta AK, Hevelone ND, Rosas HD. Choice reaction time performance correlates with diffusion anisotropy in white matter pathways supporting visuospatial attention. *Proc. Natl. Acad. Sci. U.S.A* 2005;102:12212–12217. [PubMed: 16103359]
- Xie S, Xiao JX, Gong GL, Zang YF, Wang YH, Wu HK, Jiang XX. Voxel-based detection of white matter abnormalities in mild Alzheimer disease. *Neurology* 2006;66:1845–1849. [PubMed: 16801648]
- Yasmin H, Nakata Y, Aoki S, Abe O, Sato N, Nemoto K, Arima K, Furuta N, Uno M, Hirai S, Masutani Y, Ohtomo K. Diffusion abnormalities of the uncinate fasciculus in Alzheimer's disease: diffusion tensor tract-specific analysis using a new method to measure the core of the tract. *Neuroradiology*. 2008
- Zhang Y, Schuff N, Jahng GH, Bayne W, Mori S, Schad L, Mueller S, Du AT, Kramer JH, Yaffe K, Chui H, Jagust WJ, Miller BL, Weiner MW. Diffusion tensor imaging of cingulum fibers in mild cognitive impairment and Alzheimer disease. *Neurology* 2007;68:13–19. [PubMed: 17200485]

Fig. 1.

Map showing inferotemporal regions of decreased FA in normal, high-AD risk subjects. The anatomic underlay used for illustration is the MNI-space registered target FA image. The registered average FA skeleton is represented in green. Warmer colors on red-orange scale indicate higher t -values for FA decrease at each voxel ($FA_{\text{high risk}} < FA_{\text{low risk}}$). Arrowheads in (A) mark the sagittal slice locations on the right (B) and on the left (D). Coronal slice level in A is indicated by single arrowheads in (C). The lineal patterns of FA decrease indicated by the arrowheads in (B) and (D) correspond to the inferior longitudinal fasciculus/inferior fronto-occipital fasciculus (Mori et al., 2005). Double arrowheads in (C) point to the posterior cingulum bundle on the left. Note that decreased FA adheres only to white matter regions.

Fig. 2.

Decreased FA in the right mid-cingulum bundle and splenium in normal, high-AD risk subjects. Display conventions are given in Fig. 1 legend. *Top row:* Arrowheads in (A) and (B) indicate the mid-portion of the cingulum on the right demonstrating decreased FA. A magnified view of the region indicated by a box in the mid-sagittal slice is illustrated in B. Arrows in (A) denotes symmetric cingulate areas with decreased FA on the left, but this region does not have the lineal spatial coherence of the right-sided counterpart. *Bottom row:* Arrowheads in (C) and (D) mark the region of decreased FA in the splenium of the corpus callosum. Double arrows in (D) indicate decreased FA in the posterior cingulate bundle on the left.

Fig. 3.

Map showing anterior regions of decreased FA in normal, high-AD risk subjects. Display conventions are given in Fig. 1 legend. Arrowheads in A mark the sagittal slice locations on the right (B) and on the left (D). Coronal slice level in A is indicated by single arrowheads in (C). The coherent patterns of FA decrease indicated by the arrows in (B) and (D) correspond to the inferior fronto-occipital fasciculus (Mori et al., 2005). Downward curvature of the FA decrease region, best seen on the left in (B), is typical of this tract.

Table 1A

Age and education of study participants

Group	N	Age, years	Education, years
High risk	42	57.7 ± 1.1	15.7 ± 0.4
Low risk	23	68.1 ^{**} ± 1.5	15.9 ± 0.6
<i>p</i> - Value	-	<0.0001	0.73

Standard Least Squares analysis using age and education as response variables and risk as regressor (Least square means ± S.E.M.).

Table 1B

Comparison of psychometric test scores between risk groups

	MMSE	LM Imm	LM Del	Block Des	VR	VR Del	BNT	Vocab
Risk high	29.3 ± 0.3	49.8 ± 1.6	32.6 ± 1.3	36.3 ± 1.7	80.1 ± 1.8	62.8 ± 3.6	14.4 ± 0.2	53.6 ± 1.2
Risk low	29.1 ± 0.2	51.4 ± 2.2	32.7 ± 1.8	43.3 ± 2.4	83.1 ± 2.5	65.9 ± 5.1	14.6 ± 0.3	56.8 ± 1.8
<i>p</i> -Values								
Risk	0.49	0.56	0.96	0.02*	0.32	0.63	0.44	0.14
Risk × age	0.61	0.04*	0.09	0.77	0.004**	0.19	0.77	0.49
Education	0.28	0.34	0.30	0.12	0.10	0.80	0.91	0.08
Age	0.58	0.44	0.62	0.0001** (-)	0.0004** (-)	0.009** (-)	0.58	0.13

	SRT	SRT 30	TA	TB	LF	SS	DS	Beck
Risk high	103.1 ± 3.4	8.5 ± 0.5	30.8 ± 1.2	70.7 ± 4.1	37.4 ± 2.5	15.5 ± 0.5	17.3 ± 0.7	7.3 ± 1.2
Risk low	106.9 ± 4.8	9.2 ± 0.7	25.0 ± 1.8	60.7 ± 5.8	45.6 ± 3.5	17.5 ± 0.7	17.3 ± 1.0	3.2 ± 1.6
<i>p</i> -Values								
Risk	0.52	0.41	0.01**	0.16	0.06	0.02*	0.99	0.05*
Risk × age	0.80	0.75	0.68	0.99	0.33	0.77	0.79	0.60
Education	0.29	0.99	0.90	0.78	0.05	0.70	0.34	0.05*
Age	0.09 (-)	0.05* (-)	0.002** (+)	0.002** (+)	0.06	0.03* (-)	0.61	0.66

High risk: presence of both family history of late-onset progressive dementia in a first-degree relative and possession of at least one APOE4 allele. Low risk: absence of both factors. Standard Least Squares analysis using psychometric raw scores as response variables and risk, education, age, and risk × age interaction as regressors. Least square means ± S.E.M. are entered in the first two rows, *p*-Values are unconnected; values between 0.05 down to 0.01 are denoted by a single asterisk (*), values 0.01 or less by double asterisks (**). Because no correction was made for the 16 multiple comparisons, we chose to conservatively interpret * as indicating marginal significance, and ** as significant. *Abbreviations*: MMSE, Mini Mental State Examination; LM Imm, Logical Memory I; LM Del, Logical memory II; Block Des, Block Design; VR, Visual Reproduction Immediate; VR Del, Visual Reproduction Delayed; BNT, 15-item Boston Naming Test; Vocab, Vocabulary; SRT, Selective Reminding Test Immediate; SRT Del, Selective Reminding Test Delayed; TA, Trails A; TB, Trails B; SS, Spatial Span; DS, Digit Symbol; LF, FAS Letter Fluency. The direction of correlation is indicated by a plus (+) or minus (-) sign when *p* < 0.05.

Table 2

Locations of voxels with decreased FA in high-AD risk participants

Location	Tract	Nearest BA
-27, -77, 2	Inferior fronto-occipital fasciculus L (posterior)	19
25, -78, 2	Inferior fronto-occipital fasciculus R (posterior)	19
-38, -63, -8	Inferior fronto-occipital fasciculus L (posterior)	19, 37
39, -64, -8	Inferior fronto-occipital fasciculus R (posterior)	19, 37
-37, -55, 0	Inferior longitudinal fasciculus L (posterior)	37
36, -59, 5	Inferior longitudinal fasciculus R (posterior)	37
-19, -36, -7	Cingulum bundle L (posterior)	27, 30
7, -8, 32	Cingulum bundle R (mid)	23, 24
-42, -25, -6	Inferior fronto-occipital fasciculus L (posterior)	21
-15, 2, 56	Cortico-pontine tract	6
-24, 23, 8	Inferior fronto-occipital fasciculus L (anterior)	-
26, 28, 2	Inferior fronto-occipital fasciculus L (anterior)	-
0, -36, 21	Splenium, corpus callosum	-
-4, 3, -10	Subcallosal white matter	25
4, 3, -10	Subcallosal white matter	25

Locations are given in Talairach coordinates in the first Column (Talairach and Tournoux, 1988, 1027), the corresponding tract associated with this anatomic location in the second Column (Mori et al., 2005 #67), and the nearest neighboring Brodmann cortical area (BA) to the location in the third Column.