

University of Kentucky

UKnowledge

---

Entomology Faculty Publications

Entomology

---

2-7-2012

## RNA Interference of NADPH-cytochrome P450 Reductase Results in Reduced Insecticide Resistance in the Bed Bug, *Cimex lectularius*

Fang Zhu

University of Kentucky, fangzhudy@uky.edu

Sarah Sams

University of Kentucky

Tim Moural

Bluegrass Community & Technical College

Kenneth F. Haynes

University of Kentucky, ken.haynes@uky.edu

Michael F. Potter

University of Kentucky, mpotter@uky.edu

See this page for additional authors [https://uknowledge.uky.edu/entomology\\_facpub](https://uknowledge.uky.edu/entomology_facpub)



Part of the [Agriculture Commons](#), and the [Entomology Commons](#)

[Right click to open a feedback form in a new tab to let us know how this document benefits you.](#)

---

### Repository Citation

Zhu, Fang; Sams, Sarah; Moural, Tim; Haynes, Kenneth F.; Potter, Michael F.; and Palli, Subba R., "RNA Interference of NADPH-cytochrome P450 Reductase Results in Reduced Insecticide Resistance in the Bed Bug, *Cimex lectularius*" (2012). *Entomology Faculty Publications*. 2.

[https://uknowledge.uky.edu/entomology\\_facpub/2](https://uknowledge.uky.edu/entomology_facpub/2)

This Article is brought to you for free and open access by the Entomology at UKnowledge. It has been accepted for inclusion in Entomology Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

---

## RNA Interference of NADPH-cytochrome P450 Reductase Results in Reduced Insecticide Resistance in the Bed Bug, *Cimex lectularius*

Digital Object Identifier (DOI)

<http://dx.doi.org/10.1371/journal.pone.0031037>

### Notes/Citation Information

Published in *PLoS ONE*. v. 7, no. 2, e31037.

© 2012 Zhu et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### Authors

Fang Zhu, Sarah Sams, Tim Moural, Kenneth F. Haynes, Michael F. Potter, and Subba R. Palli

# RNA Interference of NADPH-Cytochrome P450 Reductase Results in Reduced Insecticide Resistance in the Bed Bug, *Cimex lectularius*

Fang Zhu<sup>1</sup>, Sarah Sams<sup>1</sup>, Tim Moural<sup>2</sup>, Kenneth F. Haynes<sup>1</sup>, Michael F. Potter<sup>1</sup>, Subba R. Palli<sup>1\*</sup>

<sup>1</sup> Department of Entomology, College of Agriculture, University of Kentucky, Lexington, Kentucky, United States of America, <sup>2</sup> Division of Natural Sciences, Bluegrass Community & Technical College, Lexington, Kentucky, United States of America

## Abstract

**Background:** NADPH-cytochrome P450 reductase (CPR) plays a central role in cytochrome P450 action. The genes coding for P450s are not yet fully identified in the bed bug, *Cimex lectularius*. Hence, we decided to clone cDNA and knockdown the expression of the gene coding for CPR which is suggested to be required for the function of all P450s to determine whether or not P450s are involved in resistance of bed bugs to insecticides.

**Methodology/Principal Findings:** The full length *Cimex lectularius* CPR (*CICPR*) cDNA was isolated from a deltamethrin resistant bed bug population (CIN-1) using a combined PCR strategy. Bioinformatics and *in silico* modeling were employed to identify three conserved binding domains (FMN, FAD, NADP), a FAD binding motif, and the catalytic residues. The critical amino acids involved in FMN, FAD, NADP binding and their putative functions were also analyzed. No signal peptide but a membrane anchor domain with 21 amino acids which facilitates the localization of *CICPR* on the endoplasmic reticulum was identified in *CICPR* protein. Phylogenetic analysis showed that *CICPR* is closer to the CPR from the body louse, *Pediculus humanus corporis* than to the CPRs from the other insect species studied. The *CICPR* gene was ubiquitously expressed in all tissues tested but showed an increase in expression as immature stages develop into adults. We exploited the traumatic insemination mechanism of bed bugs to inject dsRNA and successfully knockdown the expression of the gene coding for *CICPR*. Suppression of the *CICPR* expression increased susceptibility to deltamethrin in resistant populations but not in the susceptible population of bed bugs.

**Conclusions/Significance:** These data suggest that P450-mediated metabolic detoxification may serve as one of the resistance mechanisms in bed bugs.

**Citation:** Zhu F, Sams S, Moural T, Haynes KF, Potter MF, et al. (2012) RNA Interference of NADPH-Cytochrome P450 Reductase Results in Reduced Insecticide Resistance in the Bed Bug, *Cimex lectularius*. PLoS ONE 7(2): e31037. doi:10.1371/journal.pone.0031037

**Editor:** Guy Smaghe, Ghent University, Belgium

**Received:** November 2, 2011; **Accepted:** December 31, 2011; **Published:** February 7, 2012

**Copyright:** © 2012 Zhu et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This work was funded by grants from Bayer and Kentucky Agricultural experimental station. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have read the journal's policy and have the following conflicts: Funding for this study was provided by a grant from Bayer Environmental Sciences. There are no patents, products in development or marketed product to declare. The authors' grant agreement with Bayer does not alter their adherence to all the PLoS ONE policies on sharing data and materials.

\* E-mail: RPALLI@UKY.EDU

## Introduction

During the past ten years, the bed bug has rapidly resurfaced throughout the world [1–3]. In an industry-wide survey of pest management firms in the United States, 76% considered bed bugs to be the most difficult pest to control [4]. It has been proposed that the global bed bug resurgence is partly due to the ubiquitous development of pyrethroid resistance [5–7]. Our previous studies determined a causal link between two identified knockdown resistance (*kdr*) mutations and deltamethrin resistance in bed bug populations, indicating decreased target-site sensitivity of voltage-gated sodium channels as one of the mechanisms of pyrethroid resistance. Interestingly, one population of bed bugs collected from Cincinnati (CIN-1) showed more than 12,765-fold deltamethrin resistance but no mutations were detected in the voltage-gated sodium channel gene [7]. Moreover, PBO, a P450 inhibitor, suppressed deltamethrin resistance in CIN-1 population [8]. These

data suggest that cytochrome P450-mediated metabolic detoxification might be a principal mechanism responsible for deltamethrin resistance in some bed bug populations. However, the identity of P450s involved in detoxification of pyrethroids in bed bugs remains unknown.

Cytochrome P450s constitute one of the largest superfamilies of enzymes that play important roles in detoxification of xenobiotics [9,10] as well as in biosynthesis and metabolism of endogenous compounds [11,12]. The reaction of the P450 system requires electrons transferred from Nicotinamide Adenine Dinucleotide Phosphate (NADPH) to the P450 heme center by a Cytochrome P450 partner enzyme, NADPH-Cytochrome P450 Reductase (CPR) [13]. Although, multiple P450 genes have been found in the genomes of insects (<http://drnelson.uthsc.edu/cytochromeP450.html>), typically only one *CPR* gene exists in each insect genome. CPR is a multidomain protein which belongs to the electron transfer flavoproteins family [14] containing both Flavin Adenine

Dinucleotide (FAD) and Flavin Mononucleotide (FMN) domains [15]. In addition to cytochrome P450s, CPR also serves as the electron donor protein for several oxygenase enzymes found in the endoplasmic reticulum of most eukaryotic cells [16–19]. Genes coding for CPRs have been identified and characterized from a few species of insects, such as house fly, *Musca domestica* L. [20–22], fruit fly, *Drosophila melanogaster* (Meigen) [23], silkworm, *Bombyx mori* L. [24], cabbage armyworm, *Mamestra brassicae* (L.) [25], mosquitoes, *Anopheles gambiae* Giles [26] and *Anopheles minimus* Theobald [27]. Sequences of CPR cDNAs are also available for many other insect species (Table S1).

With critical biological function associated with cytochrome P450s, insect CPRs have been placed in a vital path in metabolism-based insecticide resistance and were considered as the novel target for the development of synergists [26,28]. In the current study, the *Cimex lectularius* CPR (CICPR) cDNA was cloned and the gene coding for CPR was silenced in both deltamethrin resistant and susceptible populations of bed bugs. The data collected helped to reveal the role of P450-mediated metabolic detoxification in the deltamethrin resistance of bed bugs.

## Materials and Methods

### The bed bugs

Three bed bug colonies were used in this study. One insecticide-susceptible colony, LA-1, collected in 2006 in Los Angeles, CA [29] was maintained in the laboratory without any insecticide exposure. Two deltamethrin resistant populations of bed bugs, CIN-1 (Cincinnati, OH, contains no *kdr* mutations) and NY-1 (Plainview, NY, contains two *kdr* mutations) were collected from human dwellings in the United States during 2006–2008 [9] and maintained in the laboratory by using a parafilm-membrane feeder. Bed bugs were kept in screened containers and fed with 37°C heparinized chicken blood or rabbit blood with sodium citrate through a thinly stretched parafilm membrane [30]. Blood was purchased from Hema Resource and Supply Company (Aurora, OR). Bed bugs were reared at 27°C, 65±5% RH, and a photoperiod of 14:10 (L:D) h.

### RNA extraction, cDNA preparation and Cloning

Total RNA was isolated from 3 CIN-1 adults using the TRI reagent (Molecular Research Center Inc., Cincinnati, OH) and the RNA was treated with DNase I (Ambion Inc., Austin, TX). cDNA was synthesized using iScript cDNA synthesis kit (Bio-Rad Laboratories, Hercules, CA) with DNase I treated total RNA as a template. The PCR products were amplified using primer pair NADPHF/NADPHR (Table S2) that was designed based on a conserved amino acid region found in 10 insect CPR sequences. The PCR products were cloned into pGEM<sup>®</sup>-T Easy Vector Systems (Promega) and sequenced. Cloning and sequence analyses of P450 gene fragments were repeated at least three times with different preparations of RNAs. Three clones from each replication were sequenced.

### Rapid amplification of cDNA ends (RACE) of the putative CICPR gene fragment

RACE was carried out using the SMARTer<sup>™</sup> RACE cDNA Amplification Kit (Clontech) as described in the manufacturer's manual. The first strand cDNAs were synthesized with SMART-Scribe<sup>™</sup> Reverse Transcriptase using CIN-1 RNA as a template. The double-stranded cDNA was synthesized following the protocol described in the manufacturer's manual (Clontech). The 5' and/or 3' ends of the P450 cDNA fragments were amplified by PCR using adapter primers UPM and NUP and gene

specific primers generated based on the 5' and/or 3' end sequences of the putative *CICPR* transcript (Table S2). The full length of putative *CICPR* cDNA was subsequently generated by RT-PCR using specific primer pair of CICPRF/CICPRR (Table S2) synthesized based on the 5' and 3' end sequences of the putative *CICPR* mRNA. Cloning and sequence analyses of the *CICPR* transcript were repeated at least three times, and three clones from each replicate were verified by sequencing.

### In silico structural analysis

The Isoelectric point (pI) and Molecular Weight (MW) of CICPR were calculated by an ExPASy proteomics tool, Compute pI/Mw ([http://web.expasy.org/compute\\_pi](http://web.expasy.org/compute_pi)) from the Swiss Institute of Bioinformatics. The signal peptide and protein subcellular localization of CICPR were analyzed at the SignalP 3.0 server (<http://www.cbs.dtu.dk/services/SignalP/>) and WoLF PSORT (<http://wolfpsort.org/>). The secondary structure, binding domains, and catalytic residues were predicted by PHYRE2 Protein Fold Recognition Server (<http://www.sbg.bio.ic.ac.uk/phyre2/html/>), Pfam 25.0 (2011, <http://pfam.sanger.ac.uk/>), and a conserved domain search on the NCBI website (<http://www.ncbi.nlm.nih.gov/Structure/cdd/cdd.shtml>) [31]. The protein tertiary structure of CICPR was predicted by using the I-TASSER server (<http://zhanglab.cmb.med.umich.edu/I-TASSER/>) [32] and then the PDB coordinate file of the highest ranking model was loaded into Chimera (<http://plato.cgl.ucsf.edu/chimera/docs/credits.html>) for molecular visualization and modification. The transmembrane helices of CICPR were analyzed by TMHMM Server 2.0 (<http://www.cbs.dtu.dk/services/TMHMM-2.0/>). The hydrophobicity of CICPR was predicted by an on line molecular tool, Protein Hydrophobicity Plots (<http://www.vivo.colostate.edu/molkit/hydrophathy/>).

### Phylogenetic tree construction

All CPR sequences in insects which have the complete open reading frames (ORFs) were extracted from the National Center for Biotechnology Information (NCBI) (Bethesda, MD) (<http://www.ncbi.nlm.nih.gov/>). The insect CPR amino acid sequences were analyzed using ClustalW alignment through Molecular Evolutionary Genetic Analysis software version 5 (MEGA 5) (<http://www.megasoftware.net/>) [33]. To improve the alignments, the pair wise alignment was performed with the gap opening penalty at 10, and the gap extension penalty left at default 0.1. The multiple alignments were conducted with the gap opening penalty at 3 and the gap extension penalty at 1.8 [34]. The sites containing missing data or alignment gaps were eliminated in a pair-wise manner. A p-distance<0.8 when carrying out the compute overall mean distance suggested the alignment was acceptable [34]. Subsequently, the alignment result was converted to a MEGA file (.meg) and submitted to construct the phylogenetic tree with neighbor-joining algorithm. A total of 2,000 bootstrap replications were used to test of phylogeny. Ultimately, the selected tree was created with cut-off value of 50%.

### Quantitative real time PCR (qRT-PCR) and reference gene selection

qRT-PCR was performed in MyiQ single color real-time PCR detection system (Bio-Rad Laboratories, Hercules, CA). Total RNA was isolated from 3 female bed bugs at 5 days after RNAi treatment using the TRI reagent (Molecular Research Center Inc., Cincinnati, OH). The RNA was treated with DNase I (Ambion Inc., Austin, TX). cDNA was synthesized using iScript cDNA synthesis kit (Bio-Rad Laboratories, Hercules, CA). DNase I treated total RNA was used as a template. Each qRT-PCR

reaction (10  $\mu$ l final volume) contained 5  $\mu$ l FastStart SYBR Green Master (Roche Diagnostics, Indianapolis, IN), 1.2  $\mu$ l of cDNA, and 0.6  $\mu$ l each of forward and reverse gene specific primers (Table S2, stock 10  $\mu$ M). An initial incubation of 95°C for 3 min, followed by 40 cycles of 95°C for 10 s, 55°C for 20 s, and 72°C for 30 s settings were used. A fluorescence reading determined the extension of amplification at the end of each cycle. Each experiment was repeated at least three times using independent biological samples.

The suitability of four reference/control genes, *rpl11*, *rpl8*, *rps16* and *hsp70* was evaluated with the *Bestkeeper* software package [35,36]. This program was used not only to calculate potential reference genes, but also to assess the effects of RNAi on target genes. We designed primers for reference genes based on the EST sequences in the GenBank database (GenBank Accession Nos.: *rpl11*, EZ419774; *rpl8*, EZ419796; *rps16*, EZ419784; *hsp70*, EZ419756). Primers used for amplification reference genes are shown in Table S2. Relative expression levels for specific genes, in relation to the most reliable reference gene, were calculated by the  $2^{-\Delta\Delta CT}$  method [37].

### dsRNA injection by traumatic insemination mimicking

The dsRNA was synthesized using the MEGascript RNAi Kit (Ambion Inc., Austin, TX). Genomic DNA was isolated from CIN-1 adults using DNeasy Tissue Kit (QIAGEN). Genomic DNA and T7 promoter-containing PCR primers (Table S2, with T7 RNA promoter sequence (TAATACGACTCACTATAGGG) appended at the 5' ends of both sense and antisense specific primers) were used in a PCR reaction to obtain gene specific fragments containing T7 promoter sequence on both ends. PCR product (200–400 bp) was used as a template to synthesize dsRNA. The same length PCR fragments were obtained using either genomic DNA or cDNA as a template suggesting that there is no intron in the region of this gene used for dsRNA preparation. For the dsRNA purification, phenol/chloroform extraction followed by ethanol precipitation method was applied. dsRNA was diluted in nuclease-free water to 4–5  $\mu$ g/ $\mu$ l for injection into bed bug adults. The one week old female adults were anaesthetized with ether vapor for 10 min and placed on a glass slide covered with double-sided tape. The dsRNA (~1.25  $\mu$ g) was injected into the spermatheca of the abdomen with an injection needle pulled out from a glass capillary tube using a needle puller (Idaho Technology, Salt Lake City, Utah). The spermatheca is where the cuticle of the female is punctured during traumatic insemination. Prior to injection, the glass needles were sterilized by soaking in 100% ethanol for 12 h. Controls were injected with the dsRNA using bacterial *malE* gene as a template. After injection, insects were removed from the glass slide, allowed to recover for 3 h at room temperature, then returned to normal rearing conditions.

### Bioassays with deltamethrin after dsRNA injection

In the preliminary studies, bed bug adults were treated with serial dilutions of technical grade deltamethrin (99% active ingredient, Bayer Environmental Science, St. Louis, MO) prepared in acetone. A discriminating dose (causing approximately 50% of mortality) of deltamethrin was applied for the bioassays. Acetone was used as a control. The solution was dropped on the thorax of the bugs (1  $\mu$ l/drop) using a PB-600 repeating dispenser (Hamilton Co., Reno). The mortality was determined at 24 h after treatment. Mean and standard errors for each time point were obtained from at least three independent bioassays.

### Statistical analysis

Statistical analyses were carried out using SAS software (v9.1, SAS Institute Inc., Cary, NC). Student's *t*-test (two-tailed paired *t*-test) was

used to compare the gene expression and mortality difference between two samples. The differences among samples were analyzed by One-way ANOVA, followed by Duncan multiple mean separation techniques. The level of significance was set at  $P < 0.05$ .


## Results

### Cloning, sequence analysis, and structural modeling of ClCPR

The overall strategy of cloning the full length of ClCPR is shown in Figure S1. Briefly, a partial putative ClCPR cDNA fragment was amplified from deltamethrin resistant population, CIN-1, by multiple PCR amplifications using degenerate primers, NADPHF and NADPHR designed based on CPR sequences identified in other insect species (Table S2). BLAST analysis of the amino acid sequence predicted from the partial putative ClCPR cDNA sequence showed that the sequence encoded ClCPR and shared 81% amino acid similarity with the CPR sequence from the body louse, *Pediculus humanus corporis*. To amplify the 5' and 3' ends of this gene, 5'-RACE and 3'-RACE reactions were conducted using the adapter primers and gene specific primers designed based on the 5' and 3' end sequence of the putative ClCPR cDNA fragment, respectively (Table S2). The sequences of the 5'-RACE and 3'-RACE fragments overlapped with the ClCPR cDNA fragment sequence, identifying them as the 5' and 3' ends of the putative ClCPR gene. Subsequently, the full length ClCPR cDNA was amplified from CIN-1 bed bugs using PCR and the specific primers (ClCPRF and ClCPRR) designed based on the 5' and 3' end sequences of the putative ClCPR gene (Fig. S1, Table S2). The ClCPR amino acid sequence was aligned with the amino acid sequences of other CPRs from taxonomically diverse insect species. The sequence of ClCPR showed 65%, 68%, 66%, and 70% amino acid identity respectively with the CPR sequences of *An. gambiae*, *D. melanogaster*, *M. brassicae*, and *M. domestica*, (Table S1).

The ClCPR cDNA sequence contained an ORF of 2037 nucleotides encoding 679 amino acids. The pI and Mw were predicted as 5.55 and 77.01 kilodaltons, respectively. No signal peptide was found within ClCPR, but the membrane anchor which facilitates the localization of ClCPR on the endoplasmic reticulum was identified (Fig. 1A). All functional domains involved in the binding of cofactors, FMN, FAD and NADPH were identified in the predicted ClCPR protein primary and tertiary structures (Fig. 1). Three amino acid residues including Arginine 457, Tyrosine 459, and Serine 460 constitute a FAD binding motif which is similar to the conserved FAD binding domain [38] (Fig. 1A). The ClCPR catalytic residues (active site) comprise Serine 460, Cysteine 631, Aspartic 676, and Tryptophan 678 (Fig. 1A). These residues have been shown to be critical in the hydride transfer reaction catalyzed by rat CYP Oxidoreductase [39,40].

To understand the structure which may direct function, the three-dimensional model of ClCPR was predicted. As shown in Fig. 1B, ClCPR consists of three distinct protein binding domains, FMN binding domain, FAD binding domain, and NADP binding domain. The FAD and NADP binding domains present at the C terminus are similar to the FAD/NADP domain reported in human CPR [41]. The FAD binding pocket is composed of thirteen amino acids, and the NADP binding pocket is composed of fifteen amino acids (Fig. 1B). At the N terminus, the FMN domain with two FMN binding sites is structurally similar to flavodoxins and interacts with the redox-partner binding site of the P450s [42] (Fig. 1C). During a catalytic cycle, ClCPR is predicted to transfer a hydride ion derived from NADPH to FAD, and then FAD transfers electrons to FMN, from where they are delivered to acceptor proteins (such as P450s) [15,43].



**Figure 1. Structure of CICPR.** (A) Schematic drawing of CICPR with membrane anchor (orange bar), conserved binding domains (green bar-Flavodoxin, blue bar-FAD binding, cyan bar-NADP binding), FAD binding motif (ArgxTyrSer), and catalytic residues (Ser-Cys-Asp-Trp). (B) Predicted three-dimensional structure of CICPR with emphasis on FAD and NADP binding pockets. Three binding domains are highlighted in different colors (green-Falvodoxin, blue-FAD binding, and cyan-NADP binding) in the model. Fifteen amino acids composing the NADP binding pocket are highlighted as red spheres. Thirteen amino acids which constitute the FAD binding pocket are highlighted as yellow spheres. N- and C- termini are also labeled in the CICPR tertiary structure. (C) Sequence alignment for FMN binding sites in insect *CPRs*. Residues constituting the FMN1-binding site were labeled with red numbers, and the residues constituting the FMN2-binding site are labeled with blue numbers. The arrows show the direction from the N terminus to the C terminus. All insect *CPR* amino acid sequences were extracted from NCBI (Bethesda, MD) (<http://www.ncbi.nlm.nih.gov/>). The sequence alignment was performed using ClustalW through MEGA 5 [33]. The cDNA sequence of CICPR has been deposited in the GenBank database, accession number, JQ178363. doi:10.1371/journal.pone.0031037.g001

### Subcellular localization of CICPR

No conserved signal peptide was identified at the N-terminal end of CICPR suggesting that CICPR is retained in the cytoplasm. The *CPR* is anchored on the membrane of endoplasmic reticulum by an N-terminal hydrophobic segment [44]. A deduced hydrophobic transmembrane region consisting of 21 amino acids identified at the N-terminal end of CICPR may be involved in the membrane anchor function (Fig. S2).


### Phylogenetic relationship of CICPR with other insect *CPRs*

Phylogenetic analysis was performed based on the amino acid sequences of CICPR and 17 other insect *CPR* proteins (Table S1). The phylogenetic tree was generated by MEGA 5 with the neighbor-joining algorithm. As expected, insect *CPRs* from the same insect order were grouped together (Fig. 2). Among all the

sequences, CICPR shared the highest sequence similarity (75%) with the *CPR* of the body louse, *Pediculus humanus corporis* (Table S1). It was consistent with the result of phylogenetic analysis, in which CICPR originated from a same evolutionary root with the *CPR* in *P. humanus corporis* with the bootstrap value of 82 (Fig. 2) even though they do not belong to the same taxonomic order.

### Bed bug reference gene selection

Since very little information is available on gene expression in the bed bug, we decided to identify a housekeeping gene for qRT-PCR analysis. Results of reference gene examination RNA isolated from whole body samples across different developmental stages and populations are shown in Table S3 and Fig. S3. Based on the two most important criteria for evaluating the stability of reference genes by the *BestKeeper* program, the stability (SD value) and the relation to the *BestKeeper* index ( $r$  and  $P$ -value), all four reference



**Figure 2. The neighbor-joining consensus tree illustrates the phylogenetic relationship of CICPR with other insect CPRs.** The phylogenetic tree was generated by MEGA 5 according to the amino acid sequences. All nodes have significant bootstrap support based on 2,000 replicates. All insects CPRs were clustered into six groups which was six insect orders. The CPRs from bed bug (*Cimex lectularius*) and body louse (*Pediculus humanus corporis*) which showed the closest evolutionary relationship with *CICPR* were highlighted in black boxes. doi:10.1371/journal.pone.0031037.g002

genes, *rpl11*, *rpl8*, *rps16* and *hsp70* tested are stable across developmental stages and populations tested. The *rpl8* was chosen as the reference gene to calculate relative expression levels of *CICPR* because it showed the most stable expression among samples tested. In subsequent studies, *rpl8* expression was found to be stable across different tissues as well as dsRNA injected and control insects (data not shown). Therefore, *rpl8* was used as a reference gene in all the experiments.

#### Developmental, sexual, and spatial expression patterns of *CICPR*


The developmental, sexual, and spatial expression of *CICPR* gene was investigated using qRT-PCR. The mRNA levels of *CICPR* in 1 to 5 day pooled eggs, small nymphs (1<sup>st</sup> to 3<sup>rd</sup> instar), large nymphs (4<sup>th</sup> and 5<sup>th</sup> instar), and 1 week old female and male adults in both resistant CIN-1 and susceptible LA-1 populations were quantified. Low levels of *CICPR* mRNA were detected in eggs, small nymphs, and large nymphs in both CIN-1 and LA-1 populations (Figs. 3A and B). There were no significant differences in the *CICPR* mRNA levels among these samples. The mRNA levels of *CICPR* in female and male adults were significantly higher than those in the early stages of both populations (Figs. 3A and B). Moreover, the *CICPR* mRNA was detected in the antennae, head, thorax, and abdomen isolated from female CIN-1 adults (Fig. 3C).

#### Dose dependent and systemic *CICPR* knockdown

The bed bugs are unusual in being a gonochorist taxon with obligate traumatic insemination. During the process of copulation,

the male pierces the female's abdominal wall and transfers sperm into her haemocoel [2]. However, the female bed bugs evolved a unique anatomical organ, the spermalege, into which the male punctures the female with a needle-like paramere. Research suggested that the evolution of spermalege resulted from selection to defense against mating-associated pathogens [45]. Preliminary studies showed that injection of dsRNA through spermalege caused lower mortality compared to the injections at the other sites in the abdomen. Therefore, the dsRNAs were routinely injected through the spermalege into the body of female bed bugs (Fig. 4). Bed bugs injected with *malE* or *CICPR* dsRNA suffered similar rate of mortality within 5 days after injection (most of them died in the first one or two days) (Fig. S4). There was no other obvious negative effects caused by injecting *CICPR* dsRNA observed during the 6-day experimental period (including 5 days after injection and 24 h for bioassay).

A preliminary study showed that 1.25  $\mu$ g of *CICPR* dsRNA was sufficient to silence *CICPR* gene in each individual bed bug. In order to identify the most effective dose for silencing the *CICPR* gene, serial 10-fold dilutions of *CICPR* dsRNA were injected and the *CICPR* mRNA levels were quantified using qRT-PCR and total RNA isolated at 5 days after injection of dsRNA. As shown in Figure 5A, 0.125  $\mu$ g/individual of *CICPR* dsRNA was the most effective dose to suppress the expression of *CICPR* gene in CIN-1 population. Subsequently, we detected *CICPR* knockdown efficiency in different body parts, including head, thorax, and abdomen. RNAs extracted from these body parts of both control (injected with dsRNA of *malE*, a bacterial gene) and *CICPR* dsRNA



**Figure 3. Spatial and temporal expression of *CICPR*.** Changes in mRNA levels of the *CICPR* in CIN-1 (A) and LA-1 (B) populations. Egg; SN, small nymph (1–3 instar); LN, large nymph (4–5 instar); female and male, 1 week old. The relative mRNA levels were shown as a ratio in comparison with the levels of *rp18* mRNA. The data shown are mean+SEM (n = 3). (C) Relative mRNA levels of the *CICPR* in the antennae, head, thorax, and abdomen of the CIN-1. Tissues were dissected and total RNAs were isolated to quantify the *CICPR* mRNA levels by qRT-PCR as described in Materials and Methods. Relative mRNA levels were normalized using the expression of *rp18*. The data shown are mean+SEM (n = 4). Statistical significance of the gene expression among samples was calculated using one-way ANOVA followed by Duncan multiple mean separation techniques. There was no significant difference among relative expression within samples with the same alphabetic letter (i.e. a, b and c). doi:10.1371/journal.pone.0031037.g003

treated bed bugs were subjected to qRT-PCR analysis. The *CICPR* gene was successfully suppressed in all body parts tested, indicating that the RNAi effect in bed bugs is systemic (Fig. 5B).



**Figure 4. Bed bug dsRNA injection by mimicking traumatic insemination.** Female bed bug showing the site of dsRNA injection via the spermatheca using a sterilized glass needle. doi:10.1371/journal.pone.0031037.g004

### *CICPR* knockdown increases CIN-1 and NY-1 sensitivity to deltamethrin


Five days after injection of dsRNA, the survived bed bugs were exposed to deltamethrin through topical application. The percent survival was recorded after 24 h exposure to deltamethrin. The *CICPR* knockdown in deltamethrin resistant populations CIN-1 (no *kdr* mutation) and NY-1 (two *kdr* mutations) bed bugs showed a consistent increase in susceptibility to deltamethrin compared with control bed bugs (Figs. 6A and 6B). In contrast, there was no significant difference in the susceptibility to deltamethrin between *CICPR* knockdown and control in insecticide susceptible LA-1 bed bugs (Fig. 6C).

## Discussion

### Overview

The main goal of this study is to characterize NADPH-Cytochrome P450 reductase from the bed bug and investigate whether the P450-mediated metabolic detoxification plays any role in the deltamethrin resistance of bed bugs. To achieve the goal, the *CICPR* gene was isolated from a deltamethrin resistant population (CIN-1) with a combined PCR strategy (Fig. S1). Three conserved binding domains, a FAD binding motif, and the catalytic residues as well as the critical residues involved in FMN, FAD and NADP binding were identified (Fig. 1). The spatial






**Figure 5. Relative *CICPR* mRNA levels in control (*malE* dsRNA) and *CICPR* dsRNA injected bed bugs.** (A) The *CICPR* mRNA levels were quantified by qRT-PCR at 5 days after dsRNA injection in control (*malE* dsRNA) and *CICPR* dsRNA treated bed bugs with different doses of dsRNA. (B).Relative *CICPR* mRNA levels in different body parts in control (*malE* dsRNA) and *CICPR* dsRNA injected bed bugs. The relative *CICPR* mRNA levels are shown as a ratio in comparison with the levels of *rpl8* mRNA. The data shown are mean+SEM (n=3). doi:10.1371/journal.pone.0031037.g005

configuration and the putative functions of these conserved domains were analyzed by predicting a 3-D model of *CICPR* (Fig. 1B). The *CICPR* was predicted anchoring on the membrane of endoplasmic reticulum by a 21 amino acids transmembrane region (Figs. 1A and S2). The phylogenetic analysis showed that *CICPR* had the shortest genetic distance to the CPR from body louse (Fig. 2). The *CICPR* gene was ubiquitously expressed in all tissues tested (Fig. 3A and B) but showed an increase in expression as immature stages develop into adults (Fig. 3A). With mimicking the traumatic insemination of bed bugs, dsRNA of *CICPR* was injected into the bed bug and successfully suppressed the expression of the gene coding for *CICPR* throughout the body (Figs. 4 and 5). Similar to the previous report from *An. gambiae* [26], when the *CICPR* was suppressed through RNAi in deltamethrin resistant bed bug populations, the susceptibility of these bed bugs to deltamethrin was significantly enhanced. In this study, the

susceptibility enhancement was observed in populations containing *kdr* mutation (NY-1, Fig. 6B) as well as no-*kdr* mutations (CIN-1, Fig. 6A), but not in the susceptible population (LA-1, Fig. 6C), suggesting P450-mediated metabolic detoxification may serve as one of the resistance mechanisms employed by bed bugs.

#### *CICPR* gene discovery and analysis

As an obligatory electron donor, CPR transfers electrons from NADPH to various cytochrome P450s that play central roles in detoxification of xenobiotics [41]. Consequently, identification and characterization of CPR from insects will help to determine whether or not Cytochrome P450s are involved in response of insects to specific insecticides and other xenobiotics [26,28]. The primary structures of CPRs are highly conserved across diverse taxa, indicating the functional importance of this enzyme throughout the course of evolution [46]. The alignment of *CICPR*



**Figure 6. Knockdown in the expression of *CICPR* reduced the resistance to deltamethrin.** (A) The percent survival of dsRNA treated CIN-1 (deltamethrin resistant population without para-type sodium channel gene mutation at 419 aa and 925 aa) bed bugs at 0.06 µg deltamethrin 5 days after dsRNA injection. The mortality was recorded after 24 h exposure to deltamethrin (3 replicates, 50–60 individuals for each replicate). (B) The % survival of dsRNA treated NY-1 (deltamethrin resistant population with 2 para-type sodium channel gene mutations at 419 aa and 925 aa) bed bugs at 0.06 µg deltamethrin 5 days after dsRNA injection. The mortality was recorded after 24 h exposure to deltamethrin (3 replicates, 38–40 individuals for each replicate). (C) The % survival of dsRNA treated LA-1 (deltamethrin susceptible population without para-type sodium channel gene mutation at 419 aa and 925 aa) bed bugs at 0.0006 µg deltamethrin 5 days after dsRNA injection. The mortality was recorded after 24 h exposure to deltamethrin (3 replicates, 60 individuals for each replicate). The differences between control and CPR-KD in three bed bug populations were analyzed by Student's *t*-test. doi:10.1371/journal.pone.0031037.g006

with other CPRs showed that *CiCPR* shares 64–75% amino acid identity with other insect CPRs, and 54% with human (*Homo sapiens*), rat (*Rattus norvegicus*) and mouse (*Mus musculus*) CPRs. The structure analysis of *CiCPR* demonstrated that *CiCPR* has four functional domains (Fig. 1). The hydrophobic N-terminal membrane anchor of *CiCPR* is essential for its function in the P450 catalytic cycle. It serves to anchor the protein molecule to the endoplasmic reticulum which ensures proper spatial interaction for electron transfer between the *CiCPR* and cytochrome P450s [41,46]. Without the hydrophobic anchor, *CiCPR* is incapable of transferring electrons to cytochrome P450s. *CiCPR* also has three distinct protein binding domains. At the N-terminal, FMN binding domain consists of two FMN binding sites, FMN1 and FMN2 (Figs. 1A and C). FMN1 and FMN2 binding sites were well conserved among all insect CPRs indicating they play critical roles in the interaction with cytochrome P450s (Fig. 1C). For example, mutation of residues corresponding to Asp 211 and Thr 93 in *C. lectularius* to alanine in the yeast CPR resulted in complete loss of functional activity toward CYP51 [42].

The phylogenetic analysis showed that *CiCPR* had the closest evolutionary relationship with the CPR from the body louse. Although the bed bug and the body louse do not belong to the same taxonomic order and occupy different living habitats, these two species are both obligatory hematophagous (bloodsucking) insects which maintain close association with the human being throughout their life stages. Feeding on human blood implies that they share some of the same xenobiotic challenges, therefore may have evolved a similar function for their CPRs.

#### Tissue distribution of *CiCPR*

The primary tissue distribution of *CPR* is associated with the potential localization of P450 activities, which in turn reflect the prospective function of P450s. For example, the *D. melanogaster CPR* was expressed more abundantly in embryos and antennae as compared to adult heads, adult bodies and larvae [23] indicating the functions of *D. melanogaster* P450s in embryonic development and odorant clearance. The cabbage armyworm, *M. brassicae*, *CPR* was predominantly expressed in all tissues tested including male and female antenna, male brain, proboscis, thorax, abdomen, and legs and female ovipositors. The cellular localization of *CPR* and two P450s within antennae suggest the potential importance of cytochrome metabolism in the olfactory sensilla of the cabbage armyworm [25]. *An. gambiae CPR* was mainly localized in the antennae, mid-gut epithelia and oenocytes which are considered to be a major spot for heme biosynthesis in insects [26]. These observations suggest potential roles of *A. gambiae* P450s in odorant metabolism, insecticides metabolism, and regulating heme homeostasis. All insects are living in their environments surrounded by various chemicals, volatile or nonvolatile, natural or anthropic, useful or harmful. The pores located on the antenna allow the entrance of massive odor molecules which potentially trigger the olfactory signaling transduction in the olfactory sensillum and subsequently are metabolized by odorant-degrading enzymes (including Cytochrome P450s) in the neuroepithelium or the brain [25]. In this study, *CiCPR* was detected to be ubiquitously distributed throughout the body, including antenna, head, thorax, and abdomen (Fig. 3C). The significant expression of *CiCPR* in the antenna demonstrates the potentially functional importance of bed bug P450s in the odorant clearance and/or xenobiotics metabolism which might facilitate the host identification and localization in bed bugs. On the other hand, the broad presence of *CiCPR* in the head and abdomen may imply the roles of bed bug P450s in xenobiotics metabolism and/or endogenous compound biosynthesis.

#### Inactivation of *CiCPR*

Inactivation of *CPR* as a whole has been proposed to cause multiple developmental defects and embryonic lethality in mouse [47,48] and increase the permethrin susceptibility in *An. gambiae* [26]. In this study, RNAi was tested in the bed bug, and the RNAi effect was found to be dsRNA dose-dependent and systemic. When the *CiCPR* was suppressed through RNAi in deltamethrin resistant populations, the susceptibility of these bed bugs to deltamethrin was significantly enhanced. The susceptibility enhancement was not only observed in non-*kdir* mutation population (CIN-1, Fig. 6A) but also observed in the population that contains *kdir* mutation (NY-1, Fig. 6B) where decreased target-site sensitivity of voltage-gated sodium channels was suggested as one of the mechanisms of pyrethroid resistance. These data suggest that multiple resistance mechanisms may exist in NY-1 population. However, there was no significant difference in the susceptibility to deltamethrin between *CiCPR* knockdown and control in insecticide susceptible LA-1 bed bugs (Fig. 6C). These data suggest that P450-mediated metabolic detoxification might be another mechanism responsible for deltamethrin resistance in general bed bug populations. Recent report supports our prediction by showing higher levels of P450 mRNAs in pesticide resistant populations compared to their levels in susceptible populations of *C. lectularius* [49–50]. Taken together, P450-mediated metabolism could be added to factors such as the target site insensitivity, behavioral and penetration mechanisms involved in the deltamethrin resistance of bed bug populations.

#### Supporting Information

**Figure S1 A schematic diagram showing the strategy used to clone the full length *CiCPR*.** The top line stands for the cDNA. Other lines represent gene fragments isolated by RACE or PCR with specific PCR primer pair(s): fragment 1 (NADPHF/NADPHR), fragment 2 (CIRACER1-1/UPM, CIRACER2-1/NUP), fragment 3 (CIRACEF1-1/UPM, CIRACEF2-1/NUP), fragment 4 (CIRACEF3/UPM, CIRACEF4/NUP), and fragment 5 (*CiCPRF/CiCPRR*). (TIF)

**Figure S2 Transmembrane helix (A and B) and hydrophobicity (C and D) of *CiCPR* prediction.** The total 679 amino acids (A) and 140 N-terminal amino acids (B) were submitted into TMHMM Server 2.0 in turn. A 21-amino acid transmembrane region was predicted and highlighted in red. The total 679 amino acids (C) and 140 N-terminal amino acids (D) were also submitted into the on line molecular tool, Protein Hydrophobicity Plots. The hydrophobicity profiles of *CiCPR* were delineated using Kyte-Doolittle scale. Regions with values above 0 are hydrophobic in character. (TIF)

**Figure S3 Selection of a reference gene.** Stable expression of four reference genes, *rpl11*, *rpl8*, *rps16* and *hsp70* are shown across 8 RNA samples isolated from different developmental stages: eggs (1,5), small nymphs (1–3 instar) (2,6), large nymphs (4–5 instar) (3,7), and 1 week old female and male (4,8) adults in CIN-1 (1–4) and LA-1 (5–8). Products obtained after 40 cycles of PCR amplification under conditions described in the Materials and Methods section were resolved on an agarose gel and the gel was stained with ethidium bromide. (TIF)

**Figure S4 The average mortality of bed bugs at 5 days after dsRNA injection.** No significant difference was observed

in the mortalities between *CICPR* dsRNA and *malE* dsRNA (control) injected bed bugs ( $n = 7$ ; Student's *t*-test,  $P = 0.49$ ). (TIF)

**Table S1 CICPR homologues in insects.**  
(DOCX)

**Table S2 Primers used for cloning, RACE, qRT-PCR, RNAi, and housekeeping gene analysis.**  
(DOCX)

**Table S3 Statistical analyses of four candidate reference genes based on their threshold cycle ( $C_T$ ) value.**  
(DOCX)

## References

- Boase CJ (2001) Bedbugs-back from the brink. *Pestic Outlook* 12: 159–162.
- Carayon J (1966) Paragenital system In: Usinger R, ed. *Monograph of the Cimicidae*. Philadelphia, PA: Entomological Society of America. pp 81–167.
- Potter MF (2005) A bed bug state of mind: emerging issues in bed bug management. *Pest Control Technol* 33: 82–85.
- Potter MF, Rosenberg B, Henriksen M (2010) Bugs without borders: defining the global bed bug resurgence. *Pest World Sept./Oct*: 8–20.
- Romero A, Potter MF, Potter DA, Haynes KF (2007) Insecticide resistance in the bed bug: a factor in the pest's sudden resurgence? *J Med Entomol* 44: 175–178.
- Yoon KS, Kwon DH, Strycharz JP, Hollingsworth CS, Lee SH, et al. (2008) Biochemical and molecular analysis of deltamethrin resistance in the common bed bug (Hemiptera: Cimicidae). *J Med Entomol* 45: 1092–1101.
- Zhu F, Wigginton J, Romero A, Moore A, Ferguson K, et al. (2010) Widespread distribution of knockdown resistance mutations in the bed bug, *Cimex lectularius* (Hemiptera: Cimicidae), populations in the United States. *Arch Insect Biochem Physiol* 73: 245–257.
- Romero A, Potter MF, Haynes KF (2009) Evaluation of piperonyl butoxide as a deltamethrin synergist for pyrethroid-resistant bed bugs. *J Econ Entomol* 102: 2310–2315.
- Feyereisen R (2005) Insect cytochrome P450. In: Gilbert LI, Iatrou K, Gill S, eds. *Comprehensive Molecular Insect Science*. Oxford: Elsevier. pp 1–77.
- Zhu F, Parthasarathy R, Bai H, Woithe K, Kaussmann M, et al. (2010) A brain-specific cytochrome P450 responsible for the majority of deltamethrin resistance in the QITC279 strain of *Tribolium castaneum*. *Proc Natl Acad Sci U S A* 107: 8557–8562.
- Gilbert LI (2004) Halloween genes encode P450 enzymes that mediate steroid hormone biosynthesis in *Drosophila melanogaster*. *Mol Cell Endocrinol* 215: 1–10.
- Rewitz KF, O'Connor MB, Gilbert LI (2007) Molecular evolution of the insect Halloween family of cytochrome P450s: phylogeny, gene organization and functional conservation. *Insect Biochem Mol Biol* 37: 741–753.
- Lu AY, Junk KW, Coon MJ (1969) Resolution of the cytochrome P-450-containing  $\alpha$ -hydroxylation system of liver microsomes into three components. *J Biol Chem* 244: 3714–3721.
- Porter TD, Wilson TE, Kasper CB (1987) Expression of a functional 78,000 dalton mammalian flavoprotein, NADPH-cytochrome P-450 oxidoreductase, in *Escherichia coli*. *Arch Biochem Biophys* 254: 353–367.
- Vermilion JL, Ballou DP, Massey V, Coon MJ (1981) Separate roles for FMN and FAD in catalysis by liver microsomal NADPH-cytochrome P-450 reductase. *J Biol Chem* 256: 266–277.
- Nishino H, Ishibashi T (2000) Evidence for requirement of NADPH-cytochrome P450 oxidoreductase in the microsomal NADPH-sterol delta7-reductase system. *Arch Biochem Biophys* 374: 293–298.
- Ono T, Ozasa S, Hasegawa F, Imai Y (1977) Involvement of NADPH-cytochrome c reductase in the rat liver squalene epoxidase system. *Biochim Biophys Acta* 486: 401–407.
- Schenkman JB, Jansson I (1999) Interactions between cytochrome P450 monooxygenases and cytochrome b5. *Drug Metab Rev* 31: 351–364.
- Wang J, de Montellano PR (2003) The binding sites on human heme oxygenase-1 for cytochrome P450 reductase and biliverdin reductase. *J Biol Chem* 278: 20069–20076.
- Koener JF, Carino FA, Feyereisen R (1993) The cDNA and deduced protein sequence of house fly NADPH-cytochrome P450 reductase. *Insect Biochem Mol Biol* 23: 439–447.
- Andersen JF, Utermohlen JG, Feyereisen R (1994) Expression of house fly *CYP6A1* and NADPH-cytochrome P450 reductase in *Escherichia coli* and reconstitution of an insecticide-metabolizing P450 system. *Biochemistry* 33: 2171–2177.
- Murataliev MB, Arino A, Guzov VM, Feyereisen R (1999) Kinetic mechanism of cytochrome P450 reductase from the house fly (*Musca domestica*). *Insect Biochem Mol Biol* 29: 233–242.
- Hovemann BT, Schlmeyer F, Malz J (1997) *Drosophila melanogaster* NADPH-cytochrome P450 oxidoreductase: pronounced expression in antennae may be related to odorant clearance. *Gene* 189: 213–219.
- Horike N, Takemori H, Nonaka Y, Sonobe H, Okamoto M (2000) Molecular cloning of NADPH-cytochrome P450 oxidoreductase from silkworm eggs - Its involvement in 20-hydroxyecdysone biosynthesis during embryonic development. *Eur J Biochem* 267: 6914–6920.
- Maiheche-Coisne M, Merlin C, Francois MC, Porcheron P, Jacquin-Joly E (2005) P450 and P450 reductase cDNAs from the moth *Mamestra brassicae*: cloning and expression patterns in male antennae. *Gene* 346: 195–203.
- Lycett GJ, McLaughlin LA, Ranson H, Hemingway J, Kafatos FC, et al. (2006) *Anopheles gambiae* P450 reductase is highly expressed in oenocytes and *in vivo* knockdown increases permethrin susceptibility. *Insect Mol Biol* 15: 321–327.
- Kaewpa D, Boonsupsakul S, Rongnoparut P (2007) Functional expression of mosquito NADPH-cytochrome P450 reductase in *Escherichia coli*. *J Econ Entomol* 100: 946–953.
- Lian L-Y, Widdowson P, McLaughlin LA, Paine MJI (2011) Biochemical comparison of *Anopheles gambiae* and human NADPH P450 reductases reveals different 2'-5'-ADP and FMN binding traits. *PLoS One* 6.
- Romero A, Potter MF, Haynes KF (2009) Behavioral responses of the bed bug to insecticide residues. *J Med Entomol* 46: 51–57.
- Montes C, Cuadrillero C, Vilella D (2002) Maintenance of a laboratory colony of *Cimex lectularius* (Hemiptera: Cimicidae) using an artificial feeding technique. *J Med Entomol* 39: 675–679.
- Marchler-Bauer A, Lu S, Anderson JB, Chitsaz F, Derbyshire MK, et al. (2011) CDD: a Conserved Domain Database for the functional annotation of proteins. *Nucleic Acids Res* 39: D225–229.
- Roy A, Kucukural A, Zhang Y (2010) I-TASSER: a unified platform for automated protein structure and function prediction. *Nat Protoc* 5: 725–738.
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, et al. (2011) MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* 28: 2731–2739.
- Hall BG (2008) Phylogenetic trees made easy: a how-to manual. Sunderland-Massachusetts: Sinauer Associates, Inc.
- Pfaffl MW, Tichopad A, Prgomet C, Neuvians TP (2004) Determination of stable housekeeping genes, differentially regulated target genes and sample integrity: bestKeeper - excel-based tool using pair-wise correlations. *Biotechnol Lett* 26: 509–515.
- Huggett J, Dheda K, Bustin S, Zumula A (2005) Real-time RT-PCR normalisation, strategies and considerations. *Genes Immun* 6: 279–284.
- Livak KJ, Schmittgen DT (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2- $\Delta\Delta C_T$  method. *Methods* 25: 402–408.
- Ingelman M, Bianchi V, Eklund H (1997) The three-dimensional structure of flavodoxin reductase from *Escherichia coli* at 1.7 Å resolution. *J Mol Biol* 268: 147–157.
- Shen AL, Sem DS, Kasper CB (1999) Mechanistic studies on the reductive half-reaction of NADPH-cytochrome P450 oxidoreductase. *J Biol Chem* 274: 5391–5398.
- Hubbard PA, Shen AL, Paschke R, Kasper CB, Kim JJP (2001) NADPH-cytochrome P450 oxidoreductase - Structural basis for hydride and electron transfer. *J Biol Chem* 276: 29163–29170.
- Pandey AV, Kempna P, Hofer G, Mullis PE, Fluck CE (2007) Modulation of human *CYP19A1* activity by mutant NADPH P450 oxidoreductase. *Mol Endocrinol* 21: 2579–2595.
- Lamb DC, Kim Y, Yermalitskaya LV, Yermalitsky VN, Lepesheva GI, et al. (2006) A second FMN binding site in yeast NADPH-cytochrome P450 reductase suggests a mechanism of electron transfer by diflavin reductases. *Structure* 14: 51–61.
- Oprian DD, Coon MJ (1982) Oxidation-reduction states of FMN and FAD in NADPH-cytochrome P-450 reductase during reduction by NADPH. *J Biol Chem* 257: 8935–8944.
- Kida Y, Ohgiya S, Mihara K, Sakaguchi M (1998) Membrane topology of NADPH-cytochrome P450 reductase on the endoplasmic reticulum. *Arch Biochem Biophys* 351: 175–179.
- Reinhardt K, Naylor R, Siva-Jothy MT (2003) Reducing a cost of traumatic insemination: female bedbugs evolve a unique organ. *Proc Biol Sci* 270: 2371–2375.

## Acknowledgments

The authors are grateful to Dr. Alvaro Romero and Mark Goodman (University of Kentucky) for providing bed bug samples and technical help. We thank Dr. Michael Sharkey and Ms. Stephanie Clutts (University of Kentucky) for their help with phylogenetic analysis.

## Author Contributions

Conceived and designed the experiments: FZ KFH MFP SRP. Performed the experiments: FZ SS TM. Analyzed the data: FZ TM SRP. Contributed reagents/materials/analysis tools: FZ TM KFH MFP SRP. Wrote the paper: FZ SRP.

46. Wang M, Roberts DL, Paschke R, Shea TM, Masters BS, et al. (1997) Three-dimensional structure of NADPH-cytochrome P450 reductase: prototype for FMN- and FAD-containing enzymes. *Proc Natl Acad Sci U S A* 94: 8411–8416.
47. Shen AL, O'Leary KA, Kasper CB (2002) Association of multiple developmental defects and embryonic lethality with loss of microsomal NADPH-cytochrome P450 oxidoreductase. *J Biol Chem* 277: 6536–6541.
48. Henderson CJ, Otto DM, Carrie D, Magnuson MA, McLaren AW, et al. (2003) Inactivation of the hepatic cytochrome P450 system by conditional deletion of hepatic cytochrome P450 reductase. *J Biol Chem* 278: 13480–13486.
49. Bai X, Mamidala P, Rajarapu SP, Jones SC, Mittapalli O (2011) Transcriptomics of the bed bug (*Cimex lectularius*). *PLoS One* 6: e16336.
50. Adelman ZN, Kilcullen KA, Koganemaru R, Anderson MAE, Anderson TD, et al. (2011) Deep Sequencing of Pyrethroid-Resistant Bed Bugs Reveals Multiple Mechanisms of Resistance within a Single Population. *PLoS ONE* 6: e26228.