

3-2014

Evidence-Based Smoking Cessation Treatment for Disparate Populations

Chizimuzo T.C. Okoli
University of Kentucky, ctokol1@uky.edu

Follow this and additional works at: https://uknowledge.uky.edu/nursing_present



Part of the [Nursing Commons](#), and the [Public Health Commons](#)

[Right click to open a feedback form in a new tab to let us know how this document benefits you.](#)

Repository Citation

Okoli, Chizimuzo T.C., "Evidence-Based Smoking Cessation Treatment for Disparate Populations" (2014).
Nursing Presentations. 4.
https://uknowledge.uky.edu/nursing_present/4

This Presentation is brought to you for free and open access by the College of Nursing at UKnowledge. It has been accepted for inclusion in Nursing Presentations by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Evidence-Based Smoking Cessation Treatment for Disparate Populations

Chizimuzo Ökoli, PhD, MPH, RN
Assistant Professor and Director,
Tobacco Treatment and Prevention Division,
Tobacco Policy Research Program,
University of Kentucky College of Nursing

Overview

- Describe Research Group and Program of Research among disparate Populations
- Discuss smoking among persons with substance use and mental illness
- Present smoking cessation outcomes from evidence-based programs for persons with substance use disorders and mental illness
- Conclusion with current and future Projects

RESEARCH GROUP AND PROGRAM OF RESEARCH



Clean Indoor Air Partnership



Air Quality Monitoring

Tobacco Policy Research Program

Freedom from Radon Exposure and Smoking in the Home (FRESH)

Radon Policy Research Program



Kentucky Center for Smoke-Free Policy




Rural Smoke-Free Communities Project



Tobacco Treatment and Prevention Program




Program of Research



SMOKING AMONG PERSONS WITH SUBSTANCE USE AND MENTAL ILLNESS

...

Background and Significance




Vital^{CDC}**signs**[™]
www.cdc.gov/vitalsigns

- “Nicotine-dependent individuals with a comorbid psychiatric disorder made up 7.1% of the population yet consumed 34.2% of all cigarettes smoked in the United States”




Grant, B.F. , Hasin, D.S., Chou, S.P., Stinson, F.S.& Dawson, D.A. (2004). Nicotine dependence and psychiatric disorders in the United States: Results from a national epidemiologic survey on alcohol and related conditions. Archives of General Psychiatry,61,1107-1115.



* Difference between those with any mental illness and those with no mental illness is statistically significant at the .05 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Surveys on Drug Use and Health (NSDUHs), 2009 to 2010 (revised March 2012), 2011.

Past Month Cigarette Use among Adults Aged 18 or Older, by Any Mental Illness in the Past Year and Gender, by Age Group: 2009 to 2011



Kalman, Morissette and George (2005), Am. J. Addict., 14: 106-123

Health effects of smoking among persons with mental illness

Smokers with Mental illness :


- Die 10 years earlier
- Have more depression and depression
- Have more substance use problems
- Have more chest and heart problems
- Are more likely to commit suicide
- Have sexual problems

Nonsmokers with Mental illness :


- Have better health
- Live longer
- Need less medication
- Have less depression
- Save more money

Reasons for smoking among individuals with substance use disorders and mental illness

- Genetic



- Bio-behavioral



- Psychosocial



Smoking and substance use: Genetic



Familial Transmission of Substance Dependence: Alcohol, Marijuana, Cocaine, and Habitual Smoking

A Report From the Collaborative Study on the Genetics of Alcoholism

Laura Jean Bierut, MD; Stephen H. Dinwiddie, MD; Henri Begleiter, MD; Victor Hesselbrock, PhD; John I. Nurnberger, Jr, MD, PhD; Benjamin R. Crowley, MD; Marc A. Schuckit, MD; Theodore Reich, MD

Common Genetic Vulnerability for Nicotine and Alcohol Dependence in Men

William R. True, PhD, MPH; Hong Xian, PhD; Jeffrey F. Scherrer, MA; Pamela A. F. Madden, PhD; Kathleen K. Bucholz, PhD; Andrew C. Heath, DPhil; Seth A. Eisen, MD, MSc; Michael J. Lyons, PhD; Jack Goldberg, PhD; Ming Tsuang, MD, PhD, DSc

- Both common and specific addictive factors for alcohol, marijuana, cocaine, and habitual smoking transmitted in families
- This specificity suggested independent causative factors for the development of each substance dependence
- 68% of the association between nicotine and alcohol dependence explained by shared genetic effects.


Smoking and mental illness: Genetic



(1,566 female twin pairs) average life time daily cigarette consumption was found to be associated with life time prevalence of major depression, **suggesting that the relationship between smoking and major depression resulted solely from genes which predispose to both conditions.**

(8,169 male twins) shared genetic disorders **further predispose to major depression and nicotine dependence.**

Smoking and mental illness: Genetic



A Twin Registry Study of the Relationship Between Posttraumatic Stress Disorder and Nicotine Dependence in Men

Karestan C. Koenen, PhD; Brian Hitsman, PhD; Michael J. Lyons, PhD; Raymond Niaura, PhD; Jeanne McCaffery, PhD; Jack Goldberg, PhD; Seth A. Eisen, MD; William True, MD; Ming Tsuang, MD


- 63% of the association between post traumatic stress disorder and nicotine dependence co-morbidity explained by shared genetic effects.

A Novel Permutation Testing Method Implicates Sixteen Nicotinic Acetylcholine Receptor Genes as Risk Factors for Smoking in Schizophrenia Families

Stephen V. Faraone^{a,b} Jessica Su^b Levi Taylor^c Marsha Wilcox^c
Paul Van Eerdewegh^{c,d} Ming T. Tsuang^{a,b,c,e}

- A group of candidate genes and individual genes found among individuals with schizophrenia significantly linked to smoking behaviors.

Smoking and substance use: Bio-behavioral



Marijuana use reduces cessation of tobacco smoking in adults

Ford, Vu, Anthony. (2002) *Drug and Alcohol Dependence*; 67:243-248

Nicotine increases alcohol self-administration in non-dependent male smokers.

Barrett, Tichauer, Leyton, et al. (2006). *Drug and Alcohol Dependence*; 81:197-204

Increases in methadone dose could increase nicotine craving and cigarette consumption for individuals with **opioid-dependence**

Story & Stark. (1991). *Journal of psychoactive drugs*, 23:203-215

Cigarette smoking primes the brain to cocaine use

Levine et al. (2011). *Science translational medicine*, 3, 107, 107-109

Smoking and mental illness: Bio-behavioral



Nicotine reduces sensorimotor gating deficits in smokers with schizophrenia

Postma et al. (2006). *Psychopharmacology*, 184: 589–599

Brain levels of monoamine oxidase A (MAO-A) (an enzyme associated with depression) were reduced in smokers relative to nonsmokers; suggesting that people with affective disorders may smoke to reduce elevated MAO-A levels in the brain

Fowler, Volkow, Wang, et al. (1996). *Proceedings of the National Academy of Sciences of the United States of America*, 93:14065-14069

Smokers with a primary diagnosis of anxiety disorder reported greater levels of general anxiety, distress, and depression as compared to nonsmokers.

McCabe, Chudzick, Antony, et al. (2004). *Journal of Anxiety Disorders*, 18:7-18

Substance use and Smoking: Psychosocial



- Tobacco use may foster the use of other substances; and vice versa
 - Drug treatment centers may provide an environment that supports tobacco use or a factor for delayed tobacco use cessation.
- Factors such as neighborhood disadvantage and early exposure to substance use may present an ‘exposure opportunity’ for subsequent substance use.

Glautier S, Clements K, White JAW, et al. Alcohol and the reward value of cigarette smoking. *Behavioural Pharmacology* 1996; 7:144-154

King AC, Epstein AM. Alcohol Dose-Dependent Increases in Smoking Urge in Light Smokers. *Alcoholism: Clinical & Experimental Research* 2005; 29:547-552

Friend KB, Pagano ME. Smoking initiation among nonsmokers during and following treatment for alcohol use disorders. *Journal of Substance Abuse Treatment* 2004; 26:219-224

Bobo JK, Husten C. Sociocultural Influences on Smoking and Drinking. *Alcohol Research & Health* 2000; 24:225-232

Crum RM, Lillie-Blanton M, Anthony JC. Neighborhood environment and opportunity to use cocaine and other drugs in late childhood and early adolescence. *Drug and Alcohol Dependence* 1996; 43:155-161

Wagner FA, Anthony JC. Into the world of illegal drug use: Exposure opportunity and other mechanisms linking the use of alcohol, tobacco, marijuana, and cocaine. *Am. J. Epidemiol.* 2002; 155:918-925

Smoking mental illness: Psychosocial



- History of tobacco use as a **token economy**-- cigarettes used as a 'reward' for appropriate behavior (i.e., smoking privileges)

- Smoking among clients and staff to encourage **'socialization'**

Kawachi I, Berkman L. Social ties and mental health. *Journal of Urban Health* 2001; 78:458-467

Lawn S. Cigarette smoking in psychiatric settings: occupational health, safety, welfare and legal concerns. *Australian and New Zealand Journal of Psychiatry* 2005; 39:886-891

Keizer I, Eytan A. Variations in Smoking during Hospitalization in Psychiatric In-Patient Units and Smoking Prevalence in Patients and Health-Care Staff. *International Journal of Social Psychiatry* 2005; 51:317-328

Morisano D, Bacher I, Audrain-McGovern J, et al. Mechanisms underlying the comorbidity of tobacco use in mental health and addictive disorders. *Canadian Journal Of Psychiatry. Revue Canadienne De Psychiatrie* 2009; 54:356-367

Arguments for Not Providing Tobacco Treatment....

“these patients don’t want to quit”

- However....80% of participants in a methadone maintenance program and 75% of participants in an alcohol abuse program endorsed a desire to quit

(Richter KP et al., 2001; Ellingstad TP et al, 1999)

- In a review of 9 studies of motivation to quit smoking among individuals with psychiatric disorders at least 50% are contemplating cessation

(Siru, Hulse & Tait, 2009).

“these patients will relapse (to other substances) if they try to quit”

- Smoking cessation related to **IMPROVED QUALITY OF LIFE**
- Meta-analysis (n = 19 studies) of smoking cessation among individuals in addiction treatment and recovery found that smoking cessation efforts can **ENHANCE** rather than compromise long-term sobriety

Bobo JK, McIlvain HE, Lando HA, et al. Effect of smoking cessation counseling on recovery from alcoholism: findings from a randomized community intervention trial. *Addiction* 1998; 93:877-887

McCarthy WJ, Zhou Y, Hser YI, et al. To smoke or not to smoke: Impact on disability, quality of life, and illicit drug use in baseline polydrug users. *Journal of Addictive Diseases* 2002; 21:35-54

Lemon SC, Friedmann PD, Stein MD. The impact of smoking cessation on drug abuse treatment outcome. *Addictive Behaviors* 2003; 28:1323-1331

McCarthy WJ, Collins C, Hser YI. Does cigarette smoking affect drug abuse treatment? *Journal of Drug Issues* 2002; 2:61-80

● Prochaska JJ, Delucchi K, Hall SM. A meta-analysis of smoking cessation interventions with individuals in substance abuse treatment or recovery. *Journal of Consulting and Clinical Psychology* 2004; 72:1144-1156

“these patients are unable to quit”


- Meta-analysis (n = 19 studies) of smoking cessation among individuals in addiction treatment and recovery found **increased cessation at end of 12 weeks treatment**

(Prochaska JJ et al, 2004).

- Another recent study found end-of-treatment (between 8 to 26 weeks) smoking cessation rates of **40%** among individuals with SUD and/or PD who completed an intensive tailored smoking cessation intervention that provided no-cost pharmacotherapy combined with behavioural counseling

(Khara and Okoli, 2011)

Barriers to facilitating tobacco treatment

- **Concerns that smoking cessation will increase psychiatric symptoms or relapse among patients.**
 - Among individuals with depression, smoking cessation related to **increased depression symptomatology**, which is one of the symptoms of the nicotine withdrawal syndrome
 - individuals with anxiety disorders and depression report **more severe withdrawal symptoms**
 - smoking is associated with improvements in prepulse inhibition and sensory gating which may be affected by smoking cessation
- **Smoke-free policy and mental health facilities** 
 - a review of studies examining prohibitions of smoking in psychiatric facilities suggests that **prohibitions do not have long-term effects on behavioral unrest or noncompliance**, but **neither do they appear to effect smoking cessation**

Patten C, Martin J. Does nicotine withdrawal affect smoking cessation? Clinical and theoretical issues. *Annals of Behavioral Medicine* 1996; 18:190-200

Breslau N, Kilbey MM, Andreski P. Nicotine withdrawal symptoms and psychiatric disorders: findings from an epidemiologic study of young adults. *American Journal of Psychiatry* 1992; 149:464-469

Adams CE, Stevens KE. Evidence for a role of nicotinic acetylcholine receptors in schizophrenia. *Frontiers in Bioscience* 2007; 12:4755-4772

Kumari V, Postma P. Nicotine use in schizophrenia: The self medication hypotheses. *Neuroscience & Biobehavioral Reviews* 2005; 29:1021-1034

el-Guebaly N, Cathcart J, Currie S, et al. Public health and therapeutic aspects of smoking bans in mental health and addiction settings. *Psychiatr Serv* 2002; 53:1617-1622

Ziedonis DMA, Williams JMB. Management of smoking in people with psychiatric disorders. *Current Opinion in Psychiatry* 2003; 16:305-315

Costs associated with smoking cessation treatment

- Even though **less expensive than purchasing cigarettes**, the cost of pharmacotherapy and counseling presents an important barrier to seeking treatment
- Such cost barriers to accessing treatment and the potential cost-effectiveness of treatment have prompted guidelines about **reducing medication costs** (reduced cost or free of charge), **inclusion of medications as benefits on drug insurance plans**, and **setting up systems for reimbursement for tobacco cessation treatment** for health care providers.



Bansal MA, Cummings KM, Hyland A, et al. Stop-smoking medications: Who uses them, who misuses them, and who is misinformed about them? *Nicotine & Tobacco Research* 2004; 6:303-310

Reilly P, Murphy L, Alderton D. Challenging the smoking culture within a mental health service supportively. *International Journal of Mental Health Nursing* 2006; 15:272-278

Fiore M, Jaén C, Baker T, et al. A Clinical Practice Guideline for Treating Tobacco Use and Dependence: 2008 Update: A U.S. Public Health Service Report. *American Journal of Preventive Medicine* 2008; 35:158-176

Raw M, McNeill A, West R. Smoking cessation: evidence based recommendations for the healthcare system. *British Medical Journal* 1999; 318:182-185

Smoking Cessation Outcomes of evidence-based program for individuals with Substance Use Disorders and Mental Illness



- The Tobacco Dependence Clinic (TDC) for individuals with primarily substance use disorders and addictions
- The Butt-Out Group (Butt-Out) for individuals with primarily mental illnesses

Program Overview

- 24-26 weeks
 - **TDC** (8 weeks structured, up to 18 weeks of support)
 - **Butt Out group** (12 weeks structured, up to 12 weeks support)
- Tailored pharmacotherapy for up to 26 weeks at no cost
- Program is run with a team of nurses, counsellors, respiratory therapists, Occupational therapists, and a physician

TREATMENT PHILOSOPHY

Smoking cessation is a 'process' not an event. Hence the program has no specific 'quit date'. Also the program employs the concept of 'titration-to-effect' in which increasing doses of nicotine replacement therapy is provided until the behavior of smoking ceases.

Criteria

Eligibility:

- 19 years or older
- Tobacco dependent
- Have a history of substance use disorder (SUD) and/or psychiatric disorder (PD)
- Financially disadvantaged


Assessment:

- 1 hour evaluation of medical, psychiatric, substance and tobacco use history
- Expired air CO is determined and a treatment plan is developed in consultation with client

Phases of Treatment

Behavioral Counseling

Support Group



Behavioural Counseling (Weeks 1-8)

- **Phase 1:** engagement in the process – weeks 1-2
- **Phase 2:** planning for change – weeks 3-4
- **Phase 3:** sustaining change – weeks 5-8



Combination Pharmacotherapy

Nicotine Replacement Therapy



Patch



Gum



Lozenge




Inhaler

Oral Medications




Zyban




Champix

Gender




TDC (N =647)




Butt Out Group (N = 76)

Substance Use Disorder History (either past or current use)




TDC (N = 647)




Butt Out Group (N = 76)

Psychiatric Disorder History (either past or current, N = 647)





TDC (N = 647)



Butt Out Group (N = 76)

Co-occurring Disorders history (either past or current)




Sample Characteristics


	TDC (N = 647) Mean (SD)	Butt-Out (N =76) Mean (SD)
Age of participant (years)	47.3 (11.4)	47.2 (10.6)
Age at smoking initiation (years)	14.7 (4.9)	18.1 (6.4)
Importance of quitting (scale of 0 'low' to 10 'high')	9.0 (1.3)	9.0 (1.6)
Confidence in quitting (scale of 0 'low' to 10 'high')	7.3 (2.2)	6.9 (2.3)
Number of cigarettes smoked per day	20.9 (10.5)	22.5 (10.5)
Fagerstrom Test for Nicotine Dependence (scale of 0 'low' to 10 'high')	6.1 (2.1)	6.1 (2.0)
CO level at baseline (ppm)	20.5 (12.6)	22.8 (14.0)

Samples for Outcomes Assessment

Tobacco Dependence Clinic




Butt Out Group




Smoking cessation: 7-day point-prevalence of abstinence at end of treatment (i.e., anytime between 8 weeks to 26 weeks) verified by expired CO levels

Program Completion




TDC (N = 307/407)



Butt Out Group (N = 64/74)


TDC Smoking Cessation Outcomes at end-of-treatment*



* End-of-treatment (i.e., anytime between 8 weeks to 26 weeks) verified by expired CO levels

Butt Out group Smoking Cessation* Outcomes

at end-of-treatment



*End-of-treatment (i.e., anytime between 8 weeks to 26 weeks) verified by expired CO levels

Conclusions

- *With intensive tobacco dependence treatment provided within Mental Health and Addictions services, individuals with a history of substance use disorders and mental illness are able to achieve smoking abstinence.*
- *Tobacco treatment should include best practices using behavioural counseling combined with pharmacotherapy.*
- *Treatment durations beyond 12-weeks should be considered to maximize effectiveness of programs*

Current and Future Projects



- **Current projects:**

- Smoking Cessation Program among Individuals with Mental Illness (PARTICIPATION STATION- a peer led mental health support program)
- Understanding the behavioural effects of secondhand tobacco smoke exposure among nonsmokers
- Examining sex and gender differences in tobacco dependence outcomes among smokers with co-occurring substance use and psychiatric disorders

- **Future Projects:**

- Developing a tailored tobacco dependence treatment program for individuals with Chronic Mental Illnesses
- Examining the effect of different tobacco products on nicotine dependence, symptoms severity, and smoking cessation among individuals with mental illnesses.

Questions??

