

7-8-2008

# Identifying regulatory elements and gene boundaries with comparative genomic sequence analysis

Jim Lund  
*University of Kentucky*

**Right click to open a feedback form in a new tab to let us know how this document benefits you.**

Follow this and additional works at: [https://uknowledge.uky.edu/biology\\_present](https://uknowledge.uky.edu/biology_present)



Part of the [Biology Commons](#)

---

## Repository Citation

Lund, Jim, "Identifying regulatory elements and gene boundaries with comparative genomic sequence analysis" (2008). *Biology Presentations*. 4.

[https://uknowledge.uky.edu/biology\\_present/4](https://uknowledge.uky.edu/biology_present/4)

This Presentation is brought to you for free and open access by the Biology at UKnowledge. It has been accepted for inclusion in Biology Presentations by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

Poster presentation

## Identifying regulatory elements and gene boundaries with comparative genomic sequence analysis

Jim Lund

Address: Department of Biology, University of Kentucky, Lexington, KY 40506, USA

Email: Jim Lund - jim1@uky.edu

from UT-ORNL-KBRIN Bioinformatics Summit 2008  
Cadiz, KY, USA. 28–30 March 2008

Published: 8 July 2008

BMC Bioinformatics 2008, 9(Suppl 7):P17 doi:10.1186/1471-2105-9-S7-P17

This abstract is available from: <http://www.biomedcentral.com/1471-2105/9/S7/P17>

© 2008 Lund; licensee BioMed Central Ltd.

For several groups of organisms multiple genomes have been sequenced either completely or to the unfinished shotgun sequencing stage. Comparative sequence analysis can identify functional sequence elements for pairs of organisms that diverged far enough in the past to allow mutational drift of non-conserved sequences. While analysis of a pair of genomes identifies many of these functional elements adding additional genomes allows additional information to be elicited.

Additional conserved sequence elements are identified as additional genomes are analyzed. These conserved sequence elements are often regulatory elements although they are difficult to classify with *in silico* analysis. For an individual gene, the set of associated conserved sequence elements and the organisms they are found in provides insight into the evolutionary history of the regulation of the gene.

The eight complete and unfinished shotgun sequenced nematode genomes and the dozen informative insect genome sequences were used to analyze conserved non-coding sequences in these groups of organisms. I have developed web-based software to allow researchers to explore and visualize sequence conservation that differs from previous work in analyzing conserved sequences in each pair of organisms rather than with respect to a single reference genome.

We have used this analysis to identify gene regulatory boundaries in the nematode *C. elegans*. The genomes of *C.*

*elegans* and other nematodes have diverged enough that synteny regions are typically one or two genes long. I was able to associate conserved sequence elements to particular genes identifying the natural boundaries between genes and the extent of worm promoters.

### Acknowledgements

This work was supported by Grant Number P20RR-16481 from the National Center for Research Resources (NCRR), a component of the National Institute of Health (NIH).