

University of Kentucky

UKnowledge

---

Center of Excellence in Rural Health Faculty  
Publications

Rural Health

---

5-1-2015

## Isolation Housing Exacerbates Alzheimer's Disease-Like Pathophysiology in Aged APP/PS1 Mice

Huang Huang

*Nanjing Medical University, China*

Linmei Wang

*Nanjing Medical University, China*

Min Cao

*Nanjing Medical University, China*

Charles Marshall

*University of Kentucky, charles.marshall@uky.edu*

Junying Gao

*Nanjing Medical University, China*

*See next page for additional authors*

Follow this and additional works at: [https://uknowledge.uky.edu/ruralhealth\\_facpub](https://uknowledge.uky.edu/ruralhealth_facpub)



Part of the [Medicine and Health Sciences Commons](#)

**Right click to open a feedback form in a new tab to let us know how this document benefits you.**

---

### Repository Citation

Huang, Huang; Wang, Linmei; Cao, Min; Marshall, Charles; Gao, Junying; Xiao, Na; Hu, Gang; and Xiao, Ming, "Isolation Housing Exacerbates Alzheimer's Disease-Like Pathophysiology in Aged APP/PS1 Mice" (2015). *Center of Excellence in Rural Health Faculty Publications*. 4.

[https://uknowledge.uky.edu/ruralhealth\\_facpub/4](https://uknowledge.uky.edu/ruralhealth_facpub/4)

This Article is brought to you for free and open access by the Rural Health at UKnowledge. It has been accepted for inclusion in Center of Excellence in Rural Health Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

---

## Isolation Housing Exacerbates Alzheimer's Disease-Like Pathophysiology in Aged APP/PS1 Mice

Digital Object Identifier (DOI)

<http://dx.doi.org/10.1093/ijnp/pyu116>

### Notes/Citation Information

Published in *International Journal of Neuropsychopharmacology*, v. 18, no. 7, p. 1-10.

© The Author 2015. Published by Oxford University Press on behalf of CINP. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

### Authors

Huang Huang, Linmei Wang, Min Cao, Charles Marshall, Junying Gao, Na Xiao, Gang Hu, and Ming Xiao

## RESEARCH ARTICLE

# Isolation Housing Exacerbates Alzheimer's Disease-Like Pathophysiology in Aged APP/PS1 Mice

Huang Huang, MD; Linmei Wang, MD; Min Cao, Ms; Charles Marshall, PhD; Junying Gao, Ms; Na Xiao, Ms; Gang Hu, MD, PhD; Ming Xiao, MD, PhD

Jiangsu Key Laboratory of Neurodegeneration, Nanjing Medical University, Nanjing, Jiangsu, China (Drs Huang MD, Wang MD, Cao Ms, Gao Ms, N. Xiao Ms, Hu MD, PhD, and M. Xiao MD, PhD); Department of Rehabilitation Sciences, University of Kentucky Center for Excellence in Rural Health, Hazard, KY (Dr Marshall PhD).

Correspondence: Ming Xiao, MD, PhD, Jiangsu Key Laboratory of Neurodegeneration, Nanjing Medical University, 140 Hanzhong Road, Nanjing, Jiangsu 210029, P.R. China ([mingx@njmu.edu.cn](mailto:mingx@njmu.edu.cn)).

## Abstract

**Background:** Alzheimer's disease is a neurodegenerative disease characterized by gradual declines in social, cognitive, and emotional functions, leading to a loss of expected social behavior. Social isolation has been shown to have adverse effects on individual development and growth as well as health and aging. Previous experiments have shown that social isolation causes an early onset of Alzheimer's disease-like phenotypes in young APP695/PS1-dE9 transgenic mice. However, the interactions between social isolation and Alzheimer's disease still remain unknown.

**Methods:** Seventeen-month-old male APP695/PS1-dE9 transgenic mice were either singly housed or continued group housing for 3 months. Then, Alzheimer's disease-like pathophysiological changes were evaluated by using behavioral, biochemical, and pathological analyses.

**Results:** Isolation housing further promoted cognitive dysfunction and A $\beta$  plaque accumulation in the hippocampus of aged APP695/PS1-dE9 transgenic mice, associated with increased  $\gamma$ -secretase and decreased neprilysin expression. Furthermore, exacerbated hippocampal atrophy, synapse and myelin associated protein loss, and glial neuroinflammatory reactions were observed in the hippocampus of isolated aged APP695/PS1-dE9 transgenic mice.

**Conclusions:** The results demonstrate that social isolation exacerbates Alzheimer's disease-like pathophysiology in aged APP695/PS1-dE9 transgenic mice, highlighting the potential role of group life for delaying or counteracting the Alzheimer's disease process.

**Keywords:** Alzheimer's disease,  $\beta$ -amyloid, hippocampus, memory deficiency, social isolation, APP/PS1 mice

## Introduction

Alzheimer's disease (AD), the most common neurodegenerative disorder in the elderly, is reaching epidemic levels with tremendous social and financial burdens (Querfurth and Laferla, 2010). Unfortunately, the specific mechanisms causing AD remain unknown, and scientifically proven pharmacological treatment options are inadequate (Huang and Mucke, 2012). This urgent situation underscores the need for identifying crucial factors

and/or mechanisms involved in accelerating the progression of AD and developing nonpharmacological interventions to slow the process of cognitive decline.

Social isolation (SI) refers to a complete, or near complete, lack of contact with people and society for members within a social species. This isolation has been proven to be harmful to individual development and growth as well as physical and

Received: October 10, 2014; Revised: December 3, 2014; Accepted: December 23, 2014

© The Author 2015. Published by Oxford University Press on behalf of CINP.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License

(<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

mental health (Cacioppo et al., 2011). SI can be an issue for anyone, irrespective of their age. However, when compared with other age groups, the elderly are more vulnerable to suffer from SI because of declines in overall health, absent or uninvolved relatives or children, the abrupt end of daily work relationships after retirement, and/or economic struggles (Díez et al., 2014; Hand et al., 2014).

SI in elderly is associated with increased mortality risk and onset of several neuropsychological disorders, including schizophrenia (Jiang et al., 2013b), bipolar disorder (Miklowitz, 2011; Gilman et al., 2014), and AD (Dong and Csernansky, 2009). SI influences the onset and development of a mouse AD model via enhancing activity of  $\beta$  and  $\gamma$ -secretases and aberrant activation of p25/Cdk5, which plays an important role in the production of A $\beta$  peptide and phosphorylation of tau, respectively (Hsiao et al., 2011). Recently, several studies have indicated that SI increases oxidative stress and inflammatory reaction (Powell et al., 2013) and inhibits antiinflammatory responses (Azzinnari et al., 2014), synaptic plasticity (Djordjevic et al., 2009), and myelination (Liu et al., 2012a), all of which are involved in the pathogenesis of AD.

Although SI may contribute to the onset of AD, the added effect on the disease progression remains unclear. Furthermore, AD patients, especially those in the late stages, are more likely to suffer from SI due to cognitive and emotional impairment and loss of communication ability as well as potential neglect of communication by family members or attendants (Tanzi and Bertram, 2005; Peña-Longobardo et al., 2014; Zucchella et al., 2014). Thus, it is necessary to determine whether SI exacerbates pathology of late-stage AD. A better understanding of the interplay between the 2 disease processes will help establish appropriate and effective interventions to delay or counteract the neurodegenerative progression of AD.

In the present study, we evaluated the effect of isolation housing on AD-like pathophysiology in aged APP695/PS1-dE9 transgenic (APP/PS1) mice. Our results demonstrate that 17-month-old APP/PS1 mice housed separately for 3 months displayed increased cognitive impairment, A $\beta$  plaque burden, hippocampus atrophy, and reactive astrogliosis compared with group-housed controls.

## Materials and Methods

### Animals and Experimental Design

Male APP/PS1 mice were used in this study. On postnatal day 28, weaned, male APP/PS1 mice were separated from their female littermates and housed (4 mice per cage) in a temperature- and humidity-controlled facility on a 12-h-light/-dark cycle with food and water ad libitum. At 17 months old, one-half of the animals were randomly selected and singly housed; the other one-half continued group housing until 20 months old. All experiments were conducted in accordance with international standards on animal welfare and the guidelines of the Institute for Laboratory Animal Research of Nanjing Medical University. All efforts were made to minimize animal suffering and reduce the number of animals used.

### Open Field Test

After a 2-day acclimatization to the behavioral testing room, mice performed the open field test so we could measure exploration and anxiety-related behaviors (Chen et al., 2006). The open field box consisted of a square black Plexiglas box (60 cm  $\times$  60 cm  $\times$  25 cm), with an outlined center area (30 cm  $\times$  30 cm). Each animal was placed in the middle of box, which served as a starting point,

then allowed to move freely for 10 minutes within the box. The amount of time and distance traveled in the center area of the maze, number of entries into the center, and grooming numbers were measured. At the conclusion of the experiment period, defecation number was also counted.

### Y-Maze Test

One day after the open field test, the Y maze was performed to measure mouse short-term memory, as previously described (Wang et al., 2013). In brief, the Y maze was randomly assigned to 3 arms: novel arm (NA), starting arm (SA), and other arm (OA). The Y maze test contains two 5-minute stages with an interval of 2 hours. During the first stage, the NA was blocked by a black baffle, with mice entering the SA were allowed to move freely only between the SA and OA. During the second stage, the NA was opened and mice entering from SA could freely move throughout 3 arms. The percentage of time traveled in each arm, number of entries into each arms and travelling speed were analyzed.

### Morris Water Maze

The Morris water maze task was conducted to measure long-term learning and memory function as described previously (Liu et al., 2012b). Briefly, a black plastic pool with a diameter of 100 cm and a height of 50 cm was filled with water ( $22 \pm 2^\circ\text{C}$ ). Training was conducted for 7 consecutive days with 4 trials/d. During the first 2 days of testing, mice were trained with a visible platform. A cylindrical dark-colored platform with a diameter of 10 cm was placed 0.5 cm above the water surface and kept consistently within one of the quadrants. On day 3, hidden platform testing was performed, in which the platform was moved to the opposite quadrant and submerged 1 cm below the surface of the water. The escape latency, swim distance and speed, and swim patterns were analyzed. A probe trial was conducted on day 8; the hidden platform was removed, and mice were placed in the pool and allowed to swim for 60 seconds. The percent of total time spent in each quadrant and the number of crossing where the platform had been previously located were calculated.

Mouse activity in the above behavioral apparatuses was collected by a digital video camera connected with a computer-controlled system (Beijing Sunny Instruments Co. Ltd). All tests were performed by 2 independent experimenters who were blind to the treatment schedule.

### Section Preparation

Subsequent to deep anesthesia, mice were transcardially perfused with 0.9% saline by perfusion pump (Cole-parmer) for 5 minutes, followed by 4% paraformaldehyde for 12 minutes. Brains were dissected in the mid-sagittal plane, postfixed overnight at  $4^\circ\text{C}$ , then dehydrated in a series of graded ethanol solutions and embedded in paraffin. Both half-brains were serially cut into 5- $\mu\text{m}$  sagittal sections using a paraffin slicing machine (Leica RM2135, Nussloch, Germany). Sections from one half-brain were divided into 5 sets for HE staining and Thioflavin-S staining, while serial sections from the other half-brain were used for immunohistochemical stains.

### Quantitative Analysis of Hippocampal Volume and A $\beta$ Plaque Load

The dorsal hippocampus stained with H&E, Thioflavin-S or 6 E10 was photographed at 100 $\times$  magnification using a digital microscope (Leica Microsystems). Individual images were exported to

Image-Pro Plus 6.0 Analysis System (Media Cybernetics Inc., San Francisco, CA). For quantitative analysis of hippocampal volume, the boundary of hippocampus and gray region that consists of pyramidal cell layer and granular cell layer were manually delineated. The total hippocampal area and gray matter area per section were measured. Total hippocampal (gray matter) volume per mouse was also calculated according to Cavalieri's estimator (Pereda-Pérez et al., 2013),  $V = (S_1 + 2S_2 + 2S_3 + \dots + 2S_{n-1} + S_n) \mu\text{m}^2 \times 5 \mu\text{m} \times 5$ . The hippocampal white matter volume was obtained by the total hippocampal volume subtracting the hippocampal gray matter volume. For quantitative analysis of A $\beta$  plaque load, thioflavin-S positive signals were determined by standardized region of interest grayscale threshold analysis. The cross area of each plaque and total area of plaque coverage relative to the total hippocampus area were also measured.

### Immunohistochemistry

Immunohistochemical staining was performed as previously described (Xu et al., 2013). Briefly, after deparaffinization and rehydration, tissue sections were incubated with a primary antibody direct against glial fibrillary acidic protein (GFAP) (1:1000; Sigma-Aldrich), ionized calcium-binding adaptor molecule 1 (Iba-1) (1:500; Wako), double-cortin (DCX) (1:500; Abcam), or proliferating cell nuclear antigen (PCNA) (1:1000; Abcam) at 4°C overnight. Following incubation (1 hour 30 minutes) with biotinylated IgGs and ABC (1 hour 15 minutes), the reaction was visualized with DAB (Sigma-Aldrich). The mean integrated optical density (IOD/total area) was measured to assess the expression level of GFAP and Iba-1 in the entire hippocampus at 100 $\times$  magnification using an Image-Pro Plus 6.0 Analysis System (Media Cybernetics Inc). The number of DCX-positive cells and PCNA-positive cells per section was also counted.

### Western Blot

Hippocampal tissues were homogenized and centrifuged at 4°C and 12000rpm for 15 minute. The samples were resolved on SDS-PAGE, transferred onto PVDF membranes using a Bio-Rad miniprotein-III wet transfer unit, then blocked with 5% skim milk dissolved in TBST (pH 7.5, 10mM Tris-HCl, 150mM NaCl, and 0.1% Tween 20) at room temperature for 1 hour. Membranes were probed at 4°C overnight with a primary antibody directed against A $\beta_{1-42}$  (1:1000, Abcam),  $\beta$ -amyloid precursor protein (sAPP $\alpha$ ) (1:100; Immuno-Biological Laboratories), disintegrin and metalloproteinase10 (1:1000; Millipore),  $\beta$ -site amyloid precursor protein-cleaving enzyme 1 (BACE1) (1:2000; Millipore), presenilin1 (PS1) (1:1000; Sigma-Aldrich), neprilysin (NEP) (1:1000; Millipore), insulin degrading enzyme (IDE) (1:1000; Abcam), 6E10 (1:1000; Covance), myelin basic protein (MBP) (1:200; Santa Cruz Biotechnology), synaptophysin (SYP) (1:1500; Millipore), neuronal class III  $\beta$ -tubulin (1:3000; Sigma-Aldrich), or  $\beta$ -tubulin (1:3000; Sigma-Aldrich). Horseradish peroxidase-conjugated secondary antibodies (Vector Laboratories, Burlingame, CA) were used, and bands were visualized using ECL plus detection system.  $\beta$ -Tubulin was used as an internal control for protein loading and transfer efficiency.

### Statistical Analysis

All data were expressed as means  $\pm$  SEM. The effects of treatment, training day, and their interaction on the behavioral performance in the Morris water maze training task were evaluated by overall ANOVA for repeated measures. Other data were analyzed by 2-tailed Student's *t* test. A *P* value < .05 was considered statistically significant.

## Results

### Isolation Housing Decreased Learning and Memory of Aged APP/PS1 Mice

The Morris water maze data showed that escape latency of isolated APP/PS1 mice was extended during hidden-platform acquisition training compared with group controls (group effect:  $F_{1,115} = 2.889$ ,  $P = .049$ ) (Figure 1a). However, the difference was not observed on visible platform testing (group effect:  $F_{1,46} = 0.188$ ,  $P = .669$ ) (Figure 1a). Isolated APP/PS1 mice swam faster than group mice in both visible platform testing and hidden-platform testing, but the difference was not statistically significant (group effect:  $F_{1,46} = 0.994$ ,  $P = .329$ ;  $F_{1,115} = 0.923$ ,  $P = .470$ , respectively) (Figure 1b). These results suggest that isolation would specifically impair spatial learning ability but did not affect motor and/or visual ability. Additionally, isolated APP/PS1 mice displayed further spatial memory declines in the probe test, as revealed by decreased percentages of time spent in the target quadrant ( $P = .023$ ) (Figure 1c) and the number of target quadrant crossing ( $P = .032$ ) (Figure 1d). Swimming trial analysis showed that isolated APP/PS1 mice swam in each quadrant irregularly, but group controls had a preference for the target quadrant (Figure 1e). These results suggest that isolated APP/PS1 mice exhibited further deterioration of long-term memory.

Consistently, short-term memory declines were also more severe in isolated APP/PS1 mice during the Y maze test, as revealed by decreased percentages of time spent in the NA ( $P = .01$ ) (Figure 2a) and number of NA entrances ( $P = .012$ ) (Figure 2b). In addition, isolated mice moved slightly faster than group controls ( $P > .05$ ) (Figure 2c), suggesting the occurrence of mild hyperactivity.


### SI Decreased Exploring Behaviors of Aged APP/PS1 Mice

Results of the open field test suggest that isolated APP/PS1 mice showed a decline in exploratory activity compared with group controls (Figure 3a), as revealed by decreases in the time spent in the center area ( $P = .011$ ) (Figure 3b) and the number of crosses into the center area ( $P = .029$ ) (Figure 3c). In addition, isolated APP/PS1 mice exhibited mild anxiety-like behaviors, reflected by a tendency to increase movement speed ( $P = .484$ ) (Figure 3d) and defecation number ( $P = .203$ ) (Figure 3e).


### SI Increased A $\beta$ Accumulation via Upregulation of PS1 and Downregulation of NEP in the Hippocampus of Aged APP/PS1 Mice

Isolation housing increased amyloid deposits in the hippocampus, visualized as Thioflavin-S positive dense core plaques (Figure 4a) and 6E10-immunopositive diffuse and mature plaques (Figure 4b). Quantification data revealed that the percentage area occupied by Thioflavin-S or 6E10 labeled plaques was higher in the hippocampus of aged APP/PS1 mice than in group controls ( $P = .013$ ;  $P = .033$ , respectively; Figure 4c-d).

To further reveal the molecular mechanisms of SI effects on plaque pathogenesis, we examined the expression of proteins involved in A $\beta$  synthesis and degradation using Western-blot analysis (Figure 4e-f). There was no difference in APP levels between isolated and group APP/PS1 mice ( $P = .211$ ), suggesting that SI does not alter APP generation. However, SI altered APP hydrolysis in the hippocampus of isolated APP/PS1 mice, as revealed by prominent increases in



**Figure 1.** Morris water maze test. (a) The mean escape latency in the visible platform test (days 1–2) and hidden platform test (days 3–7). (b) Swimming speed. (c) Percentage of time spent in each quadrant in the probe test. (d) Crossing the platform number. (e) Tracings of the typical swim patterns in the probe test. Data represent means  $\pm$  SEM. \*  $P < .05$ , socially isolated (SI) aged APP/PS1 mice ( $n=12$ ) vs group (GR) controls ( $n=13$ ).




**Figure 2.** Y maze test. (a) Percentage of time spent in the novel arm (NA), starting arm (SA), and other arm (OA). (b) Number of arm entrances. (c) Travelling speed during the test. Data represent means  $\pm$  SEM. \*  $P < .05$ , socially isolated (SI) aged APP/PS1 mice ( $n=12$ ) vs group (GR) controls ( $n=13$ ).

the amyloidogenic peptide  $A\beta_{1-42}$  ( $P = .023$ ), and mild decreases in the nonamyloidogenic peptide sAPP $\alpha$  compared with their group controls ( $P = .05$ ). Consistent with altered hydrolysis products of APP, APP/PS1 isolated mice exhibited significantly higher levels of PS1 ( $\gamma$ -secretase) ( $P = .046$ ), which is responsible for cleavage of APP C-terminal fragment CTF $\beta$  to create the  $A\beta$  peptide (Donmez et al., 2010). Expression levels of disintegrin and metalloproteinase10 ( $\alpha$ -secretase) ( $P = .462$ ) and  $\beta$ -site amyloid precursor protein-cleaving enzyme 1 ( $\beta$ -secretase) ( $P = .725$ ) were not altered. These enzymes are required to cut APP to generate N-terminal soluble fragment sAPP $\alpha$  and C-terminal fragment CTF $\alpha$ , and sAPP $\beta$  and CTF $\beta$ , respectively (Obregon et al., 2012). Increased expression of CTF $\beta$  ( $P = .029$ ) and  $A\beta_{42}$  ( $P = .025$ ) and decreased expression of sAPP $\alpha$  ( $P = .04$ ) were also observed in the hippocampal samples of isolated APP/PS1 mice by Western blotting using 6E10 and densitometry analysis (Figure 4g-h). To determine whether decreased  $A\beta$  clearance is also involved in the exacerbated  $A\beta$  accumulation in isolated APP/PS1 mice, expression levels of NEP and IDE, 2 proteolytic enzymes that are mainly responsible for eliminating  $A\beta$  from the brain parenchyma (Fukami et al., 2002; Kasturirangan and Sierks, 2010), were compared between isolated and group APP/PS1 mice. We found that isolation housing triggered a decrease in NEP expression


( $P = .009$ ) but did not alter IDE expression ( $P = .129$ ) in the hippocampus (Figure 4e-f). Together, these results reveal that SI increased  $A\beta$  accumulation in the hippocampus of aged APP/PS1 mice, which was associated with upregulation of PS1 and downregulation of NEP.

### SI Decreased Hippocampal Volume and Synapse and Myelin-Related Proteins in Aged APP/PS1 Mice

The hippocampus is a vulnerable region to  $A\beta$  accumulation and is also sensitive to various AD-related risk factors such as hyperlipidemia, diabetes, hypertension, seizures, and stress (Dhikav and Anand, 2011). Indeed, hippocampal atrophy is a main neuropathological basis for irreversible impairment of spatial learning and memory in AD patients as well as in mouse AD models (Bird and Burgess, 2008). We examined whether isolation housing affects hippocampal size in aged APP/PS1 mice. As shown by HE-stained hippocampal serial sections, socially isolated APP/PS1 mice had a smaller dorsal hippocampus than group-housed controls (Figure 5a). Dorsal hippocampus section counts were reduced by 96 slices in isolated mice compared with group mice ( $P = .008$ ) (Figure 5b). Volumetric analyses consistently showed a significant decrease in hippocampal volume of isolated APP/PS1 mice ( $P = 0.034$ ) (Figure 5c). Moreover, the white matter volume



**Figure 3.** The open field test. (a) Tracing of mouse movement during the 10-minute test period. (b) Percentage of time spent in the center area. (c) Number of entries into the center area. (d) Travelling speed. (e) Defecation number during the test. Data represent means  $\pm$  SEM. \*  $P < .05$ , socially isolated (SI) aged APP/PS1 mice ( $n=12$ ) vs group (GR) controls ( $n=13$ ).




**Figure 4.** Analyses of amyloid metabolism-related indexes in the hippocampus. (a-b) Brain sections from isolated aged APP/PS1 mice and group controls were stained with Thioflavin-S (a) or immunostained with anti-6E10 antibody (b) to label plaques. Scale bar = 500  $\mu$ m. (c-d) Percentage of brain area occupied by Thioflavin-S (c) or 6E10 (d) labeled A $\beta$  deposition in the hippocampus. (e-f) Western blotting and densitometry analysis of APP and its hydrolysis products, A $\beta$ 42 and sAPP $\alpha$ , and hydrolytic enzymes a disintegrin and metalloproteinase (ADAM) family protease ADAM10,  $\beta$ -site amyloid precursor protein-cleaving enzyme 1 (BACE1) and PS1 as well as A $\beta$ -degrading enzymes IDE and NEP. (g-h) Western blotting using 6E10 and densitometry analysis showed expression levels of APP, sAPP $\alpha$ , CTF $\beta$ , A $\beta$ 40, and A $\beta$ 42. Data represent means  $\pm$  SEM from 5 mice per group for pathological analyses and from 3 mice per group in 3 independent Western blotting analysis experiments. \*  $P < .05$ , socially isolated (SI) aged APP/PS1 mice vs group (GR) controls.

decreased more than that of the gray matter ( $P = .018$ ) (Figure 5d), indicating that neurites undergo more prominent dystrophy or loss than neuronal cell bodies under AD-like pathology.

Synapses and oligodendrocytes are main components of the hippocampal white matter and are very vulnerable to the

toxicity of A $\beta$  peptides (Haroutunian et al., 2014). Previous studies have revealed that SI also impairs synapse and myelination plasticity in adult mice (Liu et al., 2012a; Pereda-Pérez et al., 2013). Thus, using Western-blot analyses, we determined whether SI affects synapse and myelin-related proteins by



**Figure 5.** Analyses of hippocampal volume and expression levels of synapse, myelin, and neuron-related proteins. (a) Series of H&E-stained sagittal brain sections showing that the cross area of the dorsal hippocampus was smaller in isolated aged APP/PS1 mice than group controls. Note from the No. 265 section (about 1.68 mm lateral to the mid-sagittal fissure), the intermediate region of the hippocampus, a transitional zone of the dorsal hippocampal region and ventral hippocampal region, was observed in isolated mice but was not present in group APP/PS1 mice until at No. 285 section. Scale bar = 500  $\mu$ m. (b) Counting the total number of sections that contain the dorsal hippocampus. (c) Quantification of the volume of the dorsal hippocampus. (d) Relative volume of the hippocampal gray matter and white matter between isolated aged APP/PS1 mice and group controls. (e-f) Western blotting and densitometry analysis of SYP, MBP, and neuronal class III  $\beta$ -tubulin (TUJ-1). Data represent means  $\pm$  SEM from 5 mice per group for pathological analyses and from 3 mice per group in 3 independent Western blotting analysis experiments. \* $P < .05$ , socially isolated (SI) aged APP/PS1 mice vs group (GR) controls. # $P < .05$ , the relative volume of white matter vs the relative volume of gray matter.

assessing SYP and MBP levels in the hippocampus. As expected, decreases in both SYP and MBP expression were observed in isolated APP/PS1 mice compared with group-housed controls (both  $P = .008$ ) (Figure 5e-f). Additionally, immunostaining for PCNA and DCX revealed extremely low levels of generation or proliferation of immature neurons in the hippocampal dentate gyrus of both isolated and group aged APP/PS1 mice (data not shown), supporting the evidence that cell proliferation and neurogenesis are diminished in the aging mouse dentate gyrus (Kannangara et al., 2011). Taken together, these results suggest that exacerbated hippocampal atrophy, along with increased loss of synapse and myelin-related proteins, further contribute to the severe cognitive declines of isolated aged APP/PS1 mice.

### SI Increased Reactive Gliosis in the Hippocampus of Aged APP/PS1 Mice


Reactive astrocytes and microglia that surround amyloid plaques are a neuropathological hallmark of AD (Allen and Barres, 2009). Furthermore, there is a close association of reactive gliosis with the AD process (Avila-Muñoz and Arias, 2014). To investigate the effect of SI on reactive gliosis in aged APP/PS1 mice, hippocampal distribution of astrocytes and microglia was examined by immunostaining for GFAP and Iba-1, respectively.

Isolated APP/PS1 mice showed increases in both GFAP and Iba-1 immunoreactivity surrounding plaques compared with group controls (Figure 6a-b). Quantitative analysis further confirmed that isolation housing increased GFAP or Iba-1-positive labeling intensity in the hippocampus of APP/PS1 mice ( $P = .001$ ;  $P = .006$ , respectively) (Figure 6c-d), confirming its deteriorating role in A $\beta$ -associated gliosis.

### Discussion

A social life where social creatures such as humans and rodents live in a community is vitally important to the overall health of the being. Correspondingly, scientific evidence reveals that SI is harmful to both physical and mental health (Seeman, 2000; Cacioppo et al., 2011; Steptoe et al., 2013). AD, a neurodegenerative disease with gradual declines in social, cognitive, and emotional functions, eventually separates the patients from social and family life (Luzzi et al., 2007; Weiss et al., 2008). However, the internal interaction between SI and AD still remains unclear. Previous experiments have shown that SI causes the early onset of AD-like phenotypes in young adult APP/PS1 mice (Hsiao et al., 2011, 2012). In this study, we demonstrate that aged APP/PS1 mice housed in a group exhibit more serious cognitive dysfunction after living in an isolated environment for 3 months. The





**Figure 6.** Analyses of reactive gliosis in the hippocampus. (a-b) Sagittal brain sections from isolated aged APP/PS1 mice and group controls were immunostained with GFAP and Iba-1 to show expression and distribution of astrocytes and microglia in the hippocampus, respectively. Scale bar = 500  $\mu$ m. (c-d) The quantification analysis of mean integrated optical density (MIOD) of GFAP (c) and Iba-1 (d). Data represent mean  $\pm$  SEM from 5 mice per group. \* $P < .05$ , socially isolated (SI) aged APP/PS1 mice vs group (GR) controls.

pathological basis of the further cognitive deterioration includes more prominent brain A $\beta$  deposit caused by increases in PS1 and decreases in NEP, which subsequently exacerbates hippocampal atrophy, synapse and myelin associated protein loss, and glial neuroinflammatory reactions. The finding suggests that AD patients' conditions will worsen significantly, even in the late stages of the disease, when they become socially isolated. In contrast, interventions via social and family life might slow down the progression of AD or arrest further cognitive deterioration.

Although the etiology of AD remains unknown, a number of risk factors including hypertension, diabetes, hyperlipidemia, unhealthy lifestyle, and adverse social conditions are associated with the onset and progression of AD (Barnes and Yaffe, 2011; Jiang et al., 2013a; Barnard et al., 2014). An early epidemiological study showed that SI in older individuals increases the risk of developing AD (Wilson et al., 2007). Corresponding experimental studies reveal that SI exacerbates memory impairment in 3 month-old APP/PS1 mice in a time-dependent manner (Hsiao et al., 2011). The present results suggest that isolation housing has detrimental effects on cognitive declines during the late stages of AD. These data offer clear evidence that group or family life is not only beneficial to prevent the AD onset but can also to delay AD progression.

Apart from memory deficits, the severity of AD is also associated with impairment of exploratory behaviors, especially

showing a loss of curiosity into novel surroundings or environment (Havins et al., 2012). In agreement with this view, isolated aged APP/PS1 mice showed low exploratory activity in the open field test compared with their group controls. In addition, anxiety or depression often occurs in the process of AD but has not been correlated with the severity of cognitive dysfunction (Panza et al., 2010; Maalouf et al., 2011). Previous studies have reported that rearing juvenile or adult rodents in SI environments elicits a variety of behavioral abnormalities, such as increased aggressiveness, anxiety, and hyperactivity (Weiss et al., 2004; Koike et al., 2009; Lukkes et al., 2009; Ouchi et al., 2013). In the present study, we found that isolated aged APP/PS1 mice showed no significant increases in water maze swimming speed and traveling speed in the Y maze, as well as the defecation number and grooming number in the open field, indicating that SI-induced hyperactivity and anxiety are not obvious in aged AD models. Together, the present behavioral data suggest that isolation housing selectively exacerbates cognitive impairment in the late stages of AD.

A considerable body of evidence indicates that an imbalance between A $\beta$  production and clearance leads to A $\beta$  accumulation, which plays a central role in the cognitive dysfunction and neurodegenerative changes that characterize AD (Mawuenyega et al., 2010; Holtzman et al., 2011). Along with more severe memory deficits, isolated aged APP/PS1 mice showed increased A $\beta$  accumulation in the hippocampus accompanied with increased

PS1 expression and decreased NEP expression. Early studies also revealed that 8-week isolated housing increases A $\beta$  levels and  $\gamma$ -secretase activity in the hippocampus of 3-month-old APP/PS1 mice (Hsiao et al., 2012). Aside from SI, other manners of chronic stresses, such as chronic immobilization stress (Jeong et al., 2006) and mild unpredictable chronic stress (Cuadrado-Tejedor et al., 2012), exacerbate amyloid production in transgenic mouse models of AD. Furthermore, unpredictable chronic stress can also alter APP metabolism toward the amyloidogenic pathway in normal, middle-aged rats (Catania et al., 2009). These studies highlight that various stresses, including SI, stimulate increases in the amyloidogenic pathway. Further evidence indicates that increased glucocorticoids (Green et al., 2006; Li et al., 2010) and oxidative stress (Hsiao et al., 2012) contribute to SI-altered APP metabolism, although the underlying mechanisms warrant further investigation.

NEP and IDE are 2 major A $\beta$ -degrading enzymes, with each playing an important role in maintaining homeostasis of A $\beta$  in the normal brain (Kasturirangan and Sierks, 2010). NEP is typically expressed in axonal and synaptic membranes (Fukami et al., 2002) and is responsible for degrading monomeric and oligomeric forms of A $\beta$  (El-Amouri et al., 2008). IDE is expressed predominantly in the cytosol of neurons but is also secreted from microglial cells and contributes to the degradation of polymeric and fibrillary forms of A $\beta$  in the brain (Eckman and Eckman, 2005; Qiu and Folstein, 2006). Decreases in NEP mRNA, protein level, and activity are observed in the postmortem AD hippocampus (Yasojima et al., 2001; Caccamo et al., 2005; Wang et al., 2005), and NEP protein expression is inversely correlated with the severity of A $\beta$  accumulation and cognitive impairment (Wang et al., 2010). In contrast, conflicting results exist for IDE in AD, and its protein expression seems not to be correlated with A $\beta$  or clinical diagnosis (Zhao et al., 2007; Wang et al., 2010). Experimental studies also revealed that intracranial injection of AAV expressing NEP, but not IDE, reduces amyloid pathology in both the hippocampus and cortex of APP/PS1 mice (Carty et al., 2013). These findings suggest that NEP, rather than IDE, is a possible therapeutic target for A $\beta$  degradation in AD. Consistent with these studies, the present results reveal that isolated housing decreases expression of NEP, but not IDE, in aged APP/PS1 mice. Taken together, isolated housing increases PS1 and decreases NEP expression, further exacerbating A $\beta$  accumulation in the hippocampus of aged APP/PS1 mice.

Consistent with additional A $\beta$  accumulation, the present results reveal that isolated housing decreases the hippocampal volume of APP/PS1 mice. The hippocampus is essential for declarative memory synthesis and hippocampal atrophy and is a core pathological substrate for cognitive impairment in AD patients (deToledo-Morrell et al., 2007; Barnes et al., 2009). Available evidence suggests that accumulation of A $\beta$ , together with its subsequent pathological events, including formation of neurofibrillary tangles, disrupted mitochondrial energy metabolism, Ca<sup>2+</sup> overload, activation of apoptotic pathways, oxidative stress, and glia-associated neuroinflammation, contribute to impairment of hippocampal structure and function (Dhikav and Anand, 2011). Besides A $\beta$  and the examined secondary damage factors, several studies have revealed that SI itself has detrimental effects on hippocampal volume, neurogenesis, synapse plasticity, and axonal myelination (Stranahan et al., 2006; Djordjevic et al., 2009, 2012; Liu et al., 2012a; Pereda-Pérez et al., 2013). Indeed, the hippocampus has long been known as a target of stress hormones and is an especially plastic and vulnerable region of the brain (Joëls et al., 2003; Blank et al., 2004). The hippocampus expresses a high number of corticosteroid receptors

and is responsive to circulating corticosteroids (Chaouloff and Groc, 2011). Elevated cortisol induces transcription of inhibitory glucocorticoid receptors, which are associated with depressed excitatory synaptic transmission and increased hippocampal atrophy (de Leon et al., 1998; Setiawan et al., 2007; Patel et al., 2008; Kamal et al., 2014). Furthermore, baseline plasma cortisol levels are elevated in the elderly compared with younger adults, which may identify cortisol as a key contributor to increased aging-associated hippocampal atrophy (Wright et al., 2005; McAuley et al., 2009). Isolated housing for 11 days does not change corticosterone in young (2-month-old) female C57BL/6J mice, but increases corticosterone significantly at the onset of the dark cycle in aged (17- to 18-month-old) mice (Kannangara et al., 2011). As mentioned above, glucocorticoids in turn increase A $\beta$  production (Green et al., 2006; Li et al., 2010). Therefore, both SI induced A $\beta$ -associated damage and SI itself are involved in the process of hippocampal atrophy in aged APP/PS1 mice.

In conclusion, the present results demonstrate that isolation housing exacerbates AD-like pathophysiology in aged APP/PS1 mice. It should be noted that the contribution of abnormal APP metabolism to SI-induced cognitive declines of aged mice has not been addressed in the present study. Furthermore, despite being one of the most widely used models, APP/PS1 mice cannot recapitulate fully all of the pathological features of AD. Thus, the internal interaction between SI and AD needs to be further confirmed by using multiple transgenic and/or natural models of AD. Clinical studies will also be necessary to investigate the potential benefits and mechanisms of group or family life in delay or counter the cognitive decline of AD patients. The potential findings will help establish useful nonpharmacological interventions, including lifestyle changes, to combat the devastating neurodegeneration.

## Acknowledgments

This work was supported by grants from the National Natural Science Foundation of China (no. 81271210) and the Natural Science Foundation of Jiangsu Educational Department (14KJA320001).

Interest Statement: None.

## References

- Allen NJ, Barres BA (2009) Neuroscience: glia-more than just brain glue. *Nature* 457:675–677.
- Avila-Muñoz, Arias C (2014) When astrocytes become harmful: functional and inflammatory responses that contribute to Alzheimer's disease. *Ageing Res Rev* 18C:29–40.
- Azzinnari D, Sigrist H, Staehli S, Palme R, Hildebrandt T, Lepar G, Hengerer B5, Seifritz E6, Pryce CR (2014) Mouse social stress induces increased fear conditioning, helplessness and fatigue to physical challenge together with markers of altered immune and dopamine function. *Neuropharmacology* 85:328–341.
- Barnard ND, Bush AI, Ceccarelli A, Cooper J, de Jager CA, Erickson KI, Fraser G, Kesler S, Levin SM, Lucey B, Morris MC, Squitti R (2014) Dietary and lifestyle guidelines for the prevention of Alzheimer's disease. *Neurobiol Aging* 35:S74–78.
- Barnes DE, Yaffe K (2011) The projected effect of risk factor reduction on Alzheimer's disease prevalence. *Lancet Neurol* 10:819–828.
- Barnes J, Bartlett JW, van de Pol LA, Loy CT, Scahill RI, Frost C, Thompson P, Fox NC (2009) A meta-analysis of hippocampal atrophy rates in Alzheimer's disease. *Neurobiol Aging* 30:1711–1723.

- Bird CM, Burgess N (2008) The hippocampus memory: insights from spatial processing. *Nat Rev Neurosci* 9:182–194.
- Blank T, Nijholt I, Spiess J (2004) Molecular determinants mediating effects of acute stress on hippocampus-dependent synaptic plasticity and learning. *Mol Neurobiol* 29:131–138.
- Caccamo A, Oddo S, Sugarman MC, Akbari Y, LaFerla FM (2005) Age- and region-dependent alterations in Abeta-degrading enzymes: implications for Abeta-induced disorders. *Neurobiol Aging* 26:645–654.
- Cacioppo JT, Hawkley LC, Norman GJ, Berntson GG (2011) Social isolation. *Ann NY Acad Sci* 1231:17–22.
- Carty N, Nash KR, Brownlow M, Cruite D, Wilcock D, Selenica ML, Lee DC, Gordon MN, Morgan D (2013) Intracranial injection of AAV expressing NEP but not IDE reduces amyloid pathology in APP/PS1 transgenic mice. *PLoS One* 8:e59626.
- Catania C, Sotiropoulos I, Silva R, Onofri C, Breen KC, Sousa N, Almeida OFX (2009) The amyloidogenic potential and behavioral correlates of stress. *Mol Psychiatry* 14:95–105.
- Chaoulouff F, Groc L (2011) Temporal modulation of hippocampal excitatory transmission by corticosteroids and stress. *Front Neuroendocrinol* 32:25–42.
- Chen H, Yang Y, Yao HH, Tang XC, Ding JH, Wang H, Hu G (2006) Protective effects of iptakalim, a novel ATP-sensitive potassium channel opener, on global cerebral ischemia-evoked insult in gerbils. *Acta Pharmacol Sin* 27:665–672.
- Cuadrado-Tejedor M, Ricobaraza A, Frechilla D, Franco R, Pérez-Mediavilla A, Garcia-Osta A (2012) Chronic mild stress accelerates the onset and progression of the Alzheimer's disease phenotype in Tg2576 mice. *J Alzheimers Dis* 28:567–578.
- de Leon M, Lupien SJ, de Santi S, Convit A, Tarshish C, Nair NP, Thakur M, McEwen BS, Hauger RL, Meaney MJ (1998) Cortisol levels during human aging predict hippocampal atrophy and memory deficits. *Nat Neurosci* 1:69–73.
- deToledo-Morrell L, Stoub TR, Wang C (2007) Hippocampal atrophy and disconnection in incipient and mild Alzheimer's disease. *Prog Brain Res* 163:741–753.
- Dhikav V, Anand K (2011) Potential predictors of hippocampal atrophy in Alzheimer's disease. *Drugs Aging* 28:1–11.
- Díez E, Daban F, Pasarín M, Artazcoz L, Fuertes C, López MJ, Calzada N (2014) Evaluation of a community program to reduce isolation in older people due to architectural barriers. *Gac Sanit* 28:386–388.
- Djordjevic A, Adzic M, Djordjevic J, Radojic MB (2009) Chronic social isolation is related to both upregulation of plasticity genes and initiation of proapoptotic signaling in Wistar rat hippocampus. *J Neural Transm* 116:1579–1589.
- Djordjevic A, Djordjevic J, Elaković I, Adzic M, Matić G, Radojic MB (2012) Fluoxetine affects hippocampal plasticity, apoptosis and depressive-like behavior of chronically isolated rats. *Prog Neuropsychopharmacol Biol Psychiatry* 36:92–100.
- Dong H, Csernansky JG (2009) Effects of stress and stress hormones on amyloid-beta protein and plaque deposition. *J Alzheimers Dis* 18:459–469.
- Donmez G, Wang D, Cohen DE, Guarente L (2010) SIRT1 suppresses beta-amyloid production by activating the alpha-secretase gene ADAM10. *Cell* 142:320–332.
- Eckman EA, Eckman CB (2005) Abeta-degrading enzymes: modulators of Alzheimer's disease pathogenesis and targets for therapeutic intervention. *Biochem Soc Trans* 33:1101–1105.
- El-Amouri SS, Zhu H, Yu J, Marr R, Verma IM, Kindy MS (2008) Nprilysin: an enzyme candidate to slow the progression of Alzheimer's disease. *Am J Pathol* 172:1342–1354.
- Fukami S, Watanabe K, Iwata N, Haraoka J, Lu B, Gerard NP, Gerard C, Fraser P, Westaway D, St George-Hyslop P, Saido TC (2002) Abeta-degrading endopeptidase, neprilysin, in mouse brain: synaptic and axonal localization inversely correlating with Abeta pathology. *Neurosci Res* 43:39–56.
- Gilman SE, Ni MY, Dunn EC, Breslau J, McLaughlin KA, Smoller JW, Perlis RH (2014) Contributions of the social environment to first-onset and recurrent mania. *Mol Psychiatry* [Epub ahead of print].
- Green KN, Billings LM, Roozendaal B, McLaughlin JL, LaFerla FM (2006) Glucocorticoids increase amyloid-beta and tau pathology in a mouse model of Alzheimer's disease. *J Neurosci* 26: 9047–9056.
- Hand C, McColl MA, Birtwhistle R, Kotecha JA, Batchelor D, Barber KH (2014) Social isolation in older adults who are frequent users of primary care services. *Can Fam Physician* 60:e322, e324–329.
- Haroutunian V, Katsel P, Roussos P, Davis KL, Altshuler LL, Bartzokis G (2014) Myelination, oligodendrocytes, and serious mental illness. *Glia* 62:1856–1877.
- Havins WN, Massman PJ, Doody R (2012) Factor structure of the Geriatric Depression Scale and relationships with cognition and function in Alzheimer's disease. *Dement Geriatr Cogn Disord* 34:360–372.
- Hsiao YH, Chen PS, Chen SH, Gean PW (2011) The involvement of Cdk5 activator p35 in social isolation-triggered onset of early Alzheimer's disease-related cognitive deficit in the transgenic mice. *Neuropsychopharmacology* 36:1848–1858.
- Hsiao YH, Kuo JR, Chen SH, Gean PW (2012) Amelioration of social isolation-triggered onset of early Alzheimer's disease-related cognitive deficit by N-acetylcysteine in a transgenic mouse model. *Neurobiol Dis* 45:1111–1120.
- Holtzman DM, Morris JC, Goate AM (2011) Alzheimer's disease: the challenge of the second century. *Sci Transl Med* 3:77sr1.
- Huang Y, Mucke L (2012) Alzheimer mechanisms and therapeutic strategies. *Cell* 148:1204–1222.
- Jeong YH, Park CH, Yoo J, Shin KY, Ahn SM, Kim HS, Lee SH, Emson PC, Suh YH (2006) Chronic stress accelerates learning and memory impairments and increases amyloid deposition in APPV717I-CT100 transgenic mice, an Alzheimer's disease model. *FASEB J* 20:729–731.
- Jiang T, Yu JT, Tian Y, Tan L (2013a) Epidemiology and etiology of Alzheimer's disease: from genetic to non-genetic factors. *Curr Alzheimer Res* 10:852–867.
- Jiang Z, Cowell RM, Nakazawa K (2013b) Convergence of genetic and environmental factors on parvalbumin-positive interneurons in schizophrenia. *Front Behav Neurosci* 7:116.
- Joëls M, Verkuyl JM, Van Riel E (2003) Hippocampal and hypothalamic function after chronic stress. *Ann N Y Acad Sci* 1007:367–378.
- Kamal A, Ramakers GM, Altinbilek B, Kas MJ (2014) Social isolation stress reduces hippocampal long-term potentiation: effect of animal strain and involvement of glucocorticoid receptors. *Neuroscience* 256:262–270.
- Kannangara TS, Lucero MJ, Gil-Mohapel J, Drapala RJ, Simpson JM, Christie BR, van Praag H (2011) Running reduces stress and enhances cell genesis in aged mice. *Neurobiol Aging* 32:2279–2286.
- Kasturirangan S, Sierks M (2010) Targeted hydrolysis of Beta-amyloid with engineered antibody fragment. *Curr Alzheimer Res* 7:214–222.
- Koike H, Ibi D, Mizoguchi H, Nagai T, Nitta A, Takuma K, Nabeshima T, Yoneda Y, Yamada K (2009) Behavioral abnormality and pharmacologic response in social isolation-reared mice. *Behav Brain Res* 202:114–121.
- Li WZ, Li WP, Yao YY, Zhang W, Yin YY, Wu GC, Gong HL (2010) Glucocorticoids increase impairments in learning and mem-

- ory due to elevated amyloid precursor protein expression and neuronal apoptosis in 12-month old mice. *Eur J Pharmacol* 628:108–115.
- Liu J, Dietz K, DeLooyt JM, Pedre X, Kelkar D, Kaur J, Vialou V, Lobo MK, Dietz DM, Nestler EJ, Dupree J, Casaccia P (2012a) Impaired adult myelination in the prefrontal cortex of socially isolated mice. *Nat Neurosci* 15:1621–1623.
- Liu L, Lu Y, Kong H, Li L, Marshall C, Xiao M, Ding J, Gao J, Hu G (2012b) Aquaporin-4 deficiency exacerbates brain oxidative damage and memory deficits induced by long-term ovarian hormone deprivation and D-galactose injection. *Int J Neuropsychopharmacol* 15:55–68.
- Lozano L, Lara-Lemus R, Zenteno E, Alvarado-Vásquez N (2014) The mitochondrial O-linked N-acetylglucosamine transferase (mOGT) in the diabetic patient could be the initial trigger to develop Alzheimer disease. *Exp Gerontol* 58C:198–202.
- Lukkes JL, Mokin MV, Scholl JL, Forster GL (2009) Adult rats exposed to early-life social isolation exhibit increased anxiety and conditioned fear behavior, and altered hormonal stress responses. *Horm Behav* 55:248–256.
- Luzzi S, Piccirilli M, Provinciali L (2007) Perception of emotions on happy/sad chimeric faces in Alzheimer disease: relationship with cognitive functions. *Alzheimer Dis Assoc Disord* 21:130–135.
- Maalouf M, Ringman JM, Shi J (2011) An update on the diagnosis and management of dementing conditions. *Rev Neurol Dis* 8:e68–87.
- Mawuenyega KG, Sigurdson W, Ovod V, Munsell L, Kasten T, Morris JC, Yarasheski KE, Bateman RJ (2010) Decreased clearance of CNS beta-amyloid in Alzheimer's disease. *Science* 330:1774.
- McAuley MT, Kenny RA, Kirkwood TB, Wilkinson DJ, Jones JJ, Miller VM (2009) A mathematical model of aging-related and cortisol induced hippocampal dysfunction. *BMC Neurosci* 10:26.
- Miklowitz DJ (2011) Functional impairment, stress, and psychosocial intervention in bipolar disorder. *Curr Psychiatry Rep* 13:504–512.
- Obregon D, Hou H, Deng J, Giunta B, Tian J, Darlington D, Shaha-duzzaman M, Zhu Y, Mori T, Mattson MP, Tan J (2012) Soluble amyloid precursor protein- $\alpha$  modulates  $\beta$ -secretase activity and amyloid- $\beta$  generation. *Nat Commun* 3:777.
- Ouchi H, Ono K, Murakami Y, Matsumoto K (2013) Social isolation induces deficit of latent learning performance in mice: a putative animal model of attention deficit/hyperactivity disorder. *Behav Brain Res* 238:146–153.
- Panza F, Frisardi V, Capurso C, D'Introno A, Colacicco AM, Imbimbo BP, Santamato A, Vendemiale G, Seripa D, Pilotto A, Capurso A, Solfrizzi V (2010) Late-life depression, mild cognitive impairment, and dementia: possible continuum? *Am J Geriatr Psychiatry* 18:98–116.
- Patel PD, Katz M, Karssen AM, Lyons DM (2008) Stress-induced changes in corticosteroid receptor expression in primate hippocampus and prefrontal cortex. *Psychoneuroendocrinology* 33:360–367.
- Peña-Longobardo LM, Oliva-Moreno J (2014) Caregiver Burden in Alzheimer's Disease Patients in Spain. *J Alzheimers Dis* [Epub ahead of print].
- Pereda-Pérez I, Popović N, Otalora BB, Popović M, Madrid JA, Rol MA, Venero C (2013) Long-term social isolation in the adulthood results in CA1 shrinkage and cognitive impairment. *Neurobiol Learn Mem* 106:31–39.
- Powell ND, Sloan EK, Bailey MT, Arevalo JM, Miller GE, Chen E, Kobor MS, Reader BF, Sheridan JF, Cole SW (2013) Social stress up-regulates inflammatory gene expression in the leukocyte transcriptome via  $\beta$ -adrenergic induction of myelopoiesis. *Proc Natl Acad Sci U S A* 110:16574–16579.
- Qiu WQ, Folstein MF (2006) Insulin, insulin-degrading enzyme and amyloid-beta peptide in Alzheimer's disease: review and hypothesis. *Neurobiol Aging* 27:190–198.
- Querfurth HW, Laferla FM (2010) Alzheimer's Disease. *N Engl J Med* 362:329–344.
- Seeman TE (2000) Health promoting effects of friends and family on health outcomes in older adults. *American Journal of Health Promotion*. *Am J Health Promot* 14:362–370.
- Setiawan E, Jackson MF, MacDonald JF, Matthews SG (2007) Effects of repeated prenatal glucocorticoid exposure on long-term potentiation in the juvenile guinea-pig hippocampus. *J Physiol* 581:1033–1042.
- Steptoe A, Shankar A, Demakakos P, Wardle J (2013) Social isolation, loneliness, and all-cause mortality in older men and women. *Proc Natl Acad Sci U S A* 110:5797–5801.
- Stranahan AM, Khalil D, Gould E (2006) Social isolation delays the positive effects of running on adult neurogenesis. *Nat Neurosci* 9:526–533.
- Tanzi RE, Bertram L (2005) Twenty years of the Alzheimer's disease amyloid hypothesis: a genetic perspective. *Cell* 120:545–555.
- Wang DS, Lipton RB, Katz MJ, Davies P, Buschke H, Kuslansky G, Verghese J, Younkin SG, Eckman C, Dickson DW (2005) Decreased neprilysin immunoreactivity in Alzheimer disease, but not in pathological aging. *J Neuropathol Exp Neurol* 64:378–385.
- Wang Q, Xu Z, Tang J, Sun J, Gao J, Wu T, Xiao M (2013) Voluntary exercise counteracts A $\beta$ 25–35-induced memory impairment in mice. *Behav Brain Res* 256: 618–625.
- Wang S, Wang R, Chen L, Bennett DA, Dickson DW, Wang DS (2010) Expression and functional profiling of neprilysin, insulin-degrading enzyme, and endothelin-converting enzyme in prospectively studied elderly and Alzheimer's brain. *J Neurochem* 115:47–57.
- Weiss EM, Kohler CG, Vonbank J, Stadelmann E, Kemmler G, Hinterhuber H, Marksteiner J (2008) Impairment in emotion recognition abilities in patients with mild cognitive impairment, early and moderate Alzheimer disease compared with healthy comparison subjects. *Am J Geriatr Psychiatry* 16:974–980.
- Weiss IC, Pryce CR, Jongen-Rêlo AL, Nanz-Bahr NI, Feldon J (2004) Effect of social isolation on stress-related behavioural and neuroendocrine state in the rat. *Behav Brain Res* 152:279–295.
- Wilson RS, Krueger KR, Arnold SE, Schneider JA, Kelly JF, Barnes LL, Tang Y, Bennett DA (2007) Loneliness and risk of Alzheimer disease. *Arch Gen Psychiatry* 64:234–240.
- Wright CE, Kunz-Ebrecht SR, Illife S, Foese O, Steptoe A (2005) Physiological correlates of cognitive functioning in an elderly population. *Psychoneuroendocrinology* 30:826–838.
- Xu ZQ, Zhang LQ, Wang Q, Marshall C, Xiao N, Gao JY, Wu T, Ding J, Hu G, Xiao M (2013) Aerobic exercise combined with anti-oxidative treatment does not counteract moderate or mid-stage Alzheimer-like pathophysiology of APP/PS1 mice. *CNS Neurosci Ther* 19:795–803.
- Yasojima K, Akiyama H, McGeer EG, McGeer PL (2001) Reduced neprilysin in high plaque areas of Alzheimer brain: a possible relationship to deficient degradation of beta-amyloid peptide. *Neurosci Lett* 297:97–100.
- Zhao Z, Xiang Z, Haroutunian V, Buxbaum JD, Stetka B, Pasinetti GM (2007) Insulin degrading enzyme activity selectively decreases in the hippocampal formation of cases at high risk to develop Alzheimer's disease. *Neurobiol Aging* 28:824–830.
- Zucchella C, Bartolo M, Bernini S, Picascia M, Sinforiani E (2014). Quality of Life in Alzheimer Disease: A Comparison of Patients' and Caregivers' Points of View. *Alzheimer Dis Assoc Disord* [Epub ahead of print].