University of Kentucky UKnowledge

Internal Medicine Presentations

Internal Medicine

1-21-2010

Ventricular function, myocardial delayed enhancement and patient-reported quality of life in adolescents and adults with repaired tetralogy of Fallot

Jimmy C. Lu University of Michigan - Ann Arbor

Timothy B. Cotts University of Michigan - Ann Arbor

Prachi P. Agarwal University of Michigan - Ann Arbor

Anil K. Attili University of Kentucky, anil.attili@uky.edu

Adam L. Dorfman University of Michigan - Ann Arbor

Follow this and additional works at: https://uknowledge.uky.edu/internalmedicine_present

Part of the Medicine and Health Sciences Commons Right click to open a feedback form in a new tab to let us know how this document benefits you.

Repository Citation

Lu, Jimmy C.; Cotts, Timothy B.; Agarwal, Prachi P.; Attili, Anil K.; and Dorfman, Adam L., "Ventricular function, myocardial delayed enhancement and patient-reported quality of life in adolescents and adults with repaired tetralogy of Fallot" (2010). *Internal Medicine Presentations*. 5. https://uknowledge.uky.edu/internalmedicine_present/5

This Presentation is brought to you for free and open access by the Internal Medicine at UKnowledge. It has been accepted for inclusion in Internal Medicine Presentations by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Journal of Cardiovascular Magnetic Resonance

Poster presentation

Ventricular function, myocardial delayed enhancement and patient-reported quality of life in adolescents and adults with repaired tetralogy of Fallot

Jimmy C Lu^{*1}, Timothy B Cotts¹, Prachi P Agarwal¹, Anil K Attili² and Adam L Dorfman¹

Address: ¹University of Michigan, Ann Arbor, MI, USA and ²University of Kentucky, Lexington, KY, USA * Corresponding author

from 13th Annual SCMR Scientific Sessions Phoenix, AZ, USA. 21-24 January 2010

Published: 21 January 2010

Journal of Cardiovascular Magnetic Resonance 2010, 12(Suppl 1):P11 doi:10.1186/1532-429X-12-S1-P11

This abstract is available from: http://jcmr-online.com/content/12/S1/P11 © 2010 Lu et al; licensee BioMed Central Ltd.

Introduction

In patients with repaired tetralogy of Fallot (TOF), right ventricular myocardial delayed enhancement (MDE) and diastolic dysfunction are common, and have been associated with decreased exercise capacity and increased arrhythmia. Predictors of quality of life (QOL) have not been reported in this population.

Purpose

We assessed the hypothesis that a greater degree of MDE in adolescents and adults with repaired TOF would correlate with diastolic dysfunction and decreased QOL.

Methods

Patients with repaired TOF \geq 14 years old undergoing cardiovascular magnetic resonance (CMR) had ventricular volumes measured from SSFP cine images in the short axis plane, and peak early and atrial filling rates, early filling fraction and deceleration time calculated from right ventricular inflow curves, generated by the sum of tricuspid inflow and pulmonary insufficiency measured from phase contrast imaging. MDE was scored 0-3 (none, <2 cm, 2-3 cm, >3 cm) in each of 6 regions of the right ventricle and added for a total score. Patients completed the Short Form 36, version 2, a QOL assessment validated in multiple populations with a normalized scale.

Results

Forty-seven patients (32 ± 15 years old, 62% male) underwent CMR. MDE was present in 42/46 (91%), with enddiastolic forward flow (EDFF) in the pulmonary artery in 29/45 (64%). Mean right ventricular ejection fraction (RVEF) was $45 \pm 8\%$, with mean left ventricular ejection fraction (LVEF) 54 \pm 8%. MDE inversely correlated with RVEF (r = 0.30, p = 0.04). Patients with MDE scores above the median had a lower early filling fraction (35.5 ± 11.9% vs. $44.1 \pm 14.6\%$, p = 0.04), with a trend toward higher atrial filling rates (357 ± 123 vs. 287 ± 128 ml/s, p = 0.08). MDE did not differ in patients with or without EDFF, and did not correlate with pulmonary regurgitant fraction or volume. QOL physical and mental summary scores did not differ from population-derived norms. Physical functioning subscale scores trended lower than the general population $(47.6 \pm 9.7 \text{ vs. } 50.0 \pm 10.0, \text{ p} =$ 0.10) and correlated positively with higher MDE scores (r = 0.32, p = 0.03). Role-physical subscale scores correlated with higher LVEF (r = 0.46, p = 0.001) and longer deceleration time (r = 0.31, p < 0.05) and trended with higher RVEF (r = 0.27, p = 0.07). QOL scores did not differ with EDFF, pulmonary regurgitation, or end diastolic volume.

Conclusion

Patients with greater degree of MDE demonstrated right ventricular systolic dysfunction and evidence of impaired relaxation, but did not have lower QOL scores. LVEF was the best predictor of patient-reported QOL, underscoring

Open Access

the importance of preserving LVEF in this population. The influence of pulmonary valve replacement on QOL remains unclear.

