



University of Kentucky
UKnowledge

Superfund Research Center Faculty
Publications

Superfund Research Center

3-2020

Healthful Nutrition as a Prevention and Intervention Paradigm to Decrease the Vulnerability to Environmental Toxicity or Stressors and Associated Inflammatory Disease Risks

Bernhard Hennig
University of Kentucky, bhennig@uky.edu

Pan Deng
University of Kentucky, pde233@uky.edu

Follow this and additional works at: https://uknowledge.uky.edu/superfund_facpub



Part of the [Environmental Health and Protection Commons](#), and the [Nutrition Commons](#)

Right click to open a feedback form in a new tab to let us know how this document benefits you.

Repository Citation

Hennig, Bernhard and Deng, Pan, "Healthful Nutrition as a Prevention and Intervention Paradigm to Decrease the Vulnerability to Environmental Toxicity or Stressors and Associated Inflammatory Disease Risks" (2020). *Superfund Research Center Faculty Publications*. 8.
https://uknowledge.uky.edu/superfund_facpub/8

This Article is brought to you for free and open access by the Superfund Research Center at UKnowledge. It has been accepted for inclusion in Superfund Research Center Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@sv.uky.edu.

Healthful Nutrition as a Prevention and Intervention Paradigm to Decrease the Vulnerability to Environmental Toxicity or Stressors and Associated Inflammatory Disease Risks

Digital Object Identifier (DOI)

<https://doi.org/10.1002/fft2.6>

Notes/Citation Information

Published in *Food Frontiers*, v. 1, issue 1.

© 2020 The Authors

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial License](#), which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

HIGHLIGHTS

Healthful nutrition as a prevention and intervention paradigm to decrease the vulnerability to environmental toxicity or stressors and associated inflammatory disease risks

Bernhard Hennig | Pan Deng

Superfund Research Center, University of Kentucky, Lexington, Kentucky

CorrespondenceBernhard Hennig, Superfund Research Center, University of Kentucky, Lexington, KY.
Email: bhennig@uky.edu**Funding information**

NIEHS/NIH, Grant/Award Number: P42ES007380; Kentucky Agricultural Experiment Station

Many noncommunicable diseases, such as atherosclerosis and other cardiometabolic diseases, are defined as inflammatory diseases, which suggests that multifactorial interactions linked to exacerbated disease pathology include pro-inflammatory chemical and nonchemical stressor. Recent data suggest that genetic and lifestyle factors are independently associated with susceptibility to cardiovascular diseases (Khera et al., 2016; Satija et al., 2018; Tong, 2016). Thus, potential biological interactions between chemical and nonchemical stressors and buffers and other lifestyle factors will determine disease outcome. Chemical stressors include environmental pollutants with pro-oxidant and pro-inflammatory properties, such as air pollutants, both gaseous and particulate matter, and persistent organic pollutants, such as dioxin-like polychlorinated biphenyls (PCBs). Many persistent organic pollutants also have an affinity for the aryl hydrocarbon receptor (AhR). AhR ligands lead to induction of cytochrome P450s and free radical formation, followed by inflammation. For example, dioxin-like PCBs can increase endothelial cell dysfunction, activation, and inflammation (e.g., increased adhesion molecule expression and cytokine production) (Hennig et al., 2005).

It is unlikely that pollutant remediation alone will be sufficient to address the health impacts associated with chronic pollutant exposure, because most environmental toxicants such as persistent organic pollutants (POPs) are ubiquitous in the ecosystem. Thus, there is a need to explore preventive measures of environmental exposure and disease risk through positive lifestyle changes such as healthful nutrition (Hennig et al., 2007). Our data indicate that plant-derived diets

enriched with fiber or bioactive food components such as phytochemicals can prevent or decrease toxicant-induced inflammation (Deng et al., 2020). In fact, our data suggest that nutrition, or the type of diet we eat, can modulate environmental insults and disease outcome. For example, fats/oils high in omega-6 fatty acids can act as pro-oxidative and pro-inflammatory stressors, thus exacerbating an inflammatory response. Mechanistically, many environmental pollutants and pro-atherosclerotic nutrients/diets can both activate nuclear factor- κ B (NF- κ B) signaling leading to further increases in oxidative stress and inflammation. In contrast, many anti-inflammatory nutrients and/or bioactive phytochemicals can decrease inflammation by activating nuclear factor erythroid 2-related factor 2 (Nrf2) signaling. This is important, because foods rich in omega-3 fatty acids and plant-derived phytochemicals (e.g., polyphenols) or bioactive compounds can act as nonchemical buffers to reduce inflammation and thus prevent disease outcome associated with exposure to environmental toxicants (Hennig et al., 2018; Lim et al., 2019). Our data, including metabolomic profiling, suggest that the pathology of inflammatory diseases initiated or caused by exposure to environmental pollutants (e.g., dioxin-like PCBs) is complex and may involve disturbances in the gut microbiota, liver, and vascular tissues (Deng et al., 2019; Petriello et al., 2018). The literature, including our data, also suggest that nutritional intervention can modulate the gut microbiota (Leeming et al., 2019; Petriello et al., 2018). This suggests that positive lifestyle changes, such as healthful nutrition, can have a disease-reducing impact already at the level of the gut microbiota. Of special interest are meaningful and

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2020 The Authors. *Food Frontiers* published by John Wiley & Sons Australia, Ltd and Nanchang University, Northwest University, Jiangsu University, Zhejiang University, Fujian Agriculture and Forestry University

translational approaches of prevention/intervention to lower disease outcome linked to complex interactions of chemical stressors to modify the outcome of cardiometabolic diseases, that is, diseases linked to major causes of death especially in developed countries. Therefore, adopting healthy lifestyle changes, such as consuming healthy diets rich in plant-derived bioactive nutrients, may reduce the vulnerability to diseases linked to environmental toxic insults (Hennig et al., 2007). This nutritional paradigm in environmental toxicology requires further study in order to improve our understanding of the relationship between nutrition or other lifestyle modifications and toxicant-induced diseases.

ACKNOWLEDGMENT

This study is supported in part by NIEHS/NIH grant P42ES007380 and the Kentucky Agricultural Experiment Station.

REFERENCES

- Deng, P., Barney, J., Petriello, M. C., Morris, A. J., Wahlang, B., & Hennig, B. (2019). Hepatic metabolomics reveals that liver injury increases PCB 126-induced oxidative stress and metabolic dysfunction. *Chemosphere*, 217, 140–149.
- Deng, P., Hoffman, J. B., Petriello, M. C., Wang, C. Y., Li, X. S., Kraemer, M. P., ... Hennig, B. (2020). Dietary inulin decreases circulating ceramides by suppressing neutral sphingomyelinase expression and activity in mice. *Journal of Lipid Research*, 61, 45–53.
- Hennig, B., Ettinger, A. S., Jandacek, R. J., Koo, S., McClain, C., Seifried, H., ... Suk, W. A. (2007). Using nutrition for intervention and prevention against environmental chemical toxicity and associated diseases. *Environmental Health Perspectives*, 115, 493–495.
- Hennig, B., Petriello, M. C., Gamble, M. V., Surh, Y. J., Kresty, L. A., Frank, N., ... Suk, W. A. (2018). The role of nutrition in influencing mechanisms involved in environmentally mediated diseases. *Reviews on Environmental Health*, 33, 87–97.
- Hennig, B., Reiterer, G., Majkova, Z., Oesterling, E., Meerarani, P., & Toborek, M. (2005). Modification of environmental toxicity by nutrients: Implications in atherosclerosis. *Cardiovascular Toxicology*, 5, 153–160.
- Khera, A. V., Emdin, C. A., Drake, I., Natarajan, P., Bick, A. G., Cook, N. R., ... Kathiresan, S. (2016). Genetic risk, adherence to a healthy lifestyle, and coronary disease. *New England Journal of Medicine*, 375, 2349–2358.
- Leeming, E. R., Johnson, A. J., Spector, T. D., & Le Roy, C. I. (2019). Effect of diet on the gut microbiota: Rethinking intervention duration. *Nutrients*, 11. <https://doi.org/10.3390/nu11122862>
- Lim, C. C., Hayes, R. B., Ahn, J., Shao, Y., Silverman, D. T., Jones, R. R., & Thurston, G. D. (2019). Mediterranean diet and the association between air pollution and cardiovascular disease mortality risk. *Circulation*, 139, 1766–1775.
- Petriello, M. C., Hoffman, J. B., Vsevolozhskaya, O., Morris, A. J., & Hennig, B. (2018). Dioxin-like PCB 126 increases intestinal inflammation and disrupts gut microbiota and metabolic homeostasis. *Environmental Pollution*, 242, 1022–1032.
- Satija, A., & Hu, F. B. (2018). Plant-based diets and cardiovascular health. *Trends in Cardiovascular Medicine*, 28, 437–441.
- Tong, H. (2016). Dietary and pharmacological intervention to mitigate the cardiopulmonary effects of air pollution toxicity. *Biochimica et Biophysica Acta*, 1860, 2891–2898.

How to cite this article: Hennig B, Deng P. Healthful nutrition as a prevention and intervention paradigm to decrease the vulnerability to environmental toxicity or stressors and associated inflammatory disease risks. *Food Front*. 2020;1:13–14. <https://doi.org/10.1002/fft2.6>