

5-2013

# 5,5'-Dichloro-2,2'-dimethoxybiphenyl

Hans-Joachim Lehmler  
*University of Iowa*

Huimin Wu  
*University of Iowa*

Sean Parkin  
*University of Kentucky, spark2@uky.edu*

**Right click to open a feedback form in a new tab to let us know how this document benefits you.**

Follow this and additional works at: [https://uknowledge.uky.edu/chemistry\\_facpub](https://uknowledge.uky.edu/chemistry_facpub)

 Part of the [Chemistry Commons](#)

## Repository Citation

Lehmler, Hans-Joachim; Wu, Huimin; and Parkin, Sean, "5,5'-Dichloro-2,2'-dimethoxybiphenyl" (2013). *Chemistry Faculty Publications*. 10.  
[https://uknowledge.uky.edu/chemistry\\_facpub/10](https://uknowledge.uky.edu/chemistry_facpub/10)

This Article is brought to you for free and open access by the Chemistry at UKnowledge. It has been accepted for inclusion in Chemistry Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

---

**5,5'-Dichloro-2,2'-dimethoxybiphenyl**

**Notes/Citation Information**

Published in *Acta Crystallographica Section E: Crystallographic Communications*, v. 69, part 5, p. o650.

This is an open-access article distributed under the terms of the [Creative Commons Attribution Licence](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are cited.

**Digital Object Identifier (DOI)**

<https://doi.org/10.1107/S160053681300826X>

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 5,5'-Dichloro-2,2'-dimethoxybiphenyl

Hans-Joachim Lehmler,<sup>a\*</sup> Huimin Wu<sup>a</sup> and Sean Parkin<sup>b</sup>

<sup>a</sup>The University of Iowa, Department of Occupational and Environmental Health, Iowa City, IA 52242-5000, USA, and <sup>b</sup>University of Kentucky, Department of Chemistry, Lexington, KY 40506-0055, USA

Correspondence e-mail: hans-joachim-lehmler@uiowa.edu


Received 18 March 2013; accepted 25 March 2013

Key indicators: single-crystal X-ray study;  $T = 90$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.042;  $wR$  factor = 0.111; data-to-parameter ratio = 17.9.

In the title molecule,  $\text{C}_{14}\text{H}_{12}\text{Cl}_2\text{O}_2$ , the dihedral angle between the least-square planes of the benzene rings is  $62.17$  (6)°. Both methoxy groups are slightly out of the plane of the benzene rings to which they are attached, making dihedral angles of  $4.22$  (18) and  $18.82$  (16)°.

## Related literature

For background to polychlorinated biphenyls, see: Basu *et al.* (2009); Hu *et al.* (2008); Kaminsky *et al.* (1981); Kennedy *et al.* (1981); McLean *et al.* (1996); Rodenburg *et al.* (2010). For related structures, see: Chattopadhyay *et al.* (1987); Nakaema *et al.* (2008); Sun *et al.* (2001). For the synthesis of the title compound, see: Joshi *et al.* (2011).



## Experimental

## Crystal data

$\text{C}_{14}\text{H}_{12}\text{Cl}_2\text{O}_2$   
 $M_r = 283.14$   
 Monoclinic,  $P2_1/n$   
 $a = 10.9629$  (2) Å  
 $b = 7.2177$  (1) Å  
 $c = 16.7812$  (3) Å  
 $\beta = 104.7108$  (7)°

$V = 1284.32$  (4) Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.50$  mm<sup>-1</sup>  
 $T = 90$  K  
 $0.22 \times 0.20 \times 0.18$  mm

## Data collection

Nonius KappaCCD diffractometer  
 Absorption correction: multi-scan  
 (SCALEPACK; Otwinowski & Minor, 1997)  
 $T_{\min} = 0.899$ ,  $T_{\max} = 0.916$

21357 measured reflections  
 2948 independent reflections  
 2347 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.047$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$   
 $wR(F^2) = 0.111$   
 $S = 1.11$   
 2948 reflections

165 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.43$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.41$  e Å<sup>-3</sup>

Data collection: COLLECT (Nonius, 1998); cell refinement: SCALEPACK (Otwinowski & Minor, 1997); data reduction: DENZO-SMN (Otwinowski & Minor, 1997); program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: XP in SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXL97 and local procedures.

This research was supported by grant Nos. ES05605, ES013661 and ES017425 from the National Institute of Environmental Health Sciences, National Institutes of Health.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH5598).

## References

- Basu, I., Arnold, K. A., Venier, M. & Hites, R. A. (2009). *Environ. Sci. Technol.* **43**, 6488–6492.
- Chattopadhyay, D., Banerjee, T., Majumdar, S. K., Podder, G., Kashino, S. & Haisa, M. (1987). *Acta Cryst.* **C43**, 482–484.
- Hu, D., Martinez, A. & Hornbuckle, K. C. (2008). *Environ. Sci. Technol.* **42**, 7873–7877.
- Joshi, S. N., Vyas, S. M., Duffel, M. W., Parkin, S. & Lehmler, H. J. (2011). *Synthesis*, pp. 1045–1054.
- Kaminsky, L. S., Kennedy, M. W., Adams, S. M. & Guengerich, F. P. (1981). *Biochemistry*, **20**, 7379–7384.
- Kennedy, M. W., Carpentier, N. K., Dymerski, P. P. & Kaminsky, L. S. (1981). *Biochem. Pharmacol.* **30**, 577–588.
- McLean, M. R., Bauer, U., Amaro, A. R. & Robertson, L. W. (1996). *Chem. Res. Toxicol.* **9**, 158–164.
- Nakaema, K., Okamoto, A., Maruyama, S., Noguchi, K. & Yonezawa, N. (2008). *Acta Cryst.* **E64**, o1731.
- Nonius (1998). COLLECT. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Rodenburg, L. A., Guo, J., Du, S. & Cavallo, G. J. (2010). *Environ. Sci. Technol.* **44**, 2816–2821.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sun, D. L., Lindeman, S. V., Rathore, R. & Kochi, J. K. (2001). *J. Chem. Soc. Perkin Trans. 2*, pp. 1585–1594.

## supplementary materials

*Acta Cryst.* (2013). E69, o650 [doi:10.1107/S160053681300826X]

**5,5'-Dichloro-2,2'-dimethoxybiphenyl****Hans-Joachim Lehmler, Huimin Wu and Sean Parkin****Comment**

3,3'-Dichlorobiphenyl (CB11) is a polychlorinated biphenyl (PCB) congener found at comparatively high concentrations in water and air (Basu *et al.*, 2009; Hu *et al.*, 2008; Rodenburg *et al.*, 2010). It is probably produced during the manufacturing of paint pigments and/or the degradation of higher chlorinated PCBs (Basu *et al.*, 2009). Early metabolism studies with CB11 demonstrate its oxidation to mono-hydroxylated metabolites by hepatic cytochrome P450 enzymes (Kaminsky *et al.*, 1981; Kennedy *et al.*, 1981). Similar to other lower chlorinated PCB congeners (McLean *et al.*, 1996), it is likely that these mono-hydroxylated metabolites are further oxidized to dihydroxylated metabolites, such as 2,2'-dihydroxy PCB 11. Here we report the crystal structure of the title compound, a dimethylated derivative of 2,2'-dihydroxy PCB 11.

There is one molecule of the title compound in the asymmetric unit (Fig. 1). The dihedral angle between the least-square planes of the two aromatic rings is 62.17 (6)°. The corresponding dihedral angles of structurally related biphenyls, 2,2'-dimethoxybiphenyls, 4,4'-dimethyl-2,2',5,5'-dimethoxybiphenyl and 2,2',4,4',5,5'-hexamethoxybiphenyl, are comparable with 66.94 (7)°, 69.1° (no s.u. available) and 81.1 (1)°, respectively (Nakaema *et al.*, 2008; Sun *et al.*, 2001; Chattopadhyay *et al.*, 1987). In the title compound, the methoxy group at C7' has an almost coplanar arrangement with aromatic ring system, with a dihedral angle of 4.22 (18)°. In contrast, the methoxy group at C7, with a dihedral angle of 18.82 (16)°, is slightly out of the plane of the aromatic ring system. The twist of this methoxy group out of the plane of the aromatic ring system is relatively large compared to related crystal structures, which display dihedral angles ranging from 1.0° to 12.2° (Nakaema *et al.*, 2008; Sun *et al.*, 2001; Chattopadhyay *et al.*, 1987).

**Experimental**

The title compound was prepared by the Suzuki coupling of 2-iodo-4-chloroanisole and 2-methoxy-5-chlorobenzeneboronic acid as described previously (Joshi *et al.*, 2011). Crystals suitable for crystal structure analysis were obtained by slow evaporation of a methanolic solution.

**Refinement**

H atoms were found in difference Fourier maps and subsequently placed in idealized positions with constrained distances of 0.98 Å (RCH<sub>3</sub>), 0.95 Å (C<sub>Ar</sub>H), and with  $U_{\text{iso}}(\text{H})$  values set to either 1.2 $U_{\text{eq}}$  or 1.5 $U_{\text{eq}}$  (RCH<sub>3</sub>) of the attached atom.

**Computing details**

Data collection: *COLLECT* (Nonius, 1998); cell refinement: *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO-SMN* (Otwinowski & Minor, 1997); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XP* in *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97* (Sheldrick, 2008) and local procedures.




Figure 1

View of the title compound showing displacement ellipsoids drawn at the 50% probability level.

### 5,5'-Dichloro-2,2'-dimethoxybiphenyl

#### Crystal data

$C_{14}H_{12}Cl_2O_2$

$M_r = 283.14$

Monoclinic,  $P2_1/n$

Hall symbol: -P 2yn

$a = 10.9629 (2) \text{ \AA}$

$b = 7.2177 (1) \text{ \AA}$

$c = 16.7812 (3) \text{ \AA}$

$\beta = 104.7108 (7)^\circ$

$V = 1284.32 (4) \text{ \AA}^3$

$Z = 4$

$F(000) = 584$

$D_x = 1.464 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 3160 reflections

$\theta = 1.0\text{--}27.5^\circ$

$\mu = 0.50 \text{ mm}^{-1}$

$T = 90 \text{ K}$

Block, colourless

$0.22 \times 0.20 \times 0.18 \text{ mm}$

#### Data collection

Nonius KappaCCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution:  $9.1 \text{ pixels mm}^{-1}$

$\omega$  scans at fixed  $\chi = 55^\circ$

Absorption correction: multi-scan

(SCALEPACK; Otwinowski & Minor, 1997)

$T_{\min} = 0.899$ ,  $T_{\max} = 0.916$

21357 measured reflections

2948 independent reflections

2347 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.047$

$\theta_{\max} = 27.5^\circ$ ,  $\theta_{\min} = 2.0^\circ$

$h = -14 \rightarrow 14$

$k = -9 \rightarrow 9$

$l = -21 \rightarrow 21$

Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.042$	H-atom parameters constrained
$wR(F^2) = 0.111$	$w = 1/[\sigma^2(F_o^2) + (0.0529P)^2 + 0.9091P]$
$S = 1.11$	where $P = (F_o^2 + 2F_c^2)/3$
2948 reflections	$(\Delta/\sigma)_{\max} < 0.001$
165 parameters	$\Delta\rho_{\max} = 0.43 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.41 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against all reflections. The weighted  $R$ -value  $wR$  and goodness of fit  $S$  are based on  $F^2$ . Conventional  $R$ -values  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -values based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -values based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.07535 (4)	0.15602 (7)	0.32405 (3)	0.02285 (15)
O1	0.56298 (12)	0.53852 (19)	0.38856 (9)	0.0168 (3)
C1	0.43664 (17)	0.2807 (3)	0.33719 (12)	0.0137 (4)
C2	0.45037 (18)	0.4500 (3)	0.37970 (12)	0.0149 (4)
C3	0.35230 (18)	0.5203 (3)	0.40971 (12)	0.0170 (4)
H3	0.3636	0.6321	0.4406	0.020*
C4	0.23774 (18)	0.4264 (3)	0.39443 (12)	0.0171 (4)
H4	0.1709	0.4726	0.4154	0.020*
C5	0.22250 (18)	0.2655 (3)	0.34850 (12)	0.0170 (4)
C6	0.32045 (18)	0.1903 (3)	0.32061 (12)	0.0163 (4)
H6	0.3085	0.0776	0.2903	0.020*
C7	0.56729 (19)	0.7317 (3)	0.40866 (14)	0.0211 (4)
H7A	0.4958	0.7952	0.3718	0.032*
H7B	0.6463	0.7851	0.4020	0.032*
H7C	0.5628	0.7470	0.4659	0.032*
Cl1'	0.59609 (5)	0.10229 (8)	0.08207 (3)	0.02197 (15)
O1'	0.66596 (13)	0.1598 (2)	0.44014 (8)	0.0174 (3)
C1'	0.54159 (18)	0.2061 (3)	0.30496 (12)	0.0137 (4)
C2'	0.65747 (18)	0.1509 (3)	0.35747 (12)	0.0147 (4)
C3'	0.75356 (18)	0.0855 (3)	0.32448 (12)	0.0163 (4)
H3'	0.8319	0.0501	0.3603	0.020*
C4'	0.73626 (19)	0.0711 (3)	0.23964 (13)	0.0175 (4)
H4'	0.8024	0.0277	0.2173	0.021*
C5'	0.62136 (18)	0.1210 (3)	0.18870 (12)	0.0157 (4)
C6'	0.52457 (18)	0.1888 (3)	0.22031 (12)	0.0156 (4)
H6'	0.4465	0.2235	0.1839	0.019*

C7'	0.78413 (19)	0.1104 (3)	0.49485 (13)	0.0212 (5)
H7'1	0.8051	-0.0177	0.4841	0.032*
H7'2	0.7782	0.1214	0.5520	0.032*
H7'3	0.8501	0.1935	0.4860	0.032*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C11	0.0134 (2)	0.0267 (3)	0.0286 (3)	-0.00368 (19)	0.0056 (2)	0.0006 (2)
O1	0.0159 (7)	0.0120 (7)	0.0239 (8)	-0.0027 (5)	0.0075 (6)	-0.0041 (6)
C1	0.0140 (9)	0.0144 (9)	0.0130 (9)	0.0016 (7)	0.0040 (7)	0.0015 (8)
C2	0.0135 (9)	0.0158 (10)	0.0151 (9)	0.0010 (8)	0.0032 (7)	0.0029 (8)
C3	0.0188 (10)	0.0146 (10)	0.0178 (10)	0.0016 (8)	0.0048 (8)	-0.0016 (8)
C4	0.0146 (9)	0.0195 (10)	0.0186 (10)	0.0038 (8)	0.0070 (8)	0.0012 (8)
C5	0.0127 (9)	0.0195 (10)	0.0184 (10)	-0.0010 (8)	0.0031 (7)	0.0035 (8)
C6	0.0172 (10)	0.0152 (10)	0.0167 (10)	0.0009 (8)	0.0048 (8)	-0.0001 (8)
C7	0.0218 (10)	0.0122 (10)	0.0293 (12)	-0.0031 (8)	0.0065 (9)	-0.0035 (9)
C11'	0.0204 (3)	0.0295 (3)	0.0170 (3)	-0.0005 (2)	0.00670 (19)	-0.0048 (2)
O1'	0.0169 (7)	0.0198 (7)	0.0153 (7)	0.0038 (6)	0.0036 (5)	0.0010 (6)
C1'	0.0153 (9)	0.0084 (9)	0.0189 (10)	-0.0007 (7)	0.0069 (7)	-0.0015 (8)
C2'	0.0165 (9)	0.0113 (9)	0.0172 (10)	-0.0017 (7)	0.0058 (8)	-0.0009 (8)
C3'	0.0135 (9)	0.0140 (9)	0.0215 (10)	0.0008 (8)	0.0046 (8)	-0.0002 (8)
C4'	0.0169 (9)	0.0135 (10)	0.0250 (11)	0.0007 (8)	0.0109 (8)	-0.0015 (8)
C5'	0.0197 (10)	0.0146 (10)	0.0142 (9)	-0.0031 (8)	0.0069 (8)	-0.0020 (8)
C6'	0.0142 (9)	0.0127 (9)	0.0201 (10)	-0.0004 (7)	0.0047 (8)	0.0000 (8)
C7'	0.0202 (10)	0.0224 (11)	0.0177 (10)	0.0052 (9)	-0.0016 (8)	0.0004 (9)

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

C11—C5	1.749 (2)	C11'—C5'	1.745 (2)
O1—C2	1.364 (2)	O1'—C2'	1.368 (2)
O1—C7	1.432 (2)	O1'—C7'	1.430 (2)
C1—C6	1.395 (3)	C1'—C6'	1.391 (3)
C1—C2	1.404 (3)	C1'—C2'	1.407 (3)
C1—C1'	1.491 (2)	C2'—C3'	1.391 (3)
C2—C3	1.393 (3)	C3'—C4'	1.392 (3)
C3—C4	1.392 (3)	C3'—H3'	0.9500
C3—H3	0.9500	C4'—C5'	1.379 (3)
C4—C5	1.381 (3)	C4'—H4'	0.9500
C4—H4	0.9500	C5'—C6'	1.390 (3)
C5—C6	1.386 (3)	C6'—H6'	0.9500
C6—H6	0.9500	C7'—H7'1	0.9800
C7—H7A	0.9800	C7'—H7'2	0.9800
C7—H7B	0.9800	C7'—H7'3	0.9800
C7—H7C	0.9800		
C2—O1—C7	117.14 (15)	C2'—O1'—C7'	117.18 (15)
C6—C1—C2	118.90 (17)	C6'—C1'—C2'	118.68 (17)
C6—C1—C1'	120.57 (17)	C6'—C1'—C1	119.16 (17)
C2—C1—C1'	120.38 (16)	C2'—C1'—C1	122.16 (17)

O1—C2—C3	123.39 (18)	O1'—C2'—C3'	123.77 (18)
O1—C2—C1	116.12 (16)	O1'—C2'—C1'	116.10 (16)
C3—C2—C1	120.48 (17)	C3'—C2'—C1'	120.09 (18)
C4—C3—C2	119.96 (19)	C2'—C3'—C4'	120.81 (18)
C4—C3—H3	120.0	C2'—C3'—H3'	119.6
C2—C3—H3	120.0	C4'—C3'—H3'	119.6
C5—C4—C3	119.19 (18)	C5'—C4'—C3'	118.72 (18)
C5—C4—H4	120.4	C5'—C4'—H4'	120.6
C3—C4—H4	120.4	C3'—C4'—H4'	120.6
C4—C5—C6	121.58 (18)	C4'—C5'—C6'	121.41 (18)
C4—C5—C11	119.00 (15)	C4'—C5'—C11'	119.85 (15)
C6—C5—C11	119.41 (16)	C6'—C5'—C11'	118.74 (15)
C5—C6—C1	119.72 (18)	C5'—C6'—C1'	120.26 (18)
C5—C6—H6	120.1	C5'—C6'—H6'	119.9
C1—C6—H6	120.1	C1'—C6'—H6'	119.9
O1—C7—H7A	109.5	O1'—C7'—H7'1	109.5
O1—C7—H7B	109.5	O1'—C7'—H7'2	109.5
H7A—C7—H7B	109.5	H7'1—C7'—H7'2	109.5
O1—C7—H7C	109.5	O1'—C7'—H7'3	109.5
H7A—C7—H7C	109.5	H7'1—C7'—H7'3	109.5
H7B—C7—H7C	109.5	H7'2—C7'—H7'3	109.5
C7—O1—C2—C3	-17.7 (3)	C6—C1—C1'—C2'	-119.2 (2)
C7—O1—C2—C1	161.82 (17)	C2—C1—C1'—C2'	65.4 (3)
C6—C1—C2—O1	-175.20 (17)	C7'—O1'—C2'—C3'	4.5 (3)
C1'—C1—C2—O1	0.3 (3)	C7'—O1'—C2'—C1'	-177.96 (17)
C6—C1—C2—C3	4.3 (3)	C6'—C1'—C2'—O1'	-175.93 (17)
C1'—C1—C2—C3	179.85 (18)	C1—C1'—C2'—O1'	3.6 (3)
O1—C2—C3—C4	176.63 (18)	C6'—C1'—C2'—C3'	1.7 (3)
C1—C2—C3—C4	-2.9 (3)	C1—C1'—C2'—C3'	-178.75 (18)
C2—C3—C4—C5	-0.9 (3)	O1'—C2'—C3'—C4'	176.60 (18)
C3—C4—C5—C6	3.2 (3)	C1'—C2'—C3'—C4'	-0.9 (3)
C3—C4—C5—C11	-176.05 (15)	C2'—C3'—C4'—C5'	-0.7 (3)
C4—C5—C6—C1	-1.7 (3)	C3'—C4'—C5'—C6'	1.5 (3)
C11—C5—C6—C1	177.55 (15)	C3'—C4'—C5'—C11'	-179.32 (15)
C2—C1—C6—C5	-2.1 (3)	C4'—C5'—C6'—C1'	-0.6 (3)
C1'—C1—C6—C5	-177.59 (18)	C11'—C5'—C6'—C1'	-179.82 (15)
C6—C1—C1'—C6'	60.3 (3)	C2'—C1'—C6'—C5'	-1.0 (3)
C2—C1—C1'—C6'	-115.1 (2)	C1—C1'—C6'—C5'	179.49 (17)