3-3-1998

Cloning and Developmental Expression of Pea Ribulose-1,5-Bisphosphate Carboxylase/Oxygenase Large Subunit N-Methyltransferase

Robert L. Houtz
University of Kentucky, rhoutz@uky.edu

Follow this and additional works at: https://uknowledge.uky.edu/horticulture_patents

Part of the [Horticulture Commons](#)

[Right click to open a feedback form in a new tab to let us know how this document benefits you.](#)

Recommended Citation

Houtz, Robert L., "Cloning and Developmental Expression of Pea Ribulose-1,5-Bisphosphate Carboxylase/Oxygenase Large Subunit N-Methyltransferase" (1998). *Horticulture Faculty Patents*. 9.
https://uknowledge.uky.edu/horticulture_patents/9

This Patent is brought to you for free and open access by the Horticulture at UKnowledge. It has been accepted for inclusion in Horticulture Faculty Patents by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

US005723752A

United States Patent [19]**Houtz**[11] **Patent Number:** **5,723,752**[45] **Date of Patent:** **Mar. 3, 1998**

[54] **CLONING AND DEVELOPMENTAL
EXPRESSION OF PEA RIBULOSE-1,5-
BISPHOSPHATE CARBOXYLASE/
OXYGENASE LARGE SUBUNIT N-
METHYLTRANSFERASE**

[75] **Inventor:** **Robert L. Houtz**, Lexington, Ky.

[73] **Assignee:** **University Of Kentucky**, Lexington,
Ky.

[21] **Appl. No.:** **391,000**

[22] **Filed:** **Feb. 21, 1995**

[51] **Int. Cl.⁶** **A01H 5/00**; C12N 15/29;
C12N 15/54; C12N 15/82

[52] **U.S. Cl.** **800/205**; 800/DIG. 18;
800/DIG. 19; 800/DIG. 23; 800/DIG. 26;
800/DIG. 40; 800/DIG. 41; 800/DIG. 42;
800/DIG. 43; 800/DIG. 44; 536/23.2; 536/23.6;
435/69.1; 435/70.1; 435/172.3; 435/193;
435/320.1

[58] **Field of Search** 536/23.2, 23.6;
435/69.1, 70.1, 172.3, 320.1, 193; 800/205,
DIG. 18, 19, 23, 26, 40-44

[56] **References Cited**

PUBLICATIONS

Black et al., "Light-Regulated Methylation of Chloroplast Proteins". The Journal of Biological Chemistry, vol. 262, No. 20, Jul. 15, 1987, pp. 9803-9807.

Houtz et al., "Affinity Purification of Ribulose-1-5-Bisphosphate Carboxylase/Oxygenase Large Subunit ϵ N-Methyltransferase". Supplement to Plant Physiology, Aug. 1992, Abstract No. 343, vol. 99 Suppl. 1.
Houtz et al., "Identification and Specificity Studies of Ribulose-1,5-Bisphosphate Carboxylase/Oxygenase Large Subunit ϵ N-Methyltransferase", Supplement to Plant Physiology, Aug. 1992, Abstract No. 344, vol. 99 Suppl. 1.

Houtz et al., "Partial Amino Acid Sequence of Ribulose-1,5-Bisphosphate Carboxylase/Oxygenase Large Subunit ϵ N-Methyl Transferase", Supplement to Plant Physiology, vol. 102, No. 1, May 1993, Abstract No. 248.

Houtz et al., "Partial Purification and Characterization of Ribulose-1,5-bisphosphate Carboxylase/Oxygenase Large Subunit ϵ N-Methyltransferase", Plant Physiol. vol. 97 No. 3, 1991, pp. 913-920

Klein et al., "Cloning and Expression of the Rubisco Large Subunit Methyl-Transferase Gene from Pea". Supplement to Plant Physiology, vol. 105, May 1994, Abstract No. 438.

Niemi et al., "Protein Methylation in Pea Chloroplast", Plant Physiol., vol. 93, 1990, pp. 1235-1240.

Houtz et al., "Posttranslational Modifications in the Amino-Terminal Region of the Large Subunit of Ribulose-1,5-Bisphosphate Carboxylase/Oxygenase from Several Plant Species," *Plant Physiol.*, 98:1170-74 (1992).

Mulligan et al., "Reaction-Intermediate Analogue Binding by Ribulose Bisphosphate Carboxylase/Oxygenase Causes Specific Changes in Proteolytic Sensitivity: The Amino-Terminal Residue of the Large Subunit is Acetylated Proline," *Proc. Natl. Acad. Sci. USA*, 85:1513-17 (Mar. 1988).

Houtz et al., "Post-Translational Modifications in the Large Subunit of Ribulose Bisphosphate Carboxylase/Oxygenase," *Proc. Natl. Acad. Sci. USA*, 86:1855-1859 (Mar. 1989).

Napoli et al. 1990. *Plant Cell* 2:279-289.

Smith et al. 1988. *Nature* 334:724-726.

Klein et al. 1993. *Planta* 190(4): 498-510.

Eckes et al. 1986. *Mol. Gen. Genet.* 205:14-22.

Primary Examiner—David T. Fox

Attorney, Agent, or Firm—Burns, Doane, Swecker & Mathis, L.L.P.

[57] **ABSTRACT**

The gene sequence for ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco) large subunit (LS) ϵ N-methyltransferase (protein methylase III or Rubisco LSMT) is disclosed. This enzyme catalyzes methylation of the ϵ -amine of lysine-14 in the large subunit of Rubisco. In addition, a full-length cDNA clone for Rubisco LSMT is disclosed. Transgenic plants and methods of producing same which (1) have the Rubisco LSMT gene inserted into the DNA, and (2) have the Rubisco LSMT gene product or the action of the gene product deleted from the DNA are also provided. Further, methods of using the gene to selectively deliver desired agents to a plant are also disclosed.

15 Claims, 4 Drawing Sheets

FIGURE 2B

N	K	R	L	F	P	D	P	V	T	L	D	D	F	F	W	A	F	G	I	L	R	S	721
AAT	AAG	CGG	CTT	TTT	CCG	GAT	CCT	GTG	ACG	CTG	GAT	GAC	TTC	TTT	TGG	GCA	TTT	GGA	ATT	CTC	AGA	TCA	
R	A	F	S	R	L	R	N	E	N	L	V	V	V	P	M	A	D	L	I	N	H	S	790
AGG	GCG	TTT	TCT	CGC	CTT	CGC	AAT	GAA	AAT	CTG	GTT	GTG	GTT	CCA	ATG	GCA	GAC	TTG	ATT	AAC	CAC	AGT	
A	G	V	T	T	E	D	H	A	Y	E	V	K	G	A	A	G	L	F	S	W	D	Y	859
GCA	GGA	GTT	ACT	ACA	GAG	GAT	CAT	GCT	TAT	GAA	GTT	AAA	GGA	GCA	GCT	GGC	CTT	TTC	TCT	TGG	GAT	TAC	
L	F	S	L	K	S	P	L	S	V	K	A	G	E	Q	V	Y	I	Q	Y	D	L	N	928
CTA	TTT	TCC	TTA	AAG	AGC	CCC	CTT	TCC	GTC	AAG	GCC	GGA	GAA	CAG	GTA	TAT	ATA	CAA	TAT	GAT	TTG	AAC	
K	S	N	A	E	L	A	L	D	Y	G	F	I	E	P	N	E	N	R	H	A	Y	T	997
AAA	AGC	AAT	GCA	GAG	TTG	GCT	CTA	GAC	TAC	GGT	TTT	ATT	GAA	CCA	AAT	GAA	AAT	CGA	CAT	GCA	TAC	ACT	
L	T	L	E	I	S	E	S	D	P	F	F	D	D	K	L	D	V	A	E	S	N	G	1066
CTG	ACG	CTG	GAG	ATA	TCT	GAG	TCG	GAC	CCT	TTT	TTT	GAT	GAC	AAA	CTA	GAC	GTT	GCT	GAG	TCC	AAT	GGT	
F	A	Q	T	A	Y	F	D	I	F	Y	N	R	T	L	P	P	G	L	L	P	Y	L	1135
TTT	GCT	CAG	ACA	GCG	TAC	TTT	GAC	ATC	TTC	TAT	AAT	CGC	ACT	CTT	CCA	CCT	GGA	TTG	CTT	CCA	TAT	CTG	
R	L	V	A	L	G	G	T	D	A	F	L	L	E	S	L	F	R	D	T	I	W	G	1204
AGA	CTT	GTA	GCG	CTA	GGG	GGT	ACC	GAC	GCT	TTC	TTA	TTG	GAA	TCA	CTG	TTC	AGA	GAC	ACC	ATA	TGG	GGT	
H	L	E	L	S	V	S	R	D	N	E	E	L	L	C	K	A	V	R	E	A	C	K	1273
CAT	CTT	GAG	TTG	TCT	GTC	AGC	CGT	GAC	AAT	GAG	GAG	CTA	CTA	TGC	AAA	GCC	GTT	CGA	GAA	GCC	TGC	AAA	

CLONING AND DEVELOPMENTAL EXPRESSION OF PEA RIBULOSE-1,5- BISPHOSPHATE CARBOXYLASE/ OXYGENASE LARGE SUBUNIT N- METHYLTRANSFERASE

IDENTIFICATION OF FEDERAL FUNDING

The present invention was supported by U.S. Department of Energy Grant DE-FG05-92ER20075.

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention relates to ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco) large subunit (LS) ϵ N-methyltransferase (protein methylase III or Rubisco LSMT). This enzyme catalyzes methylation of the ϵ -amine of lysine-14 in the large subunit of Rubisco. In addition, the present invention relates to a gene and a full-length cDNA clone for Rubisco LSMT, which was isolated utilizing polymerase chain reaction-based technology and conventional bacteriophage library screening. The present invention further relates to transgenic plants and methods of producing same which (1) have the Rubisco LSMT gene inserted into the DNA, and (2) have the Rubisco LSMT gene product deleted. Methods of using the gene to selectively deliver desired agents to a plant are also disclosed.

2. Description of the Related Art

Protein methylation is a widespread and common post-translational modification catalyzed by several different protein methyltransferases (Paik et al., "Protein methylation," in Freedman et al (eds), *The Enzymology of Posttranslational Modifications of Proteins*, vol. 2, pp. 187-228, Academic Press, London (1985)). Proteins which contain trimethyllysyl residues include cytochrome c (Cessay et al., "The relationship between the trimethylation of lysine 77 and cytochrome c metabolism in *Saccharomyces cerevisiae*," *Int. J. Biochem.* 26(5):721-734 (1994); Cessay et al., "Further investigations regarding the role of trimethyllysine for cytochrome c uptake into mitochondria," *Int. J. Biochem.* 23(7.8):761-768 (1991); DiMaria et al., "Cytochrome c specific methylase from wheat germ," *Biochemistry* 21:1036-1044 (1982); Farooqui et al., "Effect of Methylation on the Stability of Cytochrome c of *Saccharomyces cerevisiae* in vivo," *J. Biol. Chem.* 256(10):5041-5045 (1981); and Farooqui et al., "In vivo studies on yeast cytochrome c methylation in relation to protein synthesis," *J. Biol. Chem.* 255(10):4468-4473 (1980)), calmodulin (Han et al., "Isolation and kinetic characterization of the calmodulin methyltransferase from sheep brain," *Biochemistry* 32:13974-13980 (1993); and Rowe et al., "Calmodulin N-methyltransferase," *J. Biol. Chem.* 261(15):7060-7069 (1986)), histone-H1 (Sarnow et al., "A histone H4-specific methyltransferase properties, specificity and effects on nucleosomal histones," *Biochim. Biophys. Acta* 655:349-358 (1981); and Tuck et al., "Two histone H1-specific protein-lysine N-methyltransferases from *Euglena gracilis*," *J. Biol. Chem.* 260(11):7114-7121 (1985)), and ribosomal proteins (Chang et al., "Purification and properties of a ribosomal protein methylase from *Escherichia coli* Q13," *Biochemistry* 14(22):4994-4998 (1975); Lobet et al., "Partial purification and characterization of the specific protein-lysine N-methyltransferase of YL32, a yeast ribosomal protein," *Biochim. Biophys. Acta* 997:224-231 (1989)). However, the biological function of post-translational protein methylation in all but a few systems remains obscure. Trimethyllysine can serve as a meta-

bolic precursor to carnitine (Paik et al., "Carnitine biosynthesis via protein methylation," *TIBS* 2:159-162 (1977)), while carboxyl methylation of bacterial membrane proteins plays a major role in chemotaxis (Clarke, "Protein carboxyl methyltransferases: Two distinct classes of enzymes," *Ann. Rev. Biochem.* 54:479-506 (1985)). Evidence suggests that methylation of Lys-115 in calmodulin affects certain activities including in vitro NAD kinase activation (Roberts et al., "Trimethyllysine and protein function," *J. Biol. Chem.* 261(4):1491-1494 (1986)), and in vivo susceptibility to ubiquitination (Gregori et al., "Bacterially synthesized vertebrate calmodulin is a specific substrate for ubiquitination," *J. Biol. Chem.* 262(6):2562-2567 (1987); and Gregori et al., "Specific recognition of calmodulin from *Dictyostelium discoideum* by the ATP ubiquitin-dependent degradative pathway," *J. Biol. Chem.* 260(9):5232-5235 (1985); but see also Ziegenhagen et al., "Multiple ubiquitination of calmodulin results in one polyubiquitin chain linked to calmodulin," *FEBS Lett.* 271(1.2):71-75 (1990); and Ziegenhagen et al., "Plant and fungus calmodulins are polyubiquitinated at a single site in a Ca^{2+} -dependent manner," *FEBS Lett.* 273(1.2):253-256 (1990)). Conflicting reports (Farooqui et al., "Effect of Methylation on the Stability of Cytochrome c of *Saccharomyces cerevisiae* in vivo," *J. Biol. Chem.* 256(10):5041-5045 (1981); Frost et al., "Cytochrome c methylation," *Protein methylation*, Ch. 4, pp. 59-76 (1990); and Frost et al., "Effect of enzymatic methylation of cytochrome c on its function and synthesis," *Int. J. Biochem.* 22(10):1069-1074 (1990); versus Cessay et al., "The relationship between the trimethylation of lysine 77 and cytochrome c metabolism in *Saccharomyces cerevisiae*," *Int. J. Biochem.* 26(5):721-734 (1994); Cessay et al., "Further investigations regarding the role of trimethyllysine for cytochrome c uptake into mitochondria," *Int. J. Biochem.* 23(7.8):761-768 (1991)) also implicate methylation of Lys-77 in cytochrome c as having a role in protein stability, heme incorporation, and mitochondrial transport. A major limitation to elucidating the biological role of lysine methylation in eukaryotes has been the absence of a protein methylase III gene. Hence, molecular studies of the physiological and biochemical function performed by methylation of protein bound lysyl residues have been restricted to site-directed mutational analysis of the methylation site in the target protein (Cessay et al., "The relationship between the trimethylation of lysine 77 and cytochrome c metabolism in *Saccharomyces cerevisiae*," *Int. J. Biochem.* 26(5):721-734 (1994); Cessay et al., "Further investigations regarding the role of trimethyllysine for cytochrome c uptake into mitochondria," *Int. J. Biochem.* 23(7.8):761-768 (1991); and Roberts et al., "Expression of a calmodulin methylation mutant affects the growth and development of transgenic tobacco plants," *Proc. Nat. Acad. Sci. USA* 89:8394-8398 (1992)). These studies have been inconclusive as to the exact biological role of methylation of the ϵ -amine of protein bound lysyl residues.

Ribulose-1,5-bisphosphate carboxylase-oxygenase (Rubisco) catalyzes the reduction of atmospheric CO_2 during photosynthesis. A great deal is known about the quaternary structure, catalytic mechanism, active site residues, in vivo regulatory mechanisms, and gene expression for this abundant enzyme, see, for example, Andrews et al., "Rubisco: Structure, Mechanisms, and Prospects for Improvement," in Hatch et al (eds), *The Biochemistry of Plants*, vol. 10, pp. 131-218, Academic Press, New York (1987); Dean et al., "Structure, evolution, and regulation of rbcS genes in higher plants," *Annu. Rev. Plant. Physiol. Plant Mol. Biol.* 40:415-439 (1989); and Mullet, "Chloro-

plast development and gene expression." *Annu. Rev. Plant. Physiol. Plant Mol. Biol.* 39:475-502 (1988). Higher plant Rubisco is a hexadecameric protein composed of eight chloroplast-encoded large subunits (referred to herein as "LS") and eight nuclear-encoded small subunits (referred to herein as "SS"). Synthesis of the LS is accompanied by post-translational processing of the N-terminal domain (Houtz et al. "Post-translational modifications in the large subunit of ribulose biphosphate carboxylase/oxygenase." *Proc. Natl. Acad. Sci. USA* 86:1855-1859 (1989); and Mulligan et al. "Reaction-intermediate analogue binding by ribulose biphosphate carboxylase/oxygenase causes specific changes in proteolytic sensitivity: The amino-terminal residue of the large subunit is acetylated proline." *Proc. Natl. Acad. Sci. USA* 85:1513-1517 (1988)). The N-terminal Met-1 and Ser-2 are removed and Pro-3 acetylate. Additionally, the LS of Rubisco from tobacco, muskmelon, pea, and several other species is post-translationally modified by trimethylation of the ϵ -amine of Lys-14 (Houtz et al. "Posttranslational modifications in the amino-terminal region of the large subunit of ribulose-1,5-bisphosphate carboxylase/oxygenase from several plant species." *Plant Physiol.* 98:1170-1174 (1992); Houtz et al. "Post-translational modifications in the large subunit of ribulose biphosphate carboxylase/oxygenase." *Proc. Natl. Acad. Sci. USA* 86:1855-1859 (1989)). The enzyme responsible for this later modification is a highly specific chloroplast-localized S-adenosylmethionine (AdoMet):protein (lys) eN-methyltransferase (protein methylase III, Rubisco LSMT, EC 2.1.1.43). Recently, Rubisco LSMT was affinity purified ~8000-fold from pea chloroplasts and identified as a monomeric protein with a molecular mass of ~57 kDa (Wang et al. "Affinity Purification of Ribulose-1,5-bisphosphate Carboxylase/Oxygenase Large Subunit eN-Methyltransferase." accepted by *Protein Expression and Purification* (1995)).

OBJECTS AND SUMMARY OF THE INVENTION

In view of the state of the art as previously described, there thus exists a need in the art for a better understanding of the biological function of post-translational protein methylation in higher plant systems. More specifically, a better understanding of the biological role of methylation of the ϵ -amine of protein bound lysyl residues.

It is thus an object of the present invention to provide a Rubisco LSMT gene.

It is a further object of the present invention to provide the DNA and amino acid sequence for a Rubisco LSMT enzyme.

It is a still further object of the present invention to provide a full-length cDNA clone for Rubisco LSMT.

Another object of the present invention is to determine and selectively manipulate the biological role of lysine methylation in eukaryotes.

In a first aspect, the present invention relates to a Rubisco LSMT gene which is expressed in a higher plant and which encodes Rubisco LSMT. Rubisco LSMT catalyzes methylation of the ϵ -amine of lysine-14 in the LS of Rubisco. A particularly preferred higher plant includes the pea.

In a second aspect, the present invention relates to the DNA and amino acid sequence for a Rubisco LSMT enzyme.

In a third aspect, the present invention relates to a recombinant vector including the Rubisco LSMT gene described above. The vector is suitable for transforming higher plant seed crops.

In a fourth aspect, the present invention relates to an isolated or recombinantly expressed Rubisco LSMT enzyme encoded by the Rubisco LSMT gene described above.

In a fifth aspect, the present invention relates to a method for introducing the Rubisco LSMT gene into a plant which does not possess said gene, which method comprises transforming a higher plant seed crop with the Rubisco LSMT gene vector described above such that the plant expresses the Rubisco LSMT enzyme encoded by the gene.

In a sixth aspect, the present invention relates to a method for selectively eliminating a plant which comprises the Rubisco LSMT gene by deleting the gene product, or eliminating the action of the gene product, from the plant. Without the Rubisco LSMT gene product or the action of the gene product, the plant would be unable to catalyze net CO₂ fixation during photosynthesis and would thus die.

In a seventh aspect, the present invention relates to a method for introducing agents to a plant cell which agents will selectively increase or decrease activity of Rubisco.

In a further aspect, the present invention relates to a recombinant or transgenic plant transformed with the Rubisco LSMT gene described above.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A-1B show a reverse phase-HPLC of peptic polypeptides from Rubisco LSMT. FIG. 1a shows the control of peptic digestion of Immobilon-CD membrane without Rubisco LSMT. FIG. 1b shows the peptic digestion of affinity-purified Rubisco LSMT (~30 μ g) electroblotted to Immobilon-CD membrane as described in the Examples. The asterisks identify peaks with A₂₁₄ absorbance which were collected and submitted for amino acid sequence analyses.

FIGS. 2A, 2B and 2C (SEQ. ID NO. 41) illustrate the nucleotide and predicted amino acid sequence of pea rbcMT cDNA. Nucleotide position is marked on the right. The start and stop codons are underlined and segments corresponding to peptic fragments are marked by lines above the amino acid sequence. The position of amino acids encoded by the PCR-derived partial cDNA is blocked.

FIG. 3 shows a Southern blot analysis of the rbcMT gene in pea. Ten μ g of genomic DNA from pea was digested with EcoR I, Hind III, or Dra I, and electrophoresed on an 0.8% agarose gel. The blot was probed with a 1775 bp rbcMT cDNA of pea. Approximate sizes in kbp are indicated to the left. Blots were exposed to x-ray film for 48 hours.

FIG. 4 illustrates organ-specific accumulation of rbcMT mRNA. Messenger-RNA was isolated from roots, stems, and leaves of 10 day old chamber-grown pea. Northern blots were loaded on an equal RNA basis and were probed with radiolabeled antisense RNA to rbcS, rbcL or rbcMT. Northern blots of rbcS, rbcL and rbcMT mRNA were exposed to x-ray film for 2 hours, 1 hour, and 36 hours, respectively.

In FIG. 5, light-dependent accumulation of rbcMT mRNA in etiolated pea is shown. Peas were germinated in a dark chamber in a light-tight room. After 8 days, etiolated seedlings were either harvested (treatment 1) or transferred to the light for 24 hours (treatment 2) or 72 hours (treatment 3). Control seedlings were germinated in the light and harvested after 8 days (treatment 4). RNA was isolated from leaf tissue from each treatment and Northern analyses were conducted. Northern blots of rbcS, rbcL, and rbcMT were exposed to x-ray film for 1 hour, 1 hour, and 36 hours, respectively.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to a Rubisco LSMT gene, its DNA and amino acid sequence encoding therefor, and a cDNA clone thereof.

In the present application, naturally occurring amino acid residues in peptides are abbreviated as recommended by the IUPAC OIUB Biochemical Nomenclature Commission as follows: Phenylalanine is Phe or F; Leucine is Leu or L; Isoleucine is Ile or I; Methionine is Met or M; Norleucine is Nle; Valine is Val or V; Serine is Ser or S; Proline is Pro or P; Threonine is Thr or T; Alanine is Ala or A; Tyrosine is Tyr or Y; Histidine is His or H; Glutamine is Gln or Q; Asparagine is Asn or N; Lysine is Lys or K; Aspartic Acid is Asp or D; Glutamic Acid is Glu or E; Cysteine is Cys or C; Tryptophan is Trp or W; Arginine is Arg or R; Glycine is Gly or G, and X is any amino acid.

Synthetic or non-naturally occurring amino acids refer to amino acids which do not naturally occur in vivo but which, nevertheless, can be incorporated into the peptide structures described herein. Preferred synthetic amino acids are the D-amino acids of naturally occurring L-amino acids as well as non-naturally occurring D and L amino acids represented by the formula $H_2NCH(R^1)COOH$, wherein R^1 is: (1) a lower alkyl group; (2) a cycloalkyl group of from 3 to 7 carbon atoms; (3) a heterocycle of from 3 to 7 carbon atoms and 1 to 2 heteroatoms selected from the group consisting of oxygen, sulfur, and nitrogen; (4) an aromatic or arylalkyl residue of from 6 to 15 carbon atoms optionally having from 1 to 3 substituents on the aromatic nucleus selected from the group consisting of hydroxyl, lower alkoxy, amino, and carboxyl; (5) alkylene-Y where alkylene is an alkylene group of from 1 to 7 carbon atoms and Y is selected from the group consisting of hydroxy, amino, cycloalkyl of from 3 to 7 carbon atoms, heterocyclic of from 3 to 7 carbon atoms and 1 to 2 heteroatoms selected from the group consisting of oxygen, sulfur and nitrogen, and $-C(O)R^2$ where R^2 is selected from the group consisting of hydrogen, lower alkyl, lower alkoxy, and $-NR^3R^4$ where R^3 and R^4 are independently selected from the group consisting of hydrogen and lower alkyl; (6) alkylene-S(O) $_nR^5$ where n is 1 or 2, and R^5 is a lower alkyl or lower alkylene.

Particularly preferred synthetic amino acids include, by way of example, the D-amino acids of naturally occurring L-amino acids, L-1-naphthylalanine, L-2-naphthylalanine, L-cyclohexylalanine, L-2-amino isobutyric acid, the sulfoxide and sulfone derivatives of methionine, and the lower alkoxy derivatives of methionine.

"Peptide mimetics" are also encompassed by the present invention and include peptides having one or more of the following modifications:

peptides wherein one or more of the peptidyl [$-C(O)NH-$] linkages (bonds) have been replaced by a non-peptidyl linkage such as carbamate linkage [$-OC(O)N<$], phosphonate linkage, amidate linkage, sulfonamide linkage, and secondary amine linkage or with an alkylated peptidyl linkage [$C(O)NR^6-$ where R^6 is a lower alkyl],

peptides wherein the N-terminus is derivatized to a $-NR^7R^8$ group, to a $-NC(O)R^7$ group where R^7 and R^8 are independently selected from hydrogen and lower alkyls with the proviso that R^7 and R^8 are both not hydrogen, to a succinimide group, to a benzyloxycarbonyl-NH-(CBZ-NH-) group, to a benzyloxycarbonyl-NH- group having from 1 to 3 substituents on the phenyl ring selected from the group consisting of lower alkyl, lower alkoxy, chloro, and bromo,

peptides wherein the C terminus is derivatized to $>C(O)R^9$ where R^9 is selected from the group consisting of hydrogen, lower alkyl, lower alkoxy, and $NR^{10}R^{11}$

where R and R^{11} are independently selected from the group consisting of hydrogen and lower alkyl.

Although the present invention is described with respect to peas, it will be appreciated that the techniques employed herein are applicable to other photosynthesizing plants, e.g., legumes, soybeans, *solanaceae* (tomato, potato, tobacco, pepper) and *cucurbitaceae* (cucumbers, melons, gourds). The protein methylase III of other photosynthesizing plants would be expected to exhibit homologous amino acid sequences to those described herein. As described herein, certain aspects of the present invention are applicable to plants not having the Rubisco LSMT gene, e.g., spinach, wheat, corn, lower plants such as algae, monocots (cereals) and the like.

Ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco) large subunit (LS) ϵ N-methyltransferase (referred to herein as "Rubisco LSMT") catalyzes methylation of the ϵ -amine of lysine-14 in the LS of Rubisco. Rubisco is the world's most abundant protein, and serves as the only significant link between the inorganic and organic carbon pools in the Earth's biosphere by catalyzing the reduction of atmospheric carbon dioxide to carbohydrates during photosynthesis. Perturbations of Rubisco activity translate directly into similar changes in plant growth and yield. Thus, there is significant interest in the art in the potential manipulation and control of Rubisco activity through genetic engineering.

However, the complexity and multimeric nature of Rubisco have proven to be substantial obstacles to achieving this goal, which have not yet been overcome. Rubisco LSMT provides an opportunity for the selective manipulation of Rubisco activity through changes in the structure and stability of the N-terminal region in the LS, an area known to be essential for catalytic activity. Rubisco LSMT is a highly specific enzyme which is found to interact only with Rubisco and does not interact with any other protein in the plant cell. Since Rubisco catalyzes the reduction of atmospheric CO_2 during photosynthesis, Rubisco and Rubisco LSMT are critical to the plant cell for viability. Furthermore, the exceptionally tight and specific nature of the interaction between Rubisco LSMT and des(methyl) forms of Rubisco creates the possibility for the development of novel synthetic polypeptide herbicides, whose target is the in vivo interaction between Rubisco LSMT and Rubisco, whose specificity crosses a group of plant species related only by the presence of Rubisco LSMT, and whose target protein has no homologue in the entire animal kingdom. Finally, this same affinity of Rubisco LSMT for des(methyl) forms of Rubisco also creates the possibility for the site and protein specific delivery of compounds into the chloroplast and to Rubisco, for the potential manipulation of Rubisco activity and/or stability.

With limited internal amino acid sequence information obtained from high performance liquid chromatography (HPLC)-purified peptic polypeptides from Rubisco LSMT, a full-length cDNA clone was isolated by the present inventor utilizing polymerase chain reaction (PCR)-based technology and conventional bacteriophage library screening. PCR techniques are disclosed, for example, in Klein et al. "Cloning and Developmental Expression of the Sucrose-Phosphate-Synthase Gene From Spinach," *Planta* 190:498-510 (1993); in Ampli-Taq PCR kit by Perkin Elmer-Cetus, Emeryville, Calif.; and in the manufacturer's instruction manual. Bacteriophage library screening is described, for example, in Gant et al. "Transfer of rp122 to the Nucleus Greatly Preceded its loss from the Chloroplast and Involved the Gain of an Intron," *EMBO J.* 10:3073-3078 (1991), and in the information provided by the manufacturer of the screening membrane (Stratagene, La Jolla, Calif.).

The 1802-base-pair cDNA of Rubisco LSMT encodes a 489-amino acid polypeptide with a predicted molecular mass of ~55 kDa. To the knowledge of the present inventor, this is the first reported DNA and amino acid sequence for a protein methylase III enzyme. A derived N-terminal amino acid sequence of the polypeptide with features common to chloroplast transit peptides was identified. The deduced sequence of Rubisco LSMT did not exhibit regions of significant homology with other protein methyltransferases known in the art, e.g., D-aspartyl/L-isoaspartyl protein methyltransferase (Kagan et al., "Widespread occurrence of three sequence motifs in diverse S-adenosylmethionine-dependent methyltransferases suggests a common structure for these enzymes," *Arch. Biochem. Biophys.* 310(2):417-427 (1994)). Widespread occurrence of three sequence motifs in diverse S-adenosylmethionine dependent methyltransferases suggest a common structure for these enzymes. Southern blot analysis of pea genomic DNA indicated a low gene copy number of Rubisco LSMT in pea. A "low gene copy number" indicates that Rubisco LSMT may be encoded by a single gene. Northern analysis revealed a single mRNA species of about 1.8 kb encoding for Rubisco LSMT which was predominately localized in leaf tissue. Illumination of etiolated pea seedlings showed that the accumulation of Rubisco LSMT mRNA is light-dependent. Maximum accumulation of Rubisco LSMT transcripts occurred during the initial phase of light-induced leaf development which preceded the maximum accumulation of rbcS and rbcL mRNA. Transcript levels of Rubisco LSMT in mature light-grown tissue were similar to transcript levels in etiolated tissues indicating that the light-dependent accumulation of Rubisco LSMT mRNA is transient.

A cDNA of the Rubisco LSMT gene from pea was isolated and studies of Rubisco LSMT gene expression initiated. Utilizing amino acid sequence information derived from purified peptic polypeptide fragments from proteolyzed Rubisco LSMT, a full-length cDNA of Rubisco LSMT was obtained. The cDNA of Rubisco LSMT, rbcMT, was used to examine organ-specific and developmental parameters affecting rbcMT gene expression. The expression of two well characterized gene families, rbcS (SS of Rubisco) and rbcL (LS of Rubisco), were also examined to determine if rbcMT expression is coregulated with that of the Rubisco subunit genes, particularly the LS.

The present specification details the purification of peptic fragments from pea Rubisco LSMT and a PCR-based cloning strategy for isolating a full-length cDNA. A similar strategy was previously utilized to obtain a full-length cDNA of sucrose-phosphate synthase from spinach (Klein et al., "Cloning and developmental expression of the sucrose-phosphate-synthase gene from spinach," *Planta*, 190:498-510 (1993)). The low abundance of Rubisco LSMT in pea leaves (~0.01%) prompted the use of PCR, since it would be more difficult to obtain enough protein to ensure the production of an antibody with high-titer and specificity with which to screen a library. Further, the protein sequence information obtained from peptic fragments permitted the confirmation of clones encoding for Rubisco LSMT. Hence, a molecular probe of the pea rbcMT gene was rapidly obtained thereby permitting identification of protein and nucleotide sequence, and characterization of rbcMT gene expression.

To date, the deduced amino-acid sequence of Rubisco LSMT represents the first reported example of a protein eN-methyltransferase. Thus, it is now possible to extend the comparison of known enzyme sequences to include this class of methyltransferases. Interestingly, the deduced

amino acid sequence of Rubisco LSMT does not possess any of the three sequence motifs proposed by Kagan and Clarke (Kagan et al., "Widespread occurrence of three sequence motifs in diverse S-adenosylmethionine-dependent methyltransferases suggests a common structure for these enzymes," *Arch. Biochem. Biophys.* 310(2):417-427 (1994)) for methyltransferases. However, knowledge of methyltransferase sequences is still fragmentary and no sequences are yet available for protein arginine, histidine, or N-terminal amino methyltransferases. As noted by Kagan and Clarke, methyltransferases whose sequences are available represent less than one-third of these enzymes and a number of other methyltransferases apparently do not possess the proposed motifs or any additional elements of sequence similarity. Furthermore, several lines of evidence suggest that Rubisco LSMT exclusively methylates the large subunit of Rubisco (Houtz et al., "Posttranslational modifications in the amino-terminal region of the large subunit of ribulose-1,5-bisphosphate carboxylase/oxygenase from several plant species," *Plant Physiol.* 98:1170-1174 (1992); and Houtz et al., "Partial purification and characterization of ribulose-1,5-bisphosphate carboxylase/oxygenase large subunit eN-methyltransferase," *Plant Physiol.* 97:913-920 (1991)). This high level of specificity may in part explain the lack of overall homology with other methyltransferases. Hence, sequence determination of other yet-to-be-discovered protein(lys)eN-methyltransferases may be necessary to identify conserved, functionally essential regions in this class of enzyme.

Several lines of evidence indicate that there is a low copy number of the rbcMT gene in pea. Genomic Southern blot analysis revealed simple hybridization patterns. DNA sequence information of several cDNA clones revealed an invariant nucleotide sequence in the coding and noncoding regions. Although these observations do not preclude the existence of multiple structural genes encoding Rubisco LSMT, they are consistent with a low- or even single-copy gene hypothesis.

Many plant genes are expressed in a highly regulated manner. Gene products may be present only in certain cell types, at specific stages of development or only following the application of distinct environmental stimuli (Kuhlemeier et al., "Regulation of gene expression in higher plants," *Annu. Rev. Plant Physiol.* 38:221-257 (1987)). In addition, the expression of nuclear genes encoding plasmid proteins is often coordinated with the expression of plastid-encoded protein subunits (Rapp et al., "Chloroplast transcription is required to express the nuclear genes rbcS and cab," *Plant Mol. Biol.* 17:813-823 (1991)). The present specification shows that rbcMT gene expression is regulated in an organ-specific manner at the level of transcription or mRNA stability. The organ-specific expression of rbcMT paralleled that of rbcS and rbcL being predominately localized to photosynthetic leaf tissue. Examination of transcript levels during the light-induced development of etiolated pea leaves indicated that accumulation of mRNA encoding for rbcS, rbcL, and rbcMT is light-dependent. However, the activation of rbcMT expression preceded the maximum accumulation of mRNA encoding for either of the Rubisco subunits. Maximum transcript levels for rbcMT were obtained in the first 24 hours of illumination, which corresponded with the initial, light-dependent phase of rbcS and rbcL transcript accumulation. Interestingly, the kinetics of Rubisco activase mRNA accumulation during the greening of etiolated barley was similar to that reported here for rbcMT mRNA (Zielinski et al., "Coordinate expression of rubisco activase and rubisco during barley leaf cell development," *Plant*

Physiol. 90:516-521 (1989)). The present inventor also observed that in continuously illuminated pea leaves *rbcMT* transcript levels were equal to the levels observed in dark-grown leaves (FIG. 5), while the activity of Rubisco LSMT was nearly 3-fold higher. Since the relative amounts of *rbcMT* transcripts increased dramatically during the initial phase of light-induced development of etiolated pea leaves and then declined to a level equal to those observed in the dark, changes in the level of Rubisco LSMT protein may be controlled by the level of *rbcMT* transcripts.

Finally, while a number of ϵ N-methylated lysyl residues in several proteins have been described, no unifying hypothesis with regards to the functional significance of methylated lysyl residues has been discovered. Molecular studies have approached this topic by engineering amino acid substitutions at the position of the methylation lysyl residue in calmodulin (Roberts et al. "Expression of a calmodulin methylation mutant affects the growth and development of transgenic tobacco plants," *Proc. Nat. Acad. Sci. USA* 89:8394-8398 (1992)) and cytochrome c (Cessay et al. "The relationship between the trimethylation of lysine 77 and cytochrome c metabolism in *Saccharomyces cerevisiae*," *Int. J. Biochem.* 26(5):721-734 (1994); and Cessay et al. "Further investigations regarding the role of trimethyllysine for cytochrome c uptake into mitochondria," *Int. J. Biochem.* 23(7.8):761-768 (1991)), followed by expression of these mutant proteins in transformed tobacco plants and yeast cells, respectively. While the mutated calmodulin and cytochrome c proteins were incapable of acting as substrates for methylation, these studies were inconclusive as to a clear role for site-specific methylation of the target lysyl residues by the calmodulin or cytochrome c protein specific ϵ N-methyltransferases.

The present invention also relates to a method for introducing the Rubisco LSMT gene into a plant which does not possess said gene, such as *Arabidopsis thaliana*. The methods employed for transforming the plants are generally known in the art. For example, the transformation method described in Bechtold et al, *Planta Agrobacterium Mediated Gene Transfer By Infiltration of Adult Arabidopsis Thaliana Plants*, C.R. Acad. Sci., Paris 316:1194-1199 (1993) and Valvekens et al, "Agrobacterium tumefaciens-mediated transformation of *Arabidopsis thaliana* root explants by using kanamycin selection," *Proc. Natl. Acad. Sci. USA* 85:5536-5540 (1988), may be used in the method of the present invention. More specifically, the method contemplated herein comprises transforming a plant with the Rubisco LSMT gene vector described above such that the plant expresses the Rubisco LSMT enzyme encoded by the gene. These methods produce transgenic plants, which will have Rubisco LSMT activity and Lys-14 methylation in the LS of Rubisco.

Further, the present invention provides a method for deleting the Rubisco LSMT gene product or the action of the gene product in a photosynthesizing plant which has the Rubisco LSMT gene. Knowing the DNA sequence of the Rubisco LSMT gene, transgenic plants can be constructed expressing antisense RNA to Rubisco LSMT which results in the down-regulation of the Rubisco LSMT gene product, by methods as set forth for example, in Eguchi et al, "Antisense RNA," *Annu. Rev. Biochem.* 60:631-652 (1991). Since the Rubisco LSMT enzyme is essential for Rubisco activity, the deletion of the enzyme would be expected to be lethal to the plant since it would be unable to catalyze net CO_2 fixation during photosynthesis. This method, and variations of this method, could thus be used as a herbicide to selectively eliminate photosynthesizing plants.

Due to the high specificity of Rubisco LSMT for Rubisco, knowledge of the sequence for the Rubisco LSMT gene can be used to introduce agents to a plant cell which agents will selectively increase or decrease the activity of Rubisco. Additionally, in this regard, a recombinant vector comprising the sequence of the Rubisco LSMT gene responsible for the tight interaction of Rubisco LSMT with Rubisco could be constructed. Additional agents which enhance or reduce the activity of Rubisco, for example, CA1P (carboxyarabinitol-1-phosphate), CABP (carboxyarabinitol bisphosphate), carbamates and divalent metal cations, are then conjugated to the vector. The vector is then inserted into the plant cell by methods known in the art. The agents will then be delivered to Rubisco as a result of the high specificity and strong interaction of Rubisco LSMT and Rubisco. These agents may be synthetically derived polypeptides that are direct representations of the sequence of amino acids responsible for the interaction of Rubisco LSMT with Rubisco. These synthetic polypeptides would delete Rubisco LSMT activity and result in plant death in the aforementioned manner.

Moreover, the particular sequence disclosed herein for the pea Rubisco LSMT gene may be used to determine the particular sequence in other photosynthesizing plants. The sequence of the gene may be used as a probe to screen cDNA or genomic DNA libraries from other plants and, due to the expected homology between the gene sequences in the various plant species, the particular sequence for the Rubisco LSMT gene in other species may then be found.

In a further aspect, the present invention relates to a recombinant or transgenic plant transformed with the Rubisco LSMT gene described above.

Having now generally described this invention, the same will be better understood by reference to certain specific examples, which are included herein for purposes of illustration only and are not intended to be limiting of the invention or any embodiment thereof, unless so specified.

EXAMPLES

Example 1

Plant Growth

Controlled environment-cultured peas (*Pisum sativum*) were germinated and maintained in environmental chambers as described in Wang et al. *Protein Expression and Purification*. For developmental studies, seeds were either germinated at 23° C. in a dark chamber located in a light-tight room or were grown in an illuminated chamber with a light intensity of 300 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ (incandescent plus fluorescent). After 8 days of growth in complete darkness, pea seedlings were either harvested into liquid nitrogen or were transferred to an illuminated chamber for a predetermined period prior to harvest.

Example 2

Purification and assay of Rubisco LSMT

Rubisco LSMT was affinity purified utilizing immobilized spinach Rubisco as describe in Wang et al. cited supra. Briefly, purified spinach Rubisco (McCurry et al. "Ribulose-1,5-bisphosphate carboxylase/oxygenase from spinach, tomato or tobacco leaves," *Methods in Enzymology* 90(82):515-521 (1982)) was immobilized to PVDF membranes (Millipore Corp., Bedford, Mass. USA, 60 mg Rubisco/450 cm^2) which were then incubated for 4 h at 4° C. with pea chloroplast lysates (20 ml at 20 mg/ml protein per 450 cm^2 membrane). After incubation, the PVDF membranes were

washed with 50 mM TRIS-HCl (pH 8.2), 5 mM MgCl₂, 1 mM EDTA, 0.4M NaCl and subsequently eluted with 20 ml of 50 mM TRIS-HCl (pH 8.2), 5 mM MgCl₂, 200 μM AdoMet and 50 μg/ml β-lactoglobulin per 450 cm² membrane. The eluent was concentrated by centrifugal ultrafiltration to a final volume of ~50 μl and used as a source for purified Rubisco LSMT. The yield from a single PVDF membrane containing immobilized spinach Rubisco was typically 7–10 μg of purified Rubisco LSMT. Assays of Rubisco LSMT activity were as previously described (Houtz et al., "Partial purification and characterization of ribulose-1,5-bisphosphate carboxylase/oxygenase large subunit eN-methyltransferase," *Plant Physiol.* 97:913–920 (1991)).

Example 3

Peptide profiles and sequence analysis

To separate affinity-purified Rubisco LSMT from the β-lactoglobulin carrier protein, Rubisco LSMT was resolved by SDS-PAGE (10% acrylamide) prior to electrophoretic transfer to Immobilon-CD membranes (Millipore Corp., Bedford, Mass. USA). Conditions for the electrophoretic transfer, visualization and subsequent in vitro enzymatic cleavage of Rubisco LSMT with pepsin were as described, for example, by Paik et al., "Protein methylation," in Freedman et al (eds), *The Enzymology of Posttranslational Modifications of Proteins*, vol. 2, pp. 187–228, Academic Press, London (1985). Peptic peptides released from Rubisco LSMT were isolated by reverse phase-HPLC using conditions described in Patterson et al., "High-yield recovery of electroblotted proteins and cleavage fragments from a cationic polyvinylidene fluoride-based membrane," *Anal. Biochem.* 202:193–203 (1992), with an Aquapore RP-300 7

micron particle size octyl reverse phase column (2.1 mm×220 mm, Applied Biosystems, San Jose, Calif. USA). Peptic peptides were manually collected based on absorbance at 214 nm and samples reduced in volume to ~50 μl under vacuum. Amino-acid sequence analyses were performed by the Macromolecular Structure Analysis Facility at the University of Kentucky, Lexington, using an Applied Biosystems 477A automated sequencer. For additional confirmation of amino acid sequence data, a duplicate sample of Rubisco LSMT was purified, proteolyzed, and peptic polypeptide fragments submitted for amino acid sequence analyses.

Example 4

Synthesis of first-strand cDNA and polymerase chain reaction amplification

Pools of oligonucleotide primers encoding portions of two LSMT peptic peptides, P14 and P18, were synthesized with the number of different species (degeneracies) in each pool minimized as previously described (Klein et al., "Cloning and developmental expression of the sucrose-phosphate-synthase gene from spinach," *Planta*. 190:498–510 (1993)). First strand cDNA synthesis and polymerase chain reaction (PCR) conditions were as described, for example, in Klein et al., "Cloning and developmental expression of the sucrose-phosphate-synthase gene from spinach," *Planta*. 190:498–510 (1993), except 5 μl of first strand cDNA was used as PCR-template and the PCR-annealing temperature was reduced to 48° C. The appropriate sense and antisense PCR-primers directed against LSMT peptides, P14 and P18, are shown in Table 1, as shown below.

TABLE 1

Degenerate PCR primer pools designed according to amino-acid sequence of Rubisco LSMT peptides P14 and P18. Underlined nucleotides represent degeneracy nearest 3' terminus at which pools or primers differ.

PEPTIDE P14																			
		NH ₂	Pro	Met	Ala	Asp	Leu	Ile	Asn	His	Ser	Ala	Gly	Val	Thr	Asn	Glu	Asp	COOH
Sense DNA	5'-	CCA	ATG	GCA	GAT	TTA	ATT	AAT	CAT	TCA	GCA	GGA	GTA	ACA	AAT	GAA	GAT	-3'	
		C		C	C	G	A	C	C	C	C	C	C	C	C	G	C		
		G		G		CTA	C			G	G	G	G	G	G				
		T		T		C	T			T	T	T	T	T	T				
P14s ₁ P14s ₂					5'-	GAT	TTT	ATT	AAT	CAT	ICI	GCI	GGI	GII	ACI	AAT	GAA	GAT	-3'
											G	G	G	G	C	C	G	C	
	3'-	GGG	TAC	CGT	CTA	AAT	TAA	TTA	GTA	AGT	CGT	CCT	CAT	TGT	TTA	CTT	CTA	-5'	Antisense DNA
		C		G	G	C	T	G	G	G	G	G	G	G	G	C	G		
		A		C		GAT	G			C	C	C	C	C	C				
		T		A		G				A	A	A	A	A					
						C				TCA									
						A				G									
	3'-	TAC	CGG	CTA	GAT	TAI	TTG	GTG	IGI	CGI	CCI	CAI	TGI	TTG	-5'				
			C	G					C										
			A																
			T																
PEPTIDE P18																			
		NH ₂	Try	Asn	Arg	Thr	Leu	Pro	Pro	Gly	Leu	Leu	Pro	Tyr	Leu	Arg	COOH		
Sense DNA	5'-	TAT	AAT	CGA	ACA	TTA	CCA	CCA	GGA	TTA	TTA	CCA	TAT	TTA	CGA	-3'			
		C	C	C	C	G	C	C	C	G	G	C	C	C	C				
				G	G	CTA	G	G	G	CTA	CTA	G							
				T	T	C	T	T	T	C	C	T							
P18s ₁ P18s ₂				AGA		G				G	G								
				G		T				T	T								
	5'-	AAT	CGI	ACI	TTI	CCI	CCI	GGI	TTI	TTI	CCA	TAT	TT	-3'					
			A								G	C	C						

TABLE 1-continued

Degenerate PCR primer pools designed according to amino-acid sequence of Rubisco LSMT peptides P14 and P18. Underlined nucleotides represent degeneracy nearest 3' terminus at which pools or primers differ.																
3'-	ATA	TTA	GCT	TGT	AAT	GGT	GGT	CCT	AAT	AAT	T	ATA	AAT	GCT	-5'	Antisense
	G	G	G	G	C	G	G	G	C	C	G	G	C	G		DNA
			C	C	GAT	C	C	C	GAT	GAT			GAT	C		
			A	A	G	A	A	A	G	G	A		G	A		
			TCT		C				C	C			C	TCT		
			C		A				A	A			C	C		
3'-	ATA	TTA	GCT	TGT	GAT	GGT	GGT	CCI	GAT	GAT	GGT	ATG	-5'			P18a ₁
	<u>G</u>	G	T													P18a ₂

Following amplification, the PCR product was purified and blunt-end ligated into the SK plasmid (Stratagene, La Jolla, Calif. USA) and sequenced as described, for example, in Klein et al. "Photoaffinity labeling of mature and precursor forms of the small subunit of ribulose-1,5-bisphosphate carboxylase/oxygenase after expression in *Escherichia coli*," *Plant Physiol.* 98:546-553 (1992).

Example 5

Screening of a pea cDNA library

To obtain a full-length cDNA of pea LSMT, a pea λ gt10 cDNA library (Gantt et al. "Transfer of rp122 to the nucleus greatly preceded its loss from the chloroplast and involved the gain of an intron," *EMBO J* 10:3073-3078 (1991)) was screened with the Rubisco LSMT-PCR product. Approximately 5×10^4 primary plaques were screened with a randomly labeled 360-bp PCR product of Rubisco LSMT under recommended conditions (Stratagene, La Jolla, Calif. USA). After four rounds of plaque purification, three potential positive plaques were identified. Following amplification and purification of bacteriophage DNA, Rubisco LSMT cDNAs were subcloned into SK plasmid and the complete sequence of all three clones (approximately 1600 to 1775 bp in length) was obtained.

The technique of PCR-RACE (Rapid Amplification of cDNA Ends) was used to obtain a portion of the 5'-region of LSMT essentially as described by the manufacturer (GIBCO-BRL, Gaithersburg, Md. USA) except 100-ng of poly(A) mRNA was substituted for total RNA. The gene-specific (antisense) primer used to prime synthesis of first-strand LSMT cDNA was 5'-CCAAAAGAAGTCATCCAGCGTCAC (SEQ. ID NO. 41, position 700-667 bp). Amplification by PCR used the Anchor primer (supplied by GIBCO-BRL) and a second antisense LSMT-specific primer (5'-CAUCAUCAUCCTGTGGCAGAATACCAAAATAGT) which annealed to an internal, nested site within the LSMT cDNA (SEQ. ID NO. 41, position 515-492 bp). The inclusion of the (CAU)₄ repeat sequence at the 5' terminus permitted a uracil DNA glycosylase (UDG) cloning strategy of the PCR-RACE product. PCR amplification conditions were as above except for an annealing temperature of 55°C. and an extension time of 40 seconds.

Example 6

Northern blot analyses

Polyadenylated mRNA (0.5 μ g per lane) or total RNA (2 μ g per lane) was loaded on formaldehyde gels (Sambrook et al. *Molecular cloning: A laboratory manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989)) and transferred to GeneScreen nylon membranes (DuPont-

NEN, Wilmington, Del. USA). Conditions for prehybridization and hybridization with radiolabeled antisense-RNA probes were as described in Klein et al. "Cloning and developmental expression of the sucrose-phosphate-synthase gene from spinach," *Planta*. 190:498-510 (1993). The northern probe for rbcS was as also described in Klein et al. supra; and Klein et al. "Photoaffinity labeling of mature and precursor forms of the small subunit of ribulose-1,5-bisphosphate carboxylase/oxygenase after expression in *Escherichia coli*," *Plant Physiol.* 98:546-553 (1992); the probe for rbcL was as described in Klein et al. "Light-induced Transcription of Chloroplast Genes," *J. Biol. Chem.* 265(4):1895-1902 (1990)). The probe for rbcMT was a 1750-base antisense RNA from a portion of the open reading frame and 3'-untranslated region of pea.

Example 7

Genomic Southern blot analyses

Nuclear DNA was isolated from nuclei as described in Bedbrook, "A plant nuclear DNA preparation procedure," *Plant Mol. Biol. Newslett.* 2:24 (1981). Ten μ g of high molecular weight DNA was digested to completion with EcoR I, Hind III, and Dra I (50 units each). Following digestion, DNA was ethanol precipitated, electrophoresed on 0.8% agarose gels and transferred to Nytran nylon membranes using an alkaline transfer solution as described (Turboblotter instruction manual, Schleicher and Schuell, Keene, N.H. USA). Blots were prehybridized and hybridized at 42°C. in the presence of 50% formamide and 10% dextran sulfate. The probe was a random primer-labeled 1775 bp cDNA of pea LSMT (encompassing the open reading frame and entire 3'-untranslated region).

Example 8

Computer alignment of the amino acid sequences was performed using the FastDB program (Intelligenetics Inc., Mountain View, Calif. USA). Autoradiograms were scanned with an image acquisition densitometer (Biolmage, Milligen/Bioscience, Ann Arbor, Mich. USA) to determine the relative intensity of mRNA signal and quantified on the basis of whole-band analysis.

In accordance with the present invention, rubisco LSMT has thus been purified ~8000-fold by a novel affinity purification technique from pea chloroplasts as described in Wang et al. *Protein Expression and Purification*. After affinity-purification of Rubisco LSMT, SDS-PAGE analysis showed a single polypeptide with an apparent molecular mass of ~57 kDa. Direct Edman degradative sequencing attempts followed by amino acid analyses after HCl hydrolysis of electroblotted affinity-purified Rubisco LSMT revealed that the N-terminus was blocked. Thus, subsequent

efforts were directed towards the acquisition of internal amino acid sequence as a starting point for isolating a cDNA of pea Rubisco LSMT. Reverse phase-HPLC isolation of peptic fragments from Rubisco LSMT resulted in the identification of several reliable amino acid sequences (FIG. 1, asterisks). One polypeptide peak, however, was heterogeneous and consisted of at least three subsequences which were identifiable based on differences in the relative amino acid yields after each cycle of sequencing.

Example 9

The partial amino-acid sequence of Rubisco LSMT enabled the inventor to develop a molecular probe for the Rubisco LSMT gene (*rbcMT*) using PCR. Pools of deoxynucleosine-containing primers encoding part of two peptic peptides, P14 and P18, were synthesized with the number of species in each pool minimized, as shown in Table 1, supra. Using random-hexamer-primed first strand cDNA as a template, the combination of primer pools P14-2s with P18-1a or P18-2a directed the synthesis of a single 360-bp PCR product. No other primer combinations yielded a detectable PCR product.

The fact that either antisense primer P18-1a or P18-2a (which differ by a single nucleotide near the 3' terminus) directed the synthesis of a PCR product reflects the relative tolerance of the PCR system for base-pair mismatches near the 3' terminus of the primer. The identity of the amplification product as a partial cDNA of *rbcMT* was confirmed by comparison of the deduced amino-acid sequence of the PCR product with additional peptic fragments from purified pea Rubisco LSMT protein (see FIG. 2).

The PCR-amplified fragment of *rbcMT* was used to screen a λ gt10 pea cDNA library (Gantt et al., "Transfer of *rp122* to the nucleus greatly preceded its loss from the chloroplast and involved the gain of an intron," *EMBO J* 10:3073-3078 (1991)). Three partial clones were obtained with inserts greater than 1600 bp in length. Complete sequence analysis of the three clones showed that the nucleotide sequence of all clones were identical. The sequence of the PCR-derived cDNA was identical to the λ gt10 cDNAs except for the incorrect identification of Thr-249 as an Asn during peptide sequencing of pepsin fragment P14. The longest clone (1775 bp in length) lacked only a portion of the 5'-untranslated region. The remainder of the 5'-untranslated region was obtained by PCR-RACE. The 515 bp PCR-RACE product was barely detectable on ethidium-stained gels which likely reflects the low abundance of the *rbcMT* mRNA in pea. Sequence analysis confirmed the identity of the PCR-RACE product as encoding for the predicted 5' portion of *rbcMT* including the remainder of the 5'-untranslated region. In the region where the PCR-RACE product overlapped the cloned cDNA of *rbcMT*, complete sequence identity was observed (SEQ. ID NO. 41, position 31-484 bp). Given these overlapping clones, the present inventor was able to assemble the sequence of the *rbcMT* cDNA as shown in SEQ. ID NO. 41. All of the peptic polypeptide sequences obtained from affinity-purified Rubisco LSMT were identified in the translated open-reading frame of the *rbcMT* cDNA.

The *rbcMT* cDNA of 1802 bp in length contained a 5' leader of 58-nucleotides which contained several short repeat elements and a 3'-untranslated region of 276-nucleotides. The *rbcMT* cDNA encoded for a protein of 489-amino acid residues with a predicted molecular mass of 55 kDa. Examination of the amino terminus of Rubisco LSMT revealed several motifs that commonly appear in

chloroplast transit-peptide sequences, such as an abundance of hydroxylated amino acids Ser and Thr, presence of small hydrophobic amino acids, and general lack of acidic amino acids (Keegstra et al., "Chloroplastic precursors and their transport across the envelope membranes," *Annu. Rev. Plant. Physiol. Plant. Mol. Biol.* 40:471-501 (1989); and Theg et al., "Protein import into chloroplasts," *Trends in Cell Biology* 3:186-190 (1993)). Given that N-terminal sequence information could not be obtained for Rubisco LSMT, and that there is as yet no amino acid consensus sequence or secondary structural motif which unambiguously identifies the processing site for removal of chloroplast transit sequences (von Heijne et al., "Chloroplast transit peptides: The perfect random coil?" *FEBS Lett.* 278(1):1-3 (1991)), the cleavage site between the precursor and mature forms of Rubisco LSMT could not be determined.

Comparison of the deduced amino acid sequence of *rbcMT* cDNA with protein carboxyl methyltransferases from wheat (D-aspartyl/L-isoaspartyl protein methyltransferase, Mudgett et al., "Characterization of plant L-isoaspartyl methyltransferases that may be involved in seed survival: Purification, cloning, and sequence analysis of the wheat germ enzyme," *Biochemistry* 32:11100-11111 (1993)) and *E. coli* (gamma-glutamyl carboxyl methyltransferase, Mutoh et al., "Nucleotide sequence corresponding to five chemotaxis genes in *Escherichia coli*," *J. Bacteriol.* 165:161-166 (1986)) showed a low alignment score with sequence identity on the order of 10% (gaps in the sequence were introduced to maximize alignment). Three short amino acid regions (8 to 10 residues) of sequence similarity have been reported for several protein and small-molecule AdoMet-dependent methyltransferases (Kagan et al., "Widespread occurrence of three sequence motifs in diverse S-adenosylmethionine-dependent methyltransferases suggests a common structure for these enzymes," *Arch. Biochem. Biophys.* 310(2):417-427 (1994)). Using manual alignment, none of the three proposed sequence motifs of AdoMet-dependent methyltransferases were detected in Rubisco *rbcMT*. In a search of the Swissprot and NBRF-PIR data banks, the best match for Rubisco *rbcMT* was AfsR protein of *Streptomyces coelicolor* which reflected a 23% sequence identity over the entire protein, again with considerable gaps introduced.

Example 10

DNA analysis

To obtain information on gene copy number, total pea leaf DNA was isolated and digested with several different restriction endonucleases (FIG. 3). A 1775 bp *rbcMT* cDNA probe hybridized to two EcoR I DNA fragments, approximately 5.3 kbp and 2.0 kbp (one EcoR I restriction endonuclease site is located within the sequenced cDNA). Two bands, approximately 3.5 kbp and 1.3 kbp, were observed after cleavage with Dra I, while a single band of 3.7 kbp was observed after DNA-digestion with Hind III. The simplicity of the DNA restriction digest pattern suggests that the gene copy number per haploid genome is low for *rbcMT*.

Example 11

RNA analyses

Northern blot analyses were conducted on pea tissues to examine several developmental and organ-specific parameters governing *rbcMT* gene expression. As a basis for comparison, the expression of genes encoding Rubisco small (*rbcS*) and large (*rbcL*) subunit were concomitantly examined. The *rbcS* gene family and *rbcL* gene were exam-

ined in an attempt to determine whether the expression of the Rubisco subunits and Rubisco LSMT was coordinated. Northern blot analysis indicated that the rbcMT gene encoded for a single species of mRNA of approximately 1.8 kb in length (see FIG. 4). Examination of organ-specific expression showed that accumulation of the rbcMT transcript paralleled the accumulation of rbcL and rbcS mRNA with the greatest proportion of mRNA being localized in green leaf tissue. Transcripts encoding rbcS, rbcL and rbcMT were detected in pea stems, though the level of expression was 7, 10, and 28-fold lower, respectively, than in green leaves. The quantity of rbcMT, rbcS, and rbcL mRNA in root tissue was below the level of Northern blot sensitivity. Maximum extractable Rubisco LSMT activity generally paralleled the accumulation of rbcMT mRNA, though the enzyme activity detected in stems was greater than would be predicted based on mRNA levels. Maximum extractable Rubisco LSMT activity of roots, stems, and green leaves was 2, 15, and 36 pmoles CH₃·min⁻¹·mg protein⁻¹, respectively. Finally, it should be noted that the exposure times of the rbcMT, rbcS, and rbcL Northern analyses differ considerably and hence should be considered when comparing the absolute amounts of each transcript. The exposure time of rbcMT Northern analyses were consistently 25- to 50-times longer than that of rbcL or rbcS, suggesting that rbcMT transcripts do not accumulate to the level of the Rubisco subunits.

Examination of the accumulation of rbcMT mRNA during the greening of pea leaves is shown in FIG. 5. A low level of rbcMT mRNA was detected in 8-day-old dark-grown pea leaves (lane 1). Upon illumination of etiolated peas, rbcMT transcript levels increased ~3-fold after 24 hours of illumination and then declined slightly after an additional 48 hours of development in the light (lanes 2-3). The maximum

extractable activity of Rubisco LSMT enzyme increased during the greening of dark-grown peas from 11 pmoles CH₃·min⁻¹·mg protein⁻¹ in dark-grown leaves to an apparent maximum of 32.5 pmoles CH₃·min⁻¹·mg protein⁻¹ after 72 hours illumination. This level of extractable Rubisco LSMT enzyme activity was similar to that observed (32.4 pmoles CH₃·min⁻¹·mg protein⁻¹) for peas grown eight days under continuous illumination. Interestingly, the level of rbcMT mRNA in continuous illuminated leaves was significantly lower than the levels observed during the early stages of greening of pea (lanes 2-3 vs. 4). In fact, levels of rbcMT mRNA from continuous illuminated plants was not visibly different from dark-grown leaves. As expected, rbcS and rbcL transcript levels also increased upon illumination of dark-grown seedlings. In contrast to rbcMT, transcripts of rbcS and rbcL reached an apparent maximum during the latter stages of greening (lane 3). In addition, rbcS and rbcL transcript levels remained elevated in leaves grown under continuous illumination (lane 4). These results indicate that, unlike rbcS and rbcL, transcript levels for rbcMT reach an apparent maximum during the early stages of light-induced leaf development and decline in mature light-grown leaf tissue. These changes in transcript levels would be expected for an enzyme whose function involves post-translational protein processing.

All of the references cited herein are effectively incorporated by reference to the same extent as if each individually had been incorporated by reference.

Other embodiments of the invention will be apparent to those skilled in the art from consideration of the specification and practice of the invention disclosed herein. It is intended that the specification and examples be considered as exemplary only, with the true scope and spirit of the invention being indicated by the following claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i i i) NUMBER OF SEQUENCES: 41

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(i x) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 1
- (D) OTHER INFORMATION: /note= "Amino acid 1 wherein Xaa = NH2."

(i x) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 18
- (D) OTHER INFORMATION: /note= "Amino acid 18 wherein Xaa = COOH."

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Xaa Pro Met Ala Asp Leu Ile Asn His Ser Ala Gly Val Thr Asn Glu
1 5 10 15
Asp Xaa

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 48 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:2:

CCAATGGCAG ATTTAATTAA TCATTTCAGCA GGAGTAACAA ATGAAGAT

4 8

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 48 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CCCATGGCCG ACTTGATAAA CCACTCCGCC GGCCTCACCA ACGAGGAC

4 8

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 48 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CCGATGGCGG ACCTAATCAA CCACTCGGCG GGGGTGACGA ACGAGGAC

4 8

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 48 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:5:

CCTATGGCTG ACCTCATCAA CCACTCTGCT GGTGTTACTA ACGAGGAC

4 8

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 48 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:6:

CCTATGGCTG ACCTGATCAA CCACAGTGCT GGTGTTACTA ACGAGGAC

4 8

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

-continued

(A) LENGTH: 48 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CCTATGGCTG ACCTTATCAA CCACAGCGCT GGTGTTACTA ACGAGGAC

4 8

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 39 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(i x) FEATURE:

(A) NAME/KEY: misc_feature
 (B) LOCATION: 6..30
 (D) OTHER INFORMATION: /note= "Nucleotides 6, 9, 16, 18,
 21, 24, 27 and 30 wherein N = I."

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:8:

GATTTNATNA ATCATNCNGC NGGNGTNACN AATGAAGAT

3 9

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 39 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(i x) FEATURE:

(A) NAME/KEY: misc_feature
 (B) LOCATION: 6..30
 (D) OTHER INFORMATION: /note= "Nucleotides 6, 9, 16, 18,
 21, 24, 27 and 30 wherein N = I."

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:9:

GATTTNATNA ATCATNCNGC NGGNGTNACN AACGAGGAC

3 9

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 48 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GGGTACCGTC TAAATTAATT AGTAAGTCGT CCTCATTGTT TACTTCTA

4 8

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 48 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:11:

-continued

GGCTACCGGC TGA ACTATTT GGTGAGGCGG CCGCAGTGGT TGCTCCTG

4 8

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 48 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:12:

GGATACCGCC TGGATTAGTT GGTGAGCCGC CCCCCTGCT TGCTCCTG

4 8

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 48 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GGTTACCGAC TGGAGTAGTT GGTGAGACGA CCACAATGAT TGCTCCTG

4 8

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 48 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:14:

GGTTACCGAC TGGACTAGTT GGTGTCACGA CCACAATGAT TGCTCCTG

4 8

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 48 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:15:

GGTTACCGAC TGGAATAGTT GGTGTCGCGA CCACAATGAT TGCTCCTG

4 8

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 39 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(i x) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 12..36
- (D) OTHER INFORMATION: /note= "Nucleotides 12, 15, 22, 24, 27, 30, 33 and 36 wherein N = I."

-continued

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:16:

TACCGGCTAG ANTANTTGGT GNGNCGNCCN CANTGNTTG

3 9

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 39 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(i x) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 12..36
- (D) OTHER INFORMATION: /note= "Nucleotides 12, 15, 22, 24, 27, 30, 33 and 36 wherein N = I."

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:17:

TACCGCCTCG ANTANTTGGT GNCNCGNCCN CANTGNTTG

3 9

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 39 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(i x) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 12..36
- (D) OTHER INFORMATION: /note= "Nucleotides 12, 15, 22, 24, 27, 30, 33 and 36 wherein N = I."

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:18:

TACCGACTCG ANTANTTGGT GNCNCGNCCN CANTGNTTG

3 9

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 39 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(i x) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 12..36
- (D) OTHER INFORMATION: /note= "Nucleotides 12, 15, 22, 24, 27, 30, 33 and 36 wherein N = I."

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:19:

TACCGTCTCG ANTANTTGGT GNCNCGNCCN CANTGNTTG

3 9

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(i x) FEATURE:

- (A) NAME/KEY: Modified-site

-continued

- (B) LOCATION: 1
 (D) OTHER INFORMATION: /note= "Amino acid 1 wherein Xaa = NH2."

- (i x) FEATURE:
 (A) NAME/KEY: Modified-site
 (B) LOCATION: 16
 (D) OTHER INFORMATION: /note= "Amino acid 16 wherein Xaa = COOH."

- (x i) SEQUENCE DESCRIPTION: SEQ ID NO:20:

Xaa Tyr Asn Arg Thr Leu Pro Pro Gly Leu Leu Pro Tyr Leu Arg Xaa
 1 5 10 15

- (2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 42 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

- (i i) MOLECULE TYPE: DNA (genomic)

- (x i) SEQUENCE DESCRIPTION: SEQ ID NO:21:

TATAATCGAA CATTACCACC AGGATTATTA CCATATTTAC GA

4 2

- (2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 42 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

- (i i) MOLECULE TYPE: DNA (genomic)

- (x i) SEQUENCE DESCRIPTION: SEQ ID NO:22:

TACAACCGCA CCTTGCCCC CGGCTTGTTG CCCTACTTGC GC

4 2

- (2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 42 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

- (i i) MOLECULE TYPE: DNA (genomic)

- (x i) SEQUENCE DESCRIPTION: SEQ ID NO:23:

TACAACCGGA CGCTACCGCC GGGGCTACTA CCGTACCTAC GG

4 2

- (2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 42 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

- (i i) MOLECULE TYPE: DNA (genomic)

- (x i) SEQUENCE DESCRIPTION: SEQ ID NO:24:

TACAACCGTA CTCTCCCTCC TGGTCTCCTC CCTTACCTCC GT

4 2

- (2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 42 base pairs

-continued

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:25:

TACAACAGAA CTCTGCCTCC TGGTCTGCTG CCTTACCTGA GA

4 2

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 42 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:26:

TACAACAGGA CTCCTCCTCC TGGTCTTCTT CCTTACCTTA GG

4 2

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 35 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(i x) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 6..27
- (D) OTHER INFORMATION: /note= "Nucleotides 6, 9, 12, 15, 18, 21, 24 and 27 wherein N = 1."

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:27:

AATCGNACNT TNCCNCCNGG NTNTTNCCA TATTT

3 5

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 35 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(i x) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 6..27
- (D) OTHER INFORMATION: /note= "Nucleotides 6, 9, 12, 15, 18, 21, 24 and 27 wherein N = 1."

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:28:

AATAGNACNT TNCCNCCNGG NTNTTNCCC TACCT

3 5

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 35 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(i x) FEATURE:

- (A) NAME/KEY: misc_feature

-continued

- (B) LOCATION: 6.27
- (D) OTHER INFORMATION: /note= "Nucleotides 6, 9, 12, 15, 18, 21, 24 and 27 wherein N = I."

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:29:

AATAGNACNT TNCCNCCNGG NTINTTNCCG TACCT

3 5

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 35 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(i x) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 6.27
- (D) OTHER INFORMATION: /note= "Nucleotides 6, 9, 12, 15, 18, 21, 24 and 27 wherein N = I."

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:30:

AATAGNACNT TNCCNCCNGG NTINTTNCCT TACCT

3 5

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 42 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:31:

ATATTAGCTT GTAATGGTGG TCCTAATAAT GGTATAAAATG CT

4 2

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 42 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:32:

ATGTTGGCGT GGAACGGGGG GCCGAACAAC GGGATGAACG CG

4 2

(2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 42 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:33:

ATGTTGGCCT GCGATGGCGG CCCCGATGAT GGCATGGATG CC

4 2

(2) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 42 base pairs
- (B) TYPE: nucleic acid

-continued

```

      ( C ) STRANDEDNESS: single
      ( D ) TOPOLOGY: linear

    ( i i ) MOLECULE TYPE: DNA (genomic)

    ( x i ) SEQUENCE DESCRIPTION: SEQ ID NO:34:
ATGTTGGCAT GAGAGGGAGG ACCAGAGGAG GGAATGGAGG CA                4 2

( 2 ) INFORMATION FOR SEQ ID NO:35:

  ( i ) SEQUENCE CHARACTERISTICS:
    ( A ) LENGTH: 42 base pairs
    ( B ) TYPE: nucleic acid
    ( C ) STRANDEDNESS: single
    ( D ) TOPOLOGY: linear

  ( i i ) MOLECULE TYPE: DNA (genomic)

  ( x i ) SEQUENCE DESCRIPTION: SEQ ID NO:35:
ATGTTGTCTT GAGACGGAGG ACCAGACGAC GGAATGGACT CT                4 2

( 2 ) INFORMATION FOR SEQ ID NO:36:

  ( i ) SEQUENCE CHARACTERISTICS:
    ( A ) LENGTH: 42 base pairs
    ( B ) TYPE: nucleic acid
    ( C ) STRANDEDNESS: single
    ( D ) TOPOLOGY: linear

  ( i i ) MOLECULE TYPE: DNA (genomic)

  ( x i ) SEQUENCE DESCRIPTION: SEQ ID NO:36:
ATGTTGTCCT GAGAAGGAGG ACCAGAAGAA GGAATGGAAT CC                4 2

( 2 ) INFORMATION FOR SEQ ID NO:37:

  ( i ) SEQUENCE CHARACTERISTICS:
    ( A ) LENGTH: 36 base pairs
    ( B ) TYPE: nucleic acid
    ( C ) STRANDEDNESS: single
    ( D ) TOPOLOGY: linear

  ( i i ) MOLECULE TYPE: DNA (genomic)

  ( i x ) FEATURE:
    ( A ) NAME/KEY: misc_feature
    ( B ) LOCATION: 9..33
    ( D ) OTHER INFORMATION: /note= "Nucleotides 9, 12, 15, 18,
      21, 24, 27, 30 and 33 wherein N = I."

  ( x i ) SEQUENCE DESCRIPTION: SEQ ID NO:37:
ATATTAGCNT GNGANGGNGG NCCNGANGAN GGNATG                3 6

( 2 ) INFORMATION FOR SEQ ID NO:38:

  ( i ) SEQUENCE CHARACTERISTICS:
    ( A ) LENGTH: 36 base pairs
    ( B ) TYPE: nucleic acid
    ( C ) STRANDEDNESS: single
    ( D ) TOPOLOGY: linear

  ( i i ) MOLECULE TYPE: DNA (genomic)

  ( i x ) FEATURE:
    ( A ) NAME/KEY: misc_feature
    ( B ) LOCATION: 9..33
    ( D ) OTHER INFORMATION: /note= "Nucleotides 9, 12, 15, 18,
      21, 24, 27, 30 and 33 wherein N = I."

  ( x i ) SEQUENCE DESCRIPTION: SEQ ID NO:38:
ATGTTGTCNT GNGANGGNGG NCCNGANGAN GGNATG                3 6

```

-continued

(2) INFORMATION FOR SEQ ID NO:39:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:39:

CCAAAAGAAG TCATCCAGCG TCAC

24

(2) INFORMATION FOR SEQ ID NO:40:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 36 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:40:

CAUCAUCAUC AUCCTGTGGC AGAATACCAA AATAGT

36

(2) INFORMATION FOR SEQ ID NO:41:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1801 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(i x) FEATURE:

- (A) NAME/KEY: CDS
 (B) LOCATION: 59..1528

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:41:

ACAACACAAA	AGAAAAGCGT	ATTATCACAA	AACAAAACCA	AGAACTAGAA	ACCAGAAA	58
ATG GCT ACT ATC TTT TCC GGA GGT TCA GTT TCT CCC TTT CTT TTT CAC	106					
Met Ala Thr Ile Phe Ser Gly Gly Ser Val Ser Pro Phe Leu Phe His						
1 5 10 15						
ACC AAC AAG GGT ACA TCT TTT ACA CCC AAA GCT CCA ATT CTT CAT CTC	154					
Thr Asn Lys Gly Thr Ser Phe Thr Pro Lys Ala Pro Ile Leu His Leu						
20 25 30						
AAG AGA TCT TTC TCT GCA AAA TCA GTA GCC TCT GTA GGA ACC GAA CCA	202					
Lys Arg Ser Phe Ser Ala Lys Ser Val Ala Ser Val Gly Thr Glu Pro						
35 40 45						
TCA CTG TCT CCA GCA GTT CAA ACC TTC TGG AAG TGG CTA CAG GAA GAA	250					
Ser Leu Ser Pro Ala Val Gln Thr Phe Trp Lys Trp Leu Gln Glu Glu						
50 55 60						
GGT GTC ATC ACT GCA AAG ACC CCA GTG AAA GCT AGT GTG GTC ACA GAA	298					
Gly Val Ile Thr Ala Lys Thr Pro Val Lys Ala Ser Val Val Thr Glu						
65 70 75 80						
GGT TTA GGA TTG GTT GCA CTT AAG GAC ATT TCT AGG AAT GAT GTT ATT	346					
Gly Leu Gly Leu Val Ala Leu Lys Asp Ile Ser Arg Asn Asp Val Ile						
85 90 95						
TTG CAG GTA CCA AAA AGG CTG TGG ATA AAT CCA GAT GCA GTT GCA GCT	394					
Leu Gln Val Pro Lys Arg Leu Trp Ile Asn Pro Asp Ala Val Ala Ala						
100 105 110						
TCA GAG ATT GGG AGA GTG TGC AGT GAG TTG AAG CCA TGG TTG TCT GTT	442					
Ser Glu Ile Gly Arg Val Cys Ser Glu Leu Lys Pro Trp Leu Ser Val						

-continued

115				120				125								
ATA	CTC	TTT	CTT	ATA	AGA	GAG	AGG	TCA	AGG	GAA	GAT	TCT	GTT	TGG	AAG	490
Ile	Leu	Phe	Leu	Ile	Arg	Glu	Arg	Ser	Arg	Glu	Asp	Ser	Val	Trp	Lys	
	130					135					140					
CAC	TAT	TTT	GGT	ATT	CTG	CCA	CAG	GAA	ACT	GAT	TCT	ACT	ATA	TAT	TGG	538
His	Tyr	Phe	Gly	Ile	Leu	Pro	Gln	Glu	Thr	Asp	Ser	Thr	Ile	Tyr	Trp	
	145				150					155					160	
TCA	GAG	GAA	GAG	CTT	CAA	GAG	CTT	CAA	GGT	TCT	CAA	CTT	TTG	AAA	ACA	586
Ser	Glu	Glu	Glu	Leu	Gln	Glu	Leu	Gln	Gly	Ser	Gln	Leu	Leu	Lys	Thr	
				165					170					175		
ACA	GTG	TCT	GTG	AAA	GAA	TAT	GTG	AAG	AAT	GAA	TGT	TTG	AAA	CTA	GAA	634
Thr	Val	Ser		Lys	Glu	Tyr	Val	Lys	Asn	Glu	Cys	Leu	Lys	Leu	Glu	
			180					185					190			
CAA	GAA	ATC	ATT	CTC	CCT	AAT	AAG	CGG	CTT	TTT	CCG	GAT	CCT	GTG	ACG	682
Gln	Glu	Ile	Ile	Leu	Pro	Asn	Lys	Arg	Leu	Phe	Pro	Asp	Pro	Val	Thr	
		195					200					205				
CTG	GAT	GAC	TTC	TTT	TGG	GCA	TTT	GGA	ATT	CTC	AGA	TCA	AGG	GCG	TTT	730
Leu	Asp	Asp	Phe	Phe	Trp	Ala	Phe	Gly	Ile	Leu	Arg	Ser	Arg	Ala	Phe	
	210					215					220					
TCT	CGC	CTT	CGC	AAT	GAA	AAT	CTG	GTT	GTG	GTT	CCA	ATG	GCA	GAC	TTG	778
Ser	Arg	Leu	Arg	Asn	Glu	Asn	Leu	Val	Val	Val	Pro	Met	Ala	Asp	Leu	
	225				230					235					240	
ATT	AAC	CAC	AGT	GCA	GGA	GTT	ACT	ACA	GAG	GAT	CAT	GCT	TAT	GAA	GTT	826
Ile	Asn	His	Ser	Ala	Gly	Val	Thr	Thr	Glu	Asp	His	Ala	Tyr	Glu	Val	
				245					250					255		
AAA	GGA	GCA	GCT	GGC	CTT	TTC	TCT	TGG	GAT	TAC	CTA	TTT	TCC	TTA	AAG	874
Lys	Gly	Ala	Ala	Gly	Leu	Phe	Ser	Trp	Asp	Tyr	Leu	Phe	Ser	Leu	Lys	
			260					265					270			
AGC	CCC	CTT	TCC	GTC	AAG	GCC	GGA	GAA	CAG	CTA	TAT	ATA	CAA	TAT	GAT	922
Ser	Pro	Leu	Ser	Val	Lys	Ala	Gly	Glu	Gln	Leu	Tyr	Ile	Gln	Tyr	Asp	
		275					280					285				
TTG	AAC	AAA	AGC	AAT	GCA	GAG	TTG	GCT	CTA	GAC	TAC	GGT	TTC	ATT	GAA	970
Leu	Asn	Lys	Ser	Asn	Ala	Glu	Leu	Ala	Leu	Asp	Tyr	Gly	Phe	Ile	Glu	
		290				295					300					
CCA	AAT	GAA	AAT	CGA	CAT	GCA	TAC	ACT	CTG	ACG	CTG	GAG	ATA	TCT	GAG	1018
Pro	Asn	Glu	Asn	Arg	His	Ala	Tyr	Thr	Leu	Thr	Leu	Glu	Ile	Ser	Glu	
		305			310					315					320	
TCG	GAC	CCT	TTT	TTT	GAT	GAC	AAA	CTA	GAC	GTT	GCT	GAG	TCC	AAT	GGT	1066
Ser	Asp	Pro	Phe	Phe	Asp	Asp	Lys	Leu	Asp	Val	Ala	Glu	Ser	Asn	Gly	
				325					330					335		
TTT	GCT	CAG	ACA	GCG	TAC	TTT	GAC	ATC	TTC	TAT	AAT	CGC	ACT	CTT	CCA	1114
Phe	Ala	Gln	Thr	Ala	Tyr	Phe	Asp	Ile	Phe	Tyr	Asn	Arg	Thr	Leu	Pro	
			340					345					350			
CCT	GGA	TTG	CTT	CCA	TAT	CTG	AGA	CTT	GTA	GCG	CTA	GGG	GGT	ACC	GAC	1162
Pro	Gly	Leu	Leu	Pro	Tyr	Leu	Arg	Leu	Val	Ala	Leu	Gly	Gly	Thr	Asp	
		355					360					365				
GCT	TTC	TTA	TTG	GAA	TCA	CTG	TTC	AGA	GAC	ACC	ATA	TGG	GGT	CAT	CTT	1210
Ala	Phe	Leu	Leu	Glu	Ser	Leu	Phe	Arg	Asp	Thr	Ile	Trp	Gly	His	Leu	
		370				375					380					
GAG	TTG	TCT	GTC	AGC	CGT	GAC	AAT	GAG	GAG	CTA	CTA	TGC	AAA	GCC	GTT	1258
Glu	Leu	Ser	Val	Ser	Arg	Asp	Asn	Glu	Glu	Leu	Leu	Cys	Lys	Ala	Val	
					390					395				400		
CGA	GAA	GCC	TGC	AAA	TCT	GCC	CTT	GCT	GGT	TAT	CAT	ACA	ACC	ATT	GAA	1306
Arg	Glu	Ala	Cys	Lys	Ser	Ala	Leu	Ala	Gly	Tyr	His	Thr	Thr	Ile	Glu	
				405					410					415		
CAG	GAT	CGC	GAG	TTG	AAA	GAA	GGA	AAT	CTA	GAT	TCA	AGG	CTT	GCA	ATA	1354
Gln	Asp	Arg	Glu	Leu	Lys	Glu	Gly	Asn	Leu	Asp	Ser	Arg	Leu	Ala	Ile	
			420				425						430			
GCA	GTT	GGA	ATA	AGA	GAA	GGG	GAA	AAG	ATG	GTC	CTG	CAG	CAA	ATT	GAC	1402
Ala	Val	Gly	Ile	Arg	Glu	Gly	Glu	Lys	Met	Val	Leu	Gln	Gln	Ile	Asp	

-continued

435						440					445					
GGG	ATC	TTC	GAG	CAG	AAA	GAA	TTG	GAG	TTG	GAC	CAG	TTA	GAG	TAT	TAT	1450
Gly	Ile	Phe	Glu	Gln	Lys	Glu	Leu	Glu	Leu	Asp	Gln	Leu	Glu	Tyr	Tyr	
450						455				460						
CAA	GAA	AGG	AGG	CTC	AAG	GAT	CTT	GGA	CTT	TGC	GGA	GAA	AAT	GGC	GAT	1498
Gln	Glu	Arg	Arg	Leu	Lys	Asp	Leu	Gly	Leu	Cys	Gly	Glu	Asn	Gly	Asp	
465						470				475				480		
ATC	CTT	GGA	GAC	CTA	GGA	AAA	TTC	TTC	TAA	TCTTGCAGGA	AAATTCTTCT					1548
Ile	Leu	Gly	Asp	Leu	Gly	Lys	Phe	Phe	*							
				485						490						
AATCTTGCA		G		AAGCATTTC		A		AACCTGTTAA		AGATACACTG		TTGTTTACAA		ATGGAGTCTT		1608
CTGAGACGTA		C		CGATGCCATG		A		TTTTTGCAAT		CAATCTTAAG		AGGATCGTGA		TCAATTTTGA		1668
CTCTGGAGTC		T		TGGACCAATC		C		ATTACATGC		TTGAAGTTTG		TAAAGAGGAA		AATGTAATGT		1728
GTGAAATATA		A		AATTACACTT		C		TGTACTGGT		GATTATTTAT		AAAGCAGTTG		ACCATTATTA		1788
TTACAAAAAA		A		A		A		A		A		A		A		1801

What is claimed is:

1. An isolated Rubisco LSMT gene which is expressed in a photosynthesizing plant having a large subunit of Rubisco and which encodes a Rubisco LSMT enzyme.

2. The isolated Rubisco LSMT gene of claim 1, wherein said enzyme catalyzes methylation of the ϵ -amine of lysine 14 in the large subunit of Rubisco.

3. The isolated gene of claim 1, wherein said photosynthesizing plant is selected from the group consisting of pea, soybean, tomato, potato, tobacco, pepper cucumber, melon and gourd.

4. The isolated gene of claim 3, wherein said plant is pea.

5. A recombinant vector comprising the Rubisco LSMT gene of claim 1, said vector being one which can transform a photosynthesizing plant.

6. A method for expressing a Rubisco LSMT gene in a plant comprising transforming a plant with the Rubisco LSMT gene of claim 1, said plant thereby expressing the Rubisco LSMT enzyme encoded by said Rubisco LSMT gene.

7. A recombinant photosynthesizing non-pea plant transformed with the Rubisco LSMT gene of claim 1.

8. The recombinant plant of claim 7; wherein said Rubisco LSMT gene expresses said Rubisco LSMT enzyme.

9. The recombinant plant of claim 8, wherein said Rubisco LSMT enzyme catalyzes methylation of an ϵ -amine of lysine-14 in the large subunit of Rubisco.

10. The recombinant plant of claim 9, wherein said plant is selected from the group consisting of a soybean, tomato, potato, tobacco, pepper, cucumber, melon and gourd.

11. A method for transforming a plant comprising inserting said vector of claim 5, into a plant.

12. A recombinant photosynthesizing non-pea plant transformed with the vector of claim 5.

13. The isolated gene of claim 1, wherein said photosynthesizing plant is a legume.

14. The recombinant plant of claim 9, wherein said plant is a legume.

15. An isolated cDNA having the sequence of SEQ. ID NO. 41.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 5,723,752

Page 1 of 4

DATED : March 3, 1998

INVENTOR(S) :
Robert L. HOUTZ

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

The title page, showing "4 Drawing Sheets" should be --7 Drawing Sheets--.

IN THE DRAWINGS:

Add the Drawing Sheet consisting of Figs. 3, 4 and 5, as shown on the attached pages.

Signed and Sealed this

Twentieth Day of February, 2001

Attest:

NICHOLAS P. GODICI

Attesting Officer

Acting Director of the United States Patent and Trademark Office

Figure 3

Figure 4

Light Treatment

Figure 5