

University of Kentucky

UKnowledge

Oral Health Practice Faculty Publications

Oral Health Practice

4-13-2020

Medication-Related Osteonecrosis of the Mandible

Galal Omami

University of Kentucky, galal.omami@uky.edu

Follow this and additional works at: https://uknowledge.uky.edu/ohp_facpub



Part of the [Dentistry Commons](#)

[Right click to open a feedback form in a new tab to let us know how this document benefits you.](#)

Repository Citation

Omami, Galal, "Medication-Related Osteonecrosis of the Mandible" (2020). *Oral Health Practice Faculty Publications*. 17.

https://uknowledge.uky.edu/ohp_facpub/17

This Article is brought to you for free and open access by the Oral Health Practice at UKnowledge. It has been accepted for inclusion in Oral Health Practice Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Medication-Related Osteonecrosis of the Mandible

Digital Object Identifier (DOI)

<https://doi.org/10.1177/0145561320919574>

Notes/Citation Information

Published in *Ear, Nose, & Throat Journal*.

© The Author(s) 2020

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

Medication-Related Osteonecrosis of the Mandible

Galal Omami, BDS, MSc, MDentSc, FRCD(C)¹ 

Ear, Nose & Throat Journal

1-2

© The Author(s) 2020

Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/0145561320919574

journals.sagepub.com/home/ear



Medication-related osteonecrosis of the jaw (MRONJ) is a pathologic process that rarely follows exposure to antiresorptive agents such as bisphosphonate or denosumab, which are used to treat lytic bone lesions in patients with osseous metastases related to carcinoma of the breast or prostate, multiple myeloma, Paget's disease, and osteoporosis. These drugs inhibit osteoclastic bone resorption and remodeling.

The risk of osteonecrosis of the jaw in patients receiving either bisphosphonate or denosumab ranges from 0.7% to 6.7%.¹ However, this condition occurs more commonly in patients receiving the medication intravenously and after a surgical dental procedure, such as extraction or implant placement. The diagnosis of MRONJ is made clinically when there is nonhealing exposed bone persisting for more than 8 weeks in a patient with current or previous treatment with antiresorptive agents but without a history of radiation therapy to the jaws.²

Radiologically, the bone can be normal in the early stages. In advanced cases, a mixture of bone sclerosis and rarefaction occurs. Bone sequestra are seen more commonly in the mandible and often the sequestra appear as radiopaque islands of bone in radiolucent regions (Figure 1). Large sequestrum may weaken the bone and promote pathologic fracture. In contrast to osteomyelitis, there is no periosteal bone reaction in most cases.³

Magnetic resonance imaging can detect the early stage of the disease. The bone marrow shows low signal intensity on T1-weighted images and high signal intensity on T2-weighted images.⁴

In patients with osteonecrosis of the jaw, regular preventive dental care is of foremost importance.² Conservative treatments include maintaining good oral hygiene, eliminating sources of infection, and application of antimicrobial mouth rinses. Because of the reduced healing capacity of bone, surgical intervention should be limited to a minimal debridement of the exposed necrotic bone. Systemic antibiotics and hyperbaric oxygen may be used as adjunctive therapies.^{1,2}




Figure 1. A 61-year-old woman with osteochemonecrosis of the right mandible. The patient was treated with intravenous bisphosphonates for metastatic breast cancer. A, Cone beam computed tomography (CT) reformatted thin-slice panoramic image shows large region of bone destruction in the right body of the mandible with diffuse sclerotic bone reaction and presence of multiple sequestra. Axial (B) and coronal (C) cone beam CT images show increase in density and size of the right mandible compared with the left side. Arrows (B and C) point to a large sequestrum. There is no evidence of periosteal bone formation.

¹ Division of Oral Medicine, Diagnosis and Radiology, Department of Oral Health Practice, University of Kentucky College of Dentistry, Lexington, KY, USA

Received: March 23, 2020; accepted: March 26, 2020

Corresponding Author:

Galal Omami, BDS, MSc, MDentSc, FRCD(C), Division of Oral Medicine, Diagnosis and Radiology, Department of Oral Health Practice, University of Kentucky College of Dentistry, 800 Rose Street, Room MN-320, Lexington, KY 40536, USA.

Email: galal.omami@uky.edu



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).


Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article

ORCID iD

Galal Omami  <https://orcid.org/0000-0003-1187-1816>

References

1. Ruggiero SL, Dodson TB, Fantasia J, et al. American Association of Oral and Maxillofacial Surgeons position paper on medication-related osteonecrosis of the jaw—2014 update. *J Oral Maxillofac Surg.* 2014;72(10):1938-1956.
2. Nicolatou-Galitis O, Schiødt M, Mendes RA, et al. Medication-related osteonecrosis of the jaw: definition and best practice for prevention, diagnosis, and treatment. *Oral Surg Oral Med Oral Pathol Oral Radiol.* 2019;127(2):117-135.
3. Baba A, Goto TK, Ojiri H, et al. CT imaging features of antiresorptive agent-related osteonecrosis of the jaw/medication-related osteonecrosis of the jaw. *Dentomaxillofac Radiol.* 2018;47(4):20170323.
4. Krishnan A, Arslanoglu A, Yildirim N, et al. Imaging findings of bisphosphonate-related osteonecrosis of the jaw with emphasis on early magnetic resonance imaging findings. *J Comput Assist Tomogr.* 2009;33(2):298-304.