



8-4-2010

# Diaphragm weakness and mechanical ventilation – what's the critical issue?

Leigh Ann Callahan

*University of Kentucky*, lacall2@email.uky.edu

Gerald S. Supinski

*University of Kentucky*, gsupi2@email.uky.edu

**Click here to let us know how access to this document benefits you.**

Follow this and additional works at: [https://uknowledge.uky.edu/internalmedicine\\_facpub](https://uknowledge.uky.edu/internalmedicine_facpub)

 Part of the [Medicine and Health Sciences Commons](#)

## Repository Citation

Callahan, Leigh Ann and Supinski, Gerald S., "Diaphragm weakness and mechanical ventilation – what's the critical issue?" (2010).

*Internal Medicine Faculty Publications*. 21.

[https://uknowledge.uky.edu/internalmedicine\\_facpub/21](https://uknowledge.uky.edu/internalmedicine_facpub/21)

This Response or Comment is brought to you for free and open access by the Internal Medicine at UKnowledge. It has been accepted for inclusion in Internal Medicine Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

---

**Diaphragm weakness and mechanical ventilation – what's the critical issue?**

**Notes/Citation Information**

Published in *Critical Care*, v. 14, 187.

© 2010 BioMed Central Ltd

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Digital Object Identifier (DOI)**

<http://dx.doi.org/10.1186/cc9189>

## COMMENTARY

# Diaphragm weakness and mechanical ventilation - what's the critical issue?

Leigh Ann Callahan\* and Gerald S Supinski

See related research by Hermans *et al.*, <http://ccforum.com/content/14/4/R127>

### Abstract

While animal studies indicate that controlled mechanical ventilation (MV) induces diaphragm weakness and myofiber atrophy, there are no data in humans that confirm MV *per se* produces diaphragm weakness. Whether or not diaphragm weakness results from MV, sepsis, corticosteroids, hyperglycemia, or a combination of these factors, however, is not the most important issue raised by the recent study from Hermans and colleagues. This study makes an important contribution by providing additional evidence that many critically ill patients have profound diaphragm weakness. If diaphragm weakness of this magnitude is present in most mechanically ventilated patients, a strong argument can be made that respiratory muscle weakness is a major contributor to respiratory failure.

Numerous animal studies indicate that controlled mechanical ventilation (MV) induces diaphragm weakness and myofiber atrophy, but no data in humans confirm MV *per se* produces diaphragm weakness. In their recent publication, Hermans and colleagues [1] used an objective nonvolitional technique of bilateral anterolateral magnetic stimulation of the phrenic nerves to measure twitch transdiaphragmatic pressure (TwPdi) and assessed the degree of diaphragm weakness in ten critically ill mechanically ventilated patients. Importantly, they found an average TwPdi value of  $11.5 \pm 3.9$  cm water, which represents a 70% reduction in diaphragm strength when compared to normal individuals. In their analysis, the authors suggest that duration of MV is associated with decreased diaphragm force generation. This interpretation, however, may be misleading. Importantly, nine of

the ten patients in their study were septic and eight out of ten received corticosteroids. Moreover, no information regarding glucose control is included. Since sepsis, corticosteroid use and hyperglycemia are major risk factors for acquired weakness [2,3], it seems plausible that these conditions also contributed to the diaphragm weakness observed.

Whether or not diaphragm weakness results from sepsis, respiratory muscle unloading from MV, corticosteroids, hyperglycemia, or a combination of these factors, however, is not the most important issue raised by this study. Hermans and colleagues [1] should be congratulated because their study makes an important contribution by providing additional evidence that many critically ill patients have profound diaphragm weakness [4,5]. If diaphragm weakness of this magnitude is present in most mechanically ventilated patients, a strong argument can be made that respiratory muscle weakness is a major contributor to respiratory failure.

Consider this - mechanical ventilators are not artificial lungs but simply machines that substitute for the respiratory pump. The fact is that the respiratory pump does not have an unlimited capacity; if it did, theoretically, some patients would require augmented oxygen delivery and/or end expiratory pressure but none would require MV. For patients with normal respiratory muscle function, respiratory failure usually occurs when the respiratory workload becomes too high for the normal pump to maintain ventilation. In principal, any reduction in pump function below normal should increase the propensity for respiratory failure to develop, with the level of respiratory workload required to induce respiratory failure directly related to the level of pump function. Specifically, the lower the pump function, the lower the respiratory workload required to induce respiratory failure. If this concept is correct, the level of respiratory muscle dysfunction reported by Hermans and colleagues should be a major contributor to respiratory failure.

Unless the patient has a known neuromuscular disorder, critical care physicians often overlook diaphragm weakness as an important factor contributing to respiratory failure and weaning difficulties in a significant

\*Correspondence: [lacall2@email.uky.edu](mailto:lacall2@email.uky.edu)  
Division of Pulmonary, Critical Care and Sleep Medicine, Department of Internal Medicine, University of Kentucky, 740 South Limestone, Lexington, KY 40536, USA

number of patients. We focus on improving lung function, perhaps because conceptually this is easier to understand, easier to assess using chest radiographs, and, for the most part, the treatment options are relatively straightforward. On the other hand, if we recognize that diaphragm weakness is present, what can we do? Regrettably, the current approach to diaphragm weakness in critically ill patients is similar to the approach to pulmonary hypertension 30 years ago. Physicians once believed pulmonary hypertension was extremely rare, and there were no treatments. Today, we recognize that pulmonary hypertension is more prevalent, we have better tools to diagnose this problem, and we have a growing ensemble of pharmacological agents to treat patients with this disorder. To make such progress in dealing with the problem of respiratory muscle dysfunction in critically ill patients, we need better diagnostic tools, a better understanding of the pathophysiology of this disorder and, most importantly, we need to develop rational, specific and effective treatments. Once these goals are met, we may be able to substantially shorten the duration of MV in ICU patients and improve long-term outcomes in this growing population of patients.

#### Abbreviations

MV = mechanical ventilation; TwPdi = twitch transdiaphragmatic pressure.

#### Competing interests

The authors declare that they have no competing interests.

#### Acknowledgements

Our work is supported by NIH grants R01 HL80609 (LAC), R01 HL80429 (GS), R01 HL081525 (GS), RC1 HL100239 (GS).

Published: 4 August 2010

#### References

1. Hermans G, Agten A, Testelmans D, Decramer M, Gayan-Ramirez G: **Increased duration of mechanical ventilation is associated with decreased diaphragmatic force: a prospective observational study.** *Crit Care* 2010, **14**:R127.
2. De Jonghe B, Lacherade JC, Sharshar T, Outin H: **Intensive care unit-acquired weakness: risk factors and prevention.** *Crit Care Med* 2009, **37**:S309-315.
3. Griffiths RD, Hall JB: **Intensive care unit-acquired weakness.** *Crit Care Med* 2010, **38**:779-787.
4. Watson AC, Hughes PD, Louise Harris M, Hart N, Ware RJ, Wendon J, Green M, Moxham J: **Measurement of twitch transdiaphragmatic, esophageal, and endotracheal tube pressure with bilateral anterolateral magnetic phrenic nerve stimulation in patients in the intensive care unit.** *Crit Care Med* 2001, **29**:1325-1331.
5. Laghi F, Cattapan SE, Jubran A, Parthasarathy S, Warshawsky P, Choi YS, Tobin MJ: **Is weaning failure caused by low-frequency fatigue of the diaphragm?** *Am J Respir Crit Care Med* 2003, **167**:120-127.

doi:10.1186/cc9189

**Cite this article as:** Callahan LA, Supinski GS: **Diaphragm weakness and mechanical ventilation - what's the critical issue?** *Critical Care* 2010, **14**:187.