

University of Kentucky

UKnowledge

Mathematics Faculty Publications

Mathematics

3-29-2018

Rook Placements and Jordan Forms of Upper-Triangular Nilpotent Matrices

Martha Yip

University of Kentucky, mpyi222@uky.edu

Follow this and additional works at: https://uknowledge.uky.edu/math_facpub



Part of the [Mathematics Commons](#)

[Right click to open a feedback form in a new tab to let us know how this document benefits you.](#)

Repository Citation

Yip, Martha, "Rook Placements and Jordan Forms of Upper-Triangular Nilpotent Matrices" (2018). *Mathematics Faculty Publications*. 30.

https://uknowledge.uky.edu/math_facpub/30

This Article is brought to you for free and open access by the Mathematics at UKnowledge. It has been accepted for inclusion in Mathematics Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Rook Placements and Jordan Forms of Upper-Triangular Nilpotent Matrices

Notes/Citation Information

Published in *The Electronic Journal of Combinatorics*, v. 25, issue 1, paper #P1.68, p. 1-25.

The publisher has granted the permission for posting the article here.

Rook placements and Jordan forms of upper-triangular nilpotent matrices

Martha Yip*

Department of Mathematics
University of Kentucky
Lexington, U.S.A.

martha.yip@uky.edu

Submitted: Mar 14, 2017; Accepted: Jan 17, 2018; Published: Mar 29, 2018
Mathematics Subject Classifications: 05E15, 05A19

Abstract

The set of n by n upper-triangular nilpotent matrices with entries in a finite field \mathbb{F}_q has Jordan canonical forms indexed by partitions $\lambda \vdash n$. We present a combinatorial formula for computing the number $F_\lambda(q)$ of matrices of Jordan type λ as a weighted sum over standard Young tableaux. We construct a bijection between paths in a modified version of Young's lattice and non-attacking rook placements, which leads to a refinement of the formula for $F_\lambda(q)$.

Keywords: nilpotent matrices, finite fields, Jordan form, rook placements, Young tableaux, set partitions.

1 Introduction

In the beautiful paper *Variations on the Triangular Theme* [7], Kirillov studied various structures on the set of triangular matrices. Let $G = G_n(\mathbb{F}_q)$ denote the group of n by n invertible upper-triangular matrices over the field \mathbb{F}_q having q elements, and let $\mathfrak{g} = \mathfrak{g}_n(\mathbb{F}_q) = \text{Lie}(G_n(\mathbb{F}_q))$ denote the corresponding Lie algebra of n by n upper-triangular nilpotent matrices over \mathbb{F}_q . The problem of determining the set $\mathcal{O}_n(\mathbb{F}_q)$ of adjoint G -orbits in \mathfrak{g} remains challenging, and a more tractable task is to study a decomposition of $\mathcal{O}_n(\mathbb{F}_q)$ via the Jordan canonical form. Let $\lambda \vdash n$ be a partition of n with r positive parts

*Simons Collaboration Grant 429920.

$\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_r > 0$, and let

$$J_\lambda = J_{\lambda_1} \oplus J_{\lambda_2} \oplus \dots \oplus J_{\lambda_r}, \quad \text{where} \quad J_i = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 1 & \cdots & 0 & 0 \\ 0 & 0 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & 0 & \cdots & 0 & 0 \end{bmatrix}_{i \times i}$$

is the i by i elementary Jordan matrix with all eigenvalues equal to zero. If $X \in \mathfrak{g}_n(\mathbb{F}_q)$ is similar to J_λ under $GL_n(\mathbb{F}_q)$, then X is said to have *Jordan type* λ . Each conjugacy class contains a unique Jordan matrix J_λ , so these classes are indexed by the partitions of n . Evidently, the Jordan type of X depends only on its adjoint G -orbit.

Let $\mathfrak{g}_{n,\lambda}(\mathbb{F}_q) \subseteq \mathfrak{g}_n(\mathbb{F}_q)$ be the set of upper-triangular nilpotent matrices of fixed Jordan type λ , and let

$$F_\lambda(q) = |\mathfrak{g}_{n,\lambda}(\mathbb{F}_q)|. \quad (1)$$

Springer showed that $\mathfrak{g}_{n,\lambda}(\mathbb{F}_q)$ is an algebraic manifold with f^λ irreducible components, where f^λ is the number of standard Young tableaux of shape λ , and each of which has dimension $\binom{n}{2} - n_\lambda$, where n_λ is an integer defined in Equation 10. These quantities appear in the study of $F_\lambda(q)$.

In Section 2, we show that the numbers $F_\lambda(q)$ satisfy a simple recurrence equation, and that they are polynomials in q with integer coefficients. As a consequence of the recurrence equation in Theorem 8, it follows that the coefficient of the highest degree term in $F_\lambda(q)$ is f^λ , and $\deg F_\lambda(q) = \binom{n}{2} - n_\lambda$. Equation (9) is a combinatorial formula for $F_\lambda(q)$ as a sum over standard Young tableaux of shape λ that can be derived from the recurrence equation.

The cases $F_{(1^n)}(q) = 1$ and $F_{(n)}(q) = (q-1)^{n-1}q^{\binom{n-1}{2}}$ are readily computed, since the matrix in $\mathfrak{g}_n(\mathbb{F}_q)$ of Jordan type (1^n) is the matrix of zero rank, and the matrices in $\mathfrak{g}_n(\mathbb{F}_q)$ of Jordan type (n) are the matrices of rank equal to $n-1$. Section 2 concludes with explicit formulas for $F_\lambda(q)$ in several other special cases of λ , including hook shapes, two-rowed partitions and two-columned partitions.

In Section 3, we explore a connection of $F_\lambda(q)$ with rook placements. In their study of a formula of Frobenius, Garsia and Remmel [4] introduced the *q-rook polynomial*

$$R_{B,k}(q) = \sum_{c \in \mathcal{C}(B,k)} q^{\text{inv}(c)},$$

which is a sum over the set $\mathcal{C}(B,k)$ of non-attacking placements of k rooks on a Ferrers board B , and $\text{inv}(c)$, defined in Equation (13), is the number of inversions of c . In the case when $B = B_n$ is the staircase-shaped board, Garsia and Remmel showed that $R_{B_n,k}(q) = S_{n,n-k}(q)$ is a *q-Stirling number of the second kind*. These numbers are defined by the recurrence equation

$$S_{n,k}(q) = q^{k-1}S_{n-1,k-1}(q) + [k]_q S_{n-1,k}(q) \quad \text{for} \quad 0 \leq k \leq n,$$

with initial conditions $S_{0,0}(q) = 1$, and $S_{n,k}(q) = 0$ for $k < 0$ or $k > n$.

It was shown by Solomon [12] that non-attacking placements of k rooks on rectangular $m \times n$ boards are naturally associated to m by n matrices with rank k over \mathbb{F}_q . By identifying a Ferrers board B inside an n by n grid with the entries of an n by n matrix, Haglund [5] generalized Solomon's result to the case of non-attacking placements of k rooks on Ferrers boards, and obtained a formula for the number of n by n matrices with rank k whose support is contained in the Ferrers board region. A special case of Haglund's formula shows that the number of n by n nilpotent upper-triangular matrices of rank k is

$$P_{B_n,k}(q) = (q-1)^k q^{\binom{n}{2}-k} R_{B_n,k}(q^{-1}). \quad (2)$$

Now, a matrix in $\mathfrak{g}_{n,\lambda}(\mathbb{F}_q)$ has rank $n - \ell(\lambda)$, where $\ell(\lambda)$ is the number of parts of λ , so the number of matrices in $\mathfrak{g}_n(\mathbb{F}_q)$ with rank k is

$$P_{B_n,k}(q) = \sum_{\lambda \vdash n: \ell(\lambda)=n-k} F_\lambda(q). \quad (3)$$

Given Equations 2 and 3, it would be natural to ask whether it is possible to partition the placements $\mathcal{C}(B_n, k)$ into disjoint subsets so that the sum over each subset of placements gives $F_\lambda(q)$. A central goal of this paper is to study the connection between upper-triangular nilpotent matrices over \mathbb{F}_q and non-attacking rook placements on the staircase-shaped board B_n . Theorem 28 shows that there is a weight-preserving bijection Φ between rook placements on B_n and paths in a graph \mathcal{Z} (see Figure 5), which is a multi-edged version of Young's lattice. As a result, we obtain Corollary 30, which gives a formula for $F_\lambda(q)$ as a sum over certain rook placements that can be viewed as a generalization of Haglund's formula in Equation (2).

There is a classically known bijection between rook placements in $\mathcal{C}(B_n, k)$ and set partitions of $[n]$ with $n - k$ parts, so it is logical to next study the connection between $F_\lambda(q)$ and set partitions. We do this in Section 4. Theorem 34 describes the construction of a new (weight-preserving) bijection Ψ between rook placements and set partitions. These bijections allow us to refine Equation (9) to a sum over set partitions (or rook placements). We also discuss the significance of the polynomials $F_C(q)$ indexed by rook placements in a special case.

This paper is the full version of the extended abstract [15].

2 Formulas for $F_\lambda(q)$

The recurrence equation for $F_\lambda(q)$ in Theorem 8 can be found in [1, Division Theorem], where Borodin considers the matrices as particles of a certain mass and studies the asymptotic behaviour of the formula. A preliminary version of the idea first appeared in [7]. In this section, we give an elementary proof of the formula, and investigate some of the combinatorial properties of $F_\lambda(q)$.

2.1 The recurrence equation for $F_\lambda(q)$

A *partition* λ of a nonnegative integer n , denoted by $\lambda \vdash n$, is a non-increasing sequence of nonnegative integers $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n \geq 0$ with $|\lambda| = \sum_{i=1}^n \lambda_i = n$. If λ has r positive parts, write $\ell(\lambda) = r$. A partition λ can be represented by its Ferrers diagram in the English notation, which is an array of λ_i boxes in the i th row, with the boxes justified upwards and to the left. Let λ'_j denote the size of the j th column of λ .

Young's lattice \mathcal{Y} is the lattice of partitions ordered by the inclusion of their Ferrers diagrams; that is, $\mu \leq \lambda$ if and only if $\mu_i \leq \lambda_i$ for every i . In particular, μ is covered by λ in the Hasse diagram of \mathcal{Y} and we write $\mu < \lambda$ if the Ferrers diagram of λ can be obtained by adding a box to the Ferrers diagram of μ . See Figure 1.

Example 1. The partition

$$\lambda = (4, 2, 2, 1) \vdash 9 \quad \text{has diagram} \quad \begin{array}{cccc} \square & \square & \square & \square \\ \square & \square & & \\ \square & \square & & \\ \square & & & \end{array}$$

and columns $\lambda'_1 = 4, \lambda'_2 = 3, \lambda'_3 = 1, \lambda'_4 = 1$.

Lemma 2. Let $\lambda \vdash n$ be a partition whose Ferrers diagram has r rows and c columns. The Jordan matrix J_λ satisfies

$$\text{rank}(J_\lambda^k) = \begin{cases} \lambda'_{k+1} + \dots + \lambda'_c, & \text{if } 0 \leq k < c, \\ 0, & \text{if } k \geq c. \end{cases}$$

Proof. The i by i elementary Jordan matrix J_i has $\text{rank}(J_i^k) = i - k$ if $0 \leq k \leq i$, and its rank is zero otherwise, so the Jordan matrix $J_\lambda = J_{\lambda_1} \oplus \dots \oplus J_{\lambda_r}$ has

$$\text{rank}(J_\lambda^k) = \sum_{i=1}^r \text{rank}(J_{\lambda_i}^k) = \sum_{i:\lambda_i \geq k} \text{rank}(J_{\lambda_i}^k) = \sum_{j=k+1}^c \lambda'_j,$$

for $0 \leq k < c$, which is the number of boxes in the last $c - k$ columns of λ . \square

Remark 3. Matrices which are similar have the same rank, so if $X \in \mathfrak{g}_{n,\lambda}(\mathbb{F}_q)$, then $\text{rank}(X^k) = \text{rank}(J_\lambda^k)$ for all $k \geq 0$. Conversely, let $\lambda, \nu \vdash n$. It follows from Lemma 2 that $\text{rank}(J_\lambda^k) = \text{rank}(J_\nu^k)$ for all $k \geq 0$ if and only if $\lambda = \nu$. Thus if $X \in \mathfrak{g}_n(\mathbb{F}_q)$ is a matrix such that $\text{rank}(X^k) = \text{rank}(J_\lambda^k)$ for all $k \geq 0$, then X is similar to J_λ .

Example 4. If a matrix $X \in \mathfrak{g}_n(\mathbb{F}_q)$ has Jordan type $\lambda = (4, 2, 2, 1)$, then $\text{rank}(X) = 5$, $\text{rank}(X^2) = 2$, $\text{rank}(X^3) = 1$, and $\text{rank}(X^4) = 0$.

If $X \in \mathfrak{g}_n(\mathbb{F}_q)$ is a matrix of the form

$$X = \begin{bmatrix} J_\mu & \mathbf{v} \\ \mathbf{0} & 0 \end{bmatrix},$$

where $\mu \vdash n - 1$, and $\mathbf{v} = [v_1, \dots, v_{n-1}]^T \in \mathbb{F}_q^{n-1}$, then the first order leading principal submatrix of X^k is J_μ^k , and for $1 \leq k \leq n$, we define column vectors $\mathbf{v}^k = [v_1^k, \dots, v_{n-1}^k]^T \in \mathbb{F}_q^{n-1}$ by

$$X^k = \begin{bmatrix} J_\mu^k & \mathbf{v}^k \\ \mathbf{0} & 0 \end{bmatrix}.$$

For $i \geq 1$, let $\alpha_i = \mu_1 + \dots + \mu_i$ be the sum of the first i parts of μ . The (i, j) th entry of J_μ^k is nonzero if and only if $j = i + k$, and $i, i + k \leq \alpha_b$ for all $b \geq 1$. It follows from this that

$$v_i^k = \begin{cases} v_{i+k-1}, & \text{if } i, i+k-1 \leq \alpha_b \text{ for all } b \geq 1, \\ 0, & \text{otherwise.} \end{cases} \quad (4)$$

There is a simple way to visualize the vectors \mathbf{v}^k , which we illustrate with an example.

Example 5. Let $\mu = (4, 2, 1, 1)$, so that $\alpha_1 = 4, \alpha_2 = 6, \alpha_3 = 7$, and $\alpha_4 = 8$. Let

$$X = \begin{array}{cccc|cccc} 0 & 1 & 0 & 0 & & & & v_1 \\ 0 & 0 & 1 & 0 & & & & v_2 \\ 0 & 0 & 0 & 1 & & & & v_3 \\ 0 & 0 & 0 & 0 & & & & v_4 \\ \hline & & & & 0 & 1 & & v_5 \\ & & & & 0 & 0 & & v_6 \\ \hline & & & & & & 0 & v_7 \\ \hline & & & & & & & 0 & v_8 \\ \hline & & & & & & & & 0 \end{array} \quad \text{so that} \quad X^2 = \begin{array}{cccc|cccc} 0 & 0 & 1 & 0 & & & & v_2 \\ 0 & 0 & 0 & 1 & & & & v_3 \\ 0 & 0 & 0 & 0 & & & & v_4 \\ 0 & 0 & 0 & 0 & & & & 0 \\ \hline & & & & 0 & 0 & & v_6 \\ & & & & 0 & 0 & & 0 \\ \hline & & & & & & 0 & 0 \\ \hline & & & & & & & 0 & 0 \\ \hline & & & & & & & & 0 \end{array}.$$

We may visualize the vectors \mathbf{v} and \mathbf{v}^2 as fillings of the Ferrers diagram for μ :

$$\mathbf{v} = \begin{array}{|c|c|c|c|} \hline v_4 & v_3 & v_2 & v_1 \\ \hline v_6 & v_5 & & \\ \hline v_7 & & & \\ \hline v_8 & & & \\ \hline \end{array} \quad \text{and} \quad \mathbf{v}^2 = \begin{array}{|c|c|c|c|} \hline 0 & v_4 & v_3 & v_2 \\ \hline 0 & v_6 & & \\ \hline 0 & & & \\ \hline 0 & & & \\ \hline \end{array}.$$

This way, a basis of $\ker X^k$ is the set of vectors filling the first k columns of the diagram.

Lemma 6. If $X \in \mathfrak{g}_{n,\lambda}(\mathbb{F}_q)$ and its first order leading principal submatrix $Y \in \mathfrak{g}_{n-1,\mu}(\mathbb{F}_q)$, then $\lambda \geq \mu$.

Proof. We first consider the case $Y = J_\mu$. If μ has s parts, let $\alpha_i = \mu_1 + \dots + \mu_i$ for $1 \leq i \leq s$. Then

$$\text{rank}(X^k) - \text{rank}(J_\mu^k) = \begin{cases} 0, & \text{if } v_{\alpha_i} = 0 \text{ for all } i \text{ such that } \mu_i \geq k, \\ 1, & \text{otherwise.} \end{cases} \quad (5)$$

Let $c \leq n$ be the smallest positive integer for which $\text{rank}(X^c) - \text{rank}(J_\mu^c) = 0$. Then Equation (5) implies that

$$\text{rank}(X^k) - \text{rank}(J_\mu^k) = \begin{cases} 0, & \text{if } k \geq c, \\ 1, & \text{if } k < c. \end{cases}$$

Together with Lemma 2, we deduce that

$$\lambda'_k - \mu'_k = (\text{rank}(X^{k-1}) - \text{rank}(X^k)) - (\text{rank}(J_\mu^{k-1}) - \text{rank}(J_\mu^k)) = \begin{cases} 1, & \text{if } k = c, \\ 0, & \text{if } k \neq c. \end{cases}$$

Therefore, $\lambda \succ \mu$ in the case $Y = J_\mu$.

In the general case where Y is any matrix of Jordan type μ , then $\text{rank}(Y^k) = \text{rank}(J_\mu^k)$ for all $k \geq 0$, so the argument is the same. \square

Let λ be the partition whose diagram is obtained by adding a box to the i th row and j th column of the diagram of the partition μ . Define the coefficient

$$c_{\mu,\lambda}(q) = \begin{cases} q^{|\mu| - \mu'_j}, & \text{if } j = 1, \\ q^{|\mu| - \mu'_{j-1}} (q^{\mu'_{j-1} - \mu'_j} - 1), & \text{if } j \geq 2. \end{cases} \quad (6)$$

Note that in the case $j \geq 2$, we have $\mu'_{j-1} - \mu'_j \geq 1$.

Lemma 7. *Let Y be an upper-triangular nilpotent matrix of Jordan type $\mu \vdash n - 1$. If $\mu \triangleleft \lambda$, then there are $c_{\mu,\lambda}(q)$ upper-triangular nilpotent matrices X of Jordan type λ whose first order leading principal submatrix is Y .*

Proof. By similarity, it suffices to consider the case $Y = J_\mu = J_{\mu_1} \oplus \cdots \oplus J_{\mu_m}$, where $\ell(\mu) = m$. Suppose X is a matrix of the form

$$X = \begin{bmatrix} J_\mu & \mathbf{v} \\ \mathbf{0} & 0 \end{bmatrix}$$

of Jordan type λ such that λ is obtained by adding a box to μ in the i th row and j th column.

First consider the case $j \geq 2$. Following the proof of Lemma 6, we know that j is the unique integer where $\text{rank}(X^{j-1}) = \text{rank}(J_\mu^{j-1}) + 1$, and $\text{rank}(X^j) = \text{rank}(J_\mu^j)$. In order to satisfy the first condition, the entries in the vector \mathbf{v}^{j-1} corresponding to the boxes in the $(j-1)$ th column and rows $\geq i$ must not simultaneously be zero (refer to Equation (4) and Example 5), while in order to satisfy the second condition, the entries in the vector \mathbf{v}^j corresponding to the boxes in the j th column of μ must all be zero. The remaining $n - 1 - \mu'_{j-1}$ entries of the vector \mathbf{v} are free to be any element in \mathbb{F}_q , so there are

$$q^{n-1-\mu'_{j-1}} (q^{\mu'_{j-1} - \mu'_j} - 1)$$

possible matrices X whose leading principal submatrix is J_μ .

The case $j = 1$ is simpler. The necessary and sufficient condition that X and J_μ must satisfy is that $\text{rank}(X^k) = \text{rank}(J_\mu^k)$ for all $k \geq 1$, so the entries in the vector \mathbf{v} corresponding to the boxes in the first column of the diagram for \mathbf{v}^1 must all be zero, while the remaining $n - 1 - \mu'_1$ entries are free to be any element in \mathbb{F}_q , so there are $q^{n-1-\mu'_1}$ matrices X whose leading principal submatrix is J_μ in this case. \square

λ , where $\pi^{(i)}$ is a partition of i . To simplify notation, let $\epsilon_i(q) = c_{\pi^{(i-1)}, \pi^{(i)}}(q)$. Theorem 8 is equivalently re-phrased as

$$F_\lambda(q) = \sum_{P \in \mathcal{P}_Y(\lambda)} F_P(q), \quad (7)$$

where the weight of the path P is $F_P(q) = \prod_{i=1}^n \epsilon_i(q)$.

The set of paths $\mathcal{P}_Y(\lambda)$ is in bijection with the set $\text{SYT}(\lambda)$ of standard Young tableaux of shape λ , so we can also give an equation for $F_\lambda(q)$ as a sum over standard Young tableaux.

A *standard Young tableau* T of shape λ is a filling of the Ferrers diagram of $\lambda \vdash n$ with the integers $1, \dots, n$ such that the integers increase weakly along each row and strictly along each column. For $1 \leq i \leq n$, Let $T^{(i)}$ denote the Young tableau of shape $\lambda^{(i)}$ consisting of the boxes containing $1, \dots, i$, and define weights

$$T^{(i)}(q) = \begin{cases} q^{i-\ell(\lambda^{(i)})}, & \text{if the } i\text{th box is in the first column,} \\ q^{i-\lambda_j^{(i)'} - q^{i-1-\lambda_{j-1}^{(i)'}}} & \text{if the } i\text{th box is in the } j\text{th column, } j \geq 2. \end{cases} \quad (8)$$

Then

$$F_\lambda(q) = \sum_{T \in \text{SYT}(\lambda)} F_T(q), \quad (9)$$

where the weight of the standard Young tableau T is $F_T(q) = \prod_{i=1}^n T^{(i)}(q)$.

2.2 Properties of $F_\lambda(q)$

Several properties of $F_\lambda(q)$ follow readily from Theorem 8. For $\lambda \vdash n$, let

$$n_\lambda = \sum_{i \geq 1} (i-1)\lambda_i = \sum_{b \in \lambda} \text{coleg}(b), \quad (10)$$

where if a box $b \in \lambda$ lies in the i th row of λ , then $\text{coleg}(b) = i-1$.

Corollary 10. *Let $\lambda \vdash n$. As a polynomial in q ,*

$$\deg F_\lambda(q) = \binom{n}{2} - n_\lambda.$$

Moreover, the coefficient of the highest degree term in $F_\lambda(q)$ is f^λ , the number of standard Young tableaux of shape λ .

Proof. Suppose $P = (\emptyset = \pi^{(0)}, \pi^{(1)}, \dots, \pi^{(n)} = \lambda)$ is a path in \mathcal{Y} such that $\pi^{(k)}$ is obtained by adding a box to the i th row and j th column of $\pi^{(k-1)}$. Then $\deg c_{\pi^{(k-1)}, \pi^{(k)}}(q) = k-i$, and therefore

$$\deg F_P(q) = \sum_{k=1}^n \deg c_{\pi^{(k-1)}, \pi^{(k)}}(q) = \sum_{k=1}^n k - \sum_{k \geq 1} k\lambda_k = \binom{n}{2} - n_\lambda.$$

In particular, every polynomial $F_P(q)$ arising from a path $P \in \mathcal{P}_Y(\lambda)$ has the same degree, so $\deg F_\lambda(q) = \binom{n}{2} - n_\lambda$. Moreover, each $F_P(q)$ is monic, so the coefficient of the highest degree term in $F_\lambda(q)$ is the number of paths in \mathcal{Y} from \emptyset to λ , which is f^λ . \square

Corollary 11. Let $\lambda \vdash n$. The multiplicity of the factor $q - 1$ in $F_\lambda(q)$ is $n - \ell(\lambda)$.

Proof. The weight $c_{\pi^{(k-1)}, \pi^{(k)}}(q)$ corresponding to the k th step in the path P contributes a single factor of $q - 1$ to $F_P(q)$ if and only if the k th box added is not in the first column of λ . Therefore, the multiplicity of $q - 1$ in $F_P(q)$ is $n - \ell(\lambda)$, and it follows that the multiplicity of $q - 1$ in $F_\lambda(q)$ is $n - \ell(\lambda)$. \square

Example 12. There are two partitions of 4 with two parts, namely $(3, 1)$ and $(2, 2)$.

There are three paths from \emptyset to $(3, 1)$ in \mathcal{Y} , giving

$$\begin{aligned} F_{(3,1)}(q) &= (q - 1) \cdot (q - 1)q \cdot q^2 + (q - 1) \cdot q \cdot (q - 1)q^2 + (q^2 - 1) \cdot (q - 1)q^2 \\ &= (q - 1)^2 (3q^3 + q^2), \end{aligned}$$

and there are two paths from \emptyset to $(2, 2)$ in \mathcal{Y} , giving

$$\begin{aligned} F_{(2,2)}(q) &= (q - 1) \cdot q \cdot (q - 1)q + (q^2 - 1) \cdot (q - 1)q \\ &= (q - 1)^2 (2q^2 + q). \end{aligned}$$

Summing these gives a shift of the q -Stirling polynomial $(q - 1)^2 q^4 S_{4,2}(q^{-1}) = (q - 1)^2 (3q^3 + 3q^2 + q)$.

2.3 Explicit formulas

In this section, we derive non-recursive formulas for some special cases of λ . Previously, we have noted the simple cases $F_{(1^n)} = 1$ and $F_{(n)} = q^{\binom{n-1}{2}}(q - 1)^{n-1}$.

Proposition 13 (Hook shapes). Let $n > k \geq 2$, and let $\lambda = (n - k + 1, 1^{k-1})$ be a hook-shaped partition of n with $\ell(\lambda) = k$ parts. Then

$$F_\lambda(q) = (q - 1)^{n-k} \sum_{i=0}^{k-1} \binom{n-i-1}{k-i-1} q^{\alpha-i}, \quad \text{where } \alpha = \binom{n-1}{2} - \binom{k-1}{2}.$$

Proof. We make use of Equation (7). We enumerate paths from \emptyset to λ according to the first time a box is added to the second column, so for $0 \leq r \leq k - 1$, let S_r be the set of paths in the sublattice $[\emptyset, \lambda]$ which contains the edge $((1, 1^r), (2, 1^r))$. Such paths are formed by the concatenation of the unique path between \emptyset and $\nu = (2, 1^r)$, which has weight $q^{r+1} - 1$, with any path in the sublattice $[\nu, \lambda]$. The sublattice $[\nu, \lambda]$ is the Cartesian product of a $(n - k)$ -chain and a $(k - r - 1)$ -chain, so it forms a rectangular grid, and therefore $|S_r| = \binom{n-r-1}{k-r-1}$. Notice that in any sublattice of the form

$$\begin{array}{ccc} (a, 1^{b+1}) & \xrightarrow{(q-1)q^{a+b}} & (a+1, 1^{b+1}) \\ \uparrow q^{a-1} & & \uparrow q^a \\ (a, 1^b) & \xrightarrow{(q-1)q^{a+b-1}} & (a+1, 1^b) \end{array}$$

the product of the edge weights is $(q-1)q^{2a+b-1}$ no matter which path is taken from $(a, 1^b)$ to $(a+1, 1^{b+1})$, so it follows that every path from ν to λ has the same weight. By considering the path $(\nu, 21^{r+1}, \dots, 21^{k-1}, 31^{k-1}, \dots, \lambda)$, this weight is easily seen to be $(q-1)^{n-k-1}q^{\alpha-r}$, for $\alpha = \binom{n-1}{2} - \binom{k-1}{2}$. Altogether,

$$F_\lambda(q) = (q-1)^{n-k} \sum_{r=0}^{k-1} \binom{n-r-1}{k-r-1} (q^\alpha + q^{\alpha-1} + \dots + q^{\alpha-r}).$$

For $0 \leq i \leq k-1$, the coefficient of $q^{\alpha-i}$ in $F_\lambda(q)/(q-1)^{n-k}$ is

$$\sum_{r=i}^{k-1} \binom{n-r-1}{k-r-1} = \sum_{u=0}^{k-i-1} \binom{n-k+u}{u} = \binom{n-i-1}{k-i-1},$$

since $\sum_{u=0}^M \binom{N+u}{u} = \binom{N+M+1}{M}$. Therefore,

$$F_\lambda(q) = (q-1)^{n-k} \sum_{i=0}^{k-1} \binom{n-i-1}{k-i-1} q^{\alpha-i},$$

as claimed. □

We next consider the case when λ is a partition with two parts. For $n \geq k \geq 1$, let

$$C_{n,k} = \binom{n+k}{k} - \binom{n+k}{k-1}, \tag{11}$$

and let $C_{n,0} = 1$ for all $n \geq 0$. These generalized Catalan numbers $C_{n,k}$ (see OEIS [10, A009766]) enumerate lattice paths from $(0, 0)$ to (n, k) , using the steps $(1, 0)$ and $(0, 1)$, which do not rise above the line $y = x$. In the remainder of this section, we shall refer to these as *Dyck paths*.

The generalized Catalan numbers satisfy the simple recursive formula $C_{n,k} = C_{n,k-1} + C_{n-1,k}$. Also, these are the usual Catalan numbers $C_n = \frac{1}{n+1} \binom{2n}{n} = C_{n,n} = C_{n,n-1}$ when $k = n$ or $n-1$. These facts will be used in the computations which follow.

Proposition 14 (Partitions with two parts). *If $\lambda = (r, s) \vdash n$ such that $r > s \geq 1$, then*

$$F_{(r,s)}(q) = (q-1)^{r+s-2} q^{\binom{r+s-1}{2} - 2s+1} \sum_{i=0}^s C_{r+s-i,i} q^i.$$

If $r = s$, then

$$F_{(r,r)}(q) = (q-1)^{2r-2} q^{\binom{2r-2}{2}} \sum_{i=0}^{r-1} C_{2r-1-i,i} q^i.$$

Proof. Proceed by induction on $r+s$. The base cases are $F_{(r)}(q) = (q-1)^{r-1} q^{\binom{r-1}{2}}$ for $r \geq 1$ and $F_{(1,1)}(q) = 1$.

$n \setminus k$	0	1	2	3	4	5	6
0	1						
1	1	1					
2	1	2	2				
3	1	3	5	5			
4	1	4	9	14	14		
5	1	5	14	28	42	42	
6	1	6	20	48	90	132	132

Figure 2: The Catalan triangle $C_{n,k}$.

We first handle the case $s = 1$ separately. For $r \geq 2$,

$$\begin{aligned}
 F_{(r,1)}(q) &= q^{r-1}F_{(r)}(q) + (q-1)q^{r-1}F_{(r-1,1)}(q) \\
 &= q^{r-1} \cdot (q-1)^{r-1}q^{\binom{r-1}{2}} + (q-1)q^{r-1} \cdot (q-1)^{r-2}q^{\binom{r-1}{2}-1}((r-1)q+1) \\
 &= (q-1)^{r-1}q^{\binom{r}{2}-1}(rq+1).
 \end{aligned}$$

Next, consider the case $s = r$. For $r \geq 2$,

$$\begin{aligned}
 F_{(r,r)}(q) &= (q-1)q^{2r-3}F_{(r,r-1)}(q) \\
 &= (q-1)q^{2r-3} \cdot (q-1)^{2r-3}q^{\binom{2r-2}{2}-2(r-1)+1} \sum_{i=0}^{r-1} C_{2r-1-i,i}q^i \\
 &= (q-1)^{2r-2}q^{\binom{2r-2}{2}} \sum_{i=0}^{r-1} C_{2r-1-i,i}q^i.
 \end{aligned}$$

The case $s = r - 1$ is obtained as follows. For $r \geq 3$,

$$\begin{aligned}
 F_{(r,r-1)}(q) &= (q-1)q^{2r-4}F_{(r,r-2)}(q) + (q^2-1)q^{2r-4}F_{(r-1,r-1)}(q) \\
 &= (q-1)^{2r-3}q^{2r-4}q^{\binom{2r-4}{2}} \left(q \sum_{i=0}^{r-2} C_{2r-2-i,i}q^i + (q+1) \sum_{i=0}^{r-2} C_{2r-3-i,i}q^i \right) \\
 &= (q-1)^{2r-3}q^{\binom{2r-3}{2}} \left((C_{r,r-2} + C_{r-1,r-2})q^{r-1} \right. \\
 &\quad \left. + \sum_{i=1}^{r-2} (C_{2r-1-i,i-1} + C_{2r-2-i,i-1} + C_{2r-3-i,i})q^i + C_{2r-3,0}q^0 \right).
 \end{aligned}$$

Since $C_{n,n-1} = C_{n,n}$, then $C_{r,r-2} + C_{r-1,r-2} = C_{r,r-1}$. Similarly, we obtain $C_{2r-1-i,i-1} + C_{2r-2-i,i-1} + C_{2r-3-i,i} = C_{2r-1-i,i}$ by applying the recurrence equation for the generalized Catalan numbers. Lastly, $C_{n,0} = 1$ for all $n \geq 0$, thus

$$F_{(r,r-1)}(q) = (q-1)^{2r-3}q^{\binom{2r-3}{2}} \sum_{i=0}^{r-1} C_{2r-1-i,i}q^i,$$




Figure 3: Factors of $q - 1$ are omitted from the edge weights in this sublattice of partitions with at most two rows.

which agrees with the formula for the case $s = r - 1$.

The last case to consider is the general case $r - s \geq 2$ where $s \geq 2$.

$$\begin{aligned}
 F_{(r,s)} &= (q - 1)q^{r+s-3}F_{(r,s-1)} + (q - 1)q^{r+s-2}F_{(r-1,s)} \\
 &= (q - 1)^{r+s-2}q^{r+s-3}q^{\binom{r+s-2}{2}-2s+1} \left(q^2 \sum_{i=0}^{s-1} C_{r+s-1-i,i} q^i + q \sum_{i=0}^s C_{r-1+s-i,i} q^i \right) \\
 &= (q - 1)^{r+s-2}q^{\binom{r+s-1}{2}-2s+1} \left(C_{r+s,0}q^0 + \sum_{i=1}^s (C_{r+s-i,i-1} + C_{r-1+s-i,i}) q^i \right) \\
 &= (q - 1)^{r+s-2}q^{\binom{r+s-1}{2}-2s+1} \sum_{i=0}^s C_{r+s-i,i} q^i.
 \end{aligned}$$

□

The next equation is a formula for $F_{(r,r)}(q)$ with a different factorization.

Proposition 15 (Two equal parts). *Let $\lambda = (r, r) \vdash n$, and $r \geq 1$. Then*

$$F_{(r,r)}(q) = (q - 1)^{2r-2} \sum_{i=0}^{r-1} C_{r-1,r-1-i} q^{2(r-1)^2-i} (q + 1)^i.$$

Proof. The set of paths in the sublattice $[\emptyset, \lambda]$ are in bijection with the set of lattice paths

from $(0, 0)$ to (r, r) . In any sublattice of the form

$$\begin{array}{ccc} (a, b+1) & \xrightarrow{(q-1)q^{a+b}} & (a+1, b+1) \\ \uparrow (q-1)q^{a+b-2} & & \uparrow (q-1)q^{a+b-1} \\ (a, b) & \xrightarrow{(q-1)q^{a+b-1}} & (a+1, b) \end{array}$$

where $b \leq a - 2$, the product of the edge weights is $(q - 1)^2 q^{2a+2b-2}$ no matter which path is taken from (a, b) to $(a + 1, b + 1)$. As for sublattices of the form

$$\begin{array}{ccc} (a, a) & \xrightarrow{(q^2-1)q^{2a-2}} & (a+1, a) \\ \uparrow (q-1)q^{2a-3} & & \uparrow (q-1)q^{2a-2} \\ (a, a-1) & \xrightarrow{(q-1)q^{2a-2}} & (a+1, a-1) \end{array}$$

the product of the edge weights is $(q - 1)^2 q^{4a-4}$ via the lower horizontal edge, versus $(q - 1)^2 q^{4a-5} (q + 1)$ via the upper horizontal edge. It follows that if a path P from \emptyset to λ contains i partitions of the form (a, a) , then it has the weight

$$F_p(q) = (q - 1)^{2r-2} q^{2(r-1)^2-i} (q + 1)^i.$$

Dyck paths may be enumerated according to the points at which they touch the diagonal line $y = x$, and the set of touch points are indexed by compositions $\alpha = (\alpha_1, \dots, \alpha_{i+1}) \vDash r$ where $\alpha_j \geq 1$. The number of Dyck paths from $(0, 0)$ to (r, r) which touch the diagonal exactly i times, not including the initial and the end points, is

$$\sum_{\substack{\alpha \vDash r \\ \ell(\alpha)=i+1}} \prod_{j=1}^{i+1} C_{\alpha_j-1}.$$

On the other hand, the number $C_{r-1, r-1-i}$ of Dyck paths from $(0, 0)$ to $(r - 1, r - 1 - i)$ satisfies the same recurrence equation

$$C_{r-1, r-1-i} = \sum_{\substack{\beta \vDash r-1-i \\ \ell(\beta)=i+1}} \prod_{j=1}^{i+1} C_{\beta_j},$$

but the sum is over the set of weak compositions so that $\beta_j \geq 0$. Under the appropriate shift in indices, it follows that the number of Dyck paths from $(0, 0)$ to (r, r) which touch the diagonal exactly i times is $C_{r-1, r-1-i}$. The result follows from this. \square

Corollary 16. For $k \geq m \geq 0$,

$$\sum_{j=m}^k \binom{j}{m} C_{k,j} = C_{2k+1-m,m}.$$

Proof. The two formulas for $F_{(k+1,k+1)}(q)$ yields the identity

$$\sum_{i=0}^k C_{k,k-i} q^{2k^2-i} (q+1)^i = q^{\binom{2k}{2}} \sum_{i=0}^k C_{2k-i,i} q^i.$$

Extracting the coefficient of q^{2k^2-m} in the above expressions yields the result. \square

Remark 17. The formula for $F_{(r,r)}(q)$ provided in Proposition 15 can be viewed as a sum over Dyck paths, where each Dyck path π contributes a term of the form $q^{s_1(\pi)}(q+1)^{s_2(\pi)}$ for some statistics s_1 and s_2 on the Dyck paths. This particular factorization for $F_{(r,r)}(q)$ is related to the work of Cai and Readdy on the q -Stirling numbers of the second kind, since the polynomials $F_\lambda(q)$ can be viewed as a refinement of $S_{n,k}(q)$, as explained in Section 3.

Cai and Readdy obtained a formula [2, Theorem 3.2] for $\tilde{S}_{n,k}(q)$ (they use a different recursive formula to define the q -Stirling numbers, and the two are related by $S_{n,k}(q) = q^{\binom{k}{2}} \tilde{S}_{n,k}(q)$) as a sum over allowable restricted-growth words, where each allowable word w gives rise to a term of the form $q^{a(w)}(q+1)^{b(w)}$ for some statistics $a(w)$ and $b(w)$. They also showed that this enumerative result has an interesting extension to the study of the Stirling poset of the second kind, providing a decomposition of that poset into Boolean sublattices.

For example, if we define polynomials $G_\lambda(q)$ by letting $G_\lambda(q) = F_\lambda(q)/(q-1)^{n-\ell(\lambda)}$ (see Equation (12)), then $G_{(3,1)}(q) + G_{(2,2)}(q) = q^3 \tilde{S}_{4,2}(q^{-1})$. The formula of Cai and Readdy yields $q^3 S_{4,2}(q^{-1}) = q(q+1)^2 + q^2(q+1) + q^3$, while our factorization yields $G_{(3,1)}(q) + G_{(2,2)}(q) = (q^3 + q^3 + q^2(q+1)) + (q^2 + q(q+1))$. So, the result of Proposition 15 gives a different expression for $S_{n,n-2}(q)$ as a sum with terms of the form $q^{s_1(\pi)}(q+1)^{s_2(\pi)}$, and it may be interesting to further investigate such factorizations of $F_\lambda(q)$.

Example 18. The first few $F_{(k,k)}$ are

$$\begin{aligned} F_{(1,1)} &= 1 \\ F_{(2,2)} &= (q-1)^2 (q^2 + q(q+1)) \\ &= (q-1)^2 (2q^2 + q) \\ F_{(3,3)} &= (q-1)^4 (2q^8 + 2q^7(q+1) + q^6(q+1)^2) \\ &= (q-1)^4 (5q^8 + 4q^7 + q^6) \\ F_{(4,4)} &= (q-1)^6 (5q^{18} + 5q^{17}(q+1) + 3q^{16}(q+1)^2 + q^{15}(q+1)^3) \\ &= (q-1)^6 (14q^{18} + 14q^{17} + 6q^{16} + q^{15}) \\ F_{(5,5)} &= (q-1)^8 (14q^{32} + 14q^{31}(q+1) + 9q^{30}(q+1)^2 + 4q^{29}(q+1)^3 + q^{28}(q+1)^4) \\ &= (q-1)^8 (42q^{32} + 48q^{31} + 27q^{30} + 8q^{29} + q^{28}). \end{aligned}$$

We end this section with one more closed formula for $F_\lambda(q)$ where λ is a rectangular shape with two columns. Let $\mathcal{D}(n,k)$ denote the set of Dyck paths from $(0,0)$ to (n,k) . The *coarea* of a Dyck path π is the number of whole unit squares lying between the path





Figure 4: Factors of $q - 1$ are omitted from the edge weights in this sublattice of partitions with at most two columns.

and the x -axis. For $i = 1, \dots, n$, let $\rho_i(\pi)$ be one plus the number of unit squares lying between the path and the line $y = x + 1$ in the i th row. For example, the following Dyck path π has $\text{coarea}(\pi) = 12$, and $(\rho_1(\pi), \rho_2(\pi), \rho_3(\pi), \rho_4(\pi)) = (2, 2, 1, 1)$.



For $n \geq 1$, let $[n]_q = 1 + q + \dots + q^{n-1}$.

Proposition 19. (Partitions with two columns) Let $\lambda = (2^r, 1^s) \vdash n$ such that $r, s \geq 0$. Then

$$F_{(2^r, 1^s)}(q) = (q - 1)^r q^{\binom{r}{2}} \sum_{\pi \in \mathcal{D}(r+s, r)} q^{\text{coarea}(\pi)} \prod_{i=1}^r [\rho_i(\pi)]_q.$$

Proof. By Corollary 11, we know the multiplicity of the factor $q - 1$ in $F_{(2^r, 1^s)}(q)$ is $n - \ell(\lambda) = \lambda'_2 = r$, so we focus on computing $F_{(2^r, 1^s)}(q)/(q - 1)^r$. The paths in Young's lattice from \emptyset to $(2^r, 1^s)$ are in bijection with the Dyck paths $\mathcal{D}(r + s, r)$, so we identify these paths; adding a box in the first column of a partition corresponds to a $(1, 0)$ step in the Dyck path, and adding a box in the second column of a partition corresponds to a $(0, 1)$ step in the Dyck path. As seen in Figure 4, a vertical step (i, j) to $(i, j + 1)$ has

weight $q^j[i]_q$, while a horizontal step (i, j) to $(i + 1, j)$ has weight q^j . Thus the product of the edge weights of the r vertical steps of a given Dyck path π is $q^{\binom{r}{2}} \prod_{i=1}^r [\rho_i(\pi)]_q$, while the product of the edge weights of the $r + s$ horizontal steps of a given Dyck path is $q^{\text{coarea}(\pi)}$. The result follows. \square

Example 20. The first few $F_{(2^n)}$ are

$$\begin{aligned} F_{(2)} &= (q - 1)(q + 1) \\ F_{(2^2)} &= (q - 1)^2 q (q + (q + 1)) \\ F_{(2^3)} &= (q - 1)^3 q^3 (q^3 + 2q^2(q + 1) + q(q + 1)^2 + (q^2 + q + 1)(q + 1)) \\ F_{(2^4)} &= (q - 1)^4 q^6 (q^6 + 3q^5[2] + 3q^4[2]^2 + q^3[2]^3 + 2q^3[3]! + 2q^2[2][3]! + q[3][3]! + [4]!). \end{aligned}$$

Remark 21. Kirillov and Melnikov [8] considered the number $A_n(q)$ of n by n upper-triangular matrices over \mathbb{F}_q satisfying $X^2 = 0$. In their first characterization of these polynomials, they considered the number $A_n^r(q)$ of matrices of a given rank r , so that $A_n(q) = \sum_{r \geq 0} A_n^r(q)$, and observed that $A_n^r(q)$ satisfies the recurrence equation

$$A_n^r(q) = q^r A_{n-1}^r(q) + (q^{n-r} - q^r) A_n^r(q), \quad A_n^0(q) = 1.$$

We may think of $A_n(q)$ as the sum of $F_\lambda(q)$ over $\lambda \vdash n$ with at most two columns, so Theorem 8 is a generalization of this recurrence equation.

It was also conjectured in [8] that the same sequence of polynomials arise in a number of different ways. Ekhad and Zeilberger [3] proved that one of the conjectured alternate definitions of $A_n(q)$, namely

$$C_n(q) = \sum_s c_{n+1,s} q^{\frac{n^2}{4} + \frac{1-s^2}{12}},$$

is a sum over all $s \in [-n - 1, n + 1]$ which satisfy $s \equiv n + 1 \pmod{2}$ and $s \equiv (-1)^n \pmod{3}$, and $c_{n+1,s}$ are entries in the signed Catalan triangle, is indeed the same as $A_n(q)$. It would be interesting to see what other combinatorics may arise from considering the sum of $F_\lambda(q)$ over $\lambda \vdash n$ with at most k columns for a fixed k .

3 Jordan canonical forms and q -rook placements

In light of Corollary 11, we define polynomials $G_\lambda(q) \in \mathbb{Z}[q]$ by

$$F_\lambda(q) = (q - 1)^{n-\ell(\lambda)} G_\lambda(q). \tag{12}$$

In fact, we can deduce from Corollary 11 that $G_\lambda(q) \in \mathbb{N}[q]$. In this section, we explore the connection between the nonnegative coefficients of $G_\lambda(q)$ and rook placements.

3.1 Background on rook polynomials

A board B is a subset of an n by n grid of squares. In this paper, we follow Haglund [5] and Solomon [12], and index the squares using the convention for the entries of a matrix. A Ferrers board is a board B where if a square $s \in B$, then every square lying north and/or east of s is also in B . Our Ferrers boards have squares justified upwards and to the right. Let B_n denote the staircase-shaped board with n columns of sizes $0, 1, \dots, n-1$. Let $\text{area}(B)$ be the number of squares in B , so that in particular, $\text{area}(B_n) = \binom{n}{2}$.

A placement of k rooks on a board B is *non-attacking* if there is at most one rook in each row and each column of B . Let $\mathcal{C}(B, k)$ denote the set of non-attacking placements of k rooks on B . All rook placements considered in this article are non-attacking, so from this point forward, we drop the qualifier. For a placement $C \in \mathcal{C}(B, k)$, let $\text{ne}(C)$ be the number of squares in B lying directly north or directly east of a rook. The *inversion* of the placement is the number

$$\text{inv}(C) = \text{area}(B) - k - \text{ne}(C). \quad (13)$$


As noted in [4], the statistic $\text{inv}(C)$ is a generalization of the number of inversions of a permutation, since permutations can be identified with rook placements on a square-shaped board.

For $i = 1, \dots, n$, the weight of the i th column C_i of C is

$$C_i(q) = (q-1)^{\#\text{rooks in } C_i} q^{\text{ne}(C_i)}, \quad (14)$$

and the *weight* of C is defined by $F_C(q) = \prod_{i=1}^n C_i(q)$. Alternatively, if $C \in \mathcal{C}(B, k)$, then $F_C(q) = (q-1)^k q^{\text{ne}(C)}$.

Example 22. We use \times to mark a rook and use \bullet to mark squares lying directly north or directly east of a rook (these squares shall be referred to as the north-east squares of the placement). The following illustration is a placement of four rooks on the staircase-shaped board B_7 .



This rook placement has $\text{ne}(C) = 11$, $\text{inv}(C) = 6$, and weight $F_C(q) = (q-1)^4 q^{11}$.

For $k \geq 0$, the q -rook polynomial of a Ferrers board B is defined by Garsia and Remmel [4, I.4] as

$$R_{B,k}(q) = \sum_{C \in \mathcal{C}(B,k)} q^{\text{inv}(C)}. \quad (15)$$

The following result explains the role of rook polynomials in the enumeration of matrices of given rank. The *support* of a matrix X is $\{(i, j) \mid x_{ij} \neq 0\}$. Given a Ferrers board B with n columns, we may identify the squares in B with the entries in an n by n matrix.

Theorem 23 (Haglund). *If B is a Ferrers board, then the number $P_{B,k}(q)$ of n by n matrices of rank k with support contained in B is*

$$P_{B,k}(q) = (q-1)^k q^{\text{area}(B)-k} R_{B,k}(q^{-1}).$$

Looking ahead, it will be convenient to consider Theorem 23 in the following equivalent form:

$$P_{B,k}(q) = \sum_{C \in \mathcal{C}(B,k)} (q-1)^k q^{\text{ne}(C)} = \sum_{C \in \mathcal{C}(B,k)} F_C(q). \quad (16)$$

Example 24. We list the seven rook placements on B_4 with two rooks, along with their weights.

$$(q-1)^2 q^3 \quad (q-1)^2 q^3 \quad (q-1)^2 q^3 \quad (q-1)^2 q^2 \quad (17)$$

$$(q-1)^2 q^2 \quad (q-1)^2 q^2 \quad (q-1)^2 q \quad (18)$$

Thus $P_{B_4,2}(q) = (q-1)^2(3q^3 + 3q^2 + q)$.

3.2 Rook placements and Jordan forms

The purpose of this section is to generalize Haglund's formula (16) to a formula for $F_\lambda(q)$ (Corollary 30) as a sum over a set of rook placements. We achieve this by defining a multigraph \mathcal{Z} that is related to \mathcal{Y} , and show that paths in \mathcal{Z} are equivalent to rook placements.

The multigraph \mathcal{Z} is constructed from \mathcal{Y} by replacing each edge of \mathcal{Y} by one or more edges as follows. If there is an edge from μ to λ in \mathcal{Y} of weight $q^{|\mu|-\mu'_j-1} (q^{\mu'_{j-1}-\mu'_j} - 1)$, then this edge is replaced by $\mu'_{j-1} - \mu'_j$ edges from μ to λ with weights

$$(q-1)q^{|\mu|-\mu'_j-1}, \dots, (q-1)q^{|\mu|-\mu'_{j-1}} \quad (19)$$

in \mathcal{Z} . All other edges remain as before. See Figure 5.

Let $\mathcal{P}_{\mathcal{Z}}(\lambda)$ denote the set of paths in the graph \mathcal{Z} from the empty partition \emptyset to λ . For a path $P = (\emptyset = \pi^{(0)}, \pi^{(1)}, \dots, \pi^{(n)} = \lambda)$ in $\mathcal{P}_{\mathcal{Z}}(\lambda)$, let $\epsilon_i(q)$ denote the weight of the i th edge, for $i = 1, \dots, n$. Naturally, we define the weight of the path by $F_P(q) = \prod_{i=1}^n \epsilon_i(q)$, so that

$$F_\lambda(q) = \sum_{P \in \mathcal{P}_{\mathcal{Z}}(\lambda)} F_P(q). \quad (20)$$

Lemma 25. *Let $\mu \vdash n-1$ be a partition with $\ell(\mu) = \ell$ parts. Then there are $\ell+1$ edges leaving μ in the graph \mathcal{Z} , with weights*

$$(q-1)q^{|\mu|-1}, (q-1)q^{|\mu|-2}, \dots, (q-1)q^{|\mu|-\ell}, \text{ and } q^{|\mu|-\ell}.$$




Figure 5: The multigraph \mathcal{Z} , up to $n = 4$.

Proof. If a partition $\lambda \vdash n$ is obtained by adding a box to the first column of μ , then there is a unique edge from μ to λ in \mathcal{Z} with weight $q^{|\mu|-\ell}$. Otherwise, if we consider the set of all partitions which can be obtained from μ by adding a box anywhere except in the first column, then there are a total of

$$\sum_{j \geq 2} (\mu'_{j-1} - \mu'_j) = \ell$$

edges from μ to some partition of n . Moreover, by Equation (19), these ℓ weights are $(q-1)q^{|\mu|-i}$ for $i = 1, \dots, \ell$. \square

A sequence of nonnegative integers is $\mathcal{P}_{\mathcal{Z}}$ -admissible if it is the degree sequence of a path $P = (\emptyset, \pi^{(1)}, \dots, \pi^{(n)})$ in \mathcal{Z} . That is, $(d_1, \dots, d_n) = (\deg \epsilon_1(q), \dots, \deg \epsilon_n(q))$.

Corollary 26. *A $\mathcal{P}_{\mathcal{Z}}$ -admissible sequence determines a unique path in \mathcal{Z} .*

Proof. Induct on n . When $n = 1$, the only path is the from \emptyset to (1) , and it has degree sequence (0) .

Given a $\mathcal{P}_{\mathcal{Z}}$ -admissible sequence (d_1, \dots, d_n) , the subsequence (d_1, \dots, d_{n-1}) determines a unique path $P' = (\emptyset, \pi^{(1)}, \dots, \pi^{(n-1)})$. Suppose $\mu = \pi^{(n-1)}$ has ℓ parts. Then $|\mu| - \ell + 1 \leq d_n \leq |\mu|$, and by Lemma 25, there is a unique edge leaving μ with degree d_n . \square

3.3 The construction of Φ

Let $\mathcal{P}_{\mathcal{Z}}(n, n - k)$ denote the set of paths in \mathcal{Z} from \emptyset to a partition of n with $n - k$ parts. In this section, we define a weight-preserving bijection $\Phi : \mathcal{C}(B_n, k) \rightarrow \mathcal{P}_{\mathcal{Z}}(n, n - k)$.

Proposition 27. *Let $n \geq 1$ and $k = 0, \dots, n - 1$. Let $C \in \mathcal{C}(B_n, k)$ be a rook placement with columns C_1, \dots, C_n . There exists a unique path $P \in \mathcal{P}_{\mathcal{Z}}(n, n - k)$ with edge weights $(\epsilon_1(q), \dots, \epsilon_n(q)) = (C_1(q), \dots, C_n(q))$.*

Proof. Proceed by induction on $n + k$. When $n = 1$ and $k = 0$, there is a unique rook placement on the empty board B_1 with no rooks having weight one, corresponding to the unique path $P = (\emptyset, (1))$ in \mathcal{Z} with the same weight.

Assume the result holds for all rook placements in $\mathcal{C}(B_{n-1}, k)$ and $\mathcal{C}(B_{n-1}, k - 1)$. Given a rook placement $C \in \mathcal{C}(B_n, k)$, let C' be the sub-placement consisting of the first $n - 1$ columns of C . By induction, the sequence $(C_1(q), \dots, C_{n-1}(q))$ determines a unique path $(\emptyset, \pi^{(1)}, \dots, \pi^{(n-1)})$ in \mathcal{Z} such that $\epsilon_i(q) = C_i(q)$ for $i = 1, \dots, n - 1$.

There are now two cases to consider. The first case is if $C' \in \mathcal{C}(B_{n-1}, k)$, so that $\ell(\pi^{(n-1)}) = n - k - 1$. There are k rooks in C' , so the n th column of C does not contain any rooks, and $C_n(q) = q^k$. By Lemma 25, there exists a unique edge in the graph \mathcal{Z} originating at $\pi^{(n-1)}$ with weight q^k . Thus C corresponds to the path $P = (\emptyset, \pi^{(1)}, \dots, \pi^{(n-1)}, \pi^{(n)})$ where $\pi^{(n)}$ is obtained from $\pi^{(n-1)}$ by adding a box to the first column, and $\epsilon_n(q) = q^k$. Moreover, $\ell(\pi^{(n)}) = n - k$.

The second case is if $C' \in \mathcal{C}(B_{n-1}, k - 1)$, so that $\ell(\pi^{(n-1)}) = n - k$. There must be $k - 1$ ‘northeast’ squares in the n th column of C , and there are $n - k$ remaining squares in that column where a rook may be placed. Label these available squares $a_0, a_1, \dots, a_{n-k-1}$ from the top to the bottom. Observe that $C_n(q) = (q - 1)q^{k-1+i}$ if a rook is placed in the square a_i , for $0 \leq i \leq n - k - 1$. Again by Lemma 25, there exists $n - k$ edges in the graph \mathcal{Z} originating at $\pi^{(n-1)}$ with the weights $(q - 1)q^h$ for $k - 1 \leq h \leq n - 2$. Thus if the k th rook of C is placed in the square a_i , then C corresponds to the path $P = (\emptyset, \pi^{(1)}, \dots, \pi^{(n-1)}, \pi^{(n)})$ with $\epsilon_n(q) = (q - 1)q^{k-1+i}$, and $\ell(\pi^{(n)}) = n - k$. \square

Given a rook placement $C \in \mathcal{C}(B_n, k)$, let $\Phi(C)$ be the path in $\mathcal{P}_{\mathcal{Z}}(n, n - k)$ with edge weights $(\epsilon_1(q), \dots, \epsilon_n(q)) = (C_1(q), \dots, C_n(q))$.

Theorem 28. *The map $\Phi : \mathcal{C}(B_n, k) \rightarrow \mathcal{P}_{\mathcal{Z}}(n, n - k)$ is a weight-preserving bijection.*

Proof. Proposition 27 shows that the map Φ is an injective weight-preserving map, since each column of the rook placement determines each edge of the path $\Phi(C)$:

$$F_C(q) = \prod_{i=1}^n C_i(q) = \prod_{i=1}^n \epsilon_i(q) = F_{\Phi(C)}(q).$$

In fact, the proof of the Proposition also shows that Φ is surjective because the number of possible ways to add a column to an existing rook placement is equal to the number of possible ways to extend a path in \mathcal{Z} by one edge. Therefore, Φ is a weight-preserving bijection. \square

A sequence of nonnegative integers is \mathcal{C} -admissible if it is the degree sequence of a rook placement. That is, $(d_1, \dots, d_n) = (\deg C_1(q), \dots, \deg C_n(q))$ for a $C \in \mathcal{C}(B_n, k)$. The next Corollary follows easily from Theorem 28.

Corollary 29. *A \mathcal{C} -admissible sequence determines a unique rook placement.* \square

It follows from Theorem 28 that we may associate a partition type to each rook placement on B_n . The *partition type* of a rook placement C is the partition at the endpoint of the path $\Phi(C)$ in \mathcal{Z} . Let $\mathcal{C}(\lambda) = \Phi^{-1}(P_{\mathcal{Z}}(\lambda))$ denote the set of rook placements of partition type λ .

Corollary 30. *Let $\lambda \vdash n$ be a partition with $\ell(\lambda) = n - k$ parts. Then*

$$F_\lambda(q) = \sum_{C \in \mathcal{C}(\lambda)} F_C(q) = (q - 1)^{n - \ell(\lambda)} \sum_{C \in \mathcal{C}(\lambda)} q^{\text{ne}(C)}.$$

Proof. The result follows from Equation 20 and the bijection Φ . \square

Remark 31. The polynomial $G_\lambda(q) \in \mathbb{N}[q]$ defined in Equation (12) is simply a sum over the rook placements of type λ involving the north-east statistic.

4 A connection with set partitions

The results of the previous section naturally leads to a decomposition of $F_T(q)$, indexed by some tableau T , into a sum of polynomials indexed by set partitions, which we explain below.

A *set partition* is a set $S = \{s_1, \dots, s_k\}$ of nonempty disjoint subsets of $[n]$ such that $\bigcup_{i=1}^k s_i = [n]$. The s_i 's are the *blocks* of σ . Let $\ell(S)$ denote the number of blocks of S , and let $\mathcal{S}(n, n - k)$ denote the set of set partitions of $[n]$ with $n - k$ blocks. We adopt the convention of listing the blocks in order so that

$$|s_1| \geq |s_2| \geq \dots \geq |s_k|, \text{ and } \min s_i < \min s_{i+1} \text{ if } |s_i| = |s_{i+1}|. \quad (21)$$

This allows us to represent a set partition with a diagram similar to that of a standard Young tableau; the i th row of the diagram consists of the elements in the block s_i listed in increasing order, but there are no restrictions on the entries in each column of the diagram. A set partition $S = (s_1, \dots, s_m)$ has *partition type* λ if $\lambda = (|s_1|, \dots, |s_m|)$.

For $i = 1, \dots, n$, let $S^{(i)}$ denote the sub-diagram of S consisting of the boxes containing $1, \dots, i$, with rows ordered according to the convention set forth in Equation (21). If the box containing i is not in the first column of the diagram, let u be the least element in the same row as i in $S^{(i)}$, and suppose u is in the r th row of $S^{(i-1)}$ for some $1 \leq r \leq \ell(S^{(i-1)})$. The weight arising from the i th box is

$$S^{(i)}(q) = \begin{cases} q^{i-1-\ell(S^{(i-1)})}, & \text{if the } i\text{th box is in the first column,} \\ (q-1)q^{i-1-r}, & \text{if the } i\text{th box is in the } j\text{th column, } j \geq 2. \end{cases} \quad (22)$$

We define the *weight* of S as $F_S(q) = \prod_{i=1}^n S^{(i)}(q)$.

A sequence of nonnegative integers is \mathcal{S} -*admissible* if it is the degree sequence of a set partition. That is, $(d_1, \dots, d_n) = (\deg S^{(1)}(q), \dots, \deg S^{(n)}(q))$ for a $S \in \mathcal{S}(n)$.

Lemma 32. *An \mathcal{S} -admissible sequence determines a unique set partition.*

Proof. Induct on n . When $n = 1$, the only set partition is $\{\{1\}\}$, and its degree sequence is (0) .

Given an \mathcal{S} -admissible sequence (d_1, \dots, d_n) , the subsequence (d_1, \dots, d_{n-1}) determines a unique set partition $S^{(n-1)} = (S_1^{(n-1)}, \dots, S_m^{(n-1)})$. By Equation (22), $n - 1 - m \leq d_n \leq n - 1$, and each of the $m + 1$ choices for d_n determines the block of $S^{(n-1)}$ into which n should be inserted. \square

We have already constructed a weight-preserving bijection Φ between rook placements and paths in \mathcal{Z} . We now construct a weight-preserving bijection Ψ between rook placements and set partitions, effectively showing that paths in \mathcal{Z} are equivalent to set partitions, so that $F_Z(q) = F_C(q) = F_S(q)$ if $Z \longleftrightarrow C \longleftrightarrow S$ for $Z \in \mathcal{P}_{\mathcal{Z}}(n, n - k)$, $C \in \mathcal{C}(B_n, k)$, and $S \in \mathcal{S}(n, n - k)$.

Remark 33. There is a classically known bijection (see [14]) between the set of rook placements on the staircase board B_n with k rooks and the set of set partitions of $[n] = \{1, \dots, n\}$ with $n - k$ blocks: the placement C corresponds to the set partition where the integers i and j are in the same block if and only if there is a rook in the square $(i, j) \in C$. This bijection is different from the one described in Theorem 34. For example, the classical bijection associates the rook placement

	1	2	3	4
1		×	•	•
2			•	
3				×
4				

to the set partition $(\{1, 2\}, \{3, 4\})$ and so has partition type $(2, 2)$, but as we shall see below, this placement is associated to the set partition $(\{1, 2, 4\}, \{3\})$ under the bijection in Theorem 34 and has partition type $(3, 1)$.

4.1 The construction of Ψ

Let $C \in \mathcal{C}(B_n, k)$ be a rook placement. The main idea is that the degree of $C_i(q)$ arising from the i th column of C determines the block of the set partition in which we place i . In the construction of the set partition $\Psi(C)$, we will create a sequence of intermediate set partitions $S^{(i)}$ of $[i]$ for $i = 1, \dots, n$.

The initial case is always $\deg(C_1(q)) = \deg(1) = 0$, so $S^{(1)} = \{\{1\}\}$. Assume that $S^{(i-1)} = \{S_1^{(i-1)}, \dots, S_m^{(i-1)}\}$ is the set partition which corresponds to the first $i - 1$ columns of C , so that $m = \ell(S^{(i-1)})$. Observe that there are $m + 1$ possible blocks in which to insert i to obtain $S^{(i)}$. By Corollary 26, we know that

$$i - 1 - \ell(S^{(i-1)}) \leq \deg(C_i(q)) \leq i - 1,$$

so we construct $S^{(i)}$ by placing i in the j th block of $S^{(i-1)}$, where $j = i - \deg(C_i(q))$, and then rearranging the blocks to fit the convention in Equation (21) if necessary.


Theorem 34. *The map $\Psi : \mathcal{C}(n, k) \rightarrow \mathcal{S}(n, n - k)$ is a weight-preserving bijection.*

Proof. Let $S = \Psi(C)$. The map Ψ is weight-preserving, as $C_i(q) = S^{(i)}(q)$ by construction, for each $i = 1, \dots, n$. Now, since the degrees $\deg C_i(q) = \deg S^{(i)}(q)$, and by Corollary 29 and Lemma 32 the sequences of degrees completely determine C and S respectively, then Ψ is injective. Finally, we note that $|\mathcal{C}(n, k)| = |\mathcal{S}(n, n - k)|$, so Ψ is a bijection. \square


Corollary 35. *Let $\mathcal{S}(\lambda)$ denote the set of all set partitions of partition type λ . Then*

$$F_\lambda(q) = \sum_{S \in \mathcal{S}(\lambda)} F_S(q). \quad \square$$

Example 36. Let C be the rook placement



The associated sequence of set partition diagrams associated to C is



so the set partition associated to the rook placement C is

$$S = \Psi(C) = (\{3, 8, 9\}, \{1, 5\}, \{6, 7\}, \{2\}, \{4\}).$$

Remark 37. An intriguing question is to ask for a geometric interpretation of the polynomials $F_C(q)$, indexed by rook placements (or set partitions or paths in \mathcal{Z}).

The problem of determining the number of adjoint $G_n(\mathbb{F}_q)$ orbits on $\mathfrak{g}_n(\mathbb{F}_q)$ remains open. In the case $q = 2$, this number has been computed for $n \leq 16$ by Pak and Soffer [11, Appendix B]. Let $\mathcal{O}_n(k)$ denote the orbits of rank k matrices. When $k = 1$, it turns out that the polynomials $F_C(q)$ indexed by rook placements with exactly one rook gives the sizes of the $\binom{n}{2}$ orbits in $\mathcal{O}_n(1)$. For $2 \leq i < j \leq n$, each orbit contains a unique matrix E_{ij} whose ij th entry is 1, and is zero everywhere else. The orbit containing E_{ij} is associated to the rook placement $C(i, j)$ with a single rook in the ij th square, and the size of the associated orbit is $F_{C(i,j)}(q) = (q - 1)q^{n-1-(j-i)}$.




Figure 6: Paths, rook placements, and set partitions related to the computation of $F_{(3,1)}(q) = (q - 1)^2(3q^3 + q^2)$.

In particular, the formula in Proposition 13 applied to the partition $\lambda = (2, 1^{n-2})$ gives the generating function

$$F_{(2,1^{n-2})}(q) = (q - 1) ((n - 1)q^{n-2} + (n - 2)q^{n-3} + \dots + 3q^2 + 2q + 1)$$

for rank one orbits of $G_n(\mathbb{F}_q)$ on $\mathfrak{g}_n(\mathbb{F}_q)$.

Remark 38. To close, we mention a related problem which may provide a geometric interpretation of $F_C(q)$ for every rook placement C . Let N be an $n \times n$ nilpotent matrix with entries in an algebraically closed field k containing \mathbb{F}_q , and suppose N has Jordan type $\lambda \vdash n$. A complete flag $f = (f_1, \dots, f_n)$ is a sequence of subspaces in k^n such that $f_1 \subset \dots \subset f_n$ and $\dim f_i = i$ for all i . A flag is N -stable if $N(f_i) \subseteq f_i$ for all i . Spaltenstein [13] showed that the variety X_λ of N -stable flags is a disjoint union of f^λ smooth irreducible subvarieties X_T indexed by the standard Young tableaux of shape λ . Moreover, the closures \overline{X}_T are the irreducible components of X_λ , each of which has dimension n_λ . The number of \mathbb{F}_q -rational points in X_λ is given by Green's polynomials $Q_{(1^n)}^\lambda(q)$ [9, III.7]. Evidently,

$$\left(\prod_{i \geq 1} [m_i(\lambda)]_q! \right)^{-1} Q_{(1^n)}^\lambda(q) = ((q - 1)^{n-\ell(\lambda)} q^m)^{-1} F_\lambda(q),$$

with $m = \min_{C \in \mathcal{C}(\lambda)} \text{ne}(C)$. Based on some computations for small values of n , we expect that $F_C(q)$ plays a role in counting points in certain intersections of the irreducible components \overline{X}_T .

Acknowledgements

I would like to thank Jim Haglund and Alexandre Kirillov for their invaluable guidance, and Yue Cai and Alejandro Morales for enlightening conversations.

References

- [1] A. M. Borodin, Limit Jordan normal form of large triangular matrices over a finite field. *Funct. Anal. Appl.*, 29(4):279–281, 1995.
- [2] Y. Cai and M. Readdy, q -Stirling numbers: A new view. *Adv. in Appl. Math.*, 86:50–80, 2017.
- [3] S. B. Ekhad and D. Zeilberger, The number of solutions of $X^2 = 0$ in triangular matrices over $\text{GF}(q)$. *Electron. J. Combin.*, 3(1):#R2, 1996.
- [4] A. Garsia and J. B. Remmel, q -Counting rook configurations and a formula of frobenius. *J. Combin. Theory Ser. A*, 41(2):246–275, 1986.
- [5] J. Haglund, q -Rook polynomials and matrices over finite fields. *Adv. in Appl. Math.*, 20(4):450–487, 1998.
- [6] A. Henderson, Enhancing the Jordan canonical form. *Austral. Math. Soc. Gaz.*, 38(4):206–211, 2011.
- [7] A. A. Kirillov, Variations on the triangular theme. *Amer. Math. Soc. Transl.*, 169(2):43–73, 1995.
- [8] A. A. Kirillov and A. Melnikov, On a remarkable sequence of polynomials. *Sémin. Congr. 2, Soc. Math. France*, 35–42, 1995.
- [9] I. G. Macdonald, Symmetric functions and Hall polynomials. *Oxford University Press*, 1995.
- [10] N. J. A. Sloane, *The online encyclopedia of integer sequences*.
- [11] I. Pak and A. Soffer, On Higman’s $k(U_n(\mathbb{F}_q))$ conjecture. [arXiv:1507.00411](https://arxiv.org/abs/1507.00411).
- [12] L. Solomon, The Bruhat decomposition, Tits system and Iwahori ring for the monoid of matrices over a finite field. *Geom. Dedicata.*, 36(1):15–49, 1990.
- [13] N. Spaltenstein, The fixed point set of a unipotent transformation on the flag manifold. *Proceedings of the Koninklijke Nederlandse Academie van Wetenschappen, Amsterdam, Series A 79*, 38(5):452–458, 1976.
- [14] R. P. Stanley. Enumerative combinatorics. *Cambridge University Press*, 1999.
- [15] M. Yip q -Rook placements and Jordan forms of upper-triangular matrices. DMTCS Proceedings FPSAC, 25:1017–1028, 2013.