

University of Kentucky

UKnowledge

Orthopaedic Surgery and Sports Medicine
Faculty Publications

Orthopaedic Surgery and Sports Medicine

8-2021

Unicompartmental Knee Arthroplasty Is Associated with a Lower Rate of Periprosthetic Joint Infection Compared to Total Knee Arthroplasty

Cody S. Lee
University of Chicago

Edwin P. Su
Hospital for Special Surgery

Michael B. Cross
Hospital for Special Surgery

Alberto V. Carli
Hospital for Special Surgery

David C. Landy
University of Kentucky, David.Landy@uky.edu

Follow this and additional works at: https://uknowledge.uky.edu/orthopaedicsurgery_facpub

 *next page for additional authors*
Part of the [Orthopedics Commons](#)

[Right click to open a feedback form in a new tab to let us know how this document benefits you.](#)

Repository Citation

Lee, Cody S.; Su, Edwin P.; Cross, Michael B.; Carli, Alberto V.; Landy, David C.; and Chalmers, Brian P., "Unicompartmental Knee Arthroplasty Is Associated with a Lower Rate of Periprosthetic Joint Infection Compared to Total Knee Arthroplasty" (2021). *Orthopaedic Surgery and Sports Medicine Faculty Publications*. 33.

https://uknowledge.uky.edu/orthopaedicsurgery_facpub/33

This Article is brought to you for free and open access by the Orthopaedic Surgery and Sports Medicine at UKnowledge. It has been accepted for inclusion in Orthopaedic Surgery and Sports Medicine Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Unicompartmental Knee Arthroplasty Is Associated with a Lower Rate of Periprosthetic Joint Infection Compared to Total Knee Arthroplasty

Digital Object Identifier (DOI)

<https://doi.org/10.1016/j.artd.2021.06.006>

Notes/Citation Information


Published in *Arthroplasty Today*, v. 10.

© 2021 The Authors

This is an open access article under the CC BY-NC-ND license (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

Authors

Cody S. Lee, Edwin P. Su, Michael B. Cross, Alberto V. Carli, David C. Landy, and Brian P. Chalmers



Original research

Unicompartmental Knee Arthroplasty Is Associated With a Lower Rate of Periprosthetic Joint Infection Compared to Total Knee Arthroplasty

Cody S. Lee, MD^b, Edwin P. Su, MD^a, Michael B. Cross, MD^a, Alberto V. Carli, MD^a, David C. Landy, MD, PhD^c, Brian P. Chalmers, MD^{a,*}^a Department of Orthopedic Surgery Adult Reconstruction and Joint Replacement, Hospital for Special Surgery, New York, NY^b Department of Orthopedic Surgery, University of Chicago, Chicago, IL^c Department of Orthopedic Surgery, University of Kentucky, Lexington, KY

ARTICLE INFO

Article history:

Received 16 March 2021
 Received in revised form
 10 May 2021
 Accepted 14 June 2021
 Available online 20 July 2021

Keywords:

Unicompartmental knee arthroplasty
 Infection
 Periprosthetic joint infection
 Complications
 Total knee
 UKA

ABSTRACT

Background: Several studies have reported lower perioperative complications with unicompartmental knee arthroplasty (UKA) than with total knee arthroplasty (TKA). However, there is a paucity of data analyzing the incidence of periprosthetic joint infection (PJI) in similar patients undergoing UKA and TKA. As such, we sought to analyze the incidence of UKA and TKA PJI in a large matched population.

Material and Methods: The Mariner data set of the PearlDiver database was queried for all patients undergoing UKA or TKA during 2010–2017. Included patients were required to have at least 2 years of database inclusion after surgery. Patients were then matched at a 1:3 ratio (UKA:TKA) on age, gender, Elixhauser Comorbidity Index, tobacco use, and obesity. Rates of PJI requiring operative intervention within 90 days and 1 year were calculated.

Results: In total, 5636 patients having undergone UKA were matched to 16,890 patients having undergone TKA. Fifteen (0.27%) after UKA and 79 (0.47%) after TKA had a PJI surgically managed within 90 days (risk ratio = 0.57, 95% confidence interval = 0.33–0.99, $P = .04$). Thirty (0.53%) after UKA and 136 (0.81%) after TKA had a PJI surgically managed within 1 year (risk ratio = 0.66, 95% confidence interval = 0.45–0.98, $P = .04$).

Conclusions: In a large group of rigorously matched patients, UKA was associated with a significantly lower rate of surgically managed PJI than TKA at 90 days and 1 year; however, the rate of PJI in both groups remained low at <1% at 90 days and 1 year.

© 2021 The Authors. Published by Elsevier Inc. on behalf of The American Association of Hip and Knee Surgeons. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Unicompartmental knee arthroplasty (UKA) remains a successful surgery to relieve pain and improve function in select patients with isolated compartment pathology [1–4]. While traditionally the criteria for UKA were strict [5], the indications and criteria have expanded [2,6–9], and the incidence of UKA has increased over the past decade [10,11]. Furthermore, the introduction of robotic and computer-assisted techniques has been shown to improve UKA alignment, component position, and

potentially functional outcomes and revision rates [12–14]; as such, the wider adoption of robotic-assisted UKA may further increase the incidence of UKA [14,15].

While the long-term revision rates of UKA are higher than those of total knee arthroplasty (TKA) in most series in the literature [4,16–18], there are several reported advantages of a UKA compared with TKA. These include improvement in functional outcomes, higher activity level, quicker recovery, less pain and narcotic requirements postoperatively, and lower short-term complications [17–24]. Furthermore, UKA has been touted to have a lower rate of deep periprosthetic joint infection (PJI) than TKAs, although recent data show that treatment of UKA PJIs does not have better outcomes than TKA PJI treatment [25,26]. However, owing to the low incidence of UKA compared with TKA [10,11], the relatively low incidence of deep PJI in both UKA and TKA, and selection bias of younger and healthier patients toward UKA [15–27], there is a

* Corresponding author. Department of Orthopedic Surgery, Adult Reconstruction and Joint Replacement, Hospital for Special Surgery, 535 East 70th Street, New York, NY 10021. Tel.: 816-808-6348.

E-mail address: chalmersb@hss.edu

paucity of adequately powered data comparing UKA and TKA PJI rates in similar patient populations.

As such, the goal of the present study was to compare the incidence of reoperation for deep PJI at 90 days and 1 year in a large sample of patients from an administrative claims database undergoing UKA to a matched cohort of similar patients undergoing TKA. In addition, we sought to analyze how UKA PJIs were treated at both 90 days and 1 year. We hypothesize that the rate of surgically treated PJI at 90 days and 1 year will be lower for UKA than that for TKA.

Material and methods

Data source

The Mariner data set of the PearlDiver research database (PearlDiver Technologies, Colorado Springs, CO) was used to query deidentified, administrative claims data from patients covered by private insurance, Medicare Advantage, Medicaid, and/or claims paid with cash. It includes roughly 122 million total patients and spans the years of 2010 to 2018. The present study used Common Procedural Terminology (CPT), International Classification of Diseases (ICD)-9 and ICD-10 diagnosis and procedural codes included within the data set (Appendix). The data within this database are Health Insurance Portability and Accountability Act compliant and deemed exempt from institutional review board review.

Patient selection and outcomes

Patients who underwent UKA and primary TKA were identified using codes CPT-27,446 and CPT-27,447, respectively, to preliminarily create 2 cohorts (Fig. 1). All patients were limited to analysis of their first UKA or TKA if they underwent more than one UKA or

TKA in the database. Patients who were included in both groups, with an index procedural code for TKA occurring after index UKA, were included in the UKA cohort and eliminated from the TKA cohort so that the patient could not be matched to himself/herself. Patients were then limited to only those who had continuous database inclusion for at least 2 years. In order to minimize potential bias in patient selection criteria, patients with morbid obesity (body mass index [BMI] > 40 kg.m²) were excluded as many surgeons consider morbid obesity a contraindication for UKA. Patients in each cohort were then matched on a patient to patient basis in a 1:3 ratio (UKA:TKA) on age, sex, Elixhauser Comorbidity Index, obesity (BMI > 30 kg/m²), and tobacco use, and final demographic information was reported for each matched cohort (Table 1). Outcome of infection within 90 days and 1 year of index procedure was reported for each cohort using ICD-9 and ICD-10 diagnosis codes for PJI (Table 2). In addition, the treatment modality used to address PJI in the UKA cohort was reported by stratification into explantation or treatment with debridement, antibiotics, and implant retention (DAIR) using CPT, ICD-9, and ICD-10 procedural codes (Table 2). Diagnosis codes of PJI had to be linked to the same record as the procedural code identified as the mode of treatment.

Statistical analysis

Descriptive statistics were used to describe the demographic and clinical characteristics of the matched groups. Descriptive statistics were also used to present the rates of PJI at both 90 days and 1 year in the UKA and TKA groups. The association between surgical type and PJI at both 90 days and 1 year was expressed using the relative risk of PJI in the UKA group compared with the TKA group with 95% confidence intervals (C.I.) used to assess the statistical significance of this association. Even though there was




Figure 1. Flowchart representing the total number of patients in the database and those excluded due to inadequate follow-up or exclusion criteria. ECI, Elixhauser comorbidity index.

Table 1
Characteristics of matched UKA and TKA patients.

Characteristic	UKA, N = 5636	TKA, N = 16,890
Sex		
Male	2670 (47.4%)	7993 (47.3%)
Female	2966 (52.6%)	8897 (52.7%)
Age		
30–39	20 (0.4%)	51 (0.3%)
40–49	357 (6.3%)	1063 (6.3%)
50–59	1488 (26.4%)	4463 (26.4%)
60–69	1929 (34.2%)	5786 (34.3%)
70–79	1842 (32.7%)	5527 (32.7%)
ECI		
Mean ± SD	4.84 ± 3.23	4.84 ± 3.22
Tobacco use		
Yes	1090 (19.3%)	3259 (19.3%)
No	4546 (80.7%)	13,631 (80.7%)
Obesity		
Yes	1574 (27.9%)	4711 (27.9%)
No	4062 (72.1%)	12,179 (72.1%)
Service location		
Hospital inpatient	3442 (61.1%)	16,307 (96.5%)
Hospital outpatient	2194 (38.9%)	583 (3.5%)

ECI, Elixhauser comorbidity index.

dependency structure within the data created by matching, this could not be incorporated into the statistical analysis given the manner in which data are output from the PearlDiver platform. We elected to match 1:3 UKAs to TKAs given the rarity of UKA PJIs and the significantly higher number of TKAs in the database. Descriptive statistics were also used to describe the treatment characteristics of UKA patients with PJI. No adjustment was made for multiple hypothesis testing, and an alpha threshold of 0.05 was used to assess the statistical significance of associations as reflected by the 95% level for the C.I. All analyses were performed using Stata 16.1 (StataCorp, College Station, TX).

Results

Patient matching and demographics

In total, 5636 patients having undergone UKA were matched to 16,890 patients having undergone TKA (Table 1). As would be expected based on matching, the groups were evenly matched based on demographic and clinical characteristics but not by service location, which was not included in matching. Patients undergoing UKA were more frequently treated as outpatient.

Incidence of PJI in UKA and TKA

Of patients undergoing UKA, 15 (0.27%) had a PJI surgically managed within 90 days (Table 2). Of patients undergoing TKA, 79 (0.47%) had a PJI surgically managed within 90 days. UKA was associated with a significantly lower risk of PJI within 90 days with a risk ratio of 0.57 and 95% C.I. from 0.33 to 0.99 ($P = .04$).

Table 2
Association of UKA compared to TKA with infection.

	UKA, N = 5636	TKA, N = 16,890
PJI in 90 d		
N (%)	15 (0.27%)	79 (0.47%)
RR (95% C.I.)	0.57 (0.33–0.99)	0.00 (Ref)
PJI in 1 y		
N (%)	30 (0.53%)	136 (0.81%)
RR (95% C.I.)	0.66 (0.45–0.98)	0.00 (Ref)

Of patients undergoing UKA, 30 (0.53%) had a PJI surgically managed within 1 year. Of patients undergoing TKA, 136 (0.81%) had a PJI surgically managed within 1 year. UKA was associated with a statistically significant lower risk of PJI within 1 year, with a risk ratio of 0.66 and 95% CI from 0.45 to 0.98 ($P = .04$).

UKA PJI treatment

Of the 15 patients who underwent surgical management of their UKA infection within 90 days, 10 (67%) initially underwent a DAIR. Of the 15 patients who underwent surgical management of their UKA between 91 days and 1 year of UKA, 6 (40%) initially underwent a DAIR (Table 3).

Discussion

The overall incidence of UKA is steadily increasing and expected to continue to increase in the next decade [6–11], especially with the introduction of computer navigated and robotic techniques [12–15]. Numerous studies have reported lower perioperative and short-term complications of UKA compared to TKA [4,17–24]. The low incidence of UKA compared to TKA and biased selection of younger and potentially healthier patients for UKA over TKA render such comparisons, especially for complications with a low incidence such as PJI, and lead to studies being often underpowered and biased. As such, we sought to rigorously match patients undergoing UKA to those undergoing TKA in a large non-Medicare population to compare the incidence of PJI in each cohort. We found that UKA conferred a significantly lower rate of PJI than TKA at 90 days (RR = 0.57) and 1 year (RR = 0.66).

Several studies have reported lower perioperative complication rates of UKA, including lower medical complication rates [4,18], reduced rates of venous thromboembolism [4,18], lower readmission rates [4,18], as well as decreased LOS, lower opioid requirements, and improved functional outcomes [17,19,22] compared with TKA. However, no studies to the authors' knowledge directly compare the incidence of UKA and TKA PJI in the US population. Reported rates of UKA and TKA PJI vary significantly depending on the institution and patient population, ranging from 0% to 2.5% [4,18–23,27,28–30] at early time points. Lum et al. [29] reported a 0% UKA PJI rate in 650 patients with a BMI > 35 kg/m² at a mean follow-up of 2.3 years; conversely, Nettrour et al. [28] reported a 2.2% rate of UKA PJI in patients with a BMI > 40 kg/m² at a mean follow-up of 3.4 years. While the reported rates of PJI vary, most US cohort studies report the rate of UKA PJI from 0.5% to 1% [4,27,28–30].

While there have been very few studies comparing UKA and TKA PJI rates in the United States, a few studies of European patient populations have directly or indirectly compared their rates. In a 1:3 matched cohort of 101,330 patients undergoing 25,334 UKAs and 75,996 TKAs in the National Joint Registry for England and Wales analyzing adverse outcomes as a whole, Liddle et al. [18] found a 2-fold increased risk of revision for UKA at 8 years but

Table 3
Surgical management of infection for UKA.

	UKA, N = 5636
PJI in 90 d	
Explant	5 (33%)
DAIR	10 (67%)
PJI at 91 d to 1 y	
Explant	6 (40%)
DAIR	9 (60%)

significantly reduced perioperative complications, length of stay, and rate of readmission. While not a major outcome measure, they did also find that the incidence of PJI in UKA was significantly lower than that in TKA (HR = 0.5, $P < .001$) and accounted for just ~5% of all revision UKAs compared with 25% of all revision TKAs [18]. In an unmatched observational cohort study of 679,010 knee arthroplasties from the England and Wales registry, Poisson and piecewise exponential multilevel regression models used to analyze several risk factors for PJI; cemented UKA also conferred a lower rate of PJI (RR = 0.5) compared to cemented TKA [23]. Finally, in an older study of roughly 5000 knee arthroplasties in the Norwegian registry performed between 1994 and 2004, the overall risk of revision was 2-fold higher for UKA than that for TKA, but UKAs were associated with a lower (RR = 0.28, $P = .01$) risk of PJI than TKA [31]. The results of these registry-based studies are similar to those of the present study, in which UKA conferred a lower risk of PJI in a matched patient cohort at 90 days and 1 year.

As expected, most UKA PJIs in the present study within 90 days were treated with a DAIR (67%), while 60% of UKA PJIs treated beyond 90 days were treated with a DAIR. Recent studies have shown that DAIR has equally poor, if not worse, outcomes for UKA than for TKA, even in relatively young and healthy patient populations. In the largest study on UKA PJI treatment outcomes, DAIR for acute postoperative PJI had a survivorship free from all-cause reoperation of 55% at 2 years, including a 31% failure for PJI [25]. Similarly, Hernandez et al. [26] reported that 6 of 11 (55%) patients with UKA PJI treated with DAIR underwent septic ($n = 4$) or aseptic ($m = 2$) reoperation by 5 years. In comparison, Weston et al [32] reported a 33% rate of reinfection within 5 years in patients with acute postoperative TKA PJIs treated with DAIR. As such, while UKA may be associated with slightly lower rates of deep PJI, the outcomes of PJI treatment do not appear to be superior to those of TKA; however, there is a paucity of data on their outcomes.

We acknowledge several limitations to the present study. First, it is a large database study with inherent limitations and biases. However, given the lower incidence of UKA and even lower incidence of UKA PJI, with less than 50 UKA PJIs specifically analyzed in the literature [25,26], a large database study is likely the most resource-efficient method to address this question. Furthermore, while databases rely on accurate coding of procedures and diagnoses, the main goal of this study was to compare UKA and TKA PJI, so the potential of PJI being miscoded should not bias this association differently in one group more than the other. Furthermore, it is possible despite matching on several key demographics that the study populations were not equivalent and that patient selection bias on other nonmatched factors may have occurred. Finally, we did not analyze risk factors for UKA PJI as the overall number of patients who developed a UKA PJI, even in this large database sample, was low.

Conclusions

In this large non-Medicare population study of rigorously matched cohorts of patients undergoing either UKA or TKA, UKA conferred a significantly lower risk of PJI at 90 days (RR = 0.57) and 1 year (RR = 0.66) than TKA. However, the overall incidence of PJI in both UKA and TKA patients at 90 days and 1 year was overall low at <1%. As such, surgeons should continue to balance the short-term benefits and lower perioperative complications of UKA for patients compared to the improved long-term durability of TKA when counseling patients on knee arthroplasty options.

Conflicts of interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: Edwin P. Su designed and receives royalties for a cementless unicompartmental knee arthroplasty.

References

- [1] Bosch LC, Bala A, Denduluri SK, et al. Reimbursement and complications in outpatient vs inpatient unicompartmental arthroplasty. *J Arthroplasty* 2020;35(6S):S86.
- [2] Chalmers BP, Mehrotra KG, Sierra RJ, Pagnano MW, Taunton MJ, Abdel MP. Reliable outcomes and survivorship of unicompartmental knee arthroplasty for isolated compartment osteonecrosis. *Bone Joint J* 2018;100-B(4):450.
- [3] Parratte S, Ollivier M, Lunebourg A, Abdel MP, Argenson JN. Long-term results of compartmental arthroplasties of the knee: long term results of partial knee arthroplasty. *Bone Joint J* 2015;97-B(10 Suppl A):9.
- [4] Hansen EN, Ong KL, Lau E, Kurtz SM, Lonner JH. Unicompartmental knee arthroplasty has fewer complications but higher revision rates than total knee arthroplasty in a study of large United States databases. *J Arthroplasty* 2019;34(8):1617.
- [5] Kozinn SC, Scott R. Unicompartmental knee arthroplasty. *J Bone Joint Surg Am* 1989;71(01):145.
- [6] Lombardi Jr AV, Berend KR, Berend ME, et al. Current controversies in partial knee arthroplasty. *Instr Course Lect* 2012;61:347.
- [7] Berend KR, Lombardi Jr AV, Adams JB. Obesity, young age, patellofemoral disease, and anterior knee pain: identifying the unicompartmental arthroplasty patient in the United States. *Orthopedics* 2007;30(5 Suppl):19.
- [8] Hurst JM, Berend KR, Morris MJ, Lombardi Jr AV. Abnormal preoperative MRI does not correlate with failure of UKA. *J Arthroplasty* 2013;28(9 Suppl):184.
- [9] Berend KR, Lombardi Jr AV, Morris MJ, Hurst JM, Kavolus JJ. Does preoperative patellofemoral joint state affect medial unicompartmental arthroplasty survival? *Orthopedics* 2011;34(9):e494.
- [10] Kahlenberg CA, Richardson SS, Gruskay JA, Cross MB. Trends in utilization of total and unicompartmental knee arthroplasty in the United States. *J Knee Surg* 2020.
- [11] Bolognesi MP, Greiner MA, Attarian DE, et al. Unicompartmental knee arthroplasty and total knee arthroplasty among Medicare beneficiaries, 2000 to 2009. *J Bone Joint Surg Am* 2013;95(22):e174.
- [12] Kayani B, Konan S, Tahmassebi J, Rowan FE, Haddad FS. An assessment of early functional rehabilitation and hospital discharge in conventional versus robotic-arm assisted unicompartmental knee arthroplasty: a prospective cohort study. *Bone Joint J* 2019;101-B(1):24.
- [13] Begum FA, Kayani B, Morgan SDJ, Ahmed SS, Singh S, Haddad FS. Robotic technology: current concepts, operative techniques and emerging uses in unicompartmental knee arthroplasty. *EFORT Open Rev* 2020;5(5):312.
- [14] Sousa PL, Sculco PK, Mayman DJ, Jerabek SA, Ast MP, Chalmers BP. Robots in the operating room during hip and knee arthroplasty. *Curr Rev Musculoskelet Med* 2020;13(3):309.
- [15] Vakharia RM, Sodhi N, Cohen-Levy WB, Vakharia AM, Mont MA, Roche MW. Comparison of patient demographics and utilization trends of robotic-assisted and non-robotic-assisted unicompartmental knee arthroplasty. *J Knee Surg* 2019;10.
- [16] Dyrhovden GS, Lygre SHL, Badawy M, Gøthesen Ø, Furnes O. Have the causes of revision for total and unicompartmental knee arthroplasties changed during the past two decades? *Clin Orthop Relat Res* 2017;475(7):1874.
- [17] Liddle AD, Pandit H, Judge A, Murray DW. Patient-reported outcomes after total and unicompartmental knee arthroplasty: a study of 14,076 matched patients from the National Joint Registry for England and Wales. *Bone Joint J* 2015;97-B(6):793.
- [18] Liddle AD, Judge A, Pandit H, Murray DW. Adverse outcomes after total and unicompartmental knee replacement in 101,330 matched patients: a study of data from the National Joint Registry for England and Wales. *Lancet* 2014;384(9952):1437.
- [19] Wilson HA, Middleton R, Abram SGF, et al. Patient relevant outcomes of unicompartmental versus total knee replacement: systematic review and meta-analysis. *BMJ* 2019;364:l352.
- [20] Burn E, Sanchez-Santos MT, Pandit HG, et al. Ten-year patient-reported outcomes following total and minimally invasive unicompartmental knee arthroplasty: a propensity score-matched cohort analysis. *Knee Surg Sports Traumatol Arthrosc* 2018;26(5):1455.
- [21] van der List JP, Chawla H, Villa JC, Pearle AD. The role of patient characteristics on the choice of unicompartmental versus total knee arthroplasty in patients with medial osteoarthritis. *J Arthroplasty* 2017;32(3):761.
- [22] Zuiderbaan HA, van der List JP, Khamaisy S, et al. Unicompartmental knee arthroplasty versus total knee arthroplasty: which type of artificial joint do patients forget? *Knee Surg Sports Traumatol Arthrosc* 2017;25(3):681.
- [23] Lenguerrand E, Whitehouse MR, Beswick AD, et al. Risk factors associated with revision for prosthetic joint infection following knee replacement: an observational cohort study from England and Wales. *Lancet Infect Dis* 2019;19(6):589.
- [24] Mahure SA, Feng JE, Schwarzkopf RM, Long WJ. Differences in pain, opioid use, and function following unicompartmental knee arthroplasty compared to total knee arthroplasty. *J Arthroplasty* 2020;35(9):2435.

- [25] Chalmers BP, Kapadia M, Chiu YF, Henry MW, Miller AO, Carli AV. Treatment and outcome of periprosthetic joint infection in unicompartmental knee arthroplasty. *J Arthroplasty* 2020;35(7):1917.
- [26] Hernandez NM, Petis SM, Hanssen AD, Sierra RJ, Abdel MP, Pagnano MW. Infection after unicompartmental knee arthroplasty: a high risk of subsequent complications. *Clin Orthop Relat Res* 2019;477(1):70.
- [27] Wyles CC, Vargas-Hernandez JS, Carlson SW, Carlson BC, Sierra RJ. Single-dose perioperative antibiotics do not increase the risk of surgical site infection in unicompartmental knee arthroplasty. *J Arthroplasty* 2019;34(7S):S327.
- [28] Nettrour JF, Ellis RT, Hansen BJ, Keeney JA. High failure rates for unicompartmental knee arthroplasty in morbidly obese patients: a two-year minimum follow-up study. *J Arthroplasty* 2020;35(4):989.
- [29] Lum ZC, Crawford DA, Lombardi Jr AV, et al. Early comparative outcomes of unicompartmental and total knee arthroplasty in severely obese patients. *Knee* 2018;25(1):161.
- [30] Gruskay J, Richardson S, Schairer W, et al. Incidence and safety profile of outpatient unicompartmental knee arthroplasty. *Knee* 2019;26(3):708.
- [31] Furnes O, Espehaug B, Lie SA, Vollset SE, Engesaeter LB, Havelin LI. Failure mechanisms after unicompartmental and tricompartmental primary knee replacement with cement. *J Bone Joint Surg Am* 2007;89(3):519.
- [32] Weston JT, Watts CD, Mabry TM, Hanssen AD, Berry DJ, Abdel MP. Irrigation and debridement with chronic antibiotic suppression for the management of infected total knee arthroplasty: a Contemporary Analysis. *Bone Joint J* 2018;100-B(11):1471.

Appendix**Supplementary Table 1**

Codes used.

Description	Codes used
Unicompartmental arthroplasty	CPT-27446
Total knee arthroplasty	CPT-27447
Periprosthetic joint infection	ICD-9-D-99669, ICD-10-D-T8450XA, ICD-10-D-T8450XD, ICD-10-D-T8450XS, ICD-10-D-T8459XA, ICD-10-D-T8459XD, ICD-10-D-T8459XS, ICD-10-D-T8453XA, ICD-10-D-T8453XD, ICD-10-D-T8453XS, ICD-10-D-T8454XA, ICD-10-D-T8454XD, ICD-10-D-T8454XS, ICD-9-D-99660, ICD-9-D-99666, ICD-9-D-99667
Explantation	CPT-27486, CPT-27487, CPT-27488, ICD-9-P-0080, ICD-9-P-0081, ICD-9-P-0082, ICD-9-P-0083, ICD-9-P-8006, ICD-9-P-8155, ICD-10-P-0SPC0JC, ICD-10-P-0SPC0JZ, ICD-10-P-0SPD0JC, ICD-10-P-0SPD0JZ, ICD-10-P-0SPT0JZ, ICD-10-P-0SPU0JZ, ICD-10-P-0SPV0JZ, ICD-10-P-0SPW0JZ, ICD-10-P-0SWC0JC, ICD-10-P-0SWC0JZ, ICD-10-P-0SWD0JC, ICD-10-P-0SWD0JZ, ICD-10-P-0SWT0JZ, ICD-10-P-0SWU0JZ, ICD-10-P-0SWV0JZ, ICD-10-P-0SWW0JZ
Debridement, antibiotics, and implant retention	CPT-27486, CPT-27487, CPT-27488, ICD-9-P-0080, ICD-9-P-0081, ICD-9-P-0082, ICD-9-P-0083, ICD-9-P-8006, ICD-9-P-8155, ICD-10-P-0SPC0JC, ICD-10-P-0SPC0JZ, ICD-10-P-0SPD0JC, ICD-10-P-0SPD0JZ, ICD-10-P-0SPT0JZ, ICD-10-P-0SPU0JZ, ICD-10-P-0SPV0JZ, ICD-10-P-0SPW0JZ, ICD-10-P-0SWC0JC, ICD-10-P-0SWC0JZ, ICD-10-P-0SWD0JC, ICD-10-P-0SWD0JZ, ICD-10-P-0SWT0JZ, ICD-10-P-0SWU0JZ, ICD-10-P-0SWV0JZ, ICD-10-P-0SWW0JZ