



University of Kentucky
UKnowledge

Pharmaceutical Sciences Faculty Publications

Pharmaceutical Sciences

7-1-2015

New Approach to Develop Ultra-High Inhibitory Drug Using the Power Function of the Stoichiometry of the Targeted Nanomachine or Biocomplex

Dan Shu

University of Kentucky, dan.shu@uky.edu

Fengmei Pi

University of Kentucky, fengmei.pi@uky.edu

Chi Wang

University of Kentucky, chi.wang@uky.edu

Peng Zhang

University of Michigan

See next page for additional authors

Right click to open a feedback form in a new tab to let us know how this document benefits you.

Follow this and additional works at: https://uknowledge.uky.edu/ps_facpub

 Part of the [Nanomedicine Commons](#), [Oncology Commons](#), and the [Pharmacy and Pharmaceutical Sciences Commons](#)

New Approach to Develop Ultra-High Inhibitory Drug Using the Power Function of the Stoichiometry of the Targeted Nanomachine or Biocomplex

Digital Object Identifier (DOI)

<http://dx.doi.org/10.2217/nnm.15.37>

Notes/Citation Information

Published in *Nanomedicine*, v. 10, no. 12, p. 1881-1897.

The document available for download is the authors' post-peer-review final draft of the article.

Authors

Dan Shu, Fengmei Pi, Chi Wang, Peng Zhang, and Peixuan Guo

New approach to develop ultra-high inhibitory drug using the power-function of the stoichiometry of the targeted nanomachine or biocomplex

Dan Shu^{1,2,3}, Fengmei Pi^{1,2,3}, Chi Wang,^{2,4} Peng Zhang⁵ and Peixuan Guo^{1,2,3*}

¹Nanobiotechnology Center; ²Markey Cancer Center; ³Department of Pharmaceutical Sciences; ⁴Department of Biostatistics; University of Kentucky, Lexington, KY 40536, USA. ⁵Department of Surgery, University of Michigan Health System, Ann Arbor, MI-48109, USA.

Abstract

Aims: To find a method for developing potent drugs and to prove a hypothesis that drug inhibition potency depends on the stoichiometry of the targeted biocomplex.

Methods: Phi29 DNA-packaging motor components were used to test this model for different stoichiometries. Virion assembly efficiency was assayed with Yang Hui's Triangle: $(p + q)^Z =$

$\sum_{M=0}^Z \binom{Z}{M} p^{Z-M} q^M$, where Z=stoichiometry, M=drugged subunits in each biocomplex, p and q represent the fraction of drugged and non-drugged subunits in the population.

Results: Inhibition efficiency follows a power function. When number of drugged subunits to block the function of the biocomplex K=1, the fraction of uninhibited biocomplex equals q^Z . Thus, stoichiometry has a multiplicative effect on inhibition. Targets with a thousand subunits showed the highest inhibition effect, followed by those with six and a single subunit. Complete inhibition of virus replication was found when Z = 6.

Conclusion: Drug inhibition potency depends on the stoichiometry of the targeted components of the biocomplex or nano-machine. The inhibition effect follows a power function of the stoichiometry of the target biocomplex.

Key words: Drug target, phi29 viral assembly, nanomotor, hexameric ATPase, bionanotechnology, nanobiotechnology, binomial distribution.

Introduction

Bacteria, viruses and cells contain biocomplexes and nanomachines composed of multiple subunits, such as biomotors [1,2,3,4], pumps [5], exosomes [6,7,8], valves [9,10,11], membrane pores [12,13,14,15], chaperonins[16], PCNA[17], ATPase [18,19], and tubes [20]. From a nanobiotechnological standpoint, these

nanomachines can be used and converted to build sophisticated nano-devices including molecular sensors [21,22,23], patterned arrays, tuators [24], chips, microelectromechanical systems (MEMS) [25], molecular sorters [26], single pore DNA sequencing apparatus [12,13,21,27] or other revolutionary electronic and optical devices [28,29]. From a pharmaceutical standpoint, these multi-subunit biocomplexes or nanomachines have a potential for use as drug targets for therapeutics, as well as diagnostic applications such as pathogen detection, disease diagnosis, drug delivery, and treatment of diseases [22,23,30,31]. In the ASCE (Additional Strand Catalytic E) family including the AAA+ (ATPases Associated with diverse cellular Activities) and the FtsK-HerA superfamily in bacteria, viruses and cells, there are nanomotors that perform a wide range of functions [19,32,33] critical to chromosome segregation, bacterial binary fission, DNA/RNA and cell component transportation, membrane sorting, cellular reorganization, cell division, RNA transcription, as well as DNA replication, riding, repair, and recombination [1,34,35,36]. One of the directions of NIH Roadmap is to utilize these cellular nanomachines and biocomplexes for biomedical applications.

Acquired drug resistance has become a major reason for failure treatment of a range of diseases, i.e., the chemotherapy for cancer, bacterial or viral infections. Drug resistance of cancer has escalated and has partially contributed to the ~600,000 deaths in the USA in 2012 [37]. HIV drug resistance has also become a major issue [38]. Many common pathogens have become resistant to current drug treatments, with new infectious diseases on the rise. The use of multidrug-resistant agents in biological weapons has created a previously unrealized challenge [39]. Thus, there is a need to develop new treatment strategies to combat drug resistance with new drug development methods.

The first FDA-approved drug to treat multidrug-resistant tuberculosis, bedaquiline, follows a novel mechanism of inhibiting the bacterial ATP synthase of *M. tuberculosis* and other mycobacterial species, but had little activity against other bacteria [40]. To combat multidrug resistance in cancer, several approaches have been explored. One method is to target components that are highly important for the growth of the biological entity [41,42]. Another approach uses nano-drug delivery carriers that are expected to enhance the binding efficiency of drugs to cancer cells[43,44,45,46], or cocktail therapy [47]. A third approach is to develop new combinational drugs with higher potency by acting on multiple targets [48,49]. This involves identifying multiple targets that when treated leads to a synergetic effect and optimizing the design of multi-target ligands[50].

*Address correspondence to: Peixuan Guo, peixuan.guo@uky.edu
Department of Pharmaceutical Sciences, University of Kentucky
789 S. Limestone Street, Room # 576
Lexington, KY, USA 40536-0596
Phone: 859-218-0128
Fax: 859-257-1307




Figure 1: Stoichiometry of viral DNA packaging motor. (A) Illustration of Phi29 DNA packaging motor composed of 1 copy of genomic DNA that revolves through the channel wall (left panel), 6 copies of pRNA, 6 copies of ATPase gp16 and a connector channel. (B) Yang Hui Triangle. (C) Illustration of Z=6 and K=1, drug targeting anyone subunit of the complex will block its activity. (D) AFM image of hexameric re-engineered pRNA rings. (E). 3D structure of hexameric pRNA ring top view and side view from the crystal structure of 3WJ (PDB ID: pRNA 3WJ, 4KZ2). (F) A crystal hexameric structure of AAA+ Protein CbbX with top view and side view [85] (PDB ID: CbbX, 3Zuh, <http://www.ebi.ac.uk/pdbe/emdb/>) (G) Structure of the hexameric AAA+ molecular machine ClpC with adaptor protein MecA. [86] (PDB ID: MecA-ClpC, 3PXG).

The approach of developing highly potent drugs through targeting of protein or RNA complexes with high stoichiometry has never been reported due to challenges in comparing efficacies of two drugs that can be confused by target essentiality with binding affinity. For instance, if two drugs target two stoichiometrically different targets, it becomes extremely difficult to prove whether the difference in drug efficiency is due to differences in their target binding affinity or essential level in the growth of the biological organism. In order to quantify effects from targeting biocomplexes of different stoichiometry, a well-studied multi-component system is required that allows empirical comparison of functional inhibition of individual components that are composed of different number of subunits.

An example of one nanobiomachine is the dsDNA translocation motor, for which the ATPase protein is a pivotal component that assembles into a hexameric ring structure and translates the action of ATP binding and hydrolysis into mechanical motion to translocate DNA physically. The DNA packaging motor of bacteriophage phi29 (Fig. 1A) [9,51,52,53] is composed of three essential co-axial rings:

1) a dodecameric connector ring located at the vertex of the viral procapsid; 2) a hexameric packaging RNA (pRNA) ring [52] bound to the N-terminus of the connector [54], and 3) a hexameric ring of ATPase gp16 attached to the helical region of pRNA [10,19,55], powered through the hydrolysis of ATP resulting in DNA packaging. The use of Yang Hui's Triangle (Fig. 1B) or binomial distribution to determine the stoichiometry of the pRNA was first reported in 1997 [56]. The use of similar mathematical methods to determine the stoichiometry of the protein subunits has also been reported more recently [51]. The copy number of ATP molecules required to package one full phi29 genomic dsDNA was predicted to be 10000 [57]. It has recently been shown that this hexameric motor uses a revolution mechanism without rotation to translocate its genomic DNA [10,19,33,35,36,58,59].

In this report, we hypothesize that the inhibitory efficiency of a drug is related to the stoichiometry of its targeted biocomplex; the higher the stoichiometry of the target complex, the more efficient the drug. This can lead to the development of potent therapeutics against high-stoichiometric biomachines or biocomplexes as drug targets. We

proved this hypothesis by using a mutant subunit as the drugged inactive target to calculate the theoretical inhibition efficiency via binomial distribution, and compared with experimental data from a defined *in vitro* viral assembly system. Since biomotors share certain common structures and operation mechanisms [1,36,59,60], the approach in drug development reported here should have general applications especially in developing new generations of drugs for combating the rising acquired drug resistance in viruses, bacteria, and cancers [38,61,62].

Materials and Methods

Preparation of mutant genomic dsDNA

Phi29 genomic DNA-gp3 was purified from *B. subtilis* SpoA12 cells by CsCl gradient ultracentrifugation as described previously [63]. Mutant dsDNA was prepared by digesting the phi29 genomic dsDNA with *Eco*R1 restriction enzyme in fast digest buffer (Fermentas) at 37 °C for 1 hour followed by ethanol precipitation. The mutant DNA was tested by 1% agarose gel electrophoresis, stained by ethidium bromide (Sigma) and imaged by Typhoon (GE).

Preparation of mutant pRNA

Wild-type phi29 pRNA and inactive mutant as drugged pRNA were prepared by *in vitro* transcription. In the inactive mutant pRNA, the first four bases "UUCA" located at the 5' end were mutated to "GGGG". *Bgl*III digested plasmid pRT71 was used as DNA template [64] in the PCR reaction for both RNAs. Oligonucleotide 5'-TAA TAC GAC TCA CTA TAG GGG TGG TAC-3' and 5'-TTA TCA AAG TAG CGT GCA C-3' were used as primers for mutant pRNA. RNAs were then transcribed by T7 RNA polymerase using double-stranded DNA generated from PCR, as described before [65]. The RNA from *in vitro* transcription was further purified by 8 M urea 8% polyacrylamide gel electrophoresis as described previously [64].

Preparation of mutant ATPase gp16

The purification of *wild-type* gp16 has been described previously [63]. The walker B mutant gp16 was constructed by introducing mutations in the gp16 gene. The amino acid residues D255 and E256 in walker B motif of gp16 were mutated to E255 and D256, respectively. The mutation was introduced with the Stratagene Quick Change site-directed mutagenesis kit using appropriate primers. The expression and purification of protein were carried out followed a published procedure [51].

Antisense oligonucleotides

Antisense oligonucleotides P3 and P15 were designed to be reversely complementary to different regions on the pRNA molecule and chemically synthesized by IDT. P3 oligo (5'-TTGCCATGATTGACAAC-3') targets the region of 83-99 nucleotides at the 3' end of pRNA, P15 oligo (5'-AAGTACCGTACCATTGA-3') targets the region of 1-17 nucleotides at the 5' end of pRNA. P8 oligo (5'-TAATACGACTCACTATAGGGGTGGTAC-3') was designed as a non-targeting control in the test. 1 µl of individual oligos at 100 µM were mixed with 1 µl of pRNA at 4 µM and dialyzed on a 0.025 µm mixed cellulose esters VSWP filter membrane (Millipore Corp) against TBE buffer (89 mM Tris-HCl, pH 8.3, 89 mM boric acid, 2.5

mM EDTA) at room temperature for 15 min. The purified RNA complex was used for *in vitro* phi29 assembly assay.

In vitro phi29 assembly assay

Purified components were subjected to *in vitro* viral assembly assay as described previously [66]. Briefly, 10 µg of purified procapsids were mixed with 100 ng of pRNA in 5 µl of reaction buffer (10 mM ATP, 6 mM 2-mercaptoethanol, and 3 mM spermidine in TMS buffer) at room temperature for 30 min. Purified DNA-gp3 and gp16 were then added and the reaction mixture was incubated at room temperature for one hour to initiate DNA packaging. Finally, the DNA filled procapsids were incubated with 10 µl of gp8.5-9 extract from *E. coli* containing plasmid pARgp8.5-9 and 20 µl of gp11-14 extract from *E. coli* to complete the infectious phage assembly.

The newly assembled infectious viruses were plated with inoculated *B. subtilis* bacteria Su⁴⁴ cells onto a half LB plate covered with top agar. After 12 hour incubation at 37 °C, the viral assembly efficiency (plaque-forming unit, PFU) was calculated by counting the formed plaque numbers. Mixing different ratios of mutant with *wild-type* components, while keeping all other components the same, the viral assembly efficiency (PFU) versus ratio of mutant components gave an empirical curve for vial assembly inhibition assay, and it was compared with theoretical curves from the binomial distribution equation.

In vivo viral assembly assay

Plasmid pRBwtRNA containing the pRNA coding sequence under T7 promoter was constructed by ligating the fragment coding pRNA sequence and T7 promoter into pRB381-L550 vector (modified and kindly provided by M. Wang and H. Zalkin) following a previously described method [67]. Plasmid pRBmutRNA contained mutant pRNA under its natural promoter PE1 sequence, and the mutation was changing sequence 5'UUGA-3' at its 3' end to 5'GGGG-3'. The DNA fragments coding mutant pRNA sequence and PE1 sequence were prepared by PCR as described previously [67]; and digested with *Hind*III-*Bgl*III restriction enzyme. The mutant pRNA sequence coding fragment was further ligated with a 6.0kb fragment from pRB381-L550 that was digested with *Hind*III and partially digested with *Bgl*III.

The plasmids pRBmutRNA, pRBwtRNA, and pRB381-L550 were transformed into *B. subtilis* cells following methods described previously [67]. The *B. subtilis* cells harboring transformed plasmids were incubated in 416 medium with 10 mg/ml of neomycin for 3 hours at 37 °C and then plated onto LB-neomycin (10 mg/ml) plates for plaque formation analysis.

Results

The Definition of "Stoichiometry"

The definition of the stoichiometry in this report is different from conventional definition of stoichiometry used to evaluate drug efficiency. Conventionally the concept of stoichiometry refers to the number of a drug binding to each target molecule, which is also known as B_{max} . In this study the definition of stoichiometry refers to

the copy number of subunit within a biocomplex or the nanomachine that serves as drug target.

The definition of "K value", and K = 1 is one key for ultra-high inhibition efficacy

Suppose a biocomplex drug target contains Z copies of subunits, then K is the copy number ($K \leq Z$) of drugged subunits required to inhibit the function of the complex or the nanomachine. As an analogy to the difference between the parallel circuit and the serial circuit, when the Christmas lights are arranged in a parallel circuit, any light bulbs that are burnt out will not affect other bulbs. But in a serial circuit, any one light bulb that is broken will stop the entire lighting system, which is $K=1$. Thus, the K value is the key to the probability of inactive nanomachines or biocomplexes by combination and permutation of all subunits. K equals 1 is critical for such ultra-high inhibition effect. The foundation of the approach in this report is the difference in probability of inhibited biocomplexes in systems of different K values with combination and permutation algorithms. Biological systems display complicated reactions. Many reactions involve multiple subunits to work cooperatively sequentially or processively to accomplish one essential biological function [33,68,69,70,71,72,73,74,75]. Single assembly pathways have been reported in the viral assembly system [76,77]. In most cases of the sequential, cooperative, and processive action, inactivation of any one, not necessary all, of the subunits will result in inhibition of its function, thus $K = 1$. Drug synergism was utilized in multi-target drug therapy; in short, a drug combination can simultaneously act on multiple targets in disease networks to produce a synergistic effect [50,78]. However, our design reported here is unique from the conventional synergistic approach. We suggest that using multi-subunit biocomplexes as drug target could lead to development of ultra-high potent drugs. In a conventional six-component system, for example one drug is designed to target component #3 to stop the entire system, since the drug can only target component #3, the condition fits the model of $Z=1$ and $K=1$. Thus, the inhibition efficiency and substrate targeting efficiency (p) of drug will be in linear relationship. However, in the system in this report, the entire system will be blocked when drug targets any subunit of a hexamer, which is $Z=6$ and $K=1$. Thus the probability of remaining undrugged targets will be q^6 , where q represents the fraction of untargeted hexamer subunits; in other words, the drug inhibition efficiency will be $1-q^6$, which increases following a power function compared to the linear for conventional mono-subunit approaches.

Assuming that at least K copies of drugged subunits were needed to deactivate the nanomachine or biocomplex, the probability of functional biocomplexes in the presence of various ratios of inhibited and wild-type subunits could be predicted from equation 2. When $K=1$, it implies that drug binding to one subunit will inactivate the subunit, and one drugged subunit per multi-subunit complex is sufficient to inhibit the overall function of the complex. The inhibition efficiency by drugs targeting multi-subunit biocomplexes with stoichiometry of Z will equal $1 - q^z$, as shown in table 1. An example for such a probability calculation when $Z = 6$ and $K = 1$ is as follows: since it was assumed that 6 ($Z = 6$) copies of subunits per element were required for function and one drugged subunit ($K = 1$) was sufficient to block its activity, all elements possessing 1 to 5 copies of drugged subunits would be non-functional (Fig. 1C). Only those

Table1. Predicted inhibition efficiency of drugs targeting biocomplex, K=1

Inhibited subunit (p)	Probability of the multisubunit complex is active with:											
	Z=1, K=1	Z=2, K=1	Z=3, K=1	Z=4, K=1	Z=5, K=1	Z=6, K=1	Z=7, K=1	Z=8, K=1	Z=9, K=1	Z=10, K=1	Z=11, K=1	Z=12, K=1
0	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
0.1	0.90	0.81	0.73	0.66	0.59	0.53	0.48	0.43	0.39	0.35	0.31	0.28
0.2	0.80	0.64	0.51	0.41	0.33	0.26	0.21	0.17	0.13	0.11	0.09	0.07
0.3	0.70	0.49	0.34	0.24	0.17	0.12	0.08	0.06	0.04	0.03	0.02	0.01
0.4	0.60	0.36	0.22	0.13	0.08	0.05	0.03	0.02	0.01	0.01	0.00	0.00
0.5	0.50	0.25	0.13	0.06	0.03	0.02	0.01	0.00	0.00	0.00	0.00	0.00
0.6	0.40	0.16	0.06	0.03	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00
0.7	0.30	0.09	0.03	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
0.8	0.20	0.04	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
0.9	0.10	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

complexes possessing 6 copies of normal subunits will be functional. The chance for a complex containing 6 copies of unaffected subunits in a population is q^6 and the inhibition efficiency will be $1 - q^6$.

Rationale behind selection of multi-subunit biocomplexes as efficient drug targets

Mechanisms for drug inhibition of organism growth are to block or stop an essential biological element from functioning. When a drug is designed to target the subunit of a complex with targeting efficiency p , a fraction of subunits will not interact with the drug (a percentile given as q , $p+q=1$) and will remain active and exert their function properly. Some biological elements are monomers containing only one subunit, while other biological elements, such as the bio-motors of hexameric AAA+ family, consist of multiple-subunits [19,34]. Conventional drugs are designed to inhibit pathogenesis through targeting of a single subunit molecule, such as an enzyme or a structural protein of a virus. In this situation, the inhibition efficiency is proportional to the substrate targeting efficiency p and the effect is proportional to the first order of p . As described above, in most cases of sequential action or cooperatives in multiple subunit complexes, inactivation of one, not all, of the subunits will result in inhibition of its function. Thus, if complexes containing Z copies of subunits exercise their function in a sequential and cooperative way, then $K=1$, and the fraction of the uninhibited active biocomplex will be q^z , a higher order with regards to the stoichiometry. The inhibition proportion will equal $1 - q^z$.

In this investigation, a well-defined *in vitro* phi29 viral assembly system was used to represent a multi-subunit nano-machine target, with the mutant component representing a target component that have been inactivated by an effective drug. Then, the inhibition efficiencies by targeting different elements of the phi29 DNA packaging motor with different stoichiometry were compared. The viral assembly competition assays combined with binomial distribution analysis illustrated the concept that drug targeting functional biological complexes of a higher-stoichiometry has a higher efficiency than drug acting on a single subunit target.

When the target element is a monomer containing only one subunit, the inhibition efficiency can be calculated through a binomial distribution (equation 1), where p and q are the fractions of drugged (substrate targeting efficiency) and undrugged (normal active elements) subunits, respectively ($p + q = 1$).

$$X = (p + q)^1 \quad (1)$$

However, when the target element contains multiple subunits, a high order binomial distribution (equation 2) is applied to calculate the drug inhibition effect by finding the ratio of resulted active and inactive complexes, where Z represents the total number of subunits (the stoichiometry) in one biocomplex and M represents the number of drugged subunits in one biocomplex.

$$X = (p + q)^Z = \sum_{M=0}^Z \binom{Z}{M} p^{Z-M} q^M = \sum_{M=0}^Z \left(\frac{Z!}{M!(Z-M)!} \right) p^{Z-M} q^M \quad (2)$$

For example, if Z is 3, the probability of all combinations of drugged subunits (M) and undrugged subunits (N; M + N = Z) in a given biocomplex entity can be determined by the expansion of equation 2: $(p + q)^3 = p^3 + 3p^2q + 3pq^2 + q^3 = 1$. That is, the probability of a complex element possessing three copies of drugged subunits in the population is p^3 , two copies of drugged and one copy of undrugged or *wild-type* subunit is $3p^2q$, one copy of drugged and two copies of undrugged subunits is $3pq^2$, and three copies of undrugged subunits is q^3 . Assuming there were 70% ($p=0.7$) of subunits inactivated by bound drugs, and 30% ($q=0.3$) unaffected subunits in the population, then the percentage of elements possessing at least two copies of normal subunits would be the sum of those possessing one copy of drugged and two copies of undrugged *wild-type* subunits, $3pq^2$, and those possessing three copies of native subunits is q^3 . That is $3pq^2 + q^3 = 3(0.7)(0.3)^2 + (0.3)^3 = 0.216 = 21.6\%$. In another example, if one complex contains 6 subunits, and 5 out of the 6 subunits need to remain uninhibited in order to be biologically functional, the active complex ratio in the population will be the sum of: 1) the probability of each element containing 5 undrugged subunits, and 2) the probability of each element containing 6 undrugged subunits.

The probability X in the population displaying a certain combination of undrugged versus drugged subunits can be predicted by a binomial distribution, as shown in equation 2. Table 2 shows the probability of a given element with M drugged and N undrugged subunits at increasing percentages of drugged subunits in the

population, considering that the total subunits in one element (Z) is 3 or 12. The formula, $\frac{Z!}{M!N!} p^M q^N$ (from equation 2) was used to calculate each combination probability value, the coefficient $\frac{Z!}{M!N!}$ in this equation can also be calculated using Yang Hui Triangle, which is also called Pascal's Triangle, or binomial distribution (Fig. 1B)[79].

In vitro virus assembly system used for testing the hypothesis

The highly sensitive *in vitro* phi29 assembly system was used to determine the inhibition efficiency of drugs targeting multi-subunit complexes [56,66,76,80]. Bacteriophage phi29 DNA packaging motor contains one copy of genomic dsDNA, 6 copies of packaging RNA, 6 copies of ATPase protein gp16 and more than 10000 copies of ATP. The stoichiometry of RNA in phi29 has been proven by extensive studies including single-molecule studies[81] AFM images (Fig.1D) [82,83], pRNA crystal structure determination (Fig.1E) [84], and mathematical studies [56]. The stoichiometry of gp16 in phi29 has been proven by multiple approaches including native gel binding, capillary electrophoresis assays, Hill constant determination, and by titration of mutant subunits using binomial distribution [19,33]. Many other AAA+ superfamily members have been found to be hexamers as well [85,86,87,88,89,90,91], such as a red type rubisco activase AAA+ protein CbbX (Fig. 1F) [85], MecA-ClpC molecular machine (Fig. 1G)[86]. The copy number of ATP molecules was calculated based on the fact that 6 ATP molecules are required to package one pitch of dsDNA with 10.5 basepairs (bp) [92], thus 1 ATP is used to package 1.7 bp. The entire phi29 genome is composed of 19.4 kbp, thus, it is expected that more than 10000 ATP molecules are required to package the entire phi29 genome. The phi29 DNA nanomotor which packages an entire genomic DNA into the procapsid can be treated as a disease model for drug inhibition efficiency analysis.

In vitro testing of the hypothesis using DNA element with stoichiometry of 1

Inhibited Subunits (p)	Z=3				Z=12												
	M=0, N=3	M=1, N=2	M=2, N=1	M=3, N=0	M=0, N=12	M=1, N=11	M=2, N=10	M=3, N=9	M=4, N=8	M=5, N=7	M=6, N=6	M=7, N=5	M=8, N=4	M=9, N=3	M=10, N=2	M=11, N=1	M=12, N=0
0	1.00	0.00	0.00	0.00	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
0.1	0.73	0.24	0.03	0.00	0.28	0.38	0.23	0.09	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
0.2	0.51	0.38	0.10	0.01	0.07	0.21	0.28	0.24	0.13	0.05	0.02	0.00	0.00	0.00	0.00	0.00	0.00
0.3	0.34	0.44	0.19	0.03	0.01	0.07	0.17	0.24	0.23	0.16	0.08	0.03	0.01	0.00	0.00	0.00	0.00
0.4	0.22	0.43	0.29	0.06	0.00	0.02	0.06	0.14	0.21	0.23	0.18	0.10	0.04	0.01	0.00	0.00	0.00
0.5	0.13	0.38	0.38	0.13	0.00	0.00	0.02	0.05	0.12	0.19	0.23	0.19	0.12	0.05	0.02	0.00	0.00
0.6	0.06	0.29	0.43	0.22	0.00	0.00	0.00	0.01	0.04	0.10	0.18	0.23	0.21	0.14	0.06	0.02	0.00
0.7	0.03	0.19	0.44	0.34	0.00	0.00	0.00	0.00	0.01	0.03	0.08	0.16	0.23	0.24	0.17	0.07	0.01
0.8	0.01	0.10	0.38	0.51	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.05	0.13	0.24	0.28	0.21	0.07
0.9	0.00	0.03	0.24	0.73	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.09	0.23	0.38	0.28
1	0.00	0.00	0.00	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.00




Figure 2: Theoretical plot (with variable Z) and empirical data to illustrate inhibition efficiency with drug targeting to genomic DNA (Z=1). (A) Gel showing the phi29 genome DNA treated with endonuclease EcoR1. (B) Plot of virion assembly derived from binomial distribution equation 2, which showed that the DNA has stoichiometry of 1. (C) Viral assembly inhibition effect of mutant DNA as model of drugged component with Z=1, showing the linear relationship to p with low slope.

The inhibition efficiency of drugs targeting a single subunit substrate was tested by *in vitro* phi29 assembly inhibition by mutating the genomic dsDNA (Fig. 2A). Various ratios of mutant DNA were mixed with *wild-type* DNA in *in vitro* viral assembly assays. The empirical curve of viral assembly efficiency against drugged mutant DNA ratio fits well with the theoretical curve from binomial distribution for Z = 1 and K = 1 (Fig. 2B). This suggests that when designing drug targeting the genomic DNA in phi29 nanomotor, it is expected to be a first order inhibition response. Comparing the *in vitro* phi29 assembly inhibition, by adding drugged mutant DNA, with simply diluting *wild-type* DNA concentration as a control, revealed that the drugged mutant DNA didn't cause much difference (Fig. 2C). The results showed that the inhibition effect of drugs targeting the substrate with stoichiometry of 1 is minimal.

In vitro testing of the hypothesis using RNA elements with stoichiometry of 6

The pRNA of phi29 contains two domains; a head-loop domain

essential for procapsid binding and a helix domain essential for DNA translocation (Fig. 3A, upper panel) [30,93,94]. The right-hand loop and left-hand loop of two pairing pRNA molecules can interact with each other by complementary base pairing. Extensive studies have led to the conclusion that 6 copies of pRNA form a hexameric ring which binds to the procapsid for virus activity [81,82,83,84]. Drugged mutant pRNA was constructed by mutating 4 nucleotide sequences at the 5'end region of pRNA (Fig. 3A, lower panel), which has been shown to compete with *wild-type* pRNA for procapsid binding, but was found to be deficient in allowing DNA packaging to occur [67]. The theoretical curves generated using the expansion of binomial distribution equation while total subunit number Z is 6 and varying K number from 1 to 6 are shown in Fig. 3B. Fitting the empirical data from phage assembly efficiency at different ratios of drugged mutant pRNAs into the theoretical curves, the empirical data fit into the theoretical curve of Z = 6 and K = 1. It suggested that the pRNA oligomer ring is composed of six copies of pRNA subunits and one subunit of the pRNA multimer blockage is sufficient to block the phage assembly activity. Comparing the empirical curve for viral




Figure 3: Theoretical plot (K=1 to 6) and empirical data to illustrate inhibition efficiency with drug targeting pRNA (Z=6). (A) The sequence and secondary structure of wild-type pRNA of phi29 DNA packaging motor (upper panel) and inactive mutant pRNA with 4 bases mutation at 5'end of the DNA translocation domain serving as a model of drugged inactive pRNA(lower panel). (B) Fitting the phage assembly inhibition result by inactive mutant pRNA with the theoretical plots derived from Equation 2 matched with Z=6 and K=1. (C) Comparing the viral assembly inhibition effect by drugged pRNA at different concentration with the undrugged pRNA with same dilution factor.

assembly efficiency against different ratios of drugged mutant pRNAs with the *wild-type* pRNA concentration dilution control, addition of drugged mutant pRNA showed a much stronger inhibition effect (Fig.3C).

To further prove the concept that drugs targeting biocomplex with high stoichiometry causes stronger inhibition effect, antisense oligonucleotides which can bind to pRNA molecules were designed as mock drugs in the viral assembly assay. The oligonucleotides P15, and P3 were designed to target the 5'-end and 3'-end regions on pRNA, respectively. It was confirmed that the antisense oligonucleotides can be hybridized to pRNA by gel shift assay (data not shown). When mixing the antisense oligonucleotides with *wild-type* pRNA for *in vitro* phi29 assembly assay, complete inhibition effects were shown by antisense oligonucleotides P15, and P3, but not with the non-targeting control oligonucleotide P8[95]. By mixing the non-targeting oligo P8 with pRNA, it generated plaques with 4.4×10^6 PFU on the plate.

In vivo testing of the hypothesis using RNA elements with stoichiometry of 6

Formation of the hexameric ring of pRNA in the phi29 dsDNA packaging motor has been discovered through biochemical and structural studies [52,81,84,96,97,98,99,100,101,102,103] and activity assays [94,104]. The observed high inhibition efficiency by drugged mutant pRNA on phi29 assembly *in vitro* was striking [67,105]. To test whether such a high level of inhibition was attainable *in vivo*, pRBmutRNA plasmid expressing a pRNA with 4-base mutation at the 3' end (Fig. 4A) was transformed into *B. subtilis* DE1 cells. Plasmid pRBwtRNA contained the pRNA coding sequence but do not express pRNA in *B. subtilis* DE1 cells, and vector pRB381-L550 was introduced as well as a negative control. The results showed that only cells harboring pRBmutRNA plasmid were completely resistant to plaque formation by *wild-type* phi29 virus infection. Control cells, including *B. subtilis* 12A cells alone, *B. subtilis* DE1 cells carrying vector pRB381-L550 alone, and cells carrying a *wild-type* pRNA coding sequence but no expression plasmid pRBwtRNA were all positive for plaque formation (Fig. 4B). The ability of mutant pRNAs generated in cells by plasmid pRBmutRNA completely inhibited plaque formation indicated that hexameric pRNA in DNA packaging nano-motor may be a potential target for developing potent antiviral agents [67].




Figure 4: Complete inhibition of viral assembly *in vivo* by mutant pRNA as a model of drugged complex ($Z=6$). (A) Inactivation of pRNA by introducing a 4-nucleotide mutation at the 3' end. (B) Virion production by *wild-type* phi29 infection using host cell *B. subtilis* harboring plasmid expressing mutant pRNA, *wild-type* pRNA, or plasmid only.

In vitro testing of the hypothesis using the ATPase with stoichiometry of 6

Hexameric folding of ATPase gp16 protein in the phi29 dsDNA packaging motor has been discovered [1,19,33,35]. The hexameric gp16 protein complex functions as ATPase like many other AAA+ superfamily members. ATP binding to one subunit of gp16 stimulates the ATPase to change its conformation from having a lower affinity to one having a higher affinity for dsDNA.

Determination of gp16 stoichiometry was carried out by *in vitro* phage assembly assay and based on the binomial distribution of *wild-type* and Walker B mutant gp16 [51]. Different ratios of drugged Walker B mutant gp16 were mixed with undrugged gp16 to test the inhibition efficiency of gp16 mutation on phi29 DNA packaging motor. Assuming K equals 1 and the total copy number of gp16 (Z) is between 1 and 12, twelve theoretical curves for the production of phi29 virion against the ratio of the Walker B mutant corresponding to the stoichiometry (Z) of 1 to 12 were generated according to equation 2. The empirical data nearly perfectly overlapped the theoretical curve of $Z=6$, $K=1$ [51]. This data suggested that the ATPase gp16 components of phi29 DNA packaging motor have a stoichiometry of six, and only one copy of the drugged gp16 can block the phi29 motor function. Comparing the inhibition effect of adding mutant gp16 with *wild-type* gp16 at different concentrations, it showed that adding mutant gp16 had a much stronger inhibition effect than the *wild-type* gp16 concentration dilution control (Fig. 5A). Comparing the inhibition effect of mutation on hexameric gp16 to the effect of mutation on single subunit target DNA, the gp16 mutation displayed a much stronger inhibition effect on virus assembly than the same ratio of DNA mutation, indicating the hexameric ATPase protein complex of virus assembly system should also be an efficient target for generating new anti-virus drugs with high potency.

In vitro testing of the hypothesis using ATP with stoichiometry of more than 10000

It has been reported that 6 ATP molecules are required to package one pitch of dsDNA with 10.5bp [90], thus 1 ATP is used to package 1.7bp. As the entire phi29 genomic DNA has 19,000 base pairs, it is expected that more than 10000 ATP molecules are required to package the entire phi29 genome. Since concerning ATP, the functional unit displayed in Fig 5 is the viral production expressed as plaque-forming unit (PFU), so the production of one functional unit of PFU require 10000 ATP subunits to package one genomic DNA. Thus, the ATP in one phi29 nanomotor can be regarded as a stoichiometry of 10000. One non-hydrolysable ATP analogue γ -S-ATP was treated as the drugged subunit that mixed with ATP at different ratios to test the inhibition effect of γ -S-ATP on phi29 assembly efficiency. It was found that the inhibition curve of mutant ATP fits into the theoretical curve between $Z=100$, $K=1$ and $Z=60$, $K=1$ (Fig. 5B). The empirical ATP value derived from binomial distribution assay was different from real condition, since the binomial distribution equation was based on a condition that each subunits has the same binding affinity to the biocomplex in the targeted nanomotor, but due to the change of the γ -S-ATP structure, it has a ATPase gp16 binding affinity lower than the normal ATP. Furthermore, the affinity difference in each subunit has a multiplicative effect in the nanomotor's activity. Thus, there is a big




Figure 5: Comparison of inhibition efficiency using targets with different Z values. (A) Virion production inhibition effect of mutant gp16 (Z=6) at different concentration. (B) Inhibition efficiency by γ -S-ATP with ATP with high Z value. (C) Comparing the virus assembly inhibition effect by drugged components of DNA, pRNA, gp16 and ATP with stoichiometry of 1, 6, 6, 10000 respectively.

discrepancy between the curves with predicted Z value and the empirical Z value.

Comparing virus assembly inhibition effect using different components, the γ -S-ATP showed a severe inhibition effect (Fig. 5C). Adding 20% of gamma-s-ATP nearly completely inhibited the viral assembly. Comparing the inhibition effect targeting to ATP, pRNA, ATPase gp16, and DNA with stoichiometry of 10000, 6, 6, and 1, respectively, γ -S-ATP showed the strongest inhibition effect, while drugged mutant pRNA and mutant gp16 showed stronger inhibitory effect than mutant DNA (Fig.5C). For example, adding 20% mutant DNA caused 20% inhibition effect in viral assembly, while 20% of drugged mutant pRNA exerted 74% of inhibition effect on viral assembly and 20% of γ -S-ATP almost completely inhibited the viral assembly, indicating the higher the stoichiometry, the stronger the inhibition efficacy.

Mathematical reasoning for the increase of inhibition efficacy

Using a biological complex with higher stoichiometry as drug target will substantially reduce the proportion of non-inhibited complex. For K=1, the proportion of non-inhibited complex is q^Z . Table 3 compares the proportion of non-inhibited complex from two populations with Z=6 and Z=1, respectively, with varied substrate targeting efficiency (p) when K=1. For example, when q=0.4, the proportion of non-inhibited complex is $q^Z=0.4^1=0.4$ for Z=1, K=1. Therefore, only 1-0.4=60% of complex is inhibited. In contrast, for

Table 3. Comparison of proportion of non-inhibited complex between Z=6 and Z=1 when K=1 but having equal drug targeting efficacy

Substrate targeting efficacy (p)	Proportion of non-inhibited complex from the population with Z=6	Proportion of non-inhibited complex from the population with Z=1	Ratio of the proportions of non-inhibited complex from the two populations with Z=6 and Z=1	Reduction (fold) in proportion of non-inhibited complex comparing Z=6 and Z=1
0	1	1	1	1
0.1	0.5314	0.9	0.5905	17
0.2	0.2621	0.8	0.3277	3.1
0.3	0.1176	0.7	0.1681	5.9
0.4	0.0467	0.6	0.0778	12.9
0.5	0.0156	0.5	0.0312	32
0.6	0.0041	0.4	0.0102	98
0.7	7.00E-04	0.3	0.0024	416
0.8	100E-04	0.2	0.0003	3333
0.9	100E-06	0.1	100E-05	10000

Z=6, K=1, the proportion of non-inhibited complex is $q^Z=0.4^6=0.0041$. Therefore, 1-0.0041=99.59% of complex is inhibited. The ratio of the proportions of non-inhibited complex equals 0.0041/0.4 = 0.0102, indicating a 1/0.0102=98-fold decrease in the proportion of non-inhibited complex. One more example is to use the drug targeting efficiency p = 0.9 to compare the inhibition efficiency between two groups with Z=6 and Z=1. For Z=6, K=1, the proportion of inhibited complex is $1-q^Z=1-0.1^6=0.999999$. The proportion of non-inhibited complex is $q^Z=0.1^6=1E-6$. For Z=1, K=1, the proportion of inhibited complex is $1-q^Z=1-0.1=0.9$. The proportion of non-inhibited complex is $q^Z=0.1$. The ratio of inhibition efficiency equals to $1E-6/0.1 = 1E-5$, indicating a 10000-fold increase in inhibition efficiency (Table 3).

The equation displays inhibitory effect with a power function of stoichiometry since when K=1, the percentage of uninhibited biocomplexes in the population equal to q^Z . Since $(P + q) = 1$, thus $q \leq 1$, thus the larger the Z, the smaller the value of q^Z . That is to say, the higher the stoichiometry, the smaller number of the uninhibited background will display. With the same substrate targeting efficacy, p, the inhibition efficiency is determined by z, the power of the equation component. The inhibitory effect is a power function concerning the stoichiometry. Thus, the higher the stoichiometry, the more efficient the inhibition comparing the drugs with same binding affinity.

The half maximal inhibitory concentration (IC_{50}) is commonly used to evaluate drug effect, which quantitatively indicates how much of a particular drug is needed to inhibit a given biological process by half.

If we denote p_{IC50} as the percentage of drugged subunit needed to reach to 50% inhibition in the *in vitro* assay in the defined system, thus $1 - (1 - p_{IC50})^Z = 50\%$. Solving this equation, $p_{IC50} = 1 - 0.5^{1/Z}$. Figure 6 shows the relationship between stoichiometry (Z) and drug targeting level p to reach the inhibition effect (IC), where p is the combined result of drug binding efficacy and drug concentration (dosage). When biocomplexes with stoichiometry of Z are used as drug targets, the dosage of drug or

the drug binding affinity presented by percentage of drugged subunits to reach IC_{50} , IC_{25} , or IC_{75} decreases. This clearly shows that as Z increases, PIC_{50} decreases (Fig. 6), and hence the drug is more potent.




Figure 6: Relationship between stoichiometry (Z) and drug targeting level (p , a combined result of drug binding efficacy and drug concentration) to reach the inhibition effect (IC).

Discussion

Aiming to find a method for developing drugs with ultra-high potency, we proposed that the inhibition efficiency of a given drug depends on stoichiometry of the biocomplex or bio-machine that was used as drug target. Here the definition of the stoichiometry is different from conventional definition of stoichiometry used to evaluate drug efficiency. Conventional thinking in drug development emphasizes stoichiometry which refers to the number of drug binding per target molecule, which is also known as B_{max} . In this study the definition of stoichiometry refers to the copy number of subunit within a biocomplex that serves as drug target. We used phi29 viral components with a series of variable but known stoichiometry as mock drug targets to test the hypothesis. Both *in vitro* and *in vivo* virion assembly assays were employed to compare the inhibition efficiency targeting to components with different numbers of subunit stoichiometry. Viral inhibition efficiency was analyzed with Yang Hui's (Pascal's) Triangle (or known as binomial distribution). It was found that inhibition efficiency on virus replication correlates to the component stoichiometry of nano-machine as drug target. It displayed power law inhibitory effect since when $K=1$, the percentage of uninhibited biocomplexes in the population equal to q^z . With the same q and same K value, the inhibition efficiency is determined by z , the number of subunits within the biocomplex or the bio-machine as drug target. Here z serves as the power in the equation, thus, the inhibitory effect is the power of the stoichiometry. Empirical data demonstrated that the target with thousand-subunits shows higher inhibition effect than the targets with six subunits, and in turn higher than the target with single subunit.

In evaluation of drug effect, two parameters were commonly used. One is the half maximal inhibitory concentration (IC_{50}),

which quantitatively indicates how much of a particular drug is needed to inhibit a given biological process by half. It is universally used as a measure of drug potency in pharmacological research. Another important parameter is the median lethal dose (LD_{50}), which is also known as 50% of lethal concentration (LC_{50}). LD_{50} is frequently used to indicate a substance's acute toxicity. Obviously, the usefulness of a drug will depend on the ratio of LD_{50} to IC_{50} . The larger this ratio, the safer the drug. By ways of increasing the inhibition efficiency through targeting to the components with high stoichiometry, the IC_{50} of a drug will decrease. As a result, lower concentration of drug will be required for reaching a desired effect, resulting in a reduced toxicity of the drug.

Most of current anti-cancer, anti-virus or anti-bacteria drugs target single enzymes or single proteins. Our data showed that drugs selected to target components, biocomplexes, or nano-machines with high copy numbers could lead to a much higher efficacy, and it could potentially solve the problem of low drug effect and multi-drug resistance.

Conclusions

Targeting the functional biological units with higher stoichiometries will have a higher efficiency of inhibition. The inhibition effect is power, other than proportional, and the power, is the copy number of the drug-targeted element of the machine. The new theory developed herein suggests that potent drugs can be developed by targeting biocomplex with high stoichiometry, and a complete inhibition of virus, bacterium, or cancer is possible if a bio-machine with high stoichiometry is identified. Since bio-motors share certain common structure and operation mechanism in viruses, bacteria, and cells, this approach should have general application in drug development.

Future Perspective

Living systems contain many elegant arrays, motors and nanomachines that are multi-subunit complex. As reported here, these biocomplex with high copy number of components can serve as potent drug targets. For example, most members of the AAA+ family are hexamer [19,87,88,106,107,108]. However, these machines are common in living systems therefore the specificity and toxicity is an issue. For bacteria and virus, since our goal is to kill them nonexclusively, the specificity and toxicity is not an issue as long as the target biocomplex is not identical to that in human body. For cancers drugs, as long as a mutation is found in the multiple-subunit biocomplex, it will be an ideal target for potent drug.

Executive summary

Aim

- A method for developing potent drugs is sought.

Hypothesis

- Drug inhibition potency depends on the stoichiometry of the targeted biocomplex.

Approach:

- Phi29 viral components with variable stoichiometry were used as model to prove the hypothesis
- Virion assembly efficiency was assayed and analyzed with

$$\text{Yang Hui's Triangle: } (p + q)^Z = \sum_{M=0}^Z \binom{Z}{M} p^{Z-M} q^M.$$

Results:

- Inhibition efficiency displayed a power function of the stoichiometry of the target biocomplexes. The uninhibited biocomplex in population can equals to q^Z . Thus, the inhibitory effect is a power of the stoichiometry.
- Targets with thousand-subunit showed higher inhibition effect than with six subunits, and in turn higher than target with single subunit.
- A complete inhibition of virus, bacterium, or cancer was demonstrated when targets with high stoichiometry was used as target.

Conclusion

- Drug inhibition potency depends on the stoichiometry of the targeted components of the biocomplex or nano-machine.
- The inhibition effect displayed a power function of the stoichiometry of the target biocomplex.
- Since bio-motors share certain common structure and operation mechanism in viruses, bacteria, and cells, this approach should have general application in drug development.

Acknowledgements

The authors would like to thank Timothy S. Tracy for a critical review on the manuscript and a very insightful and impressive comments of the approach reported; Esther Penni Black, Patrick McNamara, Mario Vieweger, Joseph Chappell, Brad Anderson, Chaoyuan Liu, for constructive comments on the manuscript and helpful discussion; Ashwani Sharma, Daniel Binzel, Daniel Jasinski and Jordan Surface for editing and formatting this manuscript, and Zhengyi Zhao for the drawing of figures. This publication was made possible by grants from NIH/NIBIB EB012135, EB003730, and NIH/NCI CA151648 to Peixuan Guo, and the Markey Cancer Grant P30CA177558 to B. Mark Evers. Funding to Peixuan Guo's Endowed Chair in Nanobiotechnology position is by the William Fairish Endowment Fund. Peixuan Guo is a co-founder of Kylene Therapeutic, Inc, Biomotor and RNA Nanotechnology Development Corp. Ltd, and RNA Nano, LLC..

References

1. Guo P, Zhao Z, Haak J et al. Common Mechanisms of DNA translocation motors in Bacteria and Viruses Using One-way Revolution Mechanism without Rotation. *Biotechnology Advances* 32, 853-872 (2014).

** This review reports that revolution mechanism is commonly used in bacteria and viruses which avoid DNA coiling in translocation the lengthy genomic dsDNA helix.

2. Jankowsky E, Fairman ME, Yang Q. RNA Helicases: Versatile ATP-Driven Nanomotors. *J.Nanosci.Nanotechnol.* 5, 1983-1989 (2005).

3. Guo PX, Lee TJ. Viral nanomotors for packaging of dsDNA and dsRNA. *Mol.Microbiol.* 64, 886-903 (2007).

4. Molineux IJ, Panja D. Popping the cork: mechanisms of phage genome ejection. *Nat Rev.Microbiol.* 11(3), 194-204 (2013).

5. Lee BS, Lee SC, Holliday LS. Biochemistry of mechanoenzymes: Biological motors for nanotechnology. *Biomedical Microdevices* 5(4), 269-280 (2003).

6. Pegtel DM, Cosmopoulos K, Thorley-Lawson DA et al. Functional delivery of viral miRNAs via exosomes. *Proc.Natl.Acad.Sci.U.S.A* 107(14), 6328-6333 (2010).

7. Valadi H, Ekstrom K, Bossios A, Sjostrand M, Lee JJ, Lotvall JO. Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells. *Nat Cell Biol* 9(6), 654-659 (2007).

8. Wendler F, Bota-Rabasedas N, Franch-Marro X. Cancer becomes wasteful: emerging roles of exosomes(dagger) in cell-fate determination. *J.Extracell.Vesicles.* 2 (22390), 1-9 (2013).

9. Fang H, Jing P, Haque F, Guo P. Role of channel Lysines and "Push Through a One-way Valve" Mechanism of Viral DNA packaging Motor. *Biophysical Journal* 102, 127-135 (2012).

10. Zhang H, Schwartz C, De Donatis GM, Guo P. "Push Through One-Way Valve" Mechanism of Viral DNA Packaging. *Adv.Virus Res* 83, 415-465 (2012).

11. Jing P, Haque F, Shu D, Montemagno C, Guo P. One-Way Traffic of a Viral Motor Channel for Double-Stranded DNA Translocation. *Nano Lett.* 10, 3620-3627 (2010).

12. Kasianowicz JJ, Brandin E, Branton D, Deamer DW. Characterization of individual polynucleotide molecules using a membrane channel. *Proc.Natl.Acad.Sci.U.S.A* 93(24), 13770-13773 (1996).

13. Butler TZ, Pavlenok M, Derrington IM, Niederweis M, Gundlach JH. Single-molecule DNA detection with an engineered MspA protein nanopore. *Proc.Natl.Acad.Sci.U.S.A* 105(52), 20647-20652 (2008).

14. Wendell D, Jing P, Geng J et al. Translocation of double-stranded DNA through membrane-adapted phi29 motor protein nanopores. *Nat.Nanotechnol.* 4, 765-772 (2009).

15. Pastoriza-Gallego M, Rabah L, Gibrat G et al. Dynamics of unfolded protein transport through an aerolysin pore. *J.Am.Chem.Soc.* 133(9), 2923-2931 (2011).

16. Ezzell C. How Chaperonins Monitor Their Protein Charges. *The Journal of NIH Research* 6, 31-34 (1994).

17. Maga G, Hubscher U. Proliferating cell nuclear antigen (PCNA): a dancer with many partners. *J.Cell Sci.* 116(Pt 15), 3051-3060 (2003).

18. Rubinchik S, Parris W, Gold M. The in vitro ATPases of bacteriophage I terminase and its large subunit, gene product A. *J Biol Chem* 269, 13586-13593 (1994).

19. Schwartz C, De Donatis GM, Fang H, Guo P. The ATPase of the phi29 DNA-packaging motor is a member of the hexameric AAA+ superfamily. *Virology* 443, 20-27 (2013).

* This paper confirmed the stoichiometry of ATPase in phi29 motor is a hexamer and provided data suggesting that the phi29 motor ATPase belongs to classical hexameric AAA+ superfamily.

20. Sun L, Young LN, Zhang X et al. Icosahedral bacteriophage PhiX174 forms a tail for DNA transport during infection. *Nature* 505(7483), 432-435 (2014).

21. Haque F, Li J, Wu H-C, Liang X-J, Guo P. Solid-state and biological nanopore for real-time sensing of single chemical and sequencing of DNA. *Nano Today* 8, 56-74 (2013).

22. Haque F, Lunn J, Fang H, Smithrud D, Guo P. Real-Time Sensing and Discrimination of Single Chemicals Using the Channel of Phi29 DNA Packaging Nanomotor. *ACS Nano* 6, 3251-3261 (2012).

23. Wang S, Haque F, Rychahou PG, Evers BM, Guo P. Engineered Nanopore of Phi29 DNA-Packaging Motor for Real-Time Detection of Single Colon Cancer Specific Antibody in Serum. *ACS Nano* 7, 9814-9822 (2013).

24. Fennimore AM, Yuzvinsky TD, Han WQ, Fuhrer MS, Cumings J, Zettl A. Rotational actuators based on carbon nanotubes. *Nature* 424, 408-410 (2003).

25. Craighead HG. Nanoelectromechanical systems. *Science* 290, 1532-1536 (2000).

26. Gerion D, Parak WJ, Williams SC, Zanchet D, Micheel CM, Alivisatos AP. Sorting fluorescent nanocrystals with DNA. *J Am.Chem Soc.* 124, 7070-7074 (2002).
27. Geng J, Wang S, Fang H, Guo P. Channel size conversion of Phi29 DNA-packaging nanomotor for discrimination of single- and double-stranded nucleic acids. *ACS Nano* 7(4), 3315-3323 (2013).
28. McNally B, Singer A, Yu Z, Sun Y, Weng Z, Meller A. Optical recognition of converted DNA nucleotides for single-molecule DNA sequencing using nanopore arrays. *Nano Lett.* 10(6), 2237-2244 (2010).
29. Chandler EL, Smith AL, Burden LM, Kasianowicz JJ, Burden DL. Membrane surface dynamics of DNA-threaded nanopores revealed by simultaneous single-molecule optical and ensemble electrical recording. *Langmuir* 20(3), 898-905 (2004).
30. Shu D, Shu Y, Haque F, Abdelmawla S, Guo P. Thermodynamically stable RNA three-way junctions for constructing multifunctional nanoparticles for delivery of therapeutics. *Nature Nanotechnology* 6, 658-667 (2011).
31. Haque F, Shu D, Shu Y et al. Ultrastable synergistic tetravalent RNA nanoparticles for targeting to cancers. *Nano Today* 7, 245-257 (2012).
32. Crozat E, Grainge I. FtsK DNA translocase: the fast motor that knows where it's going. *ChemBiochem.* 11, 2232-2243 (2010).
33. Schwartz C, De Donatis GM, Zhang H, Fang H, Guo P. Revolution rather than rotation of AAA+ hexameric phi29 nanomotor for viral dsDNA packaging without coiling. *Virology* 443, 28-39 (2013).
- * This paper shows how the nanomotor in phi29 virus works with a revolution mechanism using six copies of ATPase.
34. Hanson PI, Whiteheart SW. AAA+ proteins: have engine, will work. *Nat.Rev.Mol Cell Biol.* 6, 519-529 (2005).
35. Guo P, Schwartz C, Haak J, Zhao Z. Discovery of a new motion mechanism of biomotors similar to the earth revolving around the sun without rotation. *Virology* 446, 133-143 (2013).
36. De-Donatis G, Zhao Z, Wang S et al. Finding of widespread viral and bacterial revolution dsDNA translocation motors distinct from rotation motors by channel chirality and size. *Cell Biosci* 4, 30- (2014).
- ** This paper reports that the revolution motor nanomachine is widespread among biological systems, and can be distinguished from rotation motors by channel size and chirality.
37. Batra S, Adekola KUA, Rosen ST, Shanmugam M. Cancer Metabolism as a Therapeutic Target. *ONCOLOGY-NEW YORK* 27, 460-467 (2013).
38. Buskin SE, Zhang S, Thibault CS. Prevalence of and viral outcomes associated with primary HIV-1 drug resistance. *Open.AIDS J.* 6, 181-187 (2012).
39. Spellberg B, Powers JH, Brass EP, Miller LG, Edwards JE, Jr. Trends in antimicrobial drug development: implications for the future. *Clin.Infect.Dis.* 38(9), 1279-1286 (2004).
40. Lakshmanan M, Xavier AS. Bedaquiline - The first ATP synthase inhibitor against multi drug resistant tuberculosis. *J.Young.Pharm.* 5(4), 112-115 (2013).
41. Aird KM, Ding XY, Baras A et al. Trastuzumab signaling in ErbB2-overexpressing inflammatory breast cancer correlates with X-linked inhibitor of apoptosis protein expression. *Molecular Cancer Therapeutics* 7(1), 38-47 (2008).
42. Park SH, Kim H, Song BJ. Down regulation of bcl2 expression in invasive ductal carcinomas is both estrogen- and progesterone-receptor dependent and associated with poor prognostic factors. *Pathol.Oncol.Res.* 8(1), 26-30 (2002).
43. Akinc A, Querbes W, De S et al. Targeted delivery of RNAi therapeutics with endogenous and exogenous ligand-based mechanisms. *Mol.Ther.* 18(7), 1357-1364 (2010).
44. Aronov O, Horowitz AT, Gabizon A, Gibson D. Folate-targeted PEG as a potential carrier for carboplatin analogs. Synthesis and in vitro studies. *Bioconjugate Chem.* 14(3), 563-574 (2003).
45. Bae YH, Park K. Targeted drug delivery to tumors: myths, reality and possibility. *J.Control Release* 153(3), 198-205 (2011).
46. Guo P, Haque F, Hallahan B, Reif R, Li H. Uniqueness, advantages, challenges, solutions, and perspectives in therapeutics applying RNA nanotechnology. *Nucleic Acid Ther.* 22(4), 226-245 (2012).
47. Lusky K. HIV's potent cocktail. *Contemp.Longterm.Care* 22(12), 59- (1999).
48. Chatterjee A, Chattopadhyay D, Chakrabarti G. miR-17-5p Downregulation Contributes to Paclitaxel Resistance of Lung Cancer Cells through Altering Beclin1 Expression. *PLoS ONE* 9(4), e95716- (2014).
49. Sanchez C, Chan R, Bajgain P et al. Combining T-cell immunotherapy and anti-androgen therapy for prostate cancer. *Prostate Cancer Prostatic.Dis.* 16(2), 123-31 (2013).
50. Ito M, Zhao N, Zeng Z, Chang CC, Zu Y. Synergistic growth inhibition of anaplastic large cell lymphoma cells by combining cellular ALK gene silencing and a low dose of the kinase inhibitor UO126. *Cancer Gene Ther.* 17(9),633-44 (2010).
51. Fang H, Zhang P, Huang LP et al. Binomial distribution for quantification of protein subunits in biological nanoassemblies and functional nanomachines. *Nanomedicine.* 10(7),1433-40 (2014).
- **This is the first report to describe how to use the Yang Hui's Triangle (binomial distribution) to determine the stoichiometry of protein subunits in biocomplex. It precisely confirmed that phi29 motor contains six copies of ATPase gp16 and one mutant subunit would cause motor to stop.
52. Guo P, Zhang C, Chen C, Trottier M, Garver K. Inter-RNA interaction of phage phi29 pRNA to form a hexameric complex for viral DNA transportation. *Mol.Cell.* 2, 149-155 (1998).
- * This is the first paper to reveal that the pRNA of phi29 DNA packaging motor forms a hexameric ring, and prove-of -concept of RNA nanotechnology since this paper shows that by bottom-up assembly, the RNA nanoparticles of dimers, trimers and hexamers can be constructed using the reengineered RNA fragments derived from phi29 motor pRNA.
53. Hugel T, Michaelis J, Hetherington CL et al. Experimental test of connector rotation during DNA packaging into bacteriophage phi29 capsids. *Plos Biology* 5, 558-567 (2007).
54. Xiao F, Moll D, Guo S, Guo P. Binding of pRNA to the N-terminal 14 amino acids of connector protein of bacterial phage phi29. *Nucleic Acids Res* 33, 2640-2649 (2005).
55. Guo P, Peterson C, Anderson D. Prohead and DNA-gp3-dependent ATPase activity of the DNA packaging protein gp16 of bacteriophage phi29. *J Mol Biol* 197, 229-236 (1987).
56. Trottier M, Guo P. Approaches to determine stoichiometry of viral assembly components. *J.Virol.* 71, 487-494 (1997).
- *This is the first report to describe how to use the Yang Hui's Triangle (binomial distribution) to determine the stoichiometry of biocomplex or nanomachine.
57. Guo P, Peterson C, Anderson D. Initiation events in in vitro packaging of bacteriophage phi29 DNA-gp3. *J Mol Biol* 197, 219-228 (1987).
58. Zhao Z, Khisamutdinov E, Schwartz C, Guo P. Mechanism of one-way traffic of hexameric phi29 DNA packaging motor with four electropositive relaying layers facilitating anti-parallel revolution. *ACS Nano* 7, 4082-4092 (2013).
59. Guo P, Grainge I, Zhao Z, Vieweger M. Two classes of nucleic acid translocation motors: rotation and revolution without rotation. *Cell Biosci.* 4(1), 54- (2014).
60. Casjens SR. The DNA-packaging nanomotor of tailed bacteriophages. *Nat Rev.Microbiol.* 9(9), 647-657 (2011).
61. Boal AK, Ilhan F, DeRouchey JE, Thurn-Albrecht T, Russell TP, Rotello VM. Self-assembly of nanoparticles into structured spherical and network aggregates. *Nature* 404(6779), 746-748 (2000).
62. Holohan C, Van SS, Longley DB, Johnston PG. Cancer drug resistance: an evolving paradigm. *Nat.Rev.Cancer* 13(10), 714-726 (2013).
63. Guo P, Grimes S, Anderson D. A defined system for in vitro packaging of DNA-gp3 of the *Bacillus subtilis* bacteriophage phi29. *Proc.Natl.Acad.Sci.USA* 83, 3505-3509 (1986).

64. Zhang CL, Lee C-S, Guo P. The proximate 5' and 3' ends of the 120-base viral RNA (pRNA) are crucial for the packaging of bacteriophage ϕ 29 DNA. *Virology* 201, 77-85 (1994).
65. Shu D, Huang L, Hoepflich S, Guo P. Construction of ϕ 29 DNA-packaging RNA (pRNA) monomers, dimers and trimers with variable sizes and shapes as potential parts for nano-devices. *J.Nanosci.Nanotechnol.* 3, 295-302 (2003).
66. Lee CS, Guo P. A highly sensitive system for the assay of in vitro viral assembly of bacteriophage ϕ 29 of *Bacillus subtilis*. *Virology* 202, 1039-1042 (1994).
67. Trottier M, Zhang CL, Guo P. Complete inhibition of virion assembly in vivo with mutant pRNA essential for phage ϕ 29 DNA packaging. *J.Virol.* 70, 55-61 (1996).
68. Chen C, Guo P. Sequential action of six virus-encoded DNA-packaging RNAs during phage ϕ 29 genomic DNA translocation. *J.Virol.* 71(5), 3864-3871 (1997).
69. Stitt BL, Xu Y. Sequential hydrolysis of ATP molecules bound in interacting catalytic sites of *Escherichia coli* transcription termination protein Rho. *J.Biol.Chem.* 273(41), 26477-26486 (1998).
70. Kammerer RA, Schulthess T, Landwehr R, Lustig A, Fischer D, Engel J. Tenascin-C hexabrachion assembly is a sequential two-step process initiated by coiled-coil alpha-helices. *J.Biol.Chem.* 273(17), 10602-10608 (1998).
71. Lisal J, Tuma R. Cooperative mechanism of RNA packaging motor. *J Biol Chem* 280, 23157-23164 (2005).
72. Zhang Z, Lewis D, Strock C et al. Detailed characterization of the cooperative mechanism of Ca(2+) binding and catalytic activation in the Ca(2+) transport (SERC) ATPase. *Biochemistry* 39(30), 8758-8767 (2000).
73. Sun H, Squier TC. Ordered and cooperative binding of opposing globular domains of calmodulin to the plasma membrane Ca-ATPase. *J.Biol.Chem.* 275(3), 1731-1738 (2000).
74. Hiller R, Carmeli C. Kinetic analysis of cooperative interactions induced by Mn2+ binding to the chloroplast H(+)-ATPase. *Biochemistry* 29(26), 6186-6192 (1990).
75. Persechini A, Hartshorne DJ. Cooperative behavior of smooth muscle myosin. *Fed.Proc.* 41(12), 2868-2872 (1982).
76. Lee CS, Guo P. Sequential interactions of structural proteins in phage ϕ 29 procapsid assembly. *J.Virol.* 69, 5024-5032 (1995).
77. Casjens S, Hendrix R. Control mechanisms in dsDNA bacteriophage assembly. In: *The Bacteriophages Vol.1.* Calendar R (Eds.). Plenum Publishing Corp., 15-92 (1988).
78. Qian X, Ren Y, Shi Z et al. Sequence-dependent synergistic inhibition of human glioma cell lines by combined temozolomide and miR-21 inhibitor gene therapy. *Mol Pharm.* 9(9), 2636-2645 (2012).
79. Zhang H, Shu D, Huang F, Guo P. Instrumentation and metrology for single RNA counting in biological complexes or nanoparticles by a single molecule dual-view system. *RNA* 13, 1793-1802 (2007).
80. Chen C, Trottier M, Guo P. New approaches to stoichiometry determination and mechanism investigation on RNA involved in intermediate reactions. *Nucleic Acids Symposium Series* 36, 190-193 (1997).
81. Shu D, Zhang H, Jin J, Guo P. Counting of six pRNAs of ϕ 29 DNA-packaging motor with customized single molecule dual-view system. *EMBO J.* 26, 527-537 (2007).
- * This is the first report describing the use of single fluorophore photobleaching technique to count subunits in biocomplex, and documents the "seeing is believing" to confirm that ϕ 29 DNA packaging motor contains six copies of packaging pRNA.
82. Shu Y, Haque F, Shu D et al. Fabrication of 14 Different RNA Nanoparticles for Specific Tumor Targeting without Accumulation in Normal Organs. *RNA* 19, 766-777 (2013).
83. Shu Y, Shu D, Haque F, Guo P. Fabrication of pRNA nanoparticles to deliver therapeutic RNAs and bioactive compounds into tumor cells. *Nat Protoc.* 8(9), 1635-1659 (2013).
84. Zhang H, Endrizzi JA, Shu Y et al. Crystal Structure of 3WJ Core Revealing Divalent Ion-promoted Thermostability and Assembly of the ϕ 29 Hexameric Motor pRNA. *RNA* 19, 1226-1237 (2013).
85. Mueller-Cajar O, Stotz M, Wendler P, Hartl FU, Bracher A, Hayer-Hartl M. Structure and function of the AAA+ protein CbbX, a red-type Rubisco activase. *Nature* 479, 194-199 (2011).
86. Wang F, Mei Z, Qi Y et al. Structure and mechanism of the hexameric MecA-ClpC molecular machine. *Nature* 471, 331-335 (2011).
87. Aker J, Hesselink R, Engel R et al. In vivo hexamerization and characterization of the Arabidopsis AAA ATPase CDC48A complex using forster resonance energy transfer-fluorescence lifetime imaging microscopy and fluorescence correlation spectroscopy. *Plant Physiology* 145(2), 339-350 (2007).
88. Willows RD, Hansson A, Birch D, Al-Karadaghi S, Hansson M. EM single particle analysis of the ATP-dependent Bchl complex of magnesium chelatase: an AAA(+) hexamer. *J Struct Biol* 146(1-2), 227-233 (2004).
89. Chen YJ, Yu X, Egelman EH. The hexameric ring structure of the *Escherichia coli* RuvB branch migration protein. *J.Mol.Biol.* 319(3), 587-591 (2002).
90. Happonen LJ, Oksanen E, Liljeroos L, Goldman A, Kajander T, Butcher SJ. The structure of the NTPase that powers DNA packaging into *Sulfolobus turreted* icosahedral virus 2. *J.Virol.* 87, 8388-8398 (2013).
91. Snider J, Thibault G, Houry WA. The AAA+ superfamily of functionally diverse proteins. *Genome Biol* 9(4), 216-216 (2008).
92. Schwartz C, Fang H, Huang L, Guo P. Sequential action of ATPase, ATP, ADP, Pi and dsDNA in procapsid-free system to enlighten mechanism in viral dsDNA packaging. *Nucleic Acids Res.* 40, 2577-2586 (2012).
93. Trottier M, Garver K, Zhang C, Guo P. DNA-packaging pRNA as target for complete inhibition of viral assembly in vitro and in vivo. *Nucleic Acids Symposium Series* 36, 187-189 (1997).
94. Chen C, Zhang C, Guo P. Sequence requirement for hand-in-hand interaction in formation of pRNA dimers and hexamers to gear ϕ 29 DNA translocation motor. *RNA* 5, 805-818 (1999).
95. Zhang CL, Trottier M, Guo PX. Circularly permuted viral pRNA active and specific in the packaging of bacteriophage ϕ 29 DNA. *Virology* 207, 442-451 (1995).
96. Zhang F, Lemieux S, Wu X et al. Function of hexameric RNA in packaging of bacteriophage ϕ 29 DNA in vitro. *Mol.Cell.* 2, 141-147 (1998).
97. Hendrix RW. Bacteriophage DNA packaging: RNA gears in a DNA transport machine (Minireview). *Cell* 94, 147-150 (1998).
98. Bourassa N, Major F. Implication of the prohead RNA in phage ϕ 29 DNA packaging. *Biochimie* 84, 945-951 (2002).
99. Xiao F, Zhang H, Guo P. Novel mechanism of hexamer ring assembly in protein/RNA interactions revealed by single molecule imaging. *Nucleic Acids Res* 36, 6620-6632 (2008).
100. Moll D, Guo P. Grouping of Ferritin and Gold Nanoparticles Conjugated to pRNA of the Phage ϕ 29 DNA-packaging motor. *J Nanosci and Nanotech (JNN)* 7, 3257-3267 (2007).
101. Ibarra B, Caston J.R., Llorca O., Valle M, Valpuesta J.M., Carrascosa J.L. Topology of the components of the DNA packaging machinery in the phage ϕ 29 prohead. *J.Mol.Biol.* 298, 807-815 (2000).
102. Xiao F, Demeler B, Guo P. Assembly Mechanism of the Sixty-Subunit Nanoparticles via Interaction of RNA with the Reengineered Protein Connector of ϕ 29 DNA-Packaging Motor. *ACS Nano.* 4(6), 3293-3301 (2010).
103. Fang Y, Cai Q, Qin PZ. The procapsid binding domain of ϕ 29 packaging RNA has a modular architecture and requires 2'-hydroxyl groups in packaging RNA interaction. *Biochemistry* 44, 9348-9358 (2005).
104. Chen C, Sheng S, Shao Z, Guo P. A dimer as a building block in assembling RNA: A hexamer that gears bacterial virus ϕ 29 DNA-translocating machinery. *J Biol Chem* 275(23), 17510-17516 (2000).
105. Zhang CL, Garver K, Guo P. Inhibition of phage ϕ 29 assembly by antisense oligonucleotides targeting viral pRNA essential for DNA packaging. *Virology* 211, 568-576 (1995).
106. White SR, Lauring B. AAA+ ATPases: Achieving diversity of function with conserved machinery. *Traffic* 8(12), 1657-1667 (2007).
107. Iyer LM, Leipe DD, Koonin EV, Aravind L. Evolutionary history and higher order classification of AAA plus ATPases. *J Struct Biol* 146(1-2), 11-31 (2004).

108. Liu Y, Huang T, MacMorris M, Blumenthal T. Interplay between AAUAAA and the trans-splice site in processing of a *Caenorhabditis elegans* operon pre-mRNA. *RNA*. 7(2), 176-181 (2001).