University of Kentucky UKnowledge

Plant and Soil Sciences Faculty Patents

Plant and Soil Sciences

1-5-2016

Alteration of Tobacco Alkaloid Content Through Modification of Specific Cytochrome P450 Genes

Ralph E. Dewey

Balazs Siminszky University of Kentucky

Steven W. Bowen

Lily Gavilano University of Kentucky

Follow this and additional works at: https://uknowledge.uky.edu/pss_patents

Part of the Plant Sciences Commons

Right click to open a feedback form in a new tab to let us know how this document benefits you.

Recommended Citation

Dewey, Ralph E.; Siminszky, Balazs; Bowen, Steven W.; and Gavilano, Lily, "Alteration of Tobacco Alkaloid Content Through Modification of Specific Cytochrome P450 Genes" (2016). *Plant and Soil Sciences Faculty Patents*. 41.

https://uknowledge.uky.edu/pss_patents/41

This Patent is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in Plant and Soil Sciences Faculty Patents by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.



US009228194B2

(12) United States Patent

Dewey et al.

(54) ALTERATION OF TOBACCO ALKALOID CONTENT THROUGH MODIFICATION OF SPECIFIC CYTOCHROME P450 GENES

- Inventors: Ralph E. Dewey, Apex, NC (US);
 Balazs Siminszky, Neuchatel (CH);
 Steven W. Bowen, Raleigh, NC (US);
 Lily Gavilano, Raleigh, NC (US)
- (73) Assignees: North Carolina State University, Raleigh, NC (US); University of Kentucky Research Foundation, Lexington, KY (US)
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 642 days.

This patent is subject to a terminal disclaimer.

- (21) Appl. No.: 13/361,159
- (22) Filed: Jan. 30, 2012

(65) **Prior Publication Data**

US 2012/0117933 A1 May 17, 2012

Related U.S. Application Data

- (62) Division of application No. 12/269,531, filed on Nov. 12, 2008, now Pat. No. 8,124,851.
- (60) Provisional application No. 60/987,243, filed on Nov. 12, 2007.
- (51) Int. Cl.

A01H 1/06	(2006.01)
A01H 5/12	(2006.01)
C12N 15/05	(2006.01)
C12N 9/06	(2006.01)
C12N 15/82	(2006.01)
C12N 9/02	(2006.01)

- (52) U.S. CI.
 CPC C12N 15/8243 (2013.01); A01H 1/06 (2013.01); A01H 5/12 (2013.01); C12N 9/0073 (2013.01)
- (58) Field of Classification Search None

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

9/1987	Schilperoort et al.
3/1988	Federoff
8/1988	Comai
1/1989	Hiatt et al.
7/1990	Schilperoort et al.
7/1990	Sanford et al.
4/1991	Umbeck
5/1991	Dooner et al.
7/1991	Jorgensen et al.
4/1992	Saltin
4/1992	Shewmaker et al.
8/1992	Miller et al.
	9/1987 3/1988 8/1988 1/1989 7/1990 7/1990 4/1991 5/1991 4/1992 4/1992 8/1992

(10) Patent No.: US 9,228,194 B2

(45) **Date of Patent:** *Jan. 5, 2016

5,149,645	A	9/1992	Hoekema et al.
5,159,135	A	10/1992	Umbeck
5,177,010	A	1/1993	Goldman et al.
5,231,019	A	7/1993	Paszkowski et al.
5,302,523	A	4/1994	Coffee et al.
5,352,605	A	10/1994	Fraley et al.
5,378,619	A	1/1995	Rogers
5,384,253	A	1/1995	Krzyzek et al.
5,463,174	A	10/1995	Moloney et al.
5,464,763	A	11/1995	Schilperoort et al.
5,464,765	A	11/1995	Coffee et al.
5,469,976	A	11/1995	Burchell
5,472,869	A	12/1995	Krzyzek et al.
5.583.021	A	12/1996	Dougherty et al.
5.595.733	A	1/1997	Carswell et al.
5.614.399	A	3/1997	Ouail et al.
5.641.664	A	6/1997	D'Halluin et al.
5.668.295	A	9/1997	Wahab et al.
5 679 558	A	10/1997	Göbel et al
5 684 241	Δ	11/1007	Nakatani et al
5,004,241	<u>л</u>	1/1008	D'Halluin et al
5 712 376	л Л	2/1008	Bargar
5,715,570	A .	Z/1990	Shillite et el
5,700,900	A	0/1998	Smino et al.

(Continued)

FOREIGN PATENT DOCUMENTS

0	120 516	A3	10/1984
0	267 159	A3	5/1988
	(Conti	nued)

EP

EP

OTHER PUBLICATIONS

GenBank accession No. ABA07806, first available on Oct. 13, 2005.*

(Continued)

Primary Examiner — Lee A Visone

(74) Attorney, Agent, or Firm - Alston & Bird LLP

(57) **ABSTRACT**

Compositions and methods for reducing the level of nornicotine and N'-nitrosonornicotine (NNN) in Nicotiana plants and plant parts thereof are provided. The compositions comprise isolated polynucleotides and polypeptides for cytochrome P450s that are involved in the metabolic conversion of nicotine to nornicotine in these plants. Expression cassettes, vectors, plants, and plant parts thereof comprising inhibitory sequences that target expression or function of the disclosed cytochrome P450 polypeptides are also provided. Methods for the use of these novel sequences to inhibit expression or function of cytochrome P450 polypeptides involved in this metabolic conversion are also provided. The methods find use in the production of tobacco products that have reduced levels of nornicotine and its carcinogenic metabolite, NNN, and thus reduced carcinogenic potential for individuals consuming these tobacco products or exposed to secondary smoke derived from these products.

17 Claims, 4 Drawing Sheets

U.S. PATENT DOCUMENTS

5.929.304	А	7/1999	Radin et al.
6.002.070	Â	12/1999	D'Halluin et al.
6.074.877	Ā	6/2000	D'Halluin et al.
6.907.887	B2	6/2005	Conkling
6.953.040	B2	10/2005	Atchlev et al.
7.032.601	B2	4/2006	Atchlev et al.
7.700.834	B2	4/2010	Xu et al.
7.700.851	B2	4/2010	Xu
7.812.227	B2	10/2010	Xu
7.855.318	B2	12/2010	Xu
7,884,263	B2	2/2011	Dewey et al.
8,058,504	B2	11/2011	Xu
8,124,851	B2 *	2/2012	Dewey et al 800/317.3
8,319,011	B2	11/2012	Xu et al.
2002/0042934	A1	4/2002	Staub et al.
2004/0103449	A1	5/2004	Xu
2004/0111759	A1	6/2004	Xu
2004/0117869	A1	6/2004	Xu
2004/0162420	A1	8/2004	Xu
2005/0132444	A1	6/2005	Xu
2005/0160493	A9	7/2005	Ratcliffe et al.
2005/0178398	A1	8/2005	Breslin et al.
2005/0223442	A1	10/2005	Xu
2005/0244521	Al	11/2005	Strickland et al.
2006/0037096	A1	2/2006	Xu
2006/0037623	A1	2/2006	Lawrence
2006/0041949	A1	2/2006	Xu
2006/0157072	A1	7/2006	Albino et al.
2006/0185686	A1	8/2006	Lawrence
2006/0191548	A1	8/2006	Strickland et al.
2007/0149408	A1	6/2007	Thomas et al.
2007/0199097	A1*	8/2007	Xu C12N 9/0073
			800/278
2007/0292871	A1	12/2007	Xu
2008/0076126	A1	3/2008	Xu
2008/0202541	A1	8/2008	Dewey et al.
2008/0245377	A1	10/2008	Marshall et al.
2009/0119788	A1	5/2009	Mallman et al.
2009/0205072	A1	8/2009	Dewev et al.
2010/0218270	A1	8/2010	Xu et al.
2010/0235938	A1	9/2010	Xu et al.
2010/0235945	AI	9/2010	Xu et al
2010/0235952	AI	9/2010	Xu et al
2011/0048437	Al	3/2011	Xu
2011/0078817	Al	3/2011	Xu
2011/017/322	A 1	7/2011	Dewey et al
2011/01/4322	A1	10/2011	Vamashita et al
2011/0203328	A1	5/2011	Tamaonita et al. Devieu et ol
2012/011/933	A1*	5/2012	Dewey et al. 121/221
2012/0116308	ALC	J/2017	Dewey et al

FOREIGN PATENT DOCUMENTS

EP	0 292 435 B1	11/1988
EP	0 320 500 B1	6/1989
EP	0 159 418 B1	5/1990
EP	0 176 112 B1	5/1990
EP	O116 718 B1	5/1990
EP	0 131 624 B1	9/1992
EP	0 627 752 B1	7/1997
EP	1 033 405 A3	9/2000
EP	0 290 799 B9	11/2003
WO	WO 87/06614 A1	11/1987
WO	WO 92/09696 A1	6/1992
WO	WO 93/21335 A2	10/1993
WO	WO 94/01930 A1	1/1994
WO	WO 00/67558 A1	11/2000
WO	WO 02/072758 A2	9/2002
WO	WO 02/100199 A2	12/2002
WO	WO 03/078577 A2	9/2003
WO	WO 2004/035745 A2	4/2004
WO	WO 2005/038018 A2	4/2005
WO	WO 2005/038033 A2	4/2005
WO	WO 2005/046363 A2	5/2005
WO	WO 2005/111217 A2	11/2005

WO	WO 2005/116199 A2	12/2005
WO	WO 2006/022784 A1	3/2006
WO	WO 2006/091194 A1	8/2006
WO	WO 2006/120570 A2	11/2006
WO	WO 2008/070274 A2	6/2008
WO	WO 2008/076802 A2	6/2008
WO	WO 2009/064771 A2	5/2009
WO	WO 2011/088180 A1	7/2011
WO	WO 2012/118779 A1	9/2012

OTHER PUBLICATIONS

GenBank accession No. ABA07804.1, first available on Oct. 13, 2005.*

Demole et al (Helvetica Chimica Acta—vol. 55, Fasc. 6 (1972) Nbr. 176).*

Xu et al (Physiologia Plantarum 129: 307-319. 2007, first published online: Dec. 20, 2006).*

Chakrabarti et al (New Phytologist (2007) 175: 565-574, first published online: May 30, 2007).*

Uniprot Q38Q86 Cytochrome P450 monooxygenase CYP82E4v1—first available online: Nov. 22, 2005.*

Siminszky et al (PNAS (2005) 102(41) p. 14919-14924).*

Gavilano et al., "Isolation and Characterization of the Cytochrome P450 Gene CYP82E5v2 that Mediates Nicotine to Nornicotine Conversion in the Green Leaves of Tobacco," *Plant Cell Physiol.*, vol. 48(11):1567-1574.

Extended European Search Report for European Application No. 12199188.9; date of mailing Feb. 22, 2013.

U.S. Appl. No. 60/337,684, filed Nov. 12, 2001, Xu.

U.S. Appl. No. 60/347,444, filed Jan. 11, 2002, Xu.

U.S. Appl. No. 60/363,684, filed Mar. 12, 2002, Xu.

U.S. Appl. No. 60/418,933, filed Oct. 16, 2002, Xu.

U.S. Appl. No. 60/485,368, filed Jul. 8, 2003, Xu.

U.S. Appl. No. 60/503,989, filed Sep. 18, 2003, Xu.

U.S. Appl. No. 60/566,235, filed Apr. 29, 2004, Xu.

U.S. Appl. No. 60/607,357, filed Sep. 3, 2004, Xu.

U.S. Appl. No. 60/646,764, filed Jan. 25, 2005, Xu.

U.S. Appl. No. 60/665,097, filed Mar. 24, 2005, Xu.

U.S. Appl. No. 60/665,451, filed Mar. 24, 2005, Xu.

Adams et al., "Genes duplicated by polyploidy show unequal contributions to the transcriptome and organ-specific reciprocal silencing," *PNAS*, 100(8):4649-4654 (2003).

Allen et al., "RNAi-mediated replacement of morphine with the nonnarcotic alkaloid reticuline in opium poppy," *Nature Biotechnology*, 22(12):1559-1566 (2004).

Alonso et al., "A Hox gene mutation that triggers nonsense-mediated RNA decay and affects alternative splicing during Drosophila development," *Nucleic Acids Research*, 31(14):3873-3880 (2003).

Arciga-Reyes et al., "UPF1 is required for nonsense-mediated mRNA decay (NMD) and RNAi in *Arabidopsis*" The Plant Journal, 47:480-489 (2006).

Arndt et al., "Colocalization of antisense RNAs and ribozymes with their target mRNAs," *Genome*, 40:785-797 (1997).

ARS-GRIN: PI 551280, "*Nicotiana tabacum*," http://www.ars-grin.gov/cgi-bin/npgs/acc/display.pl?1446216, accessed Feb. 2009).

Bak et al., Transgenic Tobacco and *Arabidopsis* Plants Expressing the Two Multifunctional Sorghum Cytochrome P450 Enzymes, CYP79A1 and CYP71E1, Are Cyanogenic and Accumulate Metabolites Derived from Intermediates in Dhurrin.

Bartoszewski et al., "Cloning of a Wound Inducible Lycopersicon esculentum Cytochrome P450 Gene and Lack of Regeneration of Transgenic Plants with Sense or Antisense Constructs," J. Am. Soc. Hort. Sci., 127(4):535-539 (2002).

Baseggio et al., "Size and genomic location of the pMGA multigene family of *Mycoplasma gallisepticum,*" *Microbiology*, 142:1429-1435 (1996).

Batard et al., "Increasing Expression of P450 and P450-Reductase Proteins from Monocots in Heterologous Systems," *Arch. Biochem. Biophys.*, 379:161-169 (2000).

Baulcombe, "Fast Forward Genetics Based on Virus-Induced Gene Silencing," *Current Opinion in Plant Biology*, 2:109-113 (1999).

OTHER PUBLICATIONS

Bolitho et al., "Antisense apple ACC-oxidase RNA reduces ethylene production in transgenic tomato fruit," *Plant Science*, 122:91-99 (1997).

Bosher et al., "RNA interference: genetic wand and genetic watchdog," *Nat. Cell Biol.*, 2:E31-E36 (2000).

Bosl et al., "The role of noise and positive feedback in the onset of autosomal dominant diseases," *BMC Systems Biology*, 4:1-15 (2010). Boyette et al., "Results of year 2000 TSNA sampling program in flue-cured tobacco," *Recent Advances in Tobacco Science*, 27:17-22 (2001).

Branch, "A good antisense molecule is hard to find," *TIBS*, 23:45-50 (1998).

Brignetti et al., "Viral pathogenicity determinants are suppressors of transgene silencing in *Nicotiana benthamiana," EMBO J.*, 17(22):6739-6746 (1998).

Burns et al., "Large-scale analysis of gene expression, protein localization, and gene disruption in *Saccharomyces cerevisiae*," *Genes Dev.*, 8:1087-1105 (1994).

Burton et al., Changes in Chemical Composition of Burley Tobacco During Senescence and Curing. 2. Acylated Pyridine Alkaloids, American Chemical Society, pp. 579-583 (1988).

Burton et al., "Distribution of Tobacco Constituents in Tobacco Leaf Tissue. 1. Tobacco-Specific Nitrosamines, Nitrate, Nitrite, and Alkaloids," *J. Agric. Food Chem.*, 40:1050-1055.

Burton et al., "Changes in Chemical Composition of Burley Tobacco during Senescence and Curing. 2. Acylated Pyridine Alkaloids," J .Agric. Food Chem., 38(3):579-584 (1998).

Bush et al., "Formation of tobacco-specific nitrosamines in air-cured tobacco," *Rec. Adv. Tob. Sci*, 27:23-46 (2001).

Byers et al., "Killing the messenger: new insights into nonsensemediated mRNA decay" *The Journal of Clinical Investigation*, 109(1):3-6 (2002).

Byzova et al., "Transforming petals into sepaloid organs in *Arabidopsis* and oilseed rape: implementation of the hairpin RNAmediated gene silencing technology in an organ-specific manner," *Planta*, 218:379-387 (2004).

Callis et al., "Introns increase gene expression in cultured maize cells," *Genes and Dev.*, 1:1183-1200 (1987).

Carron et al., "Genetic modification of condensed tannin biosynthesis in *Lotus corniculatus*. 1. Heterologous antisense dihydroflavonol reductase down-regulates tannin accumulation in "hairy root" cultures," *Theoretical and Applied Genetics*, 87(8).

Caruthers, "Chapter 1: New Methods for Chemically Synthesizing Deoxyoligonucleotides," Methods of DNA and RNA Sequencing, Weissman (ed.), Praeger Publishers, New York, pp. 1-22 (1983).

Chai et al., "Reducing the maize amylopectin content through RNA interference manipulation," *Zhi Wu Sheng Li Yu Fen Zi Sheng Wu Xue Xue Buo*, 31:625-630 (2005) (English Abstract only).

Chakrabarti et al., "CYP82E4-mediated nicotine to nornicotine conversion in tobacco is regulated by a senescence-specific signaling pathway," *Plant Mol. Biol.*, 66: 415-427 (2008).

Chang et al., "Predicting and Testing Physical Locations of Genetically Mapped Loci on Tomato Pachytene Chromosome," *Genetics*, 176:2131-2138 (2007).

Chao et al., "A silent mutation induces exon skipping in the phenylalanine hydroxylase gene in phenylketonuria," *Hum. Genet.*, 108:14-19 (2001).

Chappell, "Biochemistry and Molecular Biology of the Isoprenoid Biosynthetic Pathway in Plants," *Annu. Rev. Plant Physiol. Plant Mol. Biol.*, 46:521-547 (1995).

Chapple, "Molecular-Genetic Analysis of Plant Cytochrome P450-Dependent Monooxygenases," *Annu. Rev. Plant Physiol. Plant Mol. Biol.*, 49:311-343 (1998).

Chelvarajan et al., "Study of Nicotine Demethylation in Nicotiana otophora," J. Agric. Food Chem., 41:858-862 (1993).

Chen et al., "Toxicological analysis of low-nicotine and nocotinefree cigarettes," Toxicology, 249: (2008). Cheung et al., "A Floral Transmitting Tissue-Specific Glycoprotein Attracts Pollen Tubes and Stimulates Their Growth," *Cell*, 82:383-393 (1995).

Cho et al., "Transcriptome Analysis and Physical Mapping of Barley Genes in Wheat-Barley Chromosome Addition Lines," *Genetics*, 172:1277-1285 (2006).

Chou et al., "Chromosome Rearrangements in Arabidopsis thaliana Generated Through Cre-lox Site Specific Recombination," Plant and Animal Genome VII Conference, Abstract No.

Chuang et al., "Specific and heritable genetic interference by doublestranded RNA in *Arabidopsis thaliana*," *PNAS*, 97(9):4985-4990 (2000).

Cogoni et al., "Post-transcriptional gene silencing across kingdoms," *Curr. Opin. Genet. Dev.*, 10:638-643 (2000).

Colbert et al., "High-throughput screening for induced point mutations," *Plant Physiology*, 126:480-484 (2001).

Collier et al., "A Method for Specific Amplification and PCR Sequencing of Individual Members of Multigene Families. Application to the Study of Steroid 21-Hydroxylase Deficiency," *PCR Methods and Applications*, 1:181-186 (1992).

Colliver et al., "Differential modification of flavonoid and isoflavonoid biosynthesis with an antisense chalcone synthase construct in transgenic *Lotus corniculatus," Plant Mol. Biol.*, 35(4):509-522 (1997).

Crookshanks et al., "The potato tuber transcriptome: analysis of 6077 expressed sequence tags," *FEBS Lett.*, 506:123-126 (2001).

Davuluri et al., "Fruit-specific RNAi-mediated suppression of DET1 enhances carotenoid and favonoid content in tomatoes," *Nat. Biotechnol.*, 23:890-895 (2005).

Dekeyser et al., "Transient Gene Expression in Intact and Organized Rice Tissues." *Plant Cell*, 2:591-602 (1990).

Dewey et al., Meeting Abstract dated Sep. 27, 2005, 1 page.

Dewey et al., Power point presentation titled "Functional characterization of the nicotine N-Demethylase gene of tobacco," Philip Morris USA, 21 pages, 2006.

Donato et al., "Fluorescence-Based Assays for Screening Nine Cytochrome P450 (P450) Activities in Intact Cells Expressing Individual Human P450 Enzymes," *Drug Metab. Dispos.*, 2004, 32(7):699-706.

D'Souza et al., "Missense and silent tau gene mutations cause frontotemporal dementia with parkinsonism-chromosome 17 type, by affecting multiple alternative RNA splicing regulatory elements" *PNAS*, 96:5598-5603 (1999).

EBI Accession AV557806, dated Jun. 16, 2000, 2 pages.

Einset, "Differential expression of antisense in regenerated tobacco plants transformed with an antisense version of a tomato ACC oxidase gene," *Plant Cell Tissue and Organ Culture*, 46(2): 137-141 (1996).

Elkind et al., "Abnormal plant development and down-regulation of phenylpropanoid biosynthesis in transgenic tobacco containing a heterologous phenylalanine ammonia-lyase gene," *PNAS*, 87(22):9057-61 (1990).

Escobar et al., "RNAi-mediated oncogene silencing confers resistance to crown gall tumorigenesis," *PNAS*, 98: 13437-13442 (2001). European Search Report completed on Feb. 10, 2010, in European Application No. EP 07 86 5628, 4 pages.

European Search Report completed on Mar. 31, 2011, in European Application No. EP 10 01 5540, 8 pages.

Falcon-Perez et al., "Functional Domain Analysis of the Yeast ABC Transporter Ycflp by Site-directed Mutagenesis," *J. Biol. Chem.*, 274(33):23584-23590 (1999).

Fang et al., "Multiple cis regulatory elements for maximal expression of the cauliflower mosaic virus 35S promoter in transgenic plants," *Plant Cell*, 1:141-150 (1989).

Fannin et al., "Nicotine demethylation in Nicotiana," Med. Sci. Res., 20:807-808 (1992).

Faske et al., "Transgenic Tobacco Plants Expressing Pea Chloroplast Nmdh cDNA in Sense and Antisense Orientation," *Plant Physiol*, 115(2): 705-715 (1997).

Fedoroff et al., "Cloning of the bronze locus in maize by a simple and generalizable procedure using the transposable controlling element Activator (Ac)," *PNAS*, 81:3825-3829 (1984).

OTHER PUBLICATIONS

Fire et al., "Potent and specific genetic interference by doublestranded RNA in *Caenorhabditis elegans*," *Nature*, 391:806-811 (1998).

Force et al., "Preservation of Duplicate Genes by Complementary, Degenerative Mutations," *Genetics*, 151:1531-1545 (1999).

Forsthoefel et al., "T-DNA Insertion Mutagenesis in Arabidopsis: Prospects and Perspectives," Aust. J. Plant Physiol., 19:353-366 (1992).

Frank et al., "Cloning of Wound-Induced Cytochrome P450 Monooxygenases Expressed in Pea," *Plant Physiol.*, 110:1035-1046 (1996).

Freeman et al., "Quantitative RT-PCR: Pitfalls and Potential," *BioTechniques*, 26:112-125 (1999).

Fromm et al., "An octopine synthase enhancer element directs tissuespecific expression and binds ASF-1, a factor from tobacco nuclear extracts," *Plant Cell*, 1:977-984 (1989).

GenBank Accession No. CAA64635, dated Sep. 12, 1996, 2 pages. GenBank Accession No. BAA35080, dated Sep. 26, 2000, 2 pages. GenBank Accession No. AAK62347, dated Jun. 14, 2001, 2 pages. GenBank Accession No. AAK62343, dated Feb. 11, 2002, 2 pages. GenBank Accession No. AAK62346, dated Feb. 11, 2002, 2 pages. GenBank Accession No. AEK08729, dated Feb. 23, 2005, 2 pages. GenBank Accession No. AAK62342, Sep. 20, 2005, 2 pages. GenBank Accession No. ABA07804, dated Oct. 13, 2005, 2 pages. GenBank Accession No. ABA07805, dated Oct. 13, 2005, 2 pages. GenBank Accession No. ABA07807, dated Oct. 13, 2005, 2 pages. GenBank Accession No. DQ131885, dated Oct. 13, 2005, 2 pages. GenBank Accession No. DQ131886, dated Oct. 13, 2005, 2 pages. GenBank Accession No. DQ131888, dated Oct. 13, 2005, 2 pages. GenBank Accession No. DQ219341, dated Oct. 1, 2006, 2 pages. GenBank Accession No. DQ219342, dated Oct. 1, 2006, 2 pages. GenBank Accession No. DQ219343, dated Oct. 1, 2006, 2 pages. GenBank Accession No. DQ219344, dated Oct. 1, 2006, 2 pages. GenBank Accession No. DQ219345, dated Oct. 1, 2006, 2 pages. GenBank Accession No. DQ219346, dated Oct. 1, 2006, 2 pages. GenBank Accession No. DQ219347, dated Oct. 1, 2006, 2 pages. GenBank Accession No. DQ219348, dated Oct. 1, 2006, 2 pages. GenBank Accession No. DQ219349, dated Oct. 1, 2006, 2 pages. GenBank Accession No. DQ219350, dated Oct. 1, 2006, 2 pages. GenBank Accession No. DQ219351, dated Oct. 1, 2006, 2 pages. GenBank Accession No. DQ219352, dated Oct. 1, 2006, 2 pages. GenBank Accession No. DQ350312, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350313, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350314, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350315, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350316, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350317, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350318, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350319, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350320, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350321, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350322, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350323, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350324, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350325, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350326, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350327, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350328, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350329, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350330, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350331, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DO350332, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350333, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350334, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350335, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350336, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350337, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350338, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350339, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350340, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DO350341, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350342, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350343, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350344, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350345, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DO350346, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350347, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350348, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350349, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350350, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350351, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350352, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350353, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350354, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350355, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DO350356, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350357, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350358, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350359, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350360, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350361, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350362, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ350363, dated Dec. 31, 2006, 2 pages. GenBank Accession No. DQ205656, dated Jan. 18, 2007, 2 pages. GenBank Accession No. ABA07806, dated Mar. 7, 2007, 2 pages. GenBank Accession No. DQ131887, dated Mar. 7, 2007, 2 pages. Ghosh, "Polyamines and plant alkaloids," Indian J. Exp. Biol., 38:1086-1091 (2000)

Goldrick et al., "Molecular Genetic Analysis of the User Group Associated with Two Mouse Light Chain Genetic Markers," J. Exp. Med., 162:713-728 (1985).

Graham-Lorence et al., "P450s: Structural similarities and functional differences," *FASEB J.*, 10:206-214 (1996).

Guo et al., "Protein Tolerance to Random Amino Acid Change," *PNAS*, 101(25):9205-9210 (2004).

Hao et al., "Mechanism of Nicotine N-Demethylation in Tobacco Cell Suspension Cultures," *Phytochemistry*, 41(2):477-482 (1995). Hao et al., "Nicotine N-Demethylase in Cell-Free Preparations from

Tobacco Cell Cultures," *Phytochemistry*, 42(2):325-329 (1996). Hao et al., "Evidence in Favour of an Oxidative N-Demethylation of Nicotine to Nornicotine in Tobacco Cell Cultures," *Journal Plant Physiology*, 152:420-426 (1998).

Haseloff et al., "Simple RNA enzymes with new and highly specific endoribonuclease activities," *Nature*, 334:585-591 (1998).

Hayes et al., "Blotting techniques for the study of DNA, RNA, and proteins," *BMJ*, 299(14):965-968 (1989).

Hecht et al., "The relevance of tobacco-specific nitrosamines to human cancer," *Cancer Surveys*, 8(2):279-294 (1989).

Hecht, "Biochemistry, Biology, and Carcinogenicity of Tobacco-Specific N-Nitrosamines," *Chemical Research in Toxicology*, 11(6):559-603 (1998).

Helene et al., "Control of Gene Expression by Triple Helix-Forming Oligonucleotides," *Ann. N.Y. Acad. Sci.*, 660:27-36 (1992).

Helene, "The anti-gene strategy: control of gene expression by triplex-forming-oligonucleotides," *Anti-Cancer Drug Des.*, 6:569-584 (1991).

Helliwell et al., "High-throughput vectors for efficient gene silencing in plants," *Funct. Plant Biol.*, 29:1217-1225 (2002).

Henikoff et al., "Single-Nucleotide Mutations for Plant Functional Genomics," *Annu. Rev. Plant Biol.*, 54:375-401 (2003).

Herbik et al., "Isolation, characterization and cDNA cloning of nicotianamine synthase from barley," *Eur J Biochem*, 265(1): 231-9 (1999).

Hibino et al., "Increase of Cinnamaldehyde Groups in Lignin of Transgenic Tobacco Plants Carrying an Antisense Gene for Cinnamyl Alcohol Dehydrogenase," *Biosci. Biotec. Biochem*, 59:929-931 (1995).

Hildering et al., "Chimeric Structure of the Tomato Plant After Seed Treatment with EMS and X-Rays," The Use of Induced Mutations in Plant Breeding, Pergamon Press, pp. 317-320 (1965).

OTHER PUBLICATIONS

Hill et al., "Functional analysis of conserved histidines in ADPglucose pyrophosphorylase from *Escerichia coli*," *Biochem. Biophys. Res. Commun.*, 244:573-577 (1998) (Abstract only).

Hoekema et al., "A binary plant vector strategy based on separation of the vir- and T-region of the *Agrobacterium tumefaciens* Ti-plasmid," *Nature*, 303:179-180 (1983).

Hoffmann et al., "Tobacco-specific N-nitrosamines and *Areca*-derived N-nitrosamines: chemistry, biochemistry, carcinogenicity, and relevance to humans," *Journal of Toxicology and Environmental Health*, 41:1-52 (1994).

Huang et al., "Insights into Regulation and Function of the Major Stress-Induced hsp70 Molecular Chaperone In Vivo: Analysis of Mice with Targeted Gene Disruption of the hsp70. 1 or hsp70.3 Gene," *Mol Cell Biol*, 21(24):8575-8591 (2001).

Ingelbrecht et al., "Posttranscriptional silencing of reporter transgenes in tobacco correlates with DNA methylation," *PNAS*, 91:10502-10506 (1994).

International Preliminary Report on Patentability in PCT/US07/087386 mailed Jun. 25, 2009, 6 pages.

International Search Report and the Written Opinion of the International Searching Authority, or the Declaration mailed on May 4, 2012, in International Application No. PCT/US2012/026864 (13 pages).

International Search Report and the Written Opinion of the International Searching Authority, or the Declaration mailed on Jul. 4, 2012, in International Application No. PCT/US2012/026795 (15 pages).

International Search Report mailed on Apr. 23, 2014, in International Patent Application No. PCT/US2014/011035.

Invitation to Pay Additional Fees issued on Jun. 23, 2014, in International Patent Application No. PCT/US2014/019381, 8 pages.

Isshiki et al., "Nonsense-mediated decay of mutant waxy mRNA in rice," *Plant Physiology*, 125:1388-1395 (2001).

Jack et al., "Relative stability of nicotine to nornicotine conversion in three burley cultivars," Coresta Congress, Kyoto, Agro-Phyto groups, Abstract AP2 (2004).

Johnston et al., "Dosage-sensitive function of retinoblastoma related and convergent epigenetic control are required during the *Arabidopsis* life cycle," *PLoS Genet*, 6(6):e1000988 (2010).

Jorgensen et al., "Chalcone synthase cosuppression phenotypes in petunia flowers: comparison of sense vs. antisense constructs and single-copy vs. complex T-DNA sequences," *Plant Mol. Biol.*, 31:957-973 (1996).

Julio et al. "Reducing the content of nornicotine in tobacco via targeted mutation breeding," *Mol. Breeding*, 21:369-381 (2008).

Julio et al., "Targeted Mutation Breeding as a tool for tobacco crop improvement," presentation made in Oct. 2008. Kafri et al., "The regulatory utilization of genetic redundancy

Kafri et al., "The regulatory utilization of genetic redundancy through responsive backup circuits," *PNAS*, 103(31):11653-11658 (2006).

Kempin et al., "Targeted disruption in Arabidopsis," Nature, 389:802-803 (1997).

Keskin et al., "A new, structurally nonredundant, diverse data set of protein-protein interfaces and its implications," *Protein Science*, 13:1043-1055 (2004).

Kim et al., "Arabidopsis CYP85A2, a Cytochrome P450, Mediates the Baeyer-Villiger Oxidation of Castasterone to Brassinolide in Brassinosteroid Biosynthesis," *Plant Cell*, 17:2397-2412 (2005).

Klahre et al., "High molecular weight RNAs and small interfering RNAs induce systemic posttranscriptional gene silencing in plants," *PNAS*, 99:11981-11986 (2002).

Klink et al., "The Efficacy of RNAi in the Study of the Plant Cytoskeleton," J. Plant Growth Regul., 19:371-384 (2000).

Koornneef, "Chapter 1: Classical mutagenesis in higher plants," *Molecular Plant Biology*, Gilmartin and Bowler, ed., Oxford University Press, pp. 1-11 (2002).

Koshinsky et al., "Cre-lox site-specific recombination between *Arabidopsis* and tobacco chromosomes," *Plant J.*, 23(6):715-722 (2000).

Kusaba et al., "Low glutelin contentl: A Dominant Mutation That Suppresses the Glutelin Multigene Family via Rna Silencing in Rice," *Plant Cell*, 15:1455-1467 (2003).

Kynast et al., "Dissecting the maize genome by using chromosome addition and radiation hybrid lines," *PNAS*, 101(26):9921-9926 (2004).

Lazar et al., "Transforming Growth Factor a Mutation of Aspartic Acid 47 and Leucine 48 Results in Different Biological Activities," *Mol. Cell. Biol.*, 8(3):1247-1252 (1988).

Levin et al., "Methods of double-stranded RNA-mediated gene inactivation in *Arabidopsis* and their use to define an essential gene in methionine biosynthesis," *Plant Mol. Biol.*, 44:759-775 (2000).

Lewis et al., "Three nicotine demethylase genes mediate nornicotine biosynthesis in *Nicotiana tabacum* L.: Functional characterization of the CYP82E10 gene," *Phytochemistry*, 71:1988-1998 (2010).

Lewis, et al. "RNA interference (RNAi)-induced suppression of nicotine demethylase activity reduces levels of a key carcinogen in cured tobacco leaves." *Plant Biotechnology Journal*, 6:1-9 (2008).

Liu et al., "High-Stearic and High-Oleic Cottonseed Oils Produced by Hairpin RNA-Mediated Post-Transcriptional Gene Silencing," *Plant Physiol.*, 129:1732-1743 (2002).

Liu et al., "Identification and characterization of HTD2: a novel gene negatively regulating tiller bud outgrowth in rice," *Planta*, 230(4):649-658 (2009).

Liu et al "Genetic and transformation studies reveal negative regulation of ERS1 ethylene receptor signaling in *Arabidopsis*," *BMC Plant Biol*, 10:60-73 (2010).

Maher, "DNA Triple-Helix Formation: An Approach to Artificial Gene Repressors?" *BioEssays*, 14(12):807-815 (1992).

Maniatis et al., "Regulation of inducible and tissue-specific gene expression," *Science*, 236:1237-1245 (1987).

Mansoor et al. "Engineering novel traits in plants through RNA interference," *Trends in Plant Science*, 11(11):1-7 (2006).

Maquat, "Nonsense-mediated mRNA decay," *Curr. Biol.*, 12(6):R196-R197 (2002).

Matthew, "RNAi for plant functional genomics," Comparative and Functional Genomics, 5:240-244 (2004).

McDougall et al., "Detection of Viral DNA and RNA by In Situ Hybridization," J. Histochem. Cvtochem., 34:33-38 (1986).

McKinney, "Sequence based identification of T-DNA insertion mutations in *Arabidopsis*: actin mutants act2-1 and act4-1," *Plant J.*, 1995, 8:613-622.

Mesnard et al., "Evidence for the involvement of tetrahydrofolate in the demethylation of nicotine by *Nicotiana plumbaginifolia* cell-suspension cultures," *Planta*, 214:911-919 (2002).

Mette et al., "Transcriptional silencing and promoter methylation triggered by double-stranded RNA," *EMBO J.*, 19(19):5194-5201 (2000).

Mol et al., "Regulation of plant gene expression by antisense RNA," *FEBS Lett.*, 268(2):427-430 (1990).

Napoli et al., "Introduction of a Chimeric Chalcone Synthase Gene into Petunia Results in Reversible Co-Suppression of Homologous Genes in trans," *Plant Cell*, 2:279-289 (1990).

Nawrath et al., "Salicylic Acid Induction-Deficient Mutants of *Arabidopsis* Express PR-2 and PR-5 and Accumulate High Levels of Camalexin after Pathogen Inoculation," *Plant Cell*, 11:1393-1404 (1999).

Needleman et al., "A General Method Applicable to the Search for Similarities in the Amino Acid Sequence of Two Proteins," *J. Mol. Biol.*, 48:443-453 (1970).

Nelson et al., "Comparative Genomics of Rice and Arabidopsis. Analysis of 727 Cytochrome P450 Genes and Pseudogenes from a Monocot and a Dicot," *Plant Physiol.*, 135:756-772.

Nelson et al., "Comparison of cytochrome P450 (CYP) genes from the mouse and human genomes, including nomenclature recommendations for genes, pseudogenes and alternative-splice variants," *Pharmacogenetics*, 14:1-18 (2004).

Ng et al., "Specific Detection and Confirmation of *Campylobacter jejuni* by DNA Hybridization and PCR," *Appl. Environ. Microbiol.*, 63(11):4558-4563 (1997).

Nishihara et al., "Flavanoid components and flower color change in transgenic tobacco plants by suppression of chalcone isomerase gene," *FEBS Lett.*, 579:6074-6078 (2005).

OTHER PUBLICATIONS

Odell et al., "Identification of DNA sequences required for activity of the cauliflower mosaic virus 35S promoter," *Nature*, 313:810-812 (1985).

Office Action mailed on Oct. 18, 2006, in U.S. Appl. No. 10/293,252. Office Action mailed on Oct. 30, 2006, in U.S. Appl. No. 10/686,947. Office Action mailed on Nov. 14, 2006, in U.S. Appl. No. 10/387,346. Office Action mailed on Nov. 14, 2006, in U.S. Appl. No. 10/340,861. Office Action mailed on May 4, 2007, in U.S. Appl. No. 10/943,507. Office Action mailed on Jun. 12, 2007, in U.S. Appl. No. 10/934,944. Ogita et al., "Application of RNAi to confirm theobromine as the major intermediate for caffeine biosynthesis in coffee plants with potential for construction of decaffeinated varieties," *Plant Mol. Biol.*, 54:931-941 (2004).

Ohshima et al., "Nucleotide sequence of the PR-1 gene of *Nicotiana* tabacum," FEBS Letters, 225:243-246 (1987).

Oliver et al., "Inhibition of tobacco NADH-hydroxypyruvate reductase by expression of a heterologous antisense RNA derived from a cucumber cDNA: Implications for the mechanism of action of antisense RNAs," *Mol Gen Genet*, 239(3):425-34 (1993).

Pearson et al., "Improved tools for biological sequence comparison," *PNAS*, 85:2444-2448 (1988).

Peele et al., "Formation of Tobacco-specific Nitrosamines in Fluecured Tobacco," CORESTA Meeting, Agro-Phyto Groups, Suzhou, China (1999).

Pickett et al., "Seeing Double: Appreciating Genetic Redundancy," *Plant Cell*, 7:1347-1356 (1995).

Plant Variety Protection Office (USDA-AMS, Beltsville, MD, http:// www.ars-grin.gov/cgi-bin/npgs/html/pvp.pl?Tobacco, accessed Feb. 2009).

Puchta et al., "Two different but related mechanisms are used in plants for the repair of genomic double-strand breaks by homologous recombination," *PNAS*, 93:5055-5060 (1996).

Qin et al., "Cre recombinase-mediated site-specific recombination between plant chromosomes," *PNAS*, 91:1706-1710 (1994).

Qiu et al. "A computational study of off-target effects of RNA interference." *Nucleic Acids Research*, 33(6)1834-1847 (2005).

Ralston et al., "Cloning, Heterologous Aristolochene-1,3-Dihydroxylase from Expression, and Functional Characterization of 5-epi-Tobacco (*Nicotiana tabacum*)," *Arch. Biochem. Biophys.*, 393(2):222-235 (2001).

Reid et al., "Studies on the Fermentation of Tobacco 1. The Microflora of Cured and Fermenting Cigar-leaf Tobacco," Bulletin 356, Pennsylvania Agricultural Experiment Station, State College, PA, 18 pages (1938).

Rodermel et al., "Nuclear-Organelle Interactions: Nuclear Antisense Gene Inhibits Ribulose Biphosphate Carboxylase Enzyme Levels in Transformed Tobacco Plants," *Cell*, 55:673-681(1988).

Rohr et al., "Tandem inverted repeat system for selection of effective transgenic RNAi strains of *Chlamydomonas*," *Plant J.*, 40:611-621 (2004).

Ruiz et al., "Nicotine-free and salt-tolerant tobacco plants obtained by grafting to salinity-resistant rootstocks of tomato," *Physiologia Plantarum*, 24:(4):465-475 (2005).

Salehuzzaman et al., "Isolation and characterization of a cDNA encoding granule-bound starch synthase in cassava (*Manihot esculenta* Crantz) and its antisense expression in potato," *Plant Mol Biol*, 23(5):947-62 (1993).

Schenk et al., "Coordinated plant defense responses in *Arabidopsis* revealed by microarray analysis," *PNAS*, 97(21):11655-11660 (2000).

Schnable et al., "Genetic recombination in plants," *Curr. Opin. Plant Biol.*, 1:123-129 (1998).

Schopfer et al., "Identification of elicitor-induced cytochrome P450s of soybean (*Glycine max* L.) using differential display of mRNA," *Mol. Gen. Genet.*, 258:315-322, (1998).

Seal et al., "Isolation of a Pseudomonas solanacearum-Specific DNA Probe by Subtraction Hybridization and Construction of Species-Specific Oligonucleotide Primers for Sensitive Detection by the Polymerase Chain Reaction," *Appl. Environ*. Sequence 6912f1 obtained from the Internet at http://mrg.pscsiken.go.ip/nicotiana/menu/069.html on Dec. 6, 2007, 1 page.

Shah et al., "Expression of Silent Mutations in Disease Phenotype," Abstract for presentation at 11^{th} International Congress of Human Genetics, 1 page, (2006).

Shen et al., "Resistance Gene Candidates Identified by PCR with Degenerate Oligonucleotide Primers Map to Clusters of Resistance Genes in Lettuce," *Molecular Plant-Microbe Interactions*, 11(8):815-823 (1998).

Shew et al. (Eds.), "Compendium of Tobacco Diseases," published by American Phytopathology Society, 99 pages (1991).

Sinvany-Villalobo et al., "Expression in Multigene Families Analysis of Chloroplast and Mitochondrial Proteases," *Plant Physiol*, 135:1336-1345 (2004).

Skarnes, "Entrapment Vectors: A New Tool for Mammalian Genetics," *Bio/Technology*, 8:827-831 (1990).

Smith et al., "Comparison of Biosequences," Adv. Appl. Math., 2:482-489 (1981).

Smith et al., "Total silencing by intron-spliced hairpin RNAs," *Nature*, 407:319-320 (2000).

Spradling et al., "Gene disruptions using P transposable elements: An integral component of the *Drosophila* genome project," *PNAS*, 92:10824-10830 (1995).

Stalberg et al., "Deletion analysis of a 2S seed storage protein promoter of *Brassica napus* in transgenic tobacco," *Plant Mol. Biol.*, 23:671-683 (1993).

Sundaresan et al., "Patterns of gene action in plant development revealed by enhancer trap and gene trap transposable elements," *Genes Dev.*, 9:1797-1810 (1995).

Sureka et al., "Positive Feedback and Noise Activate the Stringent Response Regulator Rel in Mycobacteria," *PLoS One*, 3(3):e1771 (2008).

Takeda et al., "Differentiation between Wild and Vaccine-Derived Strains of Poliovirus by Stringent Microplate Hybridization of PCR Products," *J. Clin. Microbiol.*, 32:202-204 (1994).

Takemoto et al., "Molecular Cloning of a Defense-Response-Related Cytochrome P450 Gene from Tobacco," *Plant Cell Physiol.*, 40(12):1232-1242 (1999).

Takken et al. "A functional cloning strategy, based on a binary PVXexpression vector, to isolate HR-inducing cDNAs of plant pathogens." *The Plant Journal*, 24(2): 275-283 (2000).

Tavernarakis et al., "Heritable and inducible genetic interference by double-stranded RNA encoded by transgenes," *Nat. Genet.*, 24:180-183 (2000).

Temple et al., "Modulation of glutamine synthetase gene expression in tobacco by the introduction of an alfalfa glutamine synthetase gene in sense and antisense orientation: molecular and biochemical analysis," *Mol Gen Genet*, 236(2-3):315-25 (1993).

Thomas et al., "Size constraints for targeting post-transcriptional gene silencing and for RNAdirected methylation in *Nicotiana benthamiana* using a potato virus X vector," *Plant J.*, 25(4):417-425 (2001).

Thornton et al., "From structure to function: Approaches and limitations," *Nature Structural Biology, Structural Genomics Supplement*, pp. 991-994 (2000).

Till et al., "Discovery of induced point mutations in maize genes by TILLING," *BMC Plant Biology*, 4:12 (2004).

Toscano et al., "A silent mutation (2939G>A, exon 6; CYP2D6*59) leading to impaired expression and function of CYP2D6," *Pharmacogenet. Genomics*, 16(10):767-770 (2006).

Travella, et al. "RNA Interference-Based Gene Silencing as an Efficient Tool for Functional Genomics in Hexaploid Bread Wheat." *Plant Physiology*, 142:6-20 (2006).

Trevanion et al., "NADP-Malate Dehydrogenase in the C4 Plant *Flaveria bidentis," Plant Physiol*, 113(4):1153-1165 (1997).

Turner et al., "Post-transcriptional gene-silencing and RNA interference: genetic immunity, mechanisms and applications," *J. Chem. Technol. Biotechnol.*, 75:869-882 (2000).

United States, "Tobacco in the United States," Miscellaneous Publication No. 867, U.S. Dept. of Agriculture, Agricultural Marketing Service, 27 pages (1979).

OTHER PUBLICATIONS

Vaistij et al., "Spreading of RNA Targeting and DNA Methylation in Rna Silencing Requires Transcription of the Target Gene and a Putative RNA-Dependent RNA Polymerase," *Plant Cell*, 14:857-867 (2002).

Van der Krol et al., "An anti-sense chalcone synthase gene in transgenic plants inhibits flower pigmentation," *Nature*, 333:866-869 (1988).

Van der Krol et al., "Antisense genes in plants: an overview," *Gene*, 72:45-50 (1988).

Vaucheret et al., "Post-transcriptional gene silencing in plants," J. Cell Sci., 114:3083-3091 (2001).

Veena et al., "Glyoxalase I from *Brassica juncea*: molecular cloning, regulation and its overexpression confer tolerance in transgenic tobacco under stress," *Plant Journal*, 17(4):385-395 (1999).

Verdaguer et al., "Functional organization of the cassava vein mosaic virus (CsVMV) promoter," *Plant Mol. Biol.*, 37(6):1055-1067 (1998).

Verkerk, "Chimerism of the tomato plant after seed irradiation with fast neutrons," *Neth. J. Agric. Sci.*, 19:197-203 (1971).

Voss et al., "The role of enhancers in the regulation of cell-typespecific transcriptional control," *Trends Biochem. Sci.*, 11(7):287-289 (1986).

Wang et al., "Suppression of a P450 hydroxylase gene in plant trichome glands enhances natural product-based aphid resistance," *Nat. Biotechnol.*, 19:371-374 (2001).

Wang et al., "Isolation and characterization of the CYP71D16 trichome-specific promoter from *Nicotania tabacum* L" *J. Exp. Botany*, 53(376):1891-1897 (2002).

Wang et al., "Elucidation of the functions of genes central to diterpene metabolism in tobacco trichomes using posttranscriptional gene silencing," *Planta*, 216:686-691 (2003).

Waterhouse et al., "Virus resistance and gene silencing in plants can be induced by simultaneous expression of sense and antisense RNA," *PNAS*, 95:13959-13964 (1998).

Weigel et al., "A developmental switch sufficient for flower initiation in diverse plants," *Nature*, 377:495-500 (1995).

Weising et al., "Foreign Genes in Plants: Transfer, Structure, Expression, and Applications," *Ann. Rev. Genetics*, 22:421-477 (1988).

Werck-Reichhart et al., "Cytochromes P450: a success story," *Genome Biology*, 1(6):reviews3003.1-3003.9 (2000).

Werck-Reichhart et al., "Cytochromes P450," The *Arabidopsis* Book, American Society of Plant Biologists, 28 pages (2002).

Wernsman et al., "Time and site of nicotine conversion in tobacco," *Tobacco Science*, 167(22):226-228 (1968).

Wernsman et al., "Relative Stability of Alleles at the Nicotine Conversion Locus of Tobacco," *Tobacco Science*, 14:34-36 (1970).

Wernsman et al., "Chapter Seventeen: Tobacco." Cultivar Development. Crop Species., W. H. Fehr (ed.), MacMillan Publishing Go., Inc., New York, N.Y., pp. 669-698 (1987).

Wesley et al., "Construct design for efficient, effective and highthroughput gene silencing in plants," *The Plant Journal*, 27(6): 581-590 (2001).

Wetmur, "DNA Probes: Applications of the Principles of Nucleic Acid Hybridization" *Critical Reviews in Bio. and Mol. Biol.*, 26:227-259, (1991).

Whitbred et al., "Molecular Characterization of CYP73A9 and CYP82A1 P450 Genes Involved in Plant Defense in Pea," *Plant Physiol.*, 124:47-58 (2000).

Written Opinion of the International Searching Authority mailed on Apr. 23, 2014, in International Patent Application No. PCT/US2014/011035.

Wu et al. "Herbivory Rapidly Activates MAPK Signaling in Attacked and Unattacked Leaf Regions but Not between Leaves of *Nicotiana attenuata*." *The Plant Cell*, 19:1096-1122 (2007).

Xiong et al., "Different effects on ACC oxidase gene silencing triggered by RNA interference in transgenic tomato," *Plant Cell*, 23:639-646 (2004).

Xu et al. "Computational Estimation and Experimental Verification of Off-Target Silencing during Posttranscriptional Gene Silencing in Plants," *Plant Physiology*, 142:429-440 (2006).

Xu et al., "Biochemical and molecular characterizations of nicotine demethylase in tobacco," *Physiologia Plantarum*, 129(2):307-319 (2007).

Zwart et al., "Rapid Screening for Freshwater Bacterial Groups by Using Reverse Line Blot Hybridization," *Appl. Environ. Microbiol.*, 69(10):5875-5883 (2003).

Chakrabarti, M., et al., "Inactivation of the Cytochrome P450 Gene CYP82E2 by Degenerative Mutations Was a Key Event in the Evolution of the Alkaloid Profile of Modern Tobacco," *New Phytol.*, May 30, 2007, pp. 565-574, vol. 175, No. 3.

Chintapakorn, Y., et al., "Antisense-Mediated Down-Regulation of Putrescine N-Methyltransferase Activity in Transgenic Nicotiana Tabacum L. Can Lead to Elevated Levels of Anatabine at the Expense of Nicotine," *Plant Molecular Biology*, 2003, pp. 87-105, vol. 53.

Gavilano, L.B., "Isolation, Cloning and Characterization of Novel Tobacco Cytochrome P450 Genes Involved in Secondary Metabolism," Plant Biology Meeting, Jul. 24-28, 2004, Lake Buena Vista, Florida, Abstract No. 992, Retrieved from the Internet: URL: http:// abstracts.aspb.org/pb2004/public/P75/8027.html.

Gavilano, L.B., et al., "Genetic Engineering of *Nicotiana tabacum* for Reduced Nornicotine Content," *J. Agric. Food Chem.*, Jul. 11, 2006, pp. 9071-9078, vol. 54, No. 24.

Gavilano, L.B., and B. Siminszky, "Isolation and Characterization of the Cytochrome P450 Gene CYP82E5v2 that Mediates Nicotine to Nornicotine Conversion in the Green Leaves of Tobacco," *Plant Cell Physiol.*, Oct. 8, 2007, pp. 1567-1574, vol. 48, No. 11.

Gavilano, L.B., et al., "Functional Analysis of Nicotine Demethylase Genes Reveals Insights into the Evolution of Modern Tobacco," J. Biol. Chem, Jan. 5, 2007, pp. 249-256, vol. 282, No. 1.

Siminszky, B., et al., "Conversion of Nicotine to Nornicotine in *Nicotiana tabacum* is Mediated by CYP82E4, a Cytochrome P450 Monooxygenase," *PNAS*, Oct. 11, 2005, pp. 14919-14924, vol. 102, No. 41.

Tang & Galili, "Using RNAi to Improve Plant Nutritional Value: From Mechanism to Application," *Trends in Biotechnology*, 2004, pp. 463-469, vol. 22, No. 9.

EMBL Database Report for Accession No. DQ350312, Dec. 31, 2006 (XP002511577).

EMBL Database Report for Accession No. EU182719, Dec. 2, 2007 (XP002511576).

* cited by examiner

FIG. 1A

	1				50
CYP82E4v2	MLSPIEAIVG	LVTFTFLF	LWTKKSOKPS	KPLPPKIPGG	WPVIGHLFHF
CYP82E4v6	MLSPIEAIVG	LVTFTFLF	LWTKKSÖKPS	KPLPPKIPGG	WPVIGHLFHF
CYP82E4v12	MLSPIEAIVG	LVTFTFLFEF	LWTKKSOKPS	KPLPPKIPGG	WPVIGHLFHF
58-166	MVEPIEAIVG	LVTFTFLFYF	LWTKKSÕKPS	KPLPPKIPGG	WPVIGHLFHF
CYP82E3	MVEPVEAIVG	LVTFTFLFYF	LWTKKSOKPS	KPLPPKIPGG	WPVIGHLF攀F
CYP82E2v1	MVFPIEAFVG	LVTFTFLÄYF	LWTKKSOKLP	KPLPPKIPGG	WPVIGHLFHF
CYP82E2v2	MLSPIEAEVG	LVTFTFLÄYF	LWTKKSOKLP	KPLPPKIPGG	WPVIGHLFHF
CYP82E5v2	MVSPVEAIVG	LVTETELFYF	LWPKKEOIPS	KPLPPKIPGG	WPVIGHLFSF
	1.44.4 .44.4	4,4,4,4 ,4,4,4,4	1.64.4 (4.4.6 (4.6.6		.4.4.6
	51				100
<i>CYP82E4v2</i>	NDDGDDRPLA	RKLGDLADKY	GPVFTFRLGL	PLVLVVSSYE	AVKDCFSTND
CYP82E4v6	NDDGDDRPLA	RKLGDLADKY	GPVFTFRLGL	PLVLVVSSYE	AVKDCFSTND
CYP82E4v12	NDDGDDRPLA	RKLGDLADKY	GPVFTFRLGL	PLVLVVSSYE	AVKDCFSTND
58-166	NDDGNDRPLA	RKLGDLADKY	GPVFTFRLGL	PLVLVVSSYE	AXKDCFSTND
CYP82E3	DDDGDDRPLA	RKLGDLADKY	GPVFTFRLGL	PLVLVVSSYE	AIKDCFSTND
CYP82E2v1	NNDGDDRPLA	RKLGDLADKY	GPVFTFRLGL	PLVLVVSSYE	AIKDCFSTND
CYP82E2v2	NNDGDDRPLA	RKLGDLADKY	GPVFTFRLGL	PLVLVVSSYE	AIKDCFSTND
<u>CYP82E5v2</u>	DDDGDDRPLA	RKLGDLADKY	GPVFTFRLGL	PLVLVVSSYE	AVKDCFSTND
	101				1 5 0
	101				150
<u>CYP82E4v2</u>	AIF'SNRPAF'L	YGUYLGYNNA	MLFLANYGPY	WRKNRKLVIQ	EVLSASRLEK
CYP82E4v6	AIFSNRPAFL	YGDYLGYNNA	MLFLANYGPY	WRKNRKLVIQ	EVLSASRLEK
CYP82E4v12	AIFSNRPAFL	YGQYLGYNNA	MLFLANYGPY	WRKNRKLVIQ	EVLSASRLEK
58-166	AIFSNRPAFL	YGEYLGYNNA	MLFLANYGPY	WRKNRKLVIQ	EVLSASRL
CYP82E3	AIF'SNRPAF'L	YGEYLGYXNA	MLFLANYGSY	WRKNRKLOUIQ	EVLSASRLEK
CYP82E2v1	AIF'SNRPAF'L	YGEYLGYNN	MLFLANYGPY	WRKNRKLVIQ	EVLSASRLEK
CYP82E2v2	AIFSNRPA	YGEYLGYNN	MLFLANYGPY	WRKNRKLVIQ	EVLSASRLEK
<u>CYP82E5v2</u>	AIF'SNRPAF'L	YGEYLGY蠶NA	MLFLIKKIGPI	WRKNRKLVIQ	EVLSASRLEK
	151				200
CVD82F4+2	TAT EKHNDEVDLU	STANI VTDT	DONSSTINIT	DWIEFINFCI	TURNITACKNIV
CVD82F4v6	LUINKLAKIÕ	SIKNIALBI SIKNTALBI	DGNSSTINLT	DWLEFLNFGL	TAKWIYCKNA
CVD82EAv12	LUINKLAUIÓ	ACTENILYTRT	DENSSTINLT	DWLEFLNFGL	IAKWIYCKNA
58-166	L UILAKEVELANTŐ	WO TIMUTIKI		DWLEELNEGI	TVKMTAGNNI
CVD82F3	L KHNBEVELO	TOTINUTINT	DCNSSTINIT	DWLEELNEGT	TAKWIYCKNA
CYP82E2v1	FKØVRF®RTO	TSIKNLYTRI	WGNSSTINLT	DWLEELNEGL	TARWITCHI
CYP82E2v2	FK@VRF#RTO	TSIKNLYTRI	MGNSSTINLT	DWLEELNEGL	TARWIT
OVD0202-2			**CHODITHT		TATISTICICIAL
1 1 20 28. 342	- ※KHNRE袋塗をしい	TSTK&LYTRE	DIGNISSTENUUT	I DATE ELECT. NEEDET.	

FIG. 1B

	201				250
<i>CYP82E4v2</i>	ESGKGDEQVE	RFKKAFKDFM	ILSMEFVLWD	AFPIPLFKWV	DFQGHVKAMK
CYP82E4v6	ESGKGDEÕVE	RFKKAFKDFM	ILSMEFVLWD	AFPIPLFKWV	DFÖGHVKAMK
CYP82E4v12	ESGKGDEOVE	RFKKAFKDFM	ILSMEFVLWD	AFPIPLFKWV	DFOGHVKAMK
58-166	ESGKGDEOVE	RFKKAFKDFM	IISMEFVLWD	AFPIPLFKWV	DFOGHVKAMK
CYP82E3	ESGKGDEOVE	RFKKAFKDFM	TLSMEEVLWD	AFPIPLFKWV	DFOGHVKAMK
CYP82E2v1	ESGKGDEOVE	REKNAEKDEM	WLSMEFVLWD	AFPIPLEKWV	DFOGHEKAMK
CYP82E2v2	ESGKGDEOVE	REKNAFKDEM	WISMEEVLWD	AFPIPLEKWV	DFOGHEKAMK
CYP82E5v2	ESGKGDEOVE	RERKAEKDE	TLSMEEVLWD	AFPIPLEKWV	DFOGHVKAMK
011022072	TOOLCOPP5.T	iti gattii itbi 🥁			DI QUIIVIALII
	251				300
CYP82E4v2	RTEKDIDSVE	ONWLEEHINK	REK MEVNAE	GNEODETDVV	LSKMSNEYLG
CVP82E4v6	RTEKDIDSVE	ONWLEEHINK	REK MEVNAE	GNEODEIDVV	LSKMSNEYLG
CVP82E4v12	RTEKDIDSVE	ONWLEEHINK	REK MEVNAE	GNEODEIDVV	LSKMSNEYLG
58-166	RTEKDIDSVE	ONWLEEHINK	REK MEVNAE	GNEODEIDVV	LSKMSNEYLG
CVD82F3	RTERDIDSVE	ONMI FFHT&K	DEK MEV MR	CNEODEIDVV	I SKMSNEVI C
CVD82E21	DTEKDIDSVE	ONMI FFUTNK	DEK MEVRAE	CNEODEIDVV	I CK SC C C C C C C C C C C C C C C C C C
CVD02E2VI	DTEKDIDSVI	ONWIERUINK	DEV TEVENE	CNEODEIDVV	TOKEONELLD
CIF02E2V2	DTENDIDOVE	ONWIEFIIX	DEV MEVNAG	CNEODEIDVV	LONGORLIDE
CIFOZESVZ	KILKDIDOAL	QNWLEERAAR	RER.MEVINA@	GNEQDEIDVV	LOUNDIELE®
	301				350
CVD82F4+2	FGVCDDTVIK			CMATT TNNOR	
CVD92E4VZ	EGISKDIVIK	ATVESLVLDA	ADIVALHINW	CMALLINNOR CMALLINNOR	ALENAQUEID
CVD02E4V0	EGISKDIVIK	ATVESTUDA	ADIVALIINW	CMALLINNOR CMALLINNOR	ALENAQUEID
E0 166	EGISKDIVIS	AIVESLVLDA	ADIVALIINW	CMALLINNQ	ALWRAQEEID
20-100	EGISKDIVIK	AIVESLVLDA	ADIVALHINW	GMALLINNQN CMALLINNQN	ALKKAQEEID
CIPOZES CVD02E21	EGISKDIVIK	AIVESLVLDA	ADIVALHINS	GMALLINNQN CMÆLLINNQN	ALKKAQEEID
CIPSZEZVI CVD02E22	EGISRDIVIK	AIVESLVLDA	ADIVALHINW	GMB LLINNON	ALMKAQEEID
CIP8ZEZVZ	EGISRDIVIK	AIVESLVLDA	ADIVALHINW	GM&LLINNQN	ALMKAQLEID
<u>CIP82E5V2</u>	EGISRDIVIK	AIVESLVLDA	ADIVALHMMNW	GMALLINNQ	ALKKAQEEID
	261				100
CVD07F47	JJI	PODTRDIUVI		VDDCDTTVDU	THUEDOWNOC
CVD92E4VZ	TRVGRDRWVE	ESDINDLVIL	QAIVKEVLKL	VDDCDLLVDU	ENVEDOVVOG
CIPOZE4VO	TRVGRDRWVE	ESDIKDLVIL	QAIVKEVLKL	VDDCDLLVPH	ENVEDOVVSG
CIPOZE4VIZ	TRV&RDRWVE	ESDIKDLVIL	QAIVKEVLRL	VDDCDLLVPH	ENVEDUVVSG
001-00	1 X VGKDRWVE	ESDIKDLVIL	QAIVKEVLRL	1PPGPLLVPH VDDGDLLVDU	ENEEDUVVSG
CIP8ZE3	IKVGKDRWVE	ESDIKDLVIL	QAIVKEVLRL	YPPGPLLVPH	ENVEDUVVSG
CYP8ZEZVI	TKVGKDRWVE	ESDIKDLVYL	QAIVKKVLRL	YPPGPLLVPH	ENV&DCVVSG
CYP82E2v2	TKVGKDRWVE	ESDIKDLVYL	QAIVK&VLRL	YPPGPLLVPH	ENV&DCVVSG
<u>CYP82E5v2</u>	KVGKRRWVE	ESDIKDLVYL	QAIVKEVLRL	YPPGPLLVPH	ENVEDCVVSG
	401				4 - 0
ava	401 2000	AND MUT ODDD	171 1.70 X		450
CYP82E4v2	IHIPKGIRLF	ANVMKLQRDP	KLWSEPDIFD	PERFIAIDID	FRGQXYXXIP
CYP82E4v6	YHIPKGTRLF	anvmkl <u>æ</u> rdP	KLWEDPDEFD	PERFIATDID	FRGQXYXYIP
CYP82E4v12	YHIPKGTRLF	ANVMKLQRDP	KLWSDPDIFD	PERFIATDID	FRGQXYXYIP
58-166	YXISKGTRLF	ANVMKLQRDP	KLWENPDMFD	PERFVAAGID	FRGQHYEYIP
CYP82E3	YHIPKGTRLF	ANVMKLQRDP	KLWSNPDKF	PERFIARDID	FNGQHYEYIP

FIG. 1B

(cont.)

CYP82E2v1	YHIPKGTRLF	ANVMKLQRDP	KLESNPDKFD	PERFIAGDID	FRGHHYEFIP
CYP82E2v2	YHIPKGTRLF	ANVMKLQRDP	KLLSNPDKFD	PERFIAGDID	FRGHHYEFIP
<u>CYP82E5v2</u>	YHIPKGTRLF	ANVMKLQRDP	KLWSNPDKFD	PERFFADDID	YRGQHYEFIP

FIG. 1C

	451				500
<u>CYP82E4v2</u>	FGSGRRSCPG	MTYALQVEHL	TMAHLIQGFN	YRTPNDEPLD	MKEGAGITIR
CYP82E4v6	FGSGRRSCPG	MTYALQVEHL	TMAHLIQGFN	YRTPNDEPLD	MKEGAGITIR
CYP82E4v12	FGPGRRSCPG	MTYALQVEHL	TMAHLIQGFN	YRTPNDEPLD	MKEGAGITIR
58-166	FGSGRRSCPG	MTYALQVEHL	TMAHLIQGFN	YSTPNDEPLD	MKEGAGITIR
CYP82E3	FGSGRRSCPG	MTYALQVEHL	TMAHLIQGFN	YRTPIDEPLD	MKEGAGITIR
CYP82E2v1	FGSGRRSCPG	MTYALQVEHL	TMAHLIQGFN	YKTPNDEALD	MKEGAGITIR
CYP82E2v2	FGSGRRSCPG	MTYALQVEHL	TMAHLIQGFN	YXTPNDEALD	MKEGAGITIR
<u>CYP82E5v2</u>	FGSGRRSCPG	MTYALQÆEHL	TIAHLIQGFN	YXTPNDEPLD	MKEGAGÄTIR
	501	518			
<u>CYP82E4v2</u>	501 KVNPVELIIA	518 PRLAPELY			
<u>CYP82E4v2</u> CYP82E4v6	501 KVNPVELIIA KVNPVELIIA	518 PRLAPELY PRLAPELY			
<u>CYP82E4v2</u> CYP82E4v6 CYP82E4v12	501 KVNPVELIIA KVNPVELIIA KVNPVELIIA	518 PRLAPELY PRLAPELY PRLAPELY			
<u>CYP82E4v2</u> CYP82E4v6 CYP82E4v12 58-166	501 KVNPVELIIA KVNPVELIIA KVNPVELIIA KVNPVEVIIM	518 PRLAPELY PRLAPELY PRLAPELY PRLAPELY			
<u>CYP82E4v2</u> CYP82E4v6 CYP82E4v12 58-166 CYP82E3	501 KVNPVELIIA KVNPVELIIA KVNPVEVIIA KVNPVEVIIM KVNPVKVIIT	518 PRLAPELY PRLAPELY PRLAPELY PRLAPELY PRLAPELY			
<u>CYP82E4v2</u> CYP82E4v6 CYP82E4v12 58-166 CYP82E3 CYP82E2v1	501 KVNPVELIIA KVNPVELIIA KVNPVEVIIA KVNPVEVIII KVNPVKVIIT KVNPVELIIT	518 PRLAPELY PRLAPELY PRLAPELY PRLAPELY PRLAPELY			
<u>CYP82E4v2</u> CYP82E4v6 CYP82E4v12 58-166 CYP82E3 CYP82E2v1 CYP82E2v2	501 KVNPVELIIA KVNPVELIIA KVNPVEVIIA KVNPVEVIII KVNPVELIIT KVNPVELIIT	518 PRLAPELY PRLAPELY PRLAPELY PRLAPELY PRLAPELY PRLAPELY			

ALTERATION OF TOBACCO ALKALOID CONTENT THROUGH MODIFICATION OF SPECIFIC CYTOCHROME P450 GENES

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a divisional of U.S. application Ser. No. 12/269,531, filed Nov. 12, 2008 (issued as U.S. Pat. No. 8,124,851 on Feb. 28, 2012), which claims the benefit of U.S. Provisional Application No. 60/987,243, filed Nov. 12, 2007, the contents of which are hereby incorporated in their entirety by reference herein.

REFERENCE TO SEQUENCE LISTING SUBMITTED ELECTRONICALLY

An official copy of the Sequence Listing is submitted electronically via EFS-Web as an ASCII formatted Sequence Listing with a file named "414656SEQLIST.txt," created on ²⁰ Jan. 30, 2012, and having a size of 136 kb and is filed concurrently with the specification. The Sequence Listing contained in this ASCII formatted document is part of the specification and is herein incorporated by reference in its entirety.

FIELD OF THE INVENTION

The present invention relates to compositions and methods for reducing the level of nornicotine and its metabolite, N'-nitrosonornicotine (NNN), in a plant that is a member of the ³⁰ genus *Nicotiana*, particularly compositions and methods for inhibiting expression or function of a cytochrome P450 polypeptide involved in the metabolic conversion of nicotine to nornicotine.

BACKGROUND OF THE INVENTION

A predominant alkaloid found in commercial tobacco varieties is nicotine, typically accounting for 90%-95% of the total alkaloid pool. The remaining alkaloid fraction is prima- 40 rily three additional pyridine alkaloids: nornicotine, anabasine and anatabine. Nornicotine is generated directly from nicotine by nicotine N-demethylase. Nornicotine usually represents less than 5% of the total pyridine alkaloid pool. However, tobacco plants that initially produce very low amounts 45 of nornicotine can give rise to progeny that metabolically "convert" a large percentage of leaf nicotine to nornicotine. This process is termed "conversion." In tobacco plants that have genetically converted (i.e., "converters"), the great majority of nornicotine production occurs during senescence 50 and curing of a mature leaf (Wernsman & Matzinger (1968) Tob. Sci. 12:226-228). Burley tobaccos are particularly prone to genetic conversion, with rates as high as 20% per generation observed in some cultivars.

During curing and processing of the tobacco leaf, a portion 55 of the nornicotine is metabolized to NNN, a tobacco-specific nitrosamine (TSNA) alleged to be carcinogenic in laboratory animals (Hecht & Hoffmann (1990) *Cancer Surveys* 8:273-294; and Hoffmann et al. (1994) *J. Toxicol. Environ. Health* 41:1-52; Hecht (1998) *Chem. Res. Toxicol.* 11:559-603). In 60 flue-cured tobaccos, TSNAs predominantly form through a reaction of alkaloids with minute amounts of nitrogen oxides present in combustion gases in a direct-fired heating systems used in traditional curing barns (Peele & Gentry (1999) "Formation of tobacco-specific nitrosamines in flue-cured 65 tobacco," CORESTA Meeting, Agro-Phyto Groups, Suzhou, China). The combustion gases, however, can be eliminated

when curing barns are retrofitted with heat-exchangers, which eliminate the mixing of combustion gases with curing air, thereby reducing TSNAs in tobaccos cured in this manner (Boyette & Hamm (2001) *Rec. Adv. Tob. Sci.* 27:17-22.). In

contrast, in air-cured Burley tobaccos, TSNA formation primarily proceeds through a reaction of tobacco alkaloids with nitrite, a process catalyzed by leaf-borne microbes (Bush et al. (2001) *Rec. Adv. Tob. Sci.* 27:23-46). Thus far, attempts to reduce TSNAs through modification of curing conditions while maintaining acceptable quality standards have not been successful for air-cured tobaccos.

In Burley tobacco plants, a positive correlation exists between the nornicotine content of a leaf and an amount of NNN that accumulates in the cured leaf (Bush et al. (2001)

 Rec. Adv. Tob. Sci, 27:23-46; and Shi et al. (2000) Tob. Chem. Res. Conf 54:Abstract 27). However, keeping nornicotine levels at a minimum is difficult in Burley tobacco plants because of conversion. Plant breeders and seed producers are traditionally responsible for minimizing the number of Bur-20 ley tobacco plants that accumulate high levels of nornicotine. Though the percentage of converters that are ultimately grown in fields are reduced through roguing converters during propagation of seed stocks. Unfortunately, this process is costly, time-consuming and imperfect.

Once a plant converts, the high nornicotine trait is inherited as a single dominant gene (Griffith et al. (1955) *Science* 121:343-344; Burke & Jeffrey (1958) *Tob. Sci.* 2:139-141; and Man et al. (1964) *Crop Sci.* 4:349-353). The nature of this gene, however, is currently unknown. In the simplest of scenarios, the conversion locus may represent a nonfunctional nicotine N-demethylase gene that regains its function in converters, possibly through the mobilization of a mutation-inducting transposable element. Alternatively, the converter locus may encode a protein that initiates a cascade of events that ultimately enables converters to metabolize nicotine to nornicotine, meaning that multiple genes may be involved.

Regardless of whether there are one or many genes associated with conversion, the gene(s) encoding polypeptides having nicotine demethylase activity play a pivotal role in this process. Although the inability to purify active nicotine N-demethylase from crude extracts has impeded the isolation and identification of this enzyme, there is some evidence that a member of the cytochrome P450 superfamily of monooxygenases may be involved (Hao & Yeoman (1996) Phytochem. 41:477-482; Hao & Yeoman (1996) Phytochem. 42:325-329; Chelvarajan et al. (1993) J. Agric, Food Chem. 41:858-862; and Hao & Yeoman (1998) J. Plant Physiol. 152:420-426). Unfortunately, these studies are not conclusive, as classic P450 inhibitors, such as carbon monoxide and tetcylasis, fail to lower enzyme activity at rates comparable to other reported P450-mediated reactions (Chelvarajan et al. (1993) J. Agric. Food Chem. 41:858-862).

Furthermore, cytochrome P450s are ubiquitous, transmembrane proteins that participate in metabolizing a wide range of compounds (reviewed by Schuler (1996) *Crit. Rev. Plant Sci.* 15:235-284; and Schuler & Werck-Reichhart (2003) *Annu. Rev. Plant Biol.* 54:629-667). Examples of biochemical reactions mediated by cytochrome P450s include hydroxylations, demethylations and epoxidations. In plants, cytochrome P450 gene families are very large. For example, total genome sequence examination revealed 272 predicted cytochrome P450 genes in *Arabidopsis* and at least 455 unique cytochrome P450 genes in rice (see, e.g., Nelson et al. (2004) *Plant Physiol.* 135(2):756-772). Even though cytochrome P450s have been implicated in the conversion of nicotine to nornicotine, identification of key participating members of this protein family remains a challenge.

Aside from serving as a precursor for NNN, recent studies suggest that the nornicotine found in tobacco products has undesirable health consequences. For example, Dickerson & Janda demonstrated that nornicotine causes aberrant protein glycosylation within a cell (Dickerson & Janda (2002) *Proc.* 5 *Natl. Acad. Sci. USA* 99:15084-15088). Likewise, concentrations of nornicotine-modified proteins were much higher in plasma of smokers compared to nonsmokers. Furthermore, nornicotine can covalently modify commonly prescribed steroid drugs such as prednisone, which can alter both the effi-10 cacy and toxicity of these drugs.

In view of the difficulties associated with conversion, as well as the undesirable health effects of nornicotine accumulation, improved methods for reducing the nornicotine content in tobacco varieties, particularly Burley tobacco plants, ¹⁵ are therefore desirable. Such methods would not only help ameliorate the potential negative health consequences of the nornicotine per se as described above, but also help to reduce NNN levels.

SUMMARY OF THE INVENTION

Compositions and methods are provided for reducing the nornicotine content in plants that are members of the genus *Nicotiana*. Compositions include isolated cytochrome P450 25 polynucleotides and polypeptides that are involved in conversion of nicotine to nornicotine in plants, particularly *Nicotiana* species. Isolated polynucleotides include those that comprise a nucleic acid sequence as set forth in SEQ ID NO:1, 3 or 4, a nucleic acid sequence encoding a polypeptide as set 30 forth in SEQ ID NO:2, 5-12, 14-24, and fragments and variants thereof. Isolated polypeptides of the invention include those that comprise an amino acid sequence as set forth in SEQ ID NO:2, 5-12 or 14-24, an amino acid sequence encoded by the nucleic acid sequence set forth in SEQ ID 35 NO:1, 3 or 4, and fragments and variants thereof.

In a first aspect, the present invention is summarized as polynucleotides that can suppress expression of a nicotine demethylase involved in the metabolic conversion of nicotine to nornicotine in a plant, including the nicotine demethylases 40 of the present invention. The present invention provides an isolated polynucleotide having a promoter capable of functioning in a plant cell operably linked to a nucleic acid sequence comprising a region of between about 100 nucleic acids and about 350 nucleic acids of SEQ ID NO:1 obtained 45 from a sequence selected from the group of nucleic acids at position 253, 353, 647, 733, 1050, 1397 and combinations thereof. The present invention also provides an isolated polynucleotide comprising a nucleic acid sequence encoding a green-leaf nicotine demethylase, where the amino acid 50 sequence of the encoded nicotine demethylase has a substitution at an amino acid residue in a position selected from the group consisting of residues 235, 449, 174, 410, 224, 72, 143 and 422, where the numbering is according to SEQ ID NO:2. Also provided is an isolated polynucleotide comprising a 55 promoter capable of functioning in a plant cell operably linked to a nucleic acid sequence comprising a region of between about 100 nucleic acids and about 350 nucleic acids of SEQ ID NO:4. Further provided is an isolated polynucleotide encoding an amino acid sequence as set forth in SEQ ID 60 NO:2 with a mutation of a residue that differs from the other P450 polypeptides to a conserved residue. Alternatively provided is an isolated polynucleotide encoding an amino acid sequence as set forth in SEQ ID NO:2 with a mutation at a position selected from the group consisting of residues 85, 65 118, 216, 245 and 466, where the residue at 85 is not an isoleucine, the residue at 118 is not an asparagine, the residue

at 216 is not tyrosine, the residue at 245 is not a tyrosine, and the residue at 466 is not valine.

In a second aspect, the present invention is summarized as an expression cassette comprising a polynucleotide encoding an amino acid sequence of SEQ ID NO:2 operably linked to a promoter that is functional in a plant cell. The present invention provides an expression cassette comprises a polynucleotide comprising a nucleic acid sequence of SEQ ID NO:3, or a fragment of at least 25 contiguous nucleic acids thereof, operably linked to a promoter that is functional in a plant cell. Also provided is an isolated polynucleotide comprising at least 25 nucleotides of a nucleic acid sequence of SEQ ID NO:3. Further provided is an isolated polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:2, 5-12 and 14-24. In one embodiment of the second aspect, the isolated polypeptide comprises an amino acid sequence 99% identical to SEQ ID NO:2, such that it is capable of converting nicotine to nornicotine in green leaves of tobacco.

In a third aspect, the present invention is summarized as a plant of the genus Nicotiana or a plant part thereof comprising an expression cassette, the cassette encoding SEQ ID NO:2, a fragment thereof, or a complement of either. Also provided is a transgenic Nicotiana plant having a lower level of nicotine to nornicotine conversion rate in green leaves compared to a non-transgenic plant, the plant comprising an exogenous nucleic acid construct comprising a promoter capable of functioning in a plant cell operably linked to a polynucleotide having a first nucleic acid sequence comprising a region of between about 100 nucleic acids and about 350 nucleic acids of a green-leaf tobacco nicotine demethylase sequence encoding an amino acid sequence of SEQ ID NO:2 and a second nucleic acid sequence capable of forming a double stranded RNA with the first sequence. Further provided is a transgenic Nicotiana plant having a lower level of nicotine to nornicotine conversion rate in green leaves compared to a non-transgenic plant, the plant comprising an exogenous nucleic acid construct comprising a promoter capable of functioning in a plant cell operably linked to a polynucleotide having a first nucleic acid sequence comprising a region of between about 100 nucleic acids and about 350 nucleic acids of a green-leaf tobacco nicotine demethylase sequence having the nucleic acid sequence of SEQ ID NO:3 and a second nucleic acid sequence capable of forming a double stranded RNA with the first sequence is provided. Also provided is a transgenic Nicotiana plant having a lower level of nicotine to nornicotine conversion rate in green leaves compared to a non-transgenic plant, the plant comprising an exogenous nucleic acid construct comprising a promoter capable of functioning in a plant cell operably linked to a polynucleotide having a first nucleic acid sequence comprising a region of between about 100 nucleic acids and about 350 nucleic acids of a green-leaf tobacco nicotine demethylase sequence having a nucleic acid sequence of SEQ ID NO:4 and a second nucleic acid sequence capable of forming a double stranded RNA with the first sequence.

In a fourth aspect, the present invention is summarized as a seed of a transgenic *Nicotiana* plant having a lower level of nicotine to nornicotine conversion rate in green leaves compared to a non-transgenic plant, the plant comprising a heterologous promoter capable of functioning in a plant cell operably linked to a polynucleotide having a first nucleic acid sequence comprising a region of between about 100 nucleic acids and about 350 nucleic acids of a green-leaf tobacco nicotine demethylase sequence having an amino acid sequence of SEQ ID NO:4 and a second nucleic acid sequence capable of forming a double stranded RNA with the

first sequence. Also provided is a tissue culture of regenerable tobacco cells comprising a plant cell that comprises a first polynucleotide having a fragment of the nucleic acid sequence of SEQ ID NO:1, 3 or 4 and a second polynucle-otide capable of forming a double stranded RNA with the 5 first.

In a fifth aspect, the present invention is summarized as a tobacco product comprising a transgenic Nicotiana plant cell having a lower level of nicotine to nornicotine conversion rate in green leaves compared to a non-transgenic plant, the plant 10 cell comprising a heterologous promoter capable of functioning in a plant cell operably linked to a polynucleotide having a first nucleic acid sequence comprising a region of between about 100 nucleic acids and about 350 nucleic acids of a green-leaf tobacco nicotine demethylase sequence having a 15 nucleic acid sequence of SEQ ID NO:4 and a second polynucleotide capable of forming a double stranded RNA with the first. Also provided is a tobacco cell having a genome altered to inhibit the expression of at least a green-leaf nicotine demethylase, where the cell is homozygous for a muta- 20 tion in the gene encoding the green-leaf nicotine demethylase. Further provided is a tobacco cell comprising a transgene containing green-leaf nicotine demethylase nucleic acid sequence that flanks a selectable marker gene, where the selectable marker gene disrupts the nicotine demethylase 25 gene, thereby producing a tobacco cell where the endogenous green-leaf nicotine demethylase gene has been disrupted. Alternatively provided is a tobacco cell comprising a transgene containing green-leaf nicotine demethylase nucleic acid sequence that flanks a selectable marker gene, where the 30 selectable marker gene disrupts the nicotine demethylase gene, thereby producing a tobacco cell where the endogenous green-leaf nicotine demethylase gene has been disrupted.

In a sixth aspect, the present invention is summarized as a method for reducing nornicotine levels in a plant part derived 35 from a plant of the genus *Nicotiana*, the method comprising a) inhibiting expression of a nicotine demethylase, where the nicotine demethylase has an amino acid sequence set forth in the group consisting of SEQ ID NOs: 2 and 5-12; and b) reducing nornicotine levels in a plant part derived from a plant 40 of the genus Nicotiana. Also provided is a method for reducing nornicotine levels in a tobacco product, the method comprising a) growing a transgenic tobacco plant, where the plant having a plant part that comprises an expression cassette comprising a heterologous promoter and a nicotine demethy- 45 lase, where the nicotine demethylase has an amino acid sequence selected from the group consisting of SEQ ID NOs:2 and 5-12; and b) preparing a tobacco product from the tobacco plant part. Further provided is a method for reducing nornicotine levels in a tobacco product, the method compris- 50 ing a) growing a transgenic tobacco plant, where the plant has a plant part that comprises an antibody that specifically binds a polypeptide with an amino acid sequence selected from the group consisting of SEQ ID NOs: 2 and 5-12; and b) preparing a tobacco product from the tobacco plant part. Alternatively 55 provided is a method for reducing nornicotine levels in a tobacco product, the method comprising a) growing a transgenic tobacco plant, where the plant has a plant part that comprises a fragment of a green-leaf nicotine demethylase operably linked to a heterologous promoter and the fragment 60 has at least 25 contiguous nucleic acids from a polynucleotide encoding a polypeptide selected from the group consisting of SEQ ID NOs:2 and 5-12; and b) preparing a tobacco product from the tobacco plant part. Also provided is a method for reducing nornicotine levels in a plant part derived from a plant 65 of the genus Nicotiana, the method comprising a) modifying the functional CYP82E5v2 allele to change alleles to a non6

CYP82E5v2. where the nonfunctional functional CYP82E5v2 has a substitution at an amino acid residue in a position selected from the group consisting of residues 235, 449, 174, 410, 224, 72, 143 and 422, where the numbering is according to SEQ ID NO:2; and b) reducing the level of nornicotine in a plant part derived from a plant of the genus Nicotiana. The present invention also provides a method for reducing the carcinogenic potential of a tobacco product, the method comprising preparing the tobacco product by a) growing a transgenic tobacco plant, where the plant comprises a plant part that comprises a fragment of a green-leaf nicotine demethylase operably linked to a heterologous promoter and the fragment comprises at least 25 contiguous nucleic acids from a polynucleotide encoding a polypeptide selected from the group consisting of SEQ ID NOs:2 and 5-12; and b) preparing a tobacco product from the tobacco plant part. Further, the present invention provides a method of reducing the conversion of nicotine to nornicotine in a Nicotiana plant comprising: a) transforming a Nicotiana plant with an nucleic acid construct comprising a promoter capable of functioning in a plant cell operably linked to a polynucleotide having a first nucleic acid sequence comprising a region of between 100 nucleic acids and about 350 nucleic acids of SEQ ID NO:1, 3 or 4 and a second nucleic acid sequence capable of forming a double stranded RNA with the first sequence; and b) regenerating a transgenic Nicotiana plant.

In a seventh aspect, the present invention is summarized as a method of screening for a green-leaf nicotine demethylase sequence comprising a) obtaining a nucleic acid sequence that has at least 200 nucleic acids of sequence identity with SEO ID NO:1; and b) identifying a codon sequence encoding for a stop codon at position 422 of an encoded polypeptide, where the numbering is according to SEQ ID NO:2. Also provided is a method of screening for green-leaf nicotine demethylase sequence comprising a) obtaining a nucleic acid sequence that has at least 200 nucleic acids of sequence identity with SEQ ID NO:1; and b) identifying a codon sequence encoding for a codon that is not a proline at position 449 of an encoded polypeptide, where the numbering is according to SEQ ID NO:2. Further provided is a method for identifying a tobacco plant with low levels of nornicotine, the method comprising a) obtaining a DNA sample from a tobacco plant of interest; and b) screening the sample for a mutation in SEQ ID NO:1. The present invention also provides a method for reducing the level of nornicotine in a plant part derived from a plant of the genus Nicotiana, the method comprising a) inhibiting expression of a CYP82E4v2 nicotine demethylase and CYP82E5v2 nicotine demethylase; and b) reducing the level of nornicotine in a plant part derived from a plant of the genus Nicotiana. Also included is tobacco plant material comprising a polypeptide with an amino acid sequence of SEQ ID NO:13 having a mutation at a position selected from the group consisting of residues 458, 364, 329 and combinations thereof. Further included is tobacco plant material comprising a CYP82E4v2 having a mutation at a position selected from the group consisting of residues 458, 364, 329 and combinations thereof, where the numbering corresponds to SEQ ID NO:13. Also included is tobacco plant material comprising a CYP82E4v2 having a mutation at residue 376, where the residue is not valine and the numbering of residues corresponds to SEQ ID NO:13.

BRIEF DESCRIPTION OF DRAWINGS

FIG. 1A-C show an amino acid sequence alignment of CYP82E2 gene family members that have been assayed for nicotine demethylase activity in yeast and/or transgenic

65

plants. Sequences in italics and underlined are positive for nicotine demethylase activity (CYP82E4v2 and CYP82E5v2); sequences titled in black failed to show activity in an assay. Residues that differ among the members are shaded in grey. In FIG. **1**, CYP82E4v2 is set forth in SEQ ID 5 NO:13; CYP82E4v6 is set forth in SEQ ID NO:26; CYP82E4v12 is set forth in SEQ ID NO:27; 58-166 is set forth in SEQ ID NO:28; CYP82E3 is set forth in SEQ ID NO:29; CYP82E2v1 is set forth in SEQ ID NO:30; CYP82E2v2 is set forth in SEQ ID NO:31; and CYP82E5v2 ¹⁰ is set forth in SEQ ID NO:2.

DESCRIPTION OF THE NUCLEIC ACID SEQUENCES

SEQ ID NO: 1 sets forth a nucleic acid sequence of a coding region of CYP82E5v2.

SEQ ID NO:2 sets forth an amino acid sequence of a CYP82E5v2.

SEQ ID NO:3 sets forth a nucleic acid sequence of an ²⁰ forward primer for exon 1 of CYP82E5v2. SEQ ID NO:37 sets forth a nucleic acid sequence of an ²⁰ forward primer for exon 1 of CYP82E5v2.

SEQ ID NO:4 sets forth a nucleic acid sequence of a genomic CYP82E5v2.

SEQ ID NO:5 sets forth an amino acid sequence of a CYP82E5v2 P235S.

SEQ ID NO:6 sets forth an amino acid sequence of a CYP82E5v2 P449L.

SEQ ID NO:7 sets forth an amino acid sequence of a CYP82E5v2 S174L.

SEQ ID NO:8 sets forth an amino acid sequence of a 30 CYP82E5v2 A410V.

SEQ ID NO:9 sets forth an amino acid sequence of a CYP82E5v2 M224I.

SEQ ID NO:10 sets forth an amino acid sequence of a CYP82E5v2 P72L.

SEQ ID NO:11 sets forth an amino acid sequence of a CYP82E5v2 L143F.

SEQ ID NO:12 sets forth an amino acid sequence of a CYP82E5v2 W422Stop.

SEQ ID NO:13 sets forth an amino acid sequence of a 40 primer E5Rt_F. CYP82E4v2. SEO ID NO

SEQ ID NO:14 sets forth an amino acid sequence of a CYP82E4v2 P458S.

SEQ ID NO:15 sets forth an amino acid sequence of a CYP82E4v2 K364N.

SEQ ID NO:16 sets forth an amino acid sequence of a CYP82E4v2 P38L.

SEQ ID NO:17 sets forth an amino acid sequence of a CYP82E4v2 E201K.

SEQ ID NO:18 sets forth an amino acid sequence of a 50 CYP82E4v2 R169Q.

SEQ ID NO:19 sets forth an amino acid sequence of a CYP82E4v2 G459R.

SEQ ID NO:20 sets forth an amino acid sequence of a CYP82E4v2 E296K.

SEQ ID NO:21 sets forth an amino acid sequence of a CYP82E4v2 T427I.

SEQ ID NO:22 sets forth an amino acid sequence of a CYP82E4v2 W329Stop.

SEQ ID NO:23 sets forth an amino acid sequence of a 60 CYP82E4v2 V376M.

SEQ ID NO:24 sets forth an amino acid sequence of a CYP82E4v2 D171N.

SEQ ID NO:25 sets forth an amino acid sequence of a CYP82E4.

SEQ ID NO:26 sets forth an amino acid sequence of a CYP82E4v6.

SEQ ID NO:27 sets forth an amino acid sequence of a CYP82E4v12.

SEQ ID NO:28 sets forth an amino acid sequence of a 58-166.

SEQ ID NO:29 sets forth an amino acid sequence of a CYP82E3.

SEQ ID NO:30 sets forth an amino acid sequence of a CYP82E2v1.

SEQ ID NO:31 sets forth an amino acid sequence of a CYP82E2v2.

SEQ ID NO:32 sets forth a nucleic acid sequence of a forward primer for exon 1 of CYP82E4v2.

SEQ ID NO:33 sets forth a nucleic acid sequence of a reverse primer for exon 1 of CYP82E4v2.

SEQ ID NO:34 sets forth a nucleic acid sequence of a forward primer for exon 2 of CYP82E4v2.

SEQ ID NO:35 sets forth a nucleic acid sequence of a reverse primer for exon 2 of CYP82E4v2.

SEQ ID NO:36 sets forth a nucleic acid sequence of a forward primer for exon 1 of CYP82E5v2.

SEQ ID NO:37 sets forth a nucleic acid sequence of a reverse primer for exon 1 of CYP82E5v2.

SEQ ID NO:38 sets forth a nucleic acid sequence of a forward primer for exon 2 of CYP82E5v2.

SEQ ID NO:39 sets forth a nucleic acid sequence of a reverse primer for exon 2 of CYP82E5v2.

SEQ ID NO:40 sets forth a nucleic acid sequence of a primer E5Gen_F1.

SEQ ID NO:41 sets forth a nucleic acid sequence of a primer E5Gen_R1.

SEQ ID NO:42 sets forth a nucleic acid sequence of a primer E5Gen F2.

SEQ ID NO:43 sets forth a nucleic acid sequence of a primer E5Gen_R2.

SEQ ID NO:44 sets forth a nucleic acid sequence of a primer E4Rt_F.

SEQ ID NO:45 sets forth a nucleic acid sequence of a primer E4Rt_R.

SEQ ID NO:46 sets forth a nucleic acid sequence of a primer E5Rt F.

SEQ ID NO:47 sets forth a nucleic acid sequence of a primer E5Rt_R.

SEQ ID NO:48 sets forth a nucleic acid sequence of a primer G3PDH_F.

SEQ ID NO:49 sets forth a nucleic acid sequence of a primer G3PDH_R.

SEQ ID NO:50 sets forth a nucleic acid sequence of a coding region of CYP82E4v2 and the encoded protein (i.e., SEQ ID NO:13).

DEFINITIONS

The present invention includes compositions and methods for inhibiting expression or function of nicotine demethylase 55 polypeptides that are involved in the metabolic conversion of nicotine to nornicotine in a plant, particularly plants of the *Nicotiana* genus, including tobacco plants of various commercial varieties.

As used herein, "inhibit," "inhibition" and "inhibiting" are defined as any method known in the art or described herein, which decreases the expression or function of a gene product of interest (i.e., the target gene product).

"Inhibiting" can be in the context of a comparison between two plants, for example, a genetically altered plant versus a wild-type plant. The comparison can be between plants, one of which lacks a DNA sequence capable of reducing the agent. Inhibition of expression or function of a target gene

product also can be in the context of a comparison between plant cells, organelles, organs, tissues or plant parts within the same plant or between different plants, and includes comparisons between developmental or temporal stages within the same plant or plant part or between plants or plant parts.

"Inhibiting" can include any relative decrement of function or production of a gene product of interest, up to and including complete elimination of function or production of that gene product. When levels of an agent are compared, such a comparison is preferably carried out between organisms with 10 a similar genetic background. Preferably, a similar genetic background is a background where the organisms being compared share 50% or greater, more preferably 75% or greater, and, even more preferably 90% or greater sequence identity of nuclear genetic material. A similar genetic background is a 15 background where the organisms being compared are plants, and the plants are isogenic except for any genetic material originally introduced using plant transformation techniques or a mutation generated by human intervention. Measurement of the level or amount of an agent may be carried out by any 20 suitable method, non-limiting examples of which include, but are not limited to, comparison of mRNA transcript levels, protein or peptide levels, and/or phenotype, especially the conversion of nicotine to nornicotine. As used herein, mRNA transcripts can include processed and non-processed mRNA 25 originates from a foreign species, or, if from the same species, transcripts, and polypeptides or peptides can include polypeptides or peptides with or without any post-translational modification.

As used herein, "host cell" means a cell that comprises a heterologous nucleic acid sequence of the invention. Though 30 the nucleic acid sequences of the invention, and fragments and variants thereof, can be introduced into any cell of interest, of particular interest are plant cells, more particularly cells of a Nicotiana plant species, for example, the tobacco plant species and varieties described herein below.

As used herein, "variant" means a substantially similar sequence. A variant can have different function or a substantially similar function as a wild-type polypeptide of interest. For a nicotine demethylase, a substantially similar function is at least 99%, 98%, 97%, 95%, 90%, 85%, 80%, 75%, 60%, 40 Nicotine Demethylase Polynucleotides and Polypeptides, 50%, 25% or 15% of wild-type enzyme function of converting nicotine to nornicotine under the same conditions or in a near-isogenic line. A wild-type CYP82E5v2 is SEQ ID NO:2. A wild-type CYP82E4v2 is SEQ ID NO:13. As used herein, a "variant polynucleotide" or "variant polypeptide" means a 45 nucleic acid or amino acid sequence that is not wild-type.

A variant can have one addition, deletion or substitution; two or less additions, deletions or substitutions; three or less additions, deletions or substitutions; four or less additions, deletions or substitutions; or five or less additions, deletions 50 or substitutions. A mutation includes additions, deletions, and substitutions. Such deletions or additions can be at the C-terminus, N-terminus or both the C- and N-termini. Fusion polypeptides or epitope-tagged polypeptides are also included in the present invention. "Silent" nucleotide muta- 55 tions do not change the encoded amino acid at a given position. Amino acid substitutions can be conservative. A conservative substitution is a change in the amino acid where the change is to an amino acid within the same family of amino acids as the original amino acid. The family is defined by the 60 side chain of the individual amino acids. A family of amino acids can have basic, acidic, uncharged polar or nonpolar side chains. See, Alberts et al., (1994) Molecular biology of the cell (3rd ed., pages 56-57, Garland Publishing Inc., New York, N.Y.), incorporated herein by reference as if set forth in 65 its entirety. A deletion, substitution or addition can be to the amino acid of another CYP82E family member in that same

position. See, FIGS. 1A-C. As used herein, a "fragment" means a portion of a polynucleotide or a portion of a polypeptide and hence protein encoded thereby.

As used herein, "plant part" means plant cells, plant protoplasts, plant cell tissue cultures from which a whole plant can be regenerated, plant calli, plant clumps and plant cells that are intact in plants or parts of plants such as embryos, pollen, anthers, ovules, seeds, leaves, flowers, stems, branches, fruit, roots, root tips and the like. Progeny, variants and mutants of regenerated plants are also included within the scope of the present invention, provided that they comprise the introduced polynucleotides of the invention. As used herein, "tobacco plant material" means any portion of a plant part or any combination of plant parts.

As used herein, "operably linked" means a functional linkage between two or more elements. For example, an operable linkage between a polynucleotide of interest and a regulatory sequence (i.e., a promoter) is a functional link that allows for expression of the polynucleotide of interest. Operably linked elements may be contiguous or non-contiguous. When used to refer to the fusing of two protein coding regions, by operably linked is intended that the coding regions are in the same reading frame.

As used herein, "heterologous" means a sequence that is substantially modified from its native form in composition and/or genomic locus by deliberate human intervention. For example, a promoter operably linked to a heterologous polynucleotide is from a species different from the species from which the polynucleotide was derived, or, if from the same/ analogous species, one or both are substantially modified from their original form and/or genomic locus, or the promoter is not the native promoter for the operably linked polynucleotide. Furthermore, as used herein, "chimeric gene" means a coding sequence operably linked to a transcription initiation region that is heterologous to the coding sequence.

DETAILED DESCRIPTION OF THE INVENTION

and Variants and Fragments Thereof

Compositions of the present invention include cytochrome P450 polypeptides. Cytochrome P450 polypeptides can have nicotine demethylase activity. Such nicotine demethylase polynucleotides and polypeptides are involved in the metabolic conversion of nicotine to nornicotine in plants, including commercial varieties of tobacco plants. Also included are variants of such nicotine demethylases. In particular, compositions of the invention include isolated polypeptides comprising amino acid sequences as shown in SEQ ID NOS:2 and 5-24, isolated polynucleotides comprising the nucleic acid sequences as shown in SEQ ID NOS:1, 3 and 4, and the isolated polynucleotides encoding polypeptides comprising amino acid sequences of SEQ ID NOS:2 and 5-24. The polynucleotides of the present invention can find use in inhibiting expression of nicotine demethylase polypeptides or variants thereof that are involved in the metabolic conversion of nicotine to nornicotine in plants, particularly tobacco plants. Some of the polynucleotides of the invention have mutations that inhibit nicotine demethylase activity of the wild-type nicotine demethylase. The inhibition of polypeptides of the present invention is effective in lowering nornicotine levels in tobacco lines where genetic conversion occurs in less than 30%, 50%, 70% and 90% of the population, such as fluecured tobaccos. The inhibition of polypeptides of the present invention is effective in lowering nornicotine levels in tobacco populations where genetic conversion occurs in at

least 90%, 80%, 70%, 60% and 50% of a plant population. A population preferably contains greater than about 25, 50, 100, 500, 1,000, 5,000 or 25,000 plants where, more preferably at least about 10%, 25%, 50%, 75%, 95% or 100% of the plants comprise a polypeptide of the present invention.

The present invention further provides expression cassettes comprising all or a portion of the polynucleotides having a nucleic acid sequence set forth in SEQ ID NO:1, 3 or 4, and the isolated polynucleotides encoding polypeptides having an amino acid sequence of SEQ ID NOS: 2 and 5-24, a comple-10 ment or fragment thereof, or a sequence having substantial sequence identity to SEQ ID NO:1, 3 or 4, or the polynucleotides encoding polypeptides having an amino acid sequence of SEQ ID NOS:2 and 5-24, or a complement or fragment thereof, operably linked to a heterologous promoter that is 15 functional in a plant cell for use in expressing an inhibitory RNA transcript that interferes with expression (i.e., transcription and/or translation) of nicotine demethylase polypeptides. In some embodiments, the expression cassettes comprise the nucleotide sequence as shown in SEQ ID NO:1, 3 or 4, a 20 complement or fragment thereof, or a sequence having substantial sequence identity to SEQ ID NO:1, 3 or 4, or a complement or fragment thereof. Introduction of these expression cassettes into a Nicotiana plant of interest; particularly a tobacco plant of varieties commonly known as flue 25 or bright varieties, Burley varieties, dark varieties and oriental/Turkish varieties, results in the production of tobacco plants having reduced amounts of nornicotine and NNN. Leaf and stem material from these transgenic plants can be used to produce a variety of tobacco products having, reduced levels 30 of nornicotine, and a concomitant reduction NNN.

The nicotine demethylase polynucleotides and encoded polypeptides of the present invention include a novel cytochrome P450 gene, designated the CYP82E5v2 nicotine demethylase gene, that is newly identified as having a role in 35 the metabolic conversion of nicotine to nornicotine in tobacco plants. Suppression of the expression of the encoded polypeptide in transgenic tobacco plants results in a significant reduction in the accumulation of nornicotine in the green leaves of these transgenic plants. While not being bound by theory, the 40 metabolic role of these polypeptides may be a direct one, i.e., directly catalyzing the N-demethylation reaction, or an indirect one, i.e., in the form of production of a product that leads to the up-regulation of the nicotine demethylase activity of the leaf. Regardless of the mechanism, any means by which 45 expression and/or function of the polypeptides of the present invention are targeted for inhibition or site-directed mutagenesis within a Nicotiana plant will be effective in reducing nornicotine levels, and levels of NNN, within leaves and stems of these plants.

The invention encompasses isolated or substantially purified polynucleotide or polypeptide compositions of the present invention. An "isolated" or "purified" polynucleotide or polypeptide, or biologically active portion thereof, is substantially or essentially free from components that normally 55 accompany or interact with the polynucleotide or protein as found in its naturally occurring environment. Thus, an isolated or purified polynucleotide or polypeptide is substantially free of other cellular material or culture medium when produced by recombinant techniques, or substantially free of 60 chemical precursors or other chemicals when chemically synthesized. Optimally, an "isolated" polynucleotide is free of sequences (optimally protein encoding sequences) that naturally flank the polynucleotide (i.e., sequences located at the 5' and 3' ends of the polynucleotide) in the genomic DNA of the 65 organism from which the polynucleotide is derived. For example, in various embodiments, the isolated polynucle-

otide can contain less than about 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb or 0.1 kb of nucleotide sequence that naturally flank the polynucleotide in genomic DNA of the cell from which the polynucleotide is derived. A polypeptide that is substantially free of cellular material includes preparations having less than about 30%, 20%, 10%, 5% or 1% (by dry weight) of contaminating protein. When the polypeptide of the invention or biologically active portion thereof is recombinantly produced, optimally culture medium represents less than about 30% 20%, 10%, 5% or 1% (by dry weight) of chemical precursors or non-protein-of-interest chemicals.

Fragments of the disclosed polynucleotides and polypeptides encoded thereby are also encompassed by the present invention. Fragments of a polynucleotide may encode polypeptide fragments that retain the biological activity of the native polypeptide and hence are involved in the metabolic conversion of nicotine to nornicotine in a plant. Alternatively, fragments of a polynucleotide that are useful as hybridization probes or PCR primers using methods described below generally do not encode fragment polypeptides retaining biological activity. Furthermore, fragments of the disclosed polynucleotides include those that can be assembled within recombinant constructs for use in gene silencing with any method known in the art, including, but not limited to, sense suppression/cosuppression, antisense suppression, doublestranded RNA (dsRNA) interference, hairpin RNA interference and intron-containing hairpin RNA interference, amplicon-mediated interference, ribozymes and small interfering RNA or micro RNA, as described in the art and herein below. Thus, fragments of a polynucleotide may range from at least about 20 nucleotides, about 50 nucleotides, about 70 nucleotides, about 100 nucleotides about 150 nucleotides, about 200 nucleotides, 250 nucleotides, 300 nucleotides and up to the full-length polynucleotide encoding the polypeptides of the invention, depending upon the desired outcome. For example, the fragments of a polynucleotide can be between 100 and about 350 nucleotides, between 100 and about 325 nucleotides, between 100 and about 300 nucleotides, between about 125 and about 300 nucleotides, between about 125 and about 275 nucleotides in length, between about 200 to about 320 contiguous nucleotides, between about 200 and about 420 contiguous nucleotides in length between about 250 and about 450 contiguous nucleotides in length. Alternatively, the fragment can be between about 300 and about 450 contiguous nucleotides in length.

A fragment of a nicotine demethylase polynucleotide of the present invention that encodes a biologically active portion of a cytochrome P450 polypeptide of the present invention will encode at least 15, 25, 30, 50, 75, 100, 125, 150, 175, 200, 250, 300, 350, 400, 450 or 500 contiguous amino acids, or up to the total number of amino acids present in a fulllength nicotine demethylase polypeptide of the invention (e.g., 517 amino acids for SEQ ID NOS:2 and 5), or will encode at least 15, 25, 30, 50, 75, 100, 125, 150 or up to the total number of amino acids present in a partial-length cytochrome P450 polypeptide of the invention (e.g., 422 for SEQ ID NO:12). Preferably, the fragment comprises up to amino acid residue 330 of the encoded polypeptide. A biologically active portion of a nicotine demethylase polypeptide can be prepared by isolating a portion of one of the cytochrome P450 polynucleotides of the present invention, expressing the encoded portion of the cytochrome P450 polypeptide (e.g., by recombinant expression in vitro), and assessing the activity of the encoded portion of the cytochrome P450 polypeptide, i.e., the ability to promote conversion of nicotine to nornicotine, using assays known in the art and those provided herein below.

Polynucleotides that are fragments of a cytochrome P450 nucleotide sequence of the present invention comprise at least 16, 20, 50, 75, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, 800, 900, 950, 1000, 1050, 1100, 1150, 1200, 1250, 1300, 1350, 1400, 1450, 1500, 1550, 1600, 1650 or 1700 contiguous nucleic acids, or up to the number of nucleotides present in a full-length cytochrome P450 polynucleotide as disclosed herein (e.g., 1554 for SEQ ID NO:1; or 2608 for SEQ ID NO:4). Polynucleotides that are fragments of a cytochrome P450 nucleotide sequence of the present invention comprise fragments from about 20 to about 1700 contiguous nucleic acids, from about 50 to about 1600 contiguous nucleic acids, from about 75 to about 1500 contiguous nucleic acids, from about 100 to about 1400 nucleic 15 acids, from about 150 to about 1300 contiguous nucleic acids, from about 150 to about 1200 contiguous nucleic acids, from about 175 to about 1100 contiguous nucleic acids, from about 200 to about 1000 contiguous nucleic acids, from about 225 to about 900 contiguous nucleic acids, from about 500 to 20 about 1600 contiguous nucleic acids, from about 775 to about 1700 contiguous nucleic acids, from about 1000 to about 1700 contiguous nucleic acids, or from about 300 to about 800 contiguous nucleic acids from a cytochrome P450 polynucleotide as disclosed herein. For example, polynucleotide 25 fragment can comprise a polynucleotide sequence containing the nucleic acid sequence from the polynucleotide at about position 700 to about position 1250 of a cytochrome P450 coding sequence, at about position 700 to about position 1250 of a cytochrome P450 genomic sequence, at about position 10 to about position 900 of a cytochrome P450 intron sequence, or at about position 100 to about position 800 of a cytochrome P450 intron sequence.

Variants of the disclosed polynucleotides and polypeptides 35 encoded thereby are also encompassed by the present invention. Naturally occurring variants include those variants that share substantial sequence identity to the disclosed cytochrome P450 polynucleotides and polypeptides disclosed herein. Naturally occurring variants can share substantial 40 tion (also referred to as the reference polynucleotide) can also functional identity to the disclosed cytochrome P450 polynucleotides disclosed herein. The compositions and methods of the invention can be used to target expression or function of any naturally occurring cytochrome P450 that shares substantial sequence identity to the disclosed cytochrome P450 45 polypeptides. Such cytochrome P450 can possess the relevant cytochrome P450 activity, i.e., involvement in the metabolic conversion of nicotine to nornicotine in plants, or not. Such variants may result from, e.g., genetic polymorphism or from human manipulation as occurs with breeding and selection. 50 Biologically active variants of a cytochrome P450 protein of the invention, such as variants of the polypeptide set forth in SEQ ID NO:2 and 5-24, will have at least about 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more 55 sequence identity to the amino acid sequence for the wildtype protein as determined by sequence alignment programs and parameters described elsewhere herein, and can be characterized by a functional involvement in the metabolic conversion of nicotine to nornicotine in plants or lack thereof. A 60 biologically active variant of a polypeptide of the invention may differ by as few as 1-15 amino acid residues, as few as 10, as few as 9, as few as 8, as few as 7, as few as 6, as few as 5, as few as 4, as few as 3, as few as 2, or as few as 1 amino acid residue from the wild-type polypeptide. A biologically inac- 65 tive variant of a protein of the invention may differ from that polypeptide by as few as 1-15 amino acid residues, as few as

10, as few as 9, as few as 8, as few as 7, as few as 6, as few as 5, as few as 4, as few as 3, as few as 2, or as few as 1 amino acid residue.

Variants of the polynucleotides of the present invention include those naturally occurring polynucleotides that encode a nicotine demethylase polypeptide that is involved in the metabolic conversion of nicotine to nornicotine in plants. Such polynucleotide variants can comprise a deletion and/or addition of one or more nucleotides at one or more sites within the native polynucleotide disclosed herein and/or a substitution of one or more nucleotides at one or more sites in the native polynucleotide. Because of the degeneracy of the genetic code, conservative variants for polynucleotides include those sequences that encode the amino acid sequence of one of the cytochrome P450 polypeptides of the invention. Naturally occurring variants such as these can be identified with the use of well-known molecular biology techniques, as, e.g., with polymerase chain reaction (PCR) and hybridization techniques as are known in the art and disclosed herein. Variant polynucleotides also include synthetically derived polynucleotides, such as those generated, e.g., by using sitedirected mutagenesis but which still share substantial sequence identity to the naturally occurring sequences disclosed herein, and thus can be used in the methods of the invention to inhibit the expression or function of a nicotine demethylase that is involved in the metabolic conversion of nicotine to nornicotine, including the nicotine demethylase polypeptides set forth in SEQ ID NOS:2, 5, 7-11, 13, 16-21 and 23-24. Generally, variants of a particular polynucleotide of the invention, e.g., the polynucleotide sequence of SEQ ID NO:3 or the polynucleotide sequence encoding the amino acid sequence set forth in SEQ ID NO:2 and 5-24, will have at least about 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more sequence identity to that particular polynucleotide as determined by sequence alignment programs and parameters described elsewhere herein.

Variants of a particular polynucleotide of the present invenbe evaluated by comparison of the percent sequence identity between the polypeptide encoded by the reference polynucleotide and the polypeptide encoded by a variant polynucleotide. Percent sequence identity between any two polypeptides can be calculated using sequence alignment programs and parameters described elsewhere herein. Where any given pair of polynucleotides of the invention is evaluated by comparison of the percent sequence identity shared by the two polypeptides they encode, the percent sequence identity between the two encoded polypeptides is at least about 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more sequence identity. A variant polypeptide of the present invention can include a polypeptide having a serine at position 458, an asparagine at position 364 of the cytochrome P450 polypeptide, a stop codon at position 329 of the cytochrome P450 or any combination thereof, where the numbering corresponds to SEQ ID NO:13.

Furthermore, the polynucleotides of the invention can be used to isolate corresponding cytochrome P450 sequences from other organisms, particularly other plants, more particularly other members of the Nicotiana genus. PCR, hybridization and other like methods can be used to identify such sequences based on their sequence homology to the sequences set forth herein. Sequences isolated based on their sequence identity to the nucleotide sequences set forth herein or to variants and fragments thereof are encompassed by the

present invention. Such sequences include sequences that are orthologs of the disclosed sequences.

As used herein, "orthologs" means genes derived from a common ancestral gene that are found in different species as a result of speciation. Genes found in different species are 5 considered orthologs when their nucleotide sequences and/or their encoded protein sequences share at least 60%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or greater sequence identity. Functions of orthologs are often highly conserved among species. Thus, 10 isolated polynucleotides that encode for a nicotine demethylase polypeptide that is involved in the nicotine-to-nornicotine metabolic conversion and which hybridize under stringent conditions to the cytochrome P450 sequences disclosed herein, or to variants or fragments thereof, are encompassed 15 by the present invention. Such sequences can be used in the methods of the present invention to inhibit expression of nicotine demethylase polypeptides that are involved in the metabolic conversion of nicotine to nornicotine in plants.

Using PCR, oligonucleotide primers can be designed for 20 use in PCR reactions to amplify corresponding DNA sequences from cDNA or genomic DNA extracted from any plant of interest. Methods for designing PCR primers and PCR cloning are generally known in the art and are disclosed in Sambrook et al. (1989) Molecular cloning: a laboratory 25 manual (2d ed, Cold Spring Harbor Laboratory Press, Plainview, N.Y.). Innis et al., eds. (1990) PCR protocols: a guide to methods and applications (Academic Press, New York); Innis & Gelfand, eds. (1995) PCR strategies (Academic Press, New York); and Innis & Gelfand, eds. (1999) PCR methods 30 manual (Academic Press, New York). Known methods of PCR include, but are not limited to, methods using paired primers, nested primers, single specific primers, degenerate primers, gene-specific primers, vector-specific primers, partially mismatched primers and the like.

Hybridization techniques involve the use of all or part of a known polynucleotide as a probe that selectively hybridizes to other corresponding polynucleotides present in a population of cloned genomic DNA fragments or cDNA fragments (i.e., genomic or cDNA libraries) from a chosen organism.

Hybridization may be carried out under stringent conditions. As used herein, "stringent conditions" or "stringent hybridization conditions" means conditions under which a probe will hybridize to its target sequence to a detectably greater degree than to other sequences (e.g., at least two-fold over background). Stringent conditions are sequence-dependent and will be different in different circumstances. By controlling the stringency of the hybridization and/or washing conditions, target sequences that are 100% complementary to the probe can be identified (homologous probing). Alterna- 50 tively, stringency conditions can be adjusted to allow some mismatching in sequences so that lower degrees of similarity are detected (heterologous probing). Generally, a probe is less than about 1000 nucleic acids in length, optimally less than 500 nucleic acids in length.

Typically, stringent conditions will be those in which the salt concentration is less than about 1.5 M Na ion, typically about 0.01 to 1.0 M Na ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30° C. Stringent conditions may also be achieved with the addition of desta- 60 bilizing agents such as formamide. Exemplary low stringency conditions include hybridization with a buffer solution of 30 to 35% formamide, 1 M NaCl, 1% SDS (sodium dodecyl sulphate) at 37° C., and a wash in 1× to 2×SSC (20×SSC=3.0 M NaCl/0.3 M trisodium citrate) at 50 to 55° C. Exemplary 65 moderate stringency conditions include hybridization in 40 to 45% formamide, 1.0 M NaCl, 1% SDS at 37° C., and a wash

in 0.5× to 1×SSC at 55 to 60° C. Exemplary high stringency conditions include hybridization in 50% formamide, 1 M NaCl, 1% SDS at 37° C. and a wash in 0.1×SSC at 60 to 65° C. Optionally, wash buffers may comprise about 0.1% to about 1% SDS. Duration of hybridization is generally less than about 24 hours, usually about 4 to about 12 hours. The duration of the wash time will be at least a length of time sufficient to reach equilibrium.

Preferably, stringency conditions include hybridization in a solution containing 5×SSC, 0.5% SDS, 5×Denhardt's, 0.45 ug/ul poly A RNA, 0.45 ug/ul calf thymus DNA and 50% formamide at 42° C., and at least one post-hybridization wash in a solution comprising from about 0.01×SSC to about 1×SSC. The duration of hybridization is from about 14 to about 16 hours.

Specificity is typically the function of post-hybridization washes, the critical factors being the ionic strength and temperature of the final wash solution. For DNA-DNA hybrids, the T_m can be approximated from the equation of Meinkoth & Wahl (1984) Anal. Biochem. 138:267-284: T_m=81.5° C.+16.6 $(\log M)+0.41$ (% GC)-0.61 (% form)-500/L; where M is the molarity of monovalent cations, % GC is the percentage of guanosine and cytosine nucleotides in the DNA, % form is the percentage of formamide in the hybridization solution, and L is the length of the hybrid in base pairs. The T_m is the temperature (under defined ionic strength and pH) at which 50% of a complementary target sequence hybridizes to a perfectly matched probe. T_m is reduced by about 1° C. for each 1% of mismatching; thus, T_m, hybridization, and/or wash conditions can be adjusted to hybridize to sequences of the desired identity. For example, if sequences with $\geq 90\%$ identity are sought, the T_m can be decreased 10° C. Generally, stringent conditions are selected to be about 5° C. lower than the thermal melting point (T_m) for the specific sequence and its complement at a defined ionic strength and pH. However; severely stringent conditions can utilize a hybridization and/ or wash at 1, 2, 3 or 4° C. lower than the thermal melting point (T_m) ; moderately stringent conditions can utilize a hybridization and/or wash at 6, 7, 8, 9 or 10° C. lower than the thermal melting point (T_m) ; low stringency conditions can utilize a hybridization and/or wash at 11, 12, 13, 14, 15 or 20° C. lower than the thermal melting point (T_m) . Using the equation, hybridization and wash compositions, and desired T_m , those of ordinary skill will understand that variations in the stringency of hybridization and/or wash solutions are inherently described. If the desired degree of mismatching results in a T_m, of less than 45° C. (aqueous solution) or 32° C. (formamide solution), it is optimal to increase the SSC concentration so that a higher temperature can be used. An extensive guide to the hybridization of nucleic acids is found in Tijssen (1993) Laboratory techniques in biochemistry and molecular biology-hybridization with nucleic acid probes, Part I, Chapter 2 (Elsevier, New York); and Ausubel et al., eds. (1995) Current protocols in molecular biology, Chapter 2 (Greene Publish-55 ing and Wiley-Interscience, New York). See also, Sambrook et al. (1989) Molecular cloning: a laboratory manual (2d ed., Cold Spring Harbor Laboratory Press, Plainview, N.Y.).

Hybridization probes may be genomic DNA fragments, cDNA fragments, RNA fragments or other oligonucleotides, and may be labeled with a detectable group such as ³²P, or any other delectable marker. For example, probes for hybridization can be made by labeling synthetic oligonucleotides based on the cytochrome P450 polynucleotides sequences of the present invention. Methods for preparation of probes for hybridization and for construction of cDNA and genomic libraries are generally known in the art and are disclosed in Sambrook et al., supra.

For example, the cytochrome P430 polynucleotides disclosed herein, or one or more portions thereof, may be used as probes capable of specifically hybridizing to corresponding cytochrome P450 polynucleotides and messenger RNAs. To achieve specific hybridization under a variety of conditions, 5 such probes include sequences that are unique among cytochrome P450 polynucleotide sequences or unique to one of the cytochrome P450 polynucleotide sequences, including upstream regions 5' to the coding sequence and downstream regions 3' to the coding sequence and an intron region, and are 10 optimally at least about 10 contiguous nucleotides in length, more optimally at least about 20 contiguous nucleic acids in length, more optimally at least about 50 contiguous nucleic acids in length, more optimally at least about 75 contiguous nucleic acids in length, and more optimally at least about 100 15 contiguous nucleic acids in length. Such probes may be used to amplify corresponding cytochrome P450 polynucleotides. This technique may be used to isolate additional coding sequences or mutations from a desired plant or as a diagnostic assay to determine the presence of coding sequences in a 20 plant. Hybridization techniques include hybridization screening of plated DNA libraries (either plaques or colonies; see, e.g., Sambrook et al., supra.

As used herein, with respect to the sequence relationships between two or more polynucleotides or polypeptides, the 25 term "reference sequence" means a defined sequence used as a basis for sequence comparison. A reference sequence may be a subset or the entirety of a specified sequence; for example, as a segment of a full-length cDNA or gene sequence, or the complete cDNA or gene sequence. 30

As used herein, "comparison window" means a contiguous and specified segment of a polynucleotide sequence, where the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) compared to the reference sequence (which does not comprise additions or 35 deletions) for optimal alignment of the two polynucleotides. Generally, a comparison window is at least 20 contiguous nucleic acids in length, and optionally can be 30, 40, 50 or 100 contiguous nucleic acids or longer. Those of skill in the art understand that to avoid a high similarity to a reference 40 sequence due to inclusion of gaps in the polynucleotide sequence a gap penalty is typically introduced and is subtracted from the number of matches.

Methods of alignment of sequences for comparison are well known in the art. Thus, the determination of percent 45 sequence identity between any two sequences can be accomplished using a mathematical algorithm. Non-limiting examples of such mathematical algorithms are the algorithm of Myers & Miller (1988) *CABIOS* 4:11-17; the local alignment algorithm of Smith et al. (1981) *Adv. Appl. Math.* 2:482; 50 the global alignment algorithm of Needleman & Wunsch (1970) *J. Mol. Biol.* 48:443-453; the search-for-local alignment method of Pearson & Lipman (1988) *Proc. Natl. Acad. Sci.* 85:2444-2448; the algorithm of Karlin & Altschul (1990) *Proc. Natl. Acad. Sci. USA* 872264, modified as in Karlin & 55 Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90:5873-5877.

The BLAST programs of Altschul et al. (1990) *J. Mol. Biol.* 215:403 are based on the algorithm of Karlin & Altschul (1990) supra. BLAST nucleotide searches can be performed with the BLASTN program, score=100, wordlength=12, to 60 obtain nucleotide sequences homologous to a nucleotide sequence encoding a protein of the invention. BLAST protein searches can be performed with the BLASTX program, score=50. wordlength=3, to obtain amino acid sequences homologous to a protein or polypeptide of the invention. To 65 obtain gapped alignments for comparison purposes, Gapped BLAST (in BLAST 2.0) can be utilized as described in Alts-

chul et al. (1997) *Nucleic Acids Res.* 25:3389. Alternatively, PSI-BLAST (in BLAST 2.0) can be used to perform an iterated search that detects distant relationships between molecules. See Altschul et al. (1997) supra. When utilizing BLAST, Gapped BLAST, PSI-BLAST, the default parameters of the respective programs (e.g., BLASTN for nucleotide sequences, BLASTX for proteins) can be used. Alignment may also be performed manually by inspection.

The sequence identity/similarity values provided herein were calculated using the BLASTX (Altschul et al. (1997) supra), Clustal W (Higgins et al. (1994) *Nucleic Acids Res.* 22:4673-4680), and GAP (University of Wisconsin Genetic Computing Group software package) algorithms using default parameters. The present invention also encompasses the use of any equivalent program thereof for the analysis and comparison of nucleic acid and protein sequences. By "equivalent program" is intended any sequence comparison program that, for any two sequences in question, generates an alignment having identical nucleotide or amino acid residue matches and an identical percent sequence identity when compared to the corresponding alignment generated by BLASTX. Clustal W, or GAP.

For purposes of the foregoing discussion of variant nucleotide and polypeptide sequences encompassed by the present invention, "sequence identity" or "identity" in the context of two polynucleotides or polypeptide sequences makes reference to the residues in the two sequences that are the same when aligned for maximum correspondence over a specified comparison window. When percentage of sequence identity is used in reference to proteins it is recognized that residue positions which are not identical often differ by conservative amino acid substitutions, where amino acid residues are substituted for other amino acid residues with similar chemical properties (e.g., charge or hydrophobicity) and therefore do not change the functional properties of the molecule. When sequences differ in conservative substitutions, the percent sequence identity may be adjusted upwards to correct for the conservative nature of the substitution. Sequences that differ by such conservative substitutions are said to have "sequence similarity" or "similarity." Means for malting this adjustment are well known to those of skill in the art. Typically this involves scoring a conservative substitution as a partial rather than a full mismatch, thereby increasing the percentage sequence identity. Thus, for example, where an identical amino acid is given a score of 1 and a non-conservative substitution is given a score of zero, a conservative substitution is given a score between zero and 1. The scoring of conservative substitutions is calculated, e.g., as implemented in the program PC/GENE (Intelligenetics; Mountain View, Calif.).

As used herein, "percentage of sequence identity" means the value determined by comparing two optimally aligned sequences over a comparison window, where the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison, and multiplying the result by 100 to yield the percentage of sequence identity.

Thus, cytochrome P450 polynucleotide and polypeptide sequences can be identified using the sequences provided herein. Such methods include obtaining a polynucleotide or polypeptide sequence at least 80%, 85%, 90%, 95%, 98% or 99% sequence identity with the polynucleotide sequence of SEQ ID NO:1, 3 or 4 or a complement or fragment thereof, or a polypeptide sequence of SEQ ID NO:2 or 5-24. A preferred embodiment includes a polypeptide corresponding to SEQ ID NO:13 that has a serine at position 458 or an asparagine at position 364 of the cytochrome P450 polypeptide, or a stop codon at position 329 of the cytochrome P450, or a combination thereof, where the numbering corresponds to SEQ ID NO:13. 10

Expression Cassettes for Use in the Methods of Invention Compositions of the present invention further include expression cassettes comprising inhibitory sequences capable of inhibiting expression or function of a nicotine demethylase polypeptide involved in the conversion of nico-15 tine to nornicotine in a Nicotiana plant or plant part thereof, where the inhibitory sequences are operably linked to a promoter that is functional in a plant cell. In this manner, expression cassettes comprising all or part of the sequence set forth in SEQ ID NO:1, 3 or 4 or encoding SEQ ID NO:2 or 5-24, a 20 complement or fragment thereof, or sequences sharing substantial sequence identity to such sequences, or a complement or fragment thereof, operably linked to a promoter that is functional in a plant cell are constructed for use in the genesilencing methods of the present invention described herein 25 below. Such sequences are referred to herein as "inhibitory sequences" or "inhibitory polynucleotide sequences," as they are capable of being expressed as an RNA molecule that inhibits expression (i.e., transcription and/or translation) of the target cytochrome P450 polypeptide, for example, the 30 polypeptide set forth in SEQ ID NO:2 or 5-24 and variants thereof, where the variant polypeptides have substantial sequence identity to these disclosed cytochrome P450 polypeptides. Such variants may or may not be involved in the metabolic conversion of nicotine to nornicotine in a plant. 35 Such sequences also include fragment sequences of the target cytochrome P450 polynucleotide or polypeptide. For example, a fragment sequence can include any portion of the cytochrome P450 sequence, including coding and non-coding sequence (e.g., 5' UTR, intron, and 3' UTR sequences), 40 and can include fragments between 100 and about 350 nucleic acids, between about 125 and about 300 nucleic acids, or between about 125 and about 275 nucleic acids. Preferably, a fragment of nicotine demethylase can be between about 20 and about 420, about 30 and about 420, between about 40 and 45 about 320, between about 50 and about 200, between about 50 and about 400, between about 50 and about 420, between about 60 and about 320, about 70 and about 220, between about 100 and about 200, between about 100 and about 320, between about 150 and about 200, between about 150 and 50 about 220, between about 150 and about 400, between about 200 and about 300, or between about 300 and about 400 contiguous nucleic acids. Alternatively, a fragment of a cytochrome P450 can be about 100, about 150, about 200, about 220, about 250, about 300, about 320, or about 350 contigu- 55 Physiol. 38(7):792-803; Hansen et al. (1997) Mol. Gen. ous nucleic acids in length. Alternatively yet, a cytochrome P450 fragment can be reduced in length by about 20, about 40, about 60, about 80, about 100, about 120, about 140, about 160, about 180, about 200, about 220, about 240, about 260, about 280, about 290, about 300, about 320, about 340, about 60 360, about 380, about 400 contiguous nucleic acids compared to the full-length. For all of these cytochrome P450 fragments, the truncation or deletion can start at the 5' end, start at the 3' end, or be internal to a cytochrome P450 or a cytochrome P450 intron. For a cytochrome P450 intron fragment, 65 the entire sequence of a cytochrome P450 intron can be SEQ ID NO:3.

Furthermore, a fragment of a cytochrome P450 polynucleotide or polypeptide can contain contiguous nucleotides from about 1, 2, 5, 10, 20, 30, 40, 50, 60, 70, 80, or 90% of the entire gene. Alternatively stated, a fragment of a cytochrome P450 polynucleotide or polypeptide can be between about 5%-about 80%, between about 10%-about 70%, between about 10%-about 60%, between about 10%-about 50%, between about 25%-about 60%, between about 25%-about 50%, between about 40%-about 60%, between about 40%about 80%, between about 50%-about 90% of the length of an entire cytochrome P450.

Expression cassettes of the present invention include those that encompass additional domains that modulate the level of expression, the developmental timing of expression, or tissue type that expression occurs in (e.g., AU Patent No. AU-A-77751/94 and U.S. Pat. Nos. 5,466,785 and 5,635,618). Promoters can be selected based on the desired outcome. The nucleic acids of the present invention can be combined with inducible, constitutive, pathogen- or wound-induced, environmentally- or developmentally-regulated, cell- or tissuepreferred promoter, or other promoters for expression in plants.

Chemical-inducible promoters can be used to inhibit the expression of a cytochrome P450 that is involved in the metabolic conversion of nicotine to nornicotine in a plant through the application of an exogenous chemical regulator. Chemical-inducible promoters are known in the art and include, but are not limited to, the tobacco PR-la promoter, which is activated by salicylic acid. Other chemical-inducible promoters of interest include steroid-responsive promoters (see, e.g., the glucocorticoid-inducible promoter (Schena et al. (1991) Proc. Natl. Acad. Sci. USA 88:10421-10425; and McNellis et al. (1998) Plant J. 14(2):247-257) and tetracycline-inducible promoters (see, e.g., Gatz et al. (1991) Mol. Gen. Genet. 227:229-237; and U.S. Pat. Nos. 5,814,618 and 5,789,156), each of which is incorporated herein by reference as if set forth in its entirety.

Constitutive promoters include, e.g., the core promoter of the Rsyn7 promoter and outer constitutive promoters disclosed in U.S. Pat. No. 6,072,050; the core CaMV 35S promoter (Odell et al. (1985) Nature 313:810-812); ubiquitin (Christensen et al. (1989) Plant Mol. Biol. 12:619-632; and Christensen et al. (1992) Plant Mol. Biol. 18:675-689); pEMU (Last et al. (1991) Theor. Appl. Genet. 81:581-588); MAS (Velten et al. (1984) EMBO J. 3:2723-2730); ALS promoter (U.S. Pat. No. 5,659,026), and the like. Other constitutive promoters include, e.g., those disclosed in U.S. Pat. Nos. 5,608,149; 5,608,144; 5,604,121; 5,569,597; 5,466,785; 5,399,680; 5,268,463; 5,608,142 and 6,177,611.

Tissue-preferred promoters can be utilized to target expression of an inhibitory polynucleotide sequence of the present invention within a particular plant tissue. Tissue-preferred promoters include those disclosed in Yamamoto et al. (1997) Plant J. 12(2):255-265; Kawamata et al. (1997) Plant Cell Genet. 254(3):337-343; Russell et al. (1997) Transgenic Res. 6(2):157-168; Rinehart et al. (1996) Plant Physiol. 112(3): 1331-1341; Van Camp et al. (1996) Plant Physiol. 112(2): 525-535; Canevascini et al. (1996) Plant Physiol. 112(2): 513-524; Yamamoto et al. (1994) Plant Cell Physiol. 35(5): 773-778; Lam (1994) Results Probl. Cell Differ. 20:181-196; Orozco et al. (1993) Plant Mol. Biol. 23(6):1129-1138; Matsuoka et al. (1993) Proc Natl. Acad. Sci. USA 90(20):9586-9590; and Guevara-Garcia et al. (1993) Plant J. 4(3):495-505.

Of particular interest herein are leaf preferred promoters that provide for expression predominately in leaf tissues. See, e.g., Yamamoto et al. (1997) Plant J. 12(2):255-265; Kwon et

al. (1994) Plant Physiol. 105:357-67; Yamamoto et al. (1994) Plant Cell Physiol. 35(5):773-778; Gotor et al. (1993) Plant J. 3:509-18; Orozco et al. (1993) Plant Mol. Biol. 23(6):1129-1138; Baszczynski et al. (1988) Nucl. Acid Res. 16:4732; Mitra et al. (1994) Plant Molecular Biology 26:35-93; 5 Kayaya et al. (1995) Molecular and General Genetics 248: 668-674; and Matsuoka et al. (1993) Proc. Natl. Acad. Sci. USA 90(20):9586-9590. Senecence-regulated et al. (1998) Plant Physiol. 116:329-335); SAG 13 (Gan and Amasino (1997) Plant Physiol. 113:313-319; SAG 15 (Gan (1995) 10 "Molecular Characterization and Genetic Manipulation of Plant Senescence," Ph.D. Thesis, University of Wisconsin, Madison); SEN1 (Oh et al. (1996) Plant Mol. Biol. 30:739-754; promoter of a senescence-specific gene for expression of IPT (Gan and Amasino 91995) Science 270:1986-1988); and 15 the like (see, e.g., Or et al. (1999) Plant Cell 11:1073-1080; and McCabe et al. (2001) Plant Physiol. 127:505-516).

Expression cassettes of the present invention can include 5' leader sequences that can act to enhance translation. Translation leaders are known in the art and include, but are not 20 P450 polypeptides is reduced or eliminated by transforming a limited to, picornavirus leaders, e.g., EMCV leader (Encephalomyocarditis 5' noncoding region; Elroy-Stein et al. (1989) Proc. Natl. Acid. Sci. USA 86:6126-6130); potyvirus leaders, e.g., TEV leader (Tobacco Etch Virus; Gallie et al. (1995) Gene 165(2):233-238), MDMV leader (Maize Dwarf 25 Mosaic Virus; Virology 154:9-20), and human immunoglobulin heavy-chain binding protein (BiP; Macejak et al. (1991) Nature 353:90-94); untranslated leader from the coat protein mRNA of alfalfa mosaic virus (AMV RNA 4; Jobling et al. (1987) Nature 325:622-625); tobacco mosaic virus leader 30 (TMV; Gallie et al. (1989) in Molecular biology of RNA, ed. Cech (Liss, New York), pp. 237-256); and maize chlorotic mottle virus leader (MCMV; Lommel et al. (1991) Virology 81:382-385). See also, Della-Cioppa et al. (1987) Plant Physiol. 84:965-968. Other methods known to enhance trans- 35 lation also can be utilized.

Methods for Inhibiting Expression or Function of a Nicotine Demethylase

Methods of reducing the concentration, content and/or activity of a cytochrome P450 polypeptide of the present 40 329, 376 or 171 can be crossed with a plant having one or invention in a Nicotiana plant or plant part, particularly the leaf tissue, are provided. Many methods may be used, alone or in combination, to reduce or eliminate the activity of a cytochrome P450 polypeptide of the present invention, more particularly, the CYP82E5v2 nicotine demethylase. In addition, 45 combinations of methods may be employed to reduce or eliminate the activity of two or more different cytochrome P450 polypeptides, more particularly the CYP82E5v2 and CYP82E4v2 nicotine demethylases. Preferably, the CYP82E5v2 is a polypeptide with at least one amino acid 50 mutation in SEQ ID NO:2 that negatively affects conversion in green leaves and the CYP82E4v2 has the sequence set forth in SEQ ID NO:13 with at least one amino acid mutation that negatively affects conversion in senescent leaves.

In accordance with the present invention, the expression of 55 a cytochrome P450 polypeptide of the present invention is inhibited if the protein level of the cytochrome P450 polypeptide is statistically lower than the protein level of the same cytochrome P450 polypeptide in a plant that has not been genetically modified or mutagenized to inhibit the expression 60 of that cytochrome P450 polypeptide, and where these plants have been cultured and harvested using the same protocols. In particular embodiments of the invention, the protein level of the cytochrome P450 polypeptide in a modified plant according to the invention is less than 95%, less than 90%, less than 40%, less than 30%, less than 20%, less than 10%, or less than

5% of the protein level of the same cytochrome P450 polypeptide in a plant that is not a mutant or that has not been genetically modified to inhibit the expression of that cytochrome P450 polypeptide and which has been cultured and harvested using the same protocols. The expression level of the cytochrome P450 polypeptide may be measured directly, for example, by assaying for the level of the cytochrome P450 transcript or cytochrome P450 polypeptide expressed in the Nicotiana plant or plant part, or indirectly, e.g., by measuring the conversion of nicotine to nornicotine in the Nicotiana plant or plant part. Methods for monitoring expression level of a polypeptide are known in the art, and include, but are not limited to, Northern blot analysis and RNA differentiation assays. Methods for determining the activity of a targeted cytochrome P450 polypeptide in converting nicotine to nornicotine are known in the art and described elsewhere herein below, and include, but are not limited to, alkaloid analysis using gas chromatography.

In some instances, the activity of one or more cytochrome plant or plant part with an expression cassette comprising a polynucleotide encoding a polypeptide that inhibits the activity of one or more cytochrome P450 polypeptides of the present invention. A number of approaches have been used to combine transgenes or mutations in one plant-including sexual crossing, retransformation, co-transformation and the use of linked transgenes. A chimeric transgene with linked partial gene sequences can be used to coordinately suppress numerous plant endogenous genes. Constructs modeled on viral polyproteins can be used to simultaneously introduce multiple coding genes into plant cells. For a review, see Halpin et al., Plant Mol. Biol. 47:295-310 (2001). A plant having a mutation in CYP82E4v2 that inhibits the nicotine demethylase activity in senescent leaves can be crossed with a plant having a mutation in CYP83E5v2 that inhibits nicotine demethylase in green leaves to produce a plant with conversion levels lower than about 0.2%, 0.3%, 0.4%, 0.5%, 0.6% or 0.7%. Alternatively, a plant having one or more mutations in CYP82E4v2 at position 458, 364, 38, 201, 169, 459, 296, 427, more mutations in CYP83E5v2 at position 235, 449, 174, 410, 224, 72 or 143 to produce a plant with conversion levels lower than 0.2%, 0.3%, 0.4%, 0.5%, 0.6% or 0.7%. Alternatively still, a plant having one or more mutations in CYP82E4v2 at position 458, 364 or 329 can be crossed with a plant having one or more mutations in CYP83E5v2 at position 235, 449, 174, 410, 224, 72, or 143 to produce a plant with conversion levels lower than 0.2%, 0.3%, 0.4%, 0.5%, 0.6%, or 0.7%. A particularly preferred conversion level of nicotine to nornicotine can be between 0.05%-0.4%, between 0.1-0.6%, between 0.1%-0.3%, between 0.1%-0.5%, between 0.1%-0.4%, between 0.1%-0.7%, or between 0.1%-1.0%. Any mutation of a polynucleotide of the present invention that results in a truncation of the CYP83E4v2 or CYP83E5v2 polypeptide before a conserved heme-binding motif will inhibit the enzyme and can be used in a cross described above. The domains of cytochrome P450 proteins are known in the art. See, e.g., Xu et al. (2007) Physiologia Plantarum 129:307-319, incorporated herein by reference as if set forth in its entirety. By crossing plants having a nonfunctional or inhibited CYP82E4v2 gene, a nonfunctional or inhibited CYP82E5v2 gene, or nonfunctional or inhibited CYP82E4v2 and CYP82E5v2 genes, nornicotine levels can be reduced in a tobacco plant.

The activity of a nicotine demethylase polypeptide in converting nicotine to nornicotine in a *Nicotiana* plant or plant part is inhibited according to the present invention if this

conversion activity is statistically lower than conversion activity of the same cytochrome P450 polypeptide in a Nicotiana plant or plant part that has not been genetically modified to inhibit the conversion activity of that cytochrome P450 polypeptide and that has been cultured and harvested using the same protocols. In particular embodiments, activity of a cytochrome P450 polypeptide in converting nicotine to nornicotine in a modified Nicotiana plant or plant part according to the invention is inhibited if the activity is less than 95%, less than 90%, less than 80% less than 70%, less than 60%, less than 50%, less than 40%, less than 30%, less than 20% less than 10%, less than 5%, less than 2% or less than 1% of the conversion activity of the same cytochrome P450 polypeptide in a Nicotiana plant that that has not been genetically modified to inhibit the expression of that nicotine demethylase polypeptide and that has been cultured and harvested using the same protocols. The activity of a nicotine demethylase polypeptide in converting nicotine to nornicotine in a Nicotiana plant or plant part is eliminated according to the 20 invention when it is not detectable by the assay methods known in the art or described herein. Methods of determining the activity of a cytochrome P450 polypeptide in converting nicotine to nornicotine in a Nicotiana using gas chromatography are disclosed in the examples below.

In specific embodiments, a cytochrome P450 inhibitory polynucleotide or nicotine demethylase mutation described herein is introduced into a Nicotiana plant or plant part. Subsequently, a Nicotiana plant or plant part having the introduced inhibiting polynucleotide of the present invention is 30 selected using methods known to those of skill in the art including, but not limited to, Southern blot analysis, DNA sequencing, PCR analysis or phenotypic analysis. A polynucleotide or polypeptide sequence of the present invention includes full-length polynucleotide or polypeptide 35 sequences, truncated polynucleotide or polypeptide sequences, fragments of polynucleotide or polypeptide sequences, variants of polynucleotide or polypeptide sequences, sense-oriented nucleotide sequences, antisenseoriented nucleotide sequences, the complement of a sense- or 40 antisense-oriented nucleotide sequence, inverted regions of nucleotide sequences, hairpins of nucleotide sequences, double-stranded nucleotide sequences, single-stranded nucleotide sequences, combinations thereof, and the like. A plant or plant part altered or modified by the foregoing is 45 grown under plant-forming conditions for a time sufficient to modulate the concentration and/or activity of polypeptides of the present invention in the plant. Plant forming conditions are well known in the art and discussed briefly elsewhere herein.

A transformed tobacco plant containing a nicotine demethylase inhibitory polynucleotide sequence described herein has a reduced level of conversion of nicotine to nornicotine. In particular embodiments, conversion of nicotine to nornicotine in a tranformed tobacco plant or plant part according to 55 the invention is less than 95%, less than 90%, less than 80% less than 70%, less than 60%, less than 50%, less than 40%, less than 30%, less than 20%, less than 10%, less than 5%, less than 2% or less than 1% of the conversion in a tobacco plant that has not been genetically modified to inhibit the 60 expression of that nicotine demethylase polypeptide and with has been cultured and harvested using the same protocols. In some instances, the transformed tobacco plant is a converter tobacco plant. Alternatively, the transformed tobacco plant is a nonconverter tobacco plant. Alternatively still, the trans- 65 formed tobacco plant has a conversion rate lower than the rate observed in commercial non-converter tobacco plants.

The level and/or activity of the polypeptide may be modulated by employing a polynucleotide that is not capable of directing, in a transformed plant, the expression of a protein or an RNA. For example, the polynucleotides of the invention may be used to design polynucleotide constructs that can be employed in methods for altering or mutating a genomic nucleotide sequence in an organism. Such polynucleotide constructs include, but are not limited to, RNA:DNA vectors, RNA:DNA mutational vectors, RNA:DNA repair vectors, mixed-duplex oligonucleotides, self-complementary RNA: DNA oligonucleotides and recombinogenic oligonucleobases. Such nucleotide constructs and methods of use are known in the art. See, U.S. Pat. Nos. 5,565,350; 5,731,181; 5,756,325; 5,760,012; 5,795,972 and 5,871,984; each of which is incorporated herein by reference as if set forth in its entirety. See also, International Patent Application Publication Nos. WO 98/149350, WO 99/107865 and WO 99/125921; and Beetham et al. (1999) Proc. Natl. Acad. Sci. USA 96:8774-8778; each of which is incorporated herein by reference as if set forth in its entirety.

The methods of the present invention do not depend on the incorporation of the entire nicotine demethylase inhibitory polynucleotide into the genome, only that the Nicotiana plant or plant part thereof is altered as a result of the introduction of this inhibitory polynucleotide into a cell. As such, the genome may be altered following the introduction of the nicotine demethylase inhibitory polynucleotide into a cell. For example, the inhibitory polynucleotide, or any part thereof, may incorporate into the genome of the plant. Alterations to the genome include, but are not limited to, additions, deletions and substitutions of nucleotides into the genome. While the methods of the present invention do not depend on additions, deletions and substitutions of any particular number of nucleotides, it is recognized that such additions, deletions and/or substitutions comprise at least one nucleotide.

Further, one can reduce the level and/or activity of a nicotine demethylase sequence by eliciting the effects of the sequence only during certain developmental stages and to switch the effect off in other stages where expression is no longer desirable. Control of nicotine demethylase expression can be obtained via the use of inducible or tissue-preferred promoters. Alternatively, the gene could be inverted or deleted using site-specific recombinases, transposons or recombination systems, which would also turn on or off expression of the cytochrome P450 sequence.

According to the present invention, changes in levels, ratios, activity or distribution of cytochrome P450 polypeptides of the present invention, or changes in Nicotiana plant or plant part phenotype, particularly reduced accumulation of nornicotine and its carcinogenic metabolite, NNN, can be measured by comparing a subject plant or plant part to a control plant or plant part, where the subject plant or plant part and the control plant or plant part have been cultured and/or harvested using the same protocols. As used herein, a subject plant or plant part is one in which genetic alteration, such as transformation, has been affected as to the nicotine demethylase polypeptide of interest, or is a *Nicotiana* plant or plant part that is descended from a Nicotiana plant or plant part so altered and which comprises the alteration. A control plant or plant part provides a reference point for measuring changes in phenotype of the subject plant or plant part. The measurement of changes in phenotype can be measured at any time in a plant or plant part, including during plant development, senescence, or after curing. In other embodiments, the measurement of changes in phenotype can be measured in plants grown under any conditions, including from plants grown in growth chamber, greenhouse, or in a field. In one embodiment, changes in phenotype can be measured by determining the nicotine to nornicotine conversion rate. In a preferred embodiment, conversion can be measured by dividing the percentage of nornicotine (as a percentage of the total tissue weight) by the sum of the percentage nicotine and nornicotine 5 (as percentages of the total tissue weight) and multiplying by 100.

According to the present invention, a control plant or plant part may comprise a wild-type Nicotiana plant or plant part, i.e., of the same genotype as the starting material for the 10 genetic alteration that resulted in the subject plant or plant part. A control plant or plant part may also comprise a Nicotiana plant or plant part of the same genotype as the starting material but that has been transformed with a null construct (i.e., with a construct that has no known effect on the trait of 15 interest, such as a construct comprising a selectable marker gene). Alternatively, a control plant or plan part may comprise a Nicotiana plant or plant part that is a non-transformed segregant among progeny of a subject plant or plant part, or a Nicotiana plant or plant part genetically identical to the sub- 20 ject plant or plant part but that is not exposed to conditions or stimuli that would induce suppression of the nicotine demethylase gene of interest. Finally, a control plant or plant part may comprise the subject plant or plant part itself under conditions in which the nicotine demethylase inhibitory 25 sequence is not expressed. In all such cases, the subject plant or plant part and the control plant or plant part are cultured and harvested using the same protocols.

As described elsewhere herein, methods are provided to reduce or eliminate the activity and/or concentration of a 30 nicotine demethylase polypeptide of the present invention by introducing into a Nicotiana plant or plant part a nicotine demethylase inhibitory polynucleotide sequence than is capable of inhibiting expression or function of a nicotine demethylase polypeptide that is involved in the metabolic 35 conversion of nicotine to nornicotine. In some instances, the inhibitory sequence can be introduced by transformation of the plant or plant part such as a plant cell with an expression cassette that expresses a polynucleotide that inhibits the expression of the nicotine demethylase polypeptide. The 40 polynucleotide may inhibit the expression of a nicotine demethylase polypeptide directly, by preventing translation of the nicotine demethylase polypeptide messenger RNA, or indirectly, by encoding a polypeptide that inhibits the transcription or translation of a nicotine demethylase polypeptide gene 45 encoding a nicotine demethylase polypeptide. Methods for inhibiting or eliminating the expression of a gene product in a plant are well known in the art, and any such method may be used in the present invention to inhibit the expression of nicotine demethylase polypeptides.

In other embodiments, the activity of a nicotine demethylase polypeptide of the present invention may be reduced or eliminated by disrupting the gene encoding the nicotine demethylase polypeptide. The invention encompasses mutagenized plants that carry mutations in cytochrome P450 55 genes, where the mutations reduce expression of the nicotine demethylase gene or inhibit the activity of an encoded nicotine demethylase polypeptide of the present invention.

To obtain the desired plants, a *Nicotiana* plant or plant part can be transformed with an expression cassette that is capable 60 of expressing a polynucleotide that inhibits the expression of a nicotine demethylase sequence. Such methods may include the use of sense suppression/cosuppression, antisense suppression, double-stranded RNA (dsRNA) interference, hairpin RNA interference and intron-containing hairpin RNA 65 interference, amplicon-mediated interference, ribozymes, and small interfering RNA or micro RNA.

For cosuppression, an expression cassette is designed to express an RNA molecule corresponding to all or part of a messenger RNA encoding a cytochrome P450 polypeptide of interest (for example, a cytochrome P450 polypeptide comprising the sequence set forth in SEQ ID NO:2 or 5-24 or a sequence having substantial sequence identity to SEQ ID NO:2 or 5-24) in the "sense" orientation. Over-expression of the RNA molecule can result in reduced expression of the native gene. Multiple plant lines transformed with the cosuppression expression cassette are then screened to identify those that show the greatest inhibition of nicotine demethylase polypeptide expression.

The polynucleotide used for cosuppression may correspond to all or part of the sequence encoding a cytochrome P450 polypeptide or the present invention, all or part of the 5' and/or 3' untranslated region of a cytochrome P450 polypeptide transcript, or all or part of both the coding sequence and the untranslated regions of a transcript encoding a cytochrome P450 polypeptide. In some embodiments where the polynucleotide comprises all or part of the coding region for a cytochrome P450 polypeptide of the present invention, the expression cassette can be designed to eliminate the start codon of the polynucleotide so that no polypeptide product will be transcribed.

Cosuppression may be used to inhibit the expression of plant genes to produce plants having undetectable levels for the polypeptides encoded by these genes or may also be used to inhibit the expression of multiple proteins in the same plant (e.g., Broin et al. (2002) Plant Cell 14:1417-1432; and U.S. Pat. No. 5,942,657). Methods for using cosuppression to inhibit the expression of endogenous genes in plants are described in Flavell et al. (1994) Proc Natl. Acad. Sci. USA 91:3490-3496; Jorgensen et al. (1996) Plant Mol. Biol. 31:957-973; Johansen and Carrington (2001) Plant Physiol. 126:930-938; Brain et al. (2002) Plant Cell 14:1417-1432; Stoutjesdijk et al. (2002) Plant Physiol. 129:1723-1731; Yu et al. (2003) Phytochemistry 63:753-763; and U.S. Pat. Nos. 5,034,323; 5,283,184 and 5,942,657; each of which is incorporated herein by reference as if set forth in its entirety. The efficiency of cosuppression may be increased by including a poly-dT region in the expression cassette at a position 3' to the sense sequence and 5' of the polyadenylation signal. See, US Patent Application Publication No. 2002/0048814, incorporated herein by reference as if set forth in its entirety. Typically, such a nucleotide sequence has substantial sequence identity to the sequence of the transcript of the endogenous gene, preferably greater than about 65% sequence identity, more preferably greater than about 95% sequence identity, most preferably greater than about 99% sequence identity (e.g., U.S. Pat. Nos. 5,283,184 and 5,034,323; incorporated herein by reference as if set forth in its entirety).

Inhibition of the expression of the cytochrome P450 polypeptide of the present invention also can be obtained by antisense suppression. For antisense suppression, the expression cassette is designed to express an RNA molecule complementary to all or part of a messenger RNA encoding the cytochrome P450 polypeptide. Over-expression of the antisense RNA molecule can result in reduced expression of the native gene. Accordingly, multiple plant lines transformed with the antisense suppression expression cassette are screened to identify those that show the greatest inhibition of nicotine demethylase polypeptide expression.

The polynucleotide for use in antisense suppression may correspond to all or part of the complement of the sequence encoding the cytochrome P450 polypeptide, all or part of the complement of the 5' and/or 3' untranslated region of the cytochrome P450 polypeptide transcript, or all or part of the complement of both the coding sequence and the untranslated regions of a transcript encoding the cytochrome P450 polypeptide. In addition, the antisense polynucleotide may be fully complementary (i.e., 100% identical to the complement of the target sequence) or partially complementary (i.e., less 5 than 100% identical to the complement of the target sequence) to the target sequence.

Antisense suppression also can be used to inhibit the expression of multiple proteins in the same plant (e.g., U.S. Pat. No. 5,942,657). Furthermore, portions of the antisense 10 nucleotides may be used to disrupt the expression of the target gene. Generally, sequences of at least 50 nucleic acids, 100, 200, 300, 400, 450, 500, 550 nucleic acids, or greater may be used. Methods for using antisense suppression to inhibit the expression of endogenous genes in plants are described, e.g., 15 in Liu et al. (2002) Plant Physiol. 129:1732-1743; and U.S. Pat. Nos. 5,759,829 and 5,942,657; each of which is incorporated herein by reference as if set forth in its entirety. Efficiency of antisense suppression may be increased by including a poly-dT region in the expression cassette at a 20 position 3' to the antisense sequence and 5' of the polyadenylation signal. See, US Patent Application Publication No. 2002/0048814, incorporated herein by reference as if set forth in its entirety.

For dsRNA interference, a sense RNA molecule like that 25 described above for cosuppression and an antisense RNA molecule that is fully or partially complementary to the sense RNA molecule are expressed in the, same cell, resulting in inhibition of the expression of the corresponding endogenous messenger RNA. Expression of the sense and antisense mol- 30 ecules can be accomplished by designing the expression cassette to comprise both a sense sequence and an antisense sequence for the target cytochrome P450 sequence. Alternatively, separate expression cassettes may be used for the sense and antisense sequences. Multiple plant lines transformed 35 with the dsRNA interference expression cassette or expression cassettes are then screened to identify plant lines that show the greatest inhibition of expression of the targeted cytochrome P450 polypeptide. Methods for using dsRNA interference to inhibit the expression of endogenous plant 40 genes are described in Waterhouse et al. (1998) Pros. Natl. Acad. Sci. USA 95:13939-13964; Liu et al. (2002) Plant Physiol. 129:1732'-1743; and International Patent Application Publication Nos. WO 99/149024, WO 99/153050, WO 99/16163 1 and WO 00/149035; each of which is incorpo- 45 rated herein by reference as if set forth in its entirety.

Amplicon expression cassettes comprise a plant virus-derived sequence that contains all or part of the target gene but generally not all of the genes of the native virus. The viral sequences present in the transcription product of the expres-50 sion cassette allow the transcription product to direct its own replication. The transcripts produced by the amplicon may be either sense or antisense relative to the target sequence (i.e., the messenger RNA for a cytochrome P450 polypeptide that is involved in the metabolic conversion of nicotine to norni-55 cotine). Methods of using amplicons to inhibit the expression of endogenous plant genes are described, e.g., in Angell & Baulcombe (1997) *EMBO J.* 16:3675-3684; Angell & Baulcombe (1999) *Plant J.* 20:357-362; and U.S. Pat. No. 6,646, 805, each of which is incorporated herein by reference as if set 60 forth in its entirety.

In some instances, the polynucleotide expressed by the expression cassette of the invention is catalytic RNA or has ribozyme activity specific for the messenger RNA of a cytochrome P450 polypeptide described herein. Thus, the poly- 65 nucleotide causes the degradation of the endogenous messenger RNA, resulting in reduced expression of the cytochrome

P450 polypeptide. This method is described, for example, in U.S. Pat. No. 4,987,071, incorporated herein by reference as if set forth in its entirety.

In other instances, inhibition of the expression of one or more nicotine demethylase polypeptides may be obtained by RNA interference by expression of a gene encoding a micro RNA (miRNA). miRNAs are regulatory agents consisting of about twenty-two ribonucleotides. miRNA are highly efficient at inhibiting the expression of endogenous genes. See, e.g., Javier et al. (2003) *Nature* 425:257-263, incorporated herein by reference as if set forth in its entirety.

For miRNA interference, the expression cassette is designed to express an RNA molecule that is modeled on an endogenous miRNA gene. The miRNA gene encodes an RNA that forms a hairpin structure containing a 22-nucleotide sequence that is complementary to another endogenous gene (target sequence). For suppression of nicotine demethylase polypeptide expression, the 22-nucleotide sequence is selected from a cytochrome P450 polypeptide transcript sequence and contains 22 nucleotides encoding this cytochrome P450 polypeptide sequence in sense orientation and 21 nucleotides of a corresponding antisense sequence that is complementary to the sense sequence. miRNA molecules are highly efficient at inhibiting the expression of endogenous genes, and the RNA interference they induce is inherited by subsequent generations of plants.

Alternatively still, inhibition of the expression of one or more cytochrome P450 polypeptides may be obtained by hairpin RNA (hpRNA) interference or intron-containing hairpin RNA (ihpRNA) interference. These methods are highly efficient at inhibiting the expression of endogenous genes. See, Waterhouse & Helliwell (20013) *Nat. Rev. Genet.* 4:29-38 and the references cited therein.

The expression cassette for hpRNA interference may also be designed such that the sense sequence and the antisense sequence do not correspond to an endogenous RNA in this embodiment, the sense and antisense sequence flank a loop sequence that comprises a nucleotide sequence corresponding to all or part of the endogenous messenger RNA of the target gene. Thus, it is the loop region that determines the specificity of the RNA interference. See, e.g., International Patent Application Publication No. WO 02/00904, incorporated herein by reference as if set forth in its entirety.

Transcriptional gene silencing (TGS) may be accomplished through use of hpRNA constructs where the inverted repeat of the hairpin shares sequence identity with the promoter region of a gene to be silenced. Processing of the hpRNA into short RNAs that can interact with the homologous promoter region may trigger degradation or methylation to result in silencing (Aufsatz et al. (2002) *Proc. Natl. Acad. Sci.* 99:16499-16506; and Mette et al. (2000) *EMBO J.* 19:5194-5201).

In some instances, the polynucleotide encodes an antibody that binds to at least one cytochrome P450 polypeptide, and reduces the activity of a cytochrome P450 polypeptide of the present invention. In another embodiment, the binding of the antibody results in increased turnover of the antibody-cyto-chrome P450 polypeptide complex by cellular quality control mechanisms. The expression of antibodies in plant parts and the inhibition of molecular pathways by expression and binding of antibodies to proteins in plant parts are well known in the art. See, e.g., Conrad & Sonnewald (2003) *Nature Biotech.* 21:35-36, incorporated herein by reference as if set forth in its entirety.

The activity of a cytochrome P450 polypeptide of the present invention can be reduced or eliminated by disrupting the gene encoding the cytochrome P450 polypeptide.

The gene encoding the cytochrome P450 polypeptide may be disrupted by any method known in the art, e.g., by transposon tagging or by mutagenizing plants using random or targeted mutagenesis and selecting for plants that have reduced nicotine demethylase activity or mutations in 5 CYP82E4v2 or CYP82E5v2.

Transposon tagging may be used to reduce or eliminate the activity of one or more cytochrome P450 polypeptides of the present invention. Transposon tagging comprises inserting a transposon within an endogenous cytochrome P450 gene to 10 reduce or eliminate expression of the cytochrome P450 polypeptide.

Methods for the transposon tagging of specific genes in plants are well known in the art. See, e.g., Maes et al. (1999) Trends Plant Sci. 4:90-96; Dharmapuri & Sonti (1999) FEMS 1: Micerobiol. Lett. 179:53-59; Meissner et al. (2000) Plant J. 22:265-274; Phogat et al. (2000) J. Biosci. 25:57-63; Walbot (2000) Curr. Opin. Plant Biol. 2:103-107; Gai et al. (2000) Nucleic Acids Res. 28:94-96; and Fitzmaurice et al. (1999) Genetics 153:1919-1928).

Additional methods for decreasing or eliminating the expression of endogenous genes in plants are also known in the art and can be similarly applied to the instant invention. These methods include other forms of mutagenesis, using mutagenic or carcinogenic compounds including ethyl meth- 25 anesulfonate-induced mutagenesis, deletion mutagenesis, and fast neutron deletion mutagenesis used in a reverse genetics sense (with PCR) to identify plant lines in which the endogenous gene has been deleted. For examples of these methods see, Ohshima et al. (1998) Virology 213:472-481; 30 Okubara et al. (1994) Genetics 137:867-874; and Quesada et al. (2000) Genetics 154:421-4315; each of which is incorporated herein by reference as if set forth in its entirety. In addition, a fast and automatable method for screening for chemically induced mutations, TILLING (Targeting Induced 35 Local Lesions In Genomes), using denaturing HPLC or selective endonuclease digestion of selected PCR products is also applicable to the instant invention. See, McCallum et al. (2000) Nat. Biotechnol. 18:455-457, herein incorporated by reference. 40

Mutations that impact gene expression or that interfere with the function of the encoded cytochrome P450 protein can be determined using methods that are well known in the art. Insertional mutations in gene exons usually result in nullmutants. Mutations in conserved residues can be particularly 45 effective in inhibiting the metabolic function of the encoded protein. Conserved residues of plant cytochrome P450 polypeptides suitable for mutagenesis with the goal to eliminate activity of a cytochrome P450 polypeptide in converting nicotine to nornicotine in a Nicotiana plant or plant part have 50 been described. Conserved residues of plant cytochrome P450 polypeptides are disclosed in FIG. 1A-C, where the residues that differ from the other P450 polypeptides are shaded in grey. The conserved residue is that which is not shaded in grey at each position. Such mutants can be isolated 55 according to well-known procedures.

Dominant mutants can be used to trigger RNA silencing due to gene inversion and recombination of a duplicated gene locus. See, e.g., Kusaba et al. (2003) Plant Cell 15:1455-1467.

60

While a number of sequences are recognized in the practice of the invention, in particular sequences encoding SEQ ID NO:2 and 13 find particular use. While not intending to be bound by any particular mechanism of action, the inventors believed that these sequences encode a nicotine demethylase 65 that catalyzes the oxidative N-demethylation of nicotine to nornicotine. Thus, methods to specifically inhibit these cod30

ing sequences and not other P450 sequences may be beneficial to the recombinant plant. That is, strategies that would lead to inhibition of gene function of this individual locus may prove to be superior to those that inhibit the entire gene family. The P450 enzymes are involved in many mechanisms in the plant, the inhibition of which may prove deleterious or detrimental to the growth and development of the plant or may negatively impact factors such as the disease defense capabilities of the plant. Likewise, because the Nicotiana plant P450 enzymes have been implicated in plant metabolites such as phenylpropanoid, alkaloids, terpenoids, lipids, cyanogenic glycosides, glucosinolates, and a host of other chemical entities, disruption of P450 activity may alter components involved in tobacco flavor, texture, or other properties that would impact the commercial usefulness of the plant. Therefore, the use of the methods discussed above to inhibit expression in a manner that specifically targets the sequence coding for SEQ ID NO:2 or 13 may be preferred, including 20 targeted mutational strategies, such as chimeraplasty. See, e.g., Stewart et al. (2000) Biotechniques 24:838-843; and Graham et al. (2002) Biochim Biophys Acta 1587:1-6; each of which is incorporated herein by reference as if set forth in its entirety. In some embodiments, the methods discussed above are used to inhibit expression in a manner that specifically targets SEQ ID NO:1 (encoding SEQ ID NO:2), SEQ ID NO:50 (encoding SEQ ID NO:13), or both of these sequences.

The compositions of the invention can be used in screening methods to identify nonconverter plants for use in breeding programs. In this manner, the nucleotide sequences of the invention can be used to screen native germplasms for nonconverter plants having a stable mutation in one or more P450 genes identified herein. These nonconverter plants identified by the methods of the invention can be used to develop breeding lines.

In addition to the nucleotide sequences encoding P450 coding sequences, compositions of the invention include an intron sequence in the CYP82E5v2 sequence that can be used in screening methods. While not intending to be bound by any particular mechanism of action, the CYP82E5v2 gene(s) may represent the only member(s) of the cytochrome P450 family involved in the metabolic conversion of nicotine to nornicotine (and as stated previously there is a good likelihood that the CYP82E5v2 cDNAs originated from a single unique genetic locus) in green-leaves of tobacco. For certain applications, it would be useful to have a means of diagnostically differentiating this specific member of the cytochrome P450 gene family from the rest of the closely related sequences within this family. For example, it is possible that within the naturally existing tobacco germplasm (or in mutagenized populations), accessions may exist in which this gene is naturally dysfunctional and may therefore may be valuable as a permanently nonconverter resource. Such dysfunctional genes may include those encoding the polypeptides set forth in SEQ ID NOS:6 and 12. A method to specifically assay for such genotypes (e.g., deletion mutants, rearrangements, and the like) could serve as a powerful tool. The present invention includes primers designed to specifically amplify exon 1 and exon 2 of CYP82E5v2 or CYP82E4v2 where one of the two primer pairs corresponds to the intron between the exons. Examples of primers useful to amplify the exons of CYP82E4v2 include SEQ ID NO:32 with SEQ ID NOS:33-35. Examples of primers useful to amplify the exons of CYP82E5v2 include SEQ ID NO:36 with 37 and SEQ ID NO:38 with 39. These same primers can be used for sequence analysis of the products.

When any cDNA corresponding to a member of the CYP82E2 family is used as a hybridization probe in a Southern blotting assay of tobacco genome is DNA, a complex pattern is observed. This is expected, given that there are multiple, closely related members of this gene family. 5 Because the intron regions of genes are typically less conserved than exons, it is predicted that the use of an intronspecific probe would reduce this complexity and better enable one to distinguish the gene(s) corresponding to the CYP82E4v2 gene or the CYP82E5v2 gene from the other 10 members of the family. The use of a CYP82E4v2 or CYP82E5v2 intron-specific probe, and/or the PCR primers used to generate products provide powerful tools in assays to determine whether any naturally occurring, or mutagenized, tobacco plants possess deletions or rearrangements that may 15 render the gene inactive. Such a plant can then be used in breeding programs to create tobacco lines that are incapable of converting.

Transformed Plants, Plant Parts and Products Having Reduced Nornicotine and NNN Content

The cytochrome P450 polynucleotides of the invention, and variants and fragments thereof, can be used in the methods of the present invention to inhibit cvtochrome P450s that are involved in the metabolic conversion of nicotine to nornicotine in a plant. In this manner, inhibitory sequences that 25 target expression or function of a cytochrome P450 polypeptide disclosed herein are introduced into a plant or plant cell of interest. The expression cassettes described herein can be introduced into a plant of interest, for example, a *Nicotiana* plant as noted herein below, using any suitable transformation 30 methods known in the art including those described herein.

The methods of the invention do not depend on a particular method for introducing a sequence into a plant or plant part, only that the desired sequence gains access to the interior of at least one cell of the plant or plant part. Methods for introduc-35 ing polynucleotide sequences into plants are known in the art and include, but are not limited to, stable transformation methods, transient transformation methods and viral-mediated transformation methods.

Transformation protocols as well as protocols for introduc- 40 ing heterologous polynucleotide sequences into plants vary depending on the type of plant or plant cell targeted for transformation. Suitable methods of introducing polynucleotides into plant cells of the present invention include microinjection (Crossway et al. (1986) Biotechniques 4:320-334), 45 electroporation (Shillito et al. (1987) Meth. Enzymol. 153: 313-336; and Riggs et al. (1986) Proc. Natl. Acad. Sci. USA 83:5602-56116), Agrobacterium-mediated transformation (U.S. Pat. Nos. 5,104,310; 5,149,645; 5,177,010; 5,231,019; 5,463,174; 5,464,763; 5,469,976; 4,762,785; 5,004,863; 50 5,159,135; 5,563,055 and 5,981,840), direct gene transfer (Paszkowski et al. (1984) EMBO J. 1:2717-2722) and ballistic particle acceleration (U.S. Pat. Nos. 4,945,050; 5,141,131; 5,886,244; 5,879,918 and 5,932,782; Tomes et al. (1995) in Plant cell, tissue, and organ culture fundamental methods, ed. 55 Gamborg & Phillips (Springer-Verlag, Berlin); McCabe et al. (1988) Biotechnology 6:923-926). See also, Weissinger et al. (1988) Ann. Rev. Genet. 22:421-477; Christou et al. (1988) Plant Physiol. 87:671-674 (soybean); McCabe et al. (1988) Bio/Technology 6:923-926 (soybean); Finer & McMullen 60 (1991) In Vitro Cell Dev. Biol. 27P:175-182 (soybean); Singh et al. (1998) Theor. Appl. Genet. 96:319-324 (soybean); De Wet et al. (1985) in The experimental manipulation of ovule tissues, ed. Chapman et al. (Longman, New York.); pp. 197-209 (pollen); Kaeppler et al. (1990) Plant Cell Reports 9:415- 65 418; and Kaeppler et al. (1992) Theor. Appl. Genet, 84:560-566 (whisker-mediated transformation); D'Halluin et al.

(1992) *Plant Cell* 4:1495-1505 (electroporation); each of which is incorporated herein by reference as if set forth in its entirety.

Any plant tissue that can be subsequently propagated using clonal methods whether by organogenesis or embryogenesis, may be transformed with a recombinant construct comprising a cytochrome P450 inhibitory sequence, for example, an expression cassette of the present invention. As used herein, "organogenesis" means a process by which shoots and roots are developed sequentially from meristematic centers. As used herein, "embryogenesis" means a process by which shoots and roots develop together in a concerted fashion (not sequentially), whether from somatic cells or gametes. Exemplary tissues that are suitable for various transformation protocols described herein include, but are not limited to, callus tissue, existing meristematic tissue (e.g., apical meristems, axillary buds and root ineristems) and induced meristem tissue (e.g., cotyledon meristem and hypocotyl meristem), hypocotyls, cotyledons, leaf disks, pollen, embryos and the like.

As used herein, "stable transformation" means that the nucleotide construct of interest introduced into a plant integrates into the genome of the plant and is capable of being inherited by the progeny thereof. Likewise, "transient transformation" means that a sequence is introduced into the plant and is only temporally expressed or is only transiently present in the plant.

The inhibitory sequences of the invention can be provided to a plant using a variety of transient transformation methods. The inhibitory sequences of the invention can be transiently transformed into the plant using techniques known in the art. Such techniques include viral vector systems and the precipitation of the polynucleotide in a manner that precludes subsequent release of the DNA. Thus, the transcription from the particle-bound DNA can occur, but the frequency with which it is released to become integrated into the genome is greatly reduced. Such methods include the use of particles coated with polyethyenlimine (PEI; Sigma 4P3143).

Alternatively, the inhibitory sequence of the invention may be introduced into plants by contacting plants with a virus or viral nucleic acids. Generally, such methods involve incorporating an expression cassette of the invention within a viral DNA or RNA molecule. Promoters for use in the expression cassettes of the invention also encompass promoters utilized for transcription by viral RNA polymerases. Methods for introducing polynucleotides into plants and expressing a protein encoded therein, involving viral DNA or RNA molecules, are known in the art. See, e.g., U.S. Pat. Nos. 5,889, 191; 5,889,190; 5,866,785; 5,589,367; 5,316,931; and Porta et al. (1996) *Molecular Biotechnology* 5:209-221; each of which is incorporated herein by reference as if set forth in its entirety.

Transformed cells may be grown into *Nicotiana* plants in accordance with conventional methods. See, e.g., Vasil & Hildebrandt (1965) *Science* 150:889; Negaard & Hoffman (1989) *Biotechniques* 7(8):808-812. These plants may then be grown, and either pollinated with the same transformed line or different lines, and the resulting progeny having expression of the desired phenotypic characteristic identified, i.e., reduced expression of one or more nicotine demethylases that are involved in the metabolic conversion of nicotine to nornicotine, and thus reduced content of nornicotine, and a concomitant reduced content of its nitrosamine metabolite, NNN, in the plant, particularly in the leaf tissues. Two or more generations may be grown to ensure that expression of the desired phenotypic characteristic is stably maintained and inherited and then seeds harvested to ensure expression of the

25

desired phenotypic characteristic has been achieved. In this manner, the present invention provides transformed seed (also referred to as "transgenic seed") having a polynucleotide of the invention, for example, an expression cassette of the invention, stably incorporated into their genome.

The compositions and methods of the invention can be used to reduce the nornicotine content, particularly in the leaves and stems, of any plant of the genus Nicotiana including, but not limited to, the following species: acuminata, affinis, alata, attenuate, bigelovii, clevelandii, excelsior, forgetiana, glauca, glutinosa, langsdorffii, longiflora, obtusifolia, palmeri, paniculata, plumbaginifolia, qudrivalvis, repanda, rustica, suaveolens, sylvestris, tabacum, tomentosa, trigonophylla and x sanderae. The present invention also encompasses the transformation of any varieties of a plant of the genus Nicotiana, including, but not limited to, Nicotiana acuminata multiflora, Nicotiana alata grandiflora, Nicotiana bigelovii quadrivalvis, Nicotiana bigelovii wallacei, Nicotiana obtusifolia obtusifolia, Nicotiana obtusifolia plameri, 20 Nicotiana quadrivalvis bigelovii, Nicotiana quadrivalvis quadrivalvis, Nicotiana quadrivalvis wallacei, and Nicotiana trigonophylla palmeri, as well as varieties commonly known as flue or bright varieties, Burley varieties, dark varieties and oriental/Turkish varieties.

The transgenic plants of the genus Nicotiana as described herein are suitable for conventional growing and harvesting techniques, such as cultivation in manure rich soil or without manure, bagging the flowers or no bagging, or topping or no topping. The harvested leaves and stems may be used in any 30 traditional tobacco product including, but not limited to, pipe, cigar and cigarette tobacco, and chewing tobacco in any form including leaf tobacco, shredded tobacco, or cut tobacco.

Thus the present invention provides a Nicotiana plant, particularly leaf tissues of these plants, comprising an expres- 35 sion cassette of the invention and a reduced amount of nornicotine and N'-nitrosonornicotine. As used herein, "a reduced amount" or "a reduced level" means an amount of nornicotine and/or NNN in a treated or transgenic plant of the genus Nicotiana or a plant part or tobacco product thereof that is less 40 than what would be found in a plant of the genus Nicotiana or a plant part or tobacco product from the same variety of tobacco, processed (i.e., cultured and harvested) in the same manner, that has not been treated or was not made transgenic for reduced nornicotine and/or NNN. The amount of nornicotine may be reduced by about 10% to greater than about 90%, including greater than about 20%, about 30%, about 40%, about 50%, about 60%, about 70% and about 80%. The conversion of nicotine to nornicotine can be less than 0.3%, less than 0.5%, less than 0.7%, between 0.1%-0.5%, between 50 0.1%-0.4%, between 0.1%-0.7%, or between 0.1%-1.0% in plants, plant parts, and products of the present invention, and more specifically in plants, plant parts having mutations in CYP82E4v2 and CYP825v2.

As used herein, "tobacco products" means, but is not lim- 55 ited to, smoking materials (e.g., any cigarette, including a cigarillo, a non-ventilated or vented recess filter cigarette, a cigar, pipe tobacco), smokeless products (e.g., snuff, chewing tobacco, biodegradable inserts (e.g., gum, lozenges, dissolving strips)). See, e.g., US Patent Application Publication No. 60 2005/0019448, incorporated herein by reference as if set forth in its entirety. Tobacco product also includes blends that can be made by combining conventional tobacco with differing amounts of the low nornicotine and/or NNN tobacco described herein. Tobacco products also includes plant or 65 plant part of the genus Nicotiana as described above is cured tobacco.

The tobacco products described herein reduce the carcinogenic potential of tobacco smoke that is inhaled directly with consumption of a tobacco product such as cigars, cigarettes, or pipe tobacco or inhaled as secondary smoke (i.e., by an individual that inhales the tobacco smoke generated by an individual consuming a tobacco product such as cigars, cigarettes, or pipe tobacco). The cured tobacco described herein can be used to prepare a tobacco product, particularly one that undergoes chemical changes due to heat, comprising a reduced amount of nornicotine and/or NNN in the smoke stream that is inhaled directly or inhaled as secondary smoke. In the same manner, the tobacco products of the invention may be useful in the preparation of smokeless tobacco products such as chewing tobacco, snuff and the like.

The tobacco products derived from the transgenic tobacco plants of the present invention thus find use in methods for reducing the carcinogenic potential of these tobacco products, and reducing the exposure of humans to the carcinogenic nitrosamine, NNN, particularly for individuals that are users of these tobacco products. The following examples are offered by way of illustration and not by way of limitation.

EXPERIMENTAL

The following materials and protocols are utilized in the experiments described herein below.

Plant Materials

All plant materials were obtained from Dr. Earl Wernsman, Department of Crop Science, North Carolina State University (Raleigh, N.C.). N. tomentosiformis and N. sylvestris seed were obtained from the Nicotiana germplasm collection maintained by North Carolina State University. Plants were grown in growth chambers for two months until they were transferred to a greenhouse. Burley lines DH 98-325-5 (nonconverter) and DH 98-325-6 (converter) represent nearisogenic lines, i.e., recovered from the same maternal haploid plant. SC58 is a flue-cured tobacco variety, nonconverter individuals of which are designated $SC58(c_{\tau}c_{\tau})$. SC58 $(C_{\tau}C_{\tau})$ is a near-isogenic stable converter line that originated though the introgression of the single dominant converter locus (C_r) found in the tobacco progenitor species N. tomentosiformis into SC58 (Mann et al. (1964) Crop Sci., 4:349-353). After eight additional backcrosses to SC58, the nearisogenic SC58($C_T C_T$) line was created and was subsequently maintained via self-fertilization.

All plants were maintained in growth chambers or greenhouses using standard potting soil and fertilizer. Plants kept in a 14/10 light/dark cycle, were watered once a day as needed and fertilized with Peters Professional® All Purpose Plant Food fertilizer (20-20-20; Spectrum Brands Inc.; Madison, Wis.) once a week. To facilitate senescence, detached green leaves were treated by dipping each leaf for 30 seconds in a 0.2% ethephon solution. Leaves were cured in plastic bags under dark conditions until they turned yellow.

For the ethyl methane sulphonate (EMS) mutagensis, two grams of seed from the strong converter Burley tobacco line DH98-325-6 were surface-sterilized in 50% bleach for 12 minutes and rinsed 6 times in sterile dH2O. Approximately 80 mg aliquots were then placed in screw cap vials and imbibed in 1 ml dH₂O for 12 hours. The dH₂O was decanted, and the seed was incubated in 1 ml of 0.5% EMS (Sigma; St. Louis, Mo.) and rocked gently on a Nutator® Mixer (TCS Scientific Corp.; New Hope, Pa.) at room temperature for 12 hours in the dark. After removal of the EMS solution, the treated seeds were washed eight times in 1 ml volumes of dH₂O with gentle shaking for 5 minutes. After the final wash, seeds were collected onto a Buchner funnel for a final rinse and subsequently allowed to dry on filter paper before sowing on soil. Seedlings were grown on standard float trays for about 7 weeks, then transplanted to a field. Field-grown plants were allowed to self-pollinate, and 5-10 capsules (containing M_1 generation seed) per plant were collected and catalogued ⁵ corresponding to approximately 4000 independent M_0 individuals.

Isolation of NtabCYP82E5v2 cDNA

Total RNA was extracted from green and senescing leaves 10 of DH98-325-6 tobacco using TRIzol® (Invitrogen; Carlsbad, Calif.) following the protocol of the manufacturer. Genomic DNA was removed by treating the extracts with TURBO DNA-free™ DNase Kit (Ambion; Austin, Tex.) following the protocol of the manufacturer. cDNA synthesis was performed with the StrataScript[™] First-Strand Kit (Stratagene; Cedar Creek, Tex.) using oligo dT primers and 3 µg of DNase-free total RNA. Full-length NtabCYP82E5v2 cDNAs were amplified using the ESOrf_F forward and E5Orf_R 20 reverse primers, 10 ng tobacco leaf cDNA library as a template and Long Range™ PCR Enzyme Blend (GeneChoice, Inc.; Frederick, Md.) following the protocol of the manufacturer. DNA sequence information of the primers is listed in the sequence listing and above. PCR conditions were as fol- 25 lows: 3 minutes initial denaturation at 95° C. followed by 33 cycles of denaturation at 94° C. for 30 seconds, annealing at 60° C., for 30 seconds and extension at 68° C. for 90 seconds, followed by an incubation at 68° C. for 7 minutes. PCR products were cloned into pGEM®-T Easy cloning vector (Promega Corp.; Madison, Wis.) and sequenced according to the dideoxy method using synthetic oligonucleotides as primers (Sanger et al. (1977) Proc. Natl. Acad. Sci. U.S.A. 74: 5463-5467).

Isolation of the Genomic CYP82E5 Fragments

Genomic DNA was isolated from the green leaves of *N. tomentosiformis* and DH98-325-6 tobacco using a cetyltrimethyl ammonium bromide-based extraction procedure according to the instructions of the NucleoSpin® Plant kit (BD Bioscience Clontech; Palo Alto, Calif.).

Genomic fragments of the CYP82E5 orthologs were amplified in two consecutive PCR reactions using nested primers. The first PCR mixture contained 0.5 µM each of the 45 E5Gen_F1 forward (SEQ ID NO:40) and E5Gen R1 reverse (SEQ ID NO:41) primers, 200 µM of each dNTP, 20 ng of genomic DNA template and 2 U of Platinum® Taq polymerase (Invitrogen). PCR amplification was performed under the following conditions: 90° C. for 3 minutes; 30 cycles of denaturation at 94° C. for 30 seconds, annealing at 57° C. for 30 seconds and extension at 68° C. for 2.2 minutes followed by a final extension at 68° C. for 5 minutes. The composition and conditions of the second PCR were the same as those 55 described for the first amplification, except primers E5Gen_F2 (SEQ ID NO:42) and E5Gen_R2 (SEQ ID NO:43) were added and $1 \mu l$ of a 50-fold diluted sample of the first PCR product was used as a template. The purified products of the final PCR were ligated into the pGEM-T-Easy 60 cloning vector and subjected to DNA sequencing. Expression of the NtabCYP82E5v2 cDNA in Yeast

To facilitate the expression of the NtabCYP82E5v2 cDNA in yeast, the open reading frame of the gene was inserted 65 downstream of the galactose-inducible, glucose-repressed GAL10-CYC1 promoter of the pYeDP60 yeast expression

vector (Cullin & Pompon 1988). The recombinant plasmid was introduced into WAT11 yeast strain using the lithiumacetate-based transformation method as described by Gietz et al. (1992) *Nucleic Acids Res.* 20:1425-1425. Yeast culturing, galactose-mediated induction of gene expression and the isolation of the microsomal fractions were performed according to the protocol of Pompon et al. (1996) *Methods Enzymol.* 272:51-64.

Kinetic Analysis of NtabCYP82E5-Mediated Nicotine Metabolism

The nicotine metabolism assays were performed in a reaction mixture containing 0.75 mM NADPH, 2.5 µM [pyrrolidine-2-14C] nicotine (Moravek Biochemicals; Brea, Calif.) and 90 µg of yeast microsomes in a final volume of 150 µl Pi buffer, pH 7.1. Nicotine concentrations were adjusted by the addition of nonradiolabelled nicotine to final concentrations ranging between 0.7 and 13 µM. Microsomal preparations isolated from yeast containing an empty vector were used as negative control. Reaction mixtures were incubated at 25° C. for 7 minutes, and the reaction was arrested with 50 µl of acetone. After centrifugation at 16,000×g for 3 minutes, 50 µl the sample was spotted on a 250-µm Whatman K6F silica plate. The plates were developed in choloroform:methanol: ammonia (60:10:1) solvent system. Nicotine and its nornicotine derivative were quantified by measuring the relative abundance of their radioactive traces with a Bioscan System 400 imaging scanner. K_m and v_{max} values were calculated by fitting the Michaelis-Menten equation to nicotine demethylation measurements at different substrate concentrations using the SigmaPlot® 10.0 graphing software (Systat Software, Inc.; San Jose, Calif.).

³⁵ Quantitative Real-Time PCR Analysis of CYP82E4v2 and CYP82E5v2 Expression

Expression of the N. tomentosiformis and tobacco orthologs of CYP82E4v2 and CYP82E5v2 were analyzed by allele-specific, quantitative real-time PCR (qrt-PCR) analysis using SYBR® Green I fluorescence chemistry (Morrison et al. (1998) Biotechniques 24: 954-962). Total RNA was extracted from the green and cured leaves of 3-month-old plants using the methods already described for NtabCYP82E5v2 cDNA isolation herein. CYP82E4v2 and CYP82E5v2 transcripts were amplified using the E4Rt_F (SEQ ID NO:44) and E5Rt_F (SEQ ID NO:46) forward in conjunction with the E4Rt_R (SEQ ID NO:45) and the E5Rt_R reverse (SEQ ID NO:47) primers, respectively. The qrt-PCR mixture contained 1×iQ[™] SYBR® Green Supermix (Bio-Rad Laboratories) 0.5 µM of each primer and 0.5 µg of cDNA. Transcript measurements were obtained using standard curves generated by amplifying known amounts of target DNA and a 220-bp fragment of the glyceraldehyde-3phosphate dehydrogenase (G3PDH) gene, amplified with the G3PDH_F (SEQ ID NO:48) and G3PDH_R (SEQ ID NO:49) primers, to provide an internal standard for cDNA measurements. DNA amplifications were performed using the real time PCR system (Bio-Rad Laboratories; Hercules, Calif.) under the following conditions: 95° C. for 2 minutes; 35 cycles of 95° C. for 30 seconds, 57 C for 30 seconds, 72° C. for 50 seconds followed by final extension at 72° C. for 5 minutes. DNA sequences of the amplicons generated by qrt-PCR were verified by cloning the PCR products and sequencing 20 randomly selected clones.

Transgenic Plant Analysis

The RNAi-based gene silencing constructs are assembled in a version of the pKYL80 cloning vector (Schardl et al. (1987) *Gene* 61:1-11) that is engineered to contain a 151-bp fragment of the soybean FAD3 gene intron between the Xhol and Sacl restriction sites of the polylinker (pKYLX80I). To create a construct in which the FAD3 intron was flanked by a sense and antisense fragment of CYP82E5v2, a 400-bp intron region is cloned between the HindIII-Xhol and Sacl-Xbal ¹⁰ restriction sites of pKYLX801 in its sense and antisense orientation, respectively. The resulting HindIII-Xbal fragment containing the CYP8E5v2 sense arm, FAD3 intron, and CYP82E5v2 antisense arm is subcloned into the pKYLX71 ¹⁵ plant binary expression vector (Maiti et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:6110-6114) between the 35S CaMV promoter and a rubisco small subunit terminator.

Overexpression constructs are created by replacing the 3-glucuronidase ORF of the plant binary expression vector pBI121 (Clontech) with the full-length coding regions of the CYP82E5v2, CYP82E5v2 variants, or CYP82E4v2 variants cDNAs. This places the tobacco P450s under the transcriptional control of the 35S CaMV promoter. The pBI121- and ²⁵ pKYLX71-based constructs are transformed into *Agrobacterium tumefaciens* strain LBA 4404 and introduced into tobacco cultivars Petite Havana and DH98-325-6 (converter), respectively, using established protocols (Horsch et al. (1985) *Science* 227:1229-1231).

Northern Blot Analysis

Total cellular RNAs are isolated from tobacco leaves using the TRIZOL® method as described by the manufacturer (Invitrogen). Five to ten micrograms of RNA are size fractionated on a 1.2% agarose gel prepared in TBE buffer. RNA immobilization, probe labeling, and signal detection are carried out using the DIG nucleic acid labeling and detection kits according to the manufacturer's instructions (Roche). Alternatively, probes are synthesized using ³²P-dCTP according to protocols accompanying the Random Primed DNA Labeling kit (Roche).

Isolation of DNA from EMS-Mutagenized Plants.

Genomic DNA was isolated from young leaf material of a 45 single greenhouse-grown M₁ plant from each independent M₁ seed pool. 1.5 ml screw cap vials were used to collect a single leaf disc from each plant and the samples were stored at -80° C. until they were processed. The leaf tissue was deposited to the bottom of the vial by centrifugation, and the tubes were set in a shallow amount of liquid nitrogen to facilitate grinding using plastic pestles. Genomic DNA was isolated from the ground material by adding 320 µl extraction buffer (200 mM Tris-Cl, pH 7.5, 250 mM NaCl, 25 mM 55 EDTA, 0.5% SDS). Samples were vortexed for 20 seconds, and left at room temperature for 5 minutes, followed by a 37° C. incubation for 3 minutes. The lysed material was centrifuged 10 minutes, and the supernatant transferred to fresh 60 tube. 50 µl of protein precipitation solution (Qiagen) was added, the samples briefly vortexed, then set on ice for 5 minutes, followed by a 4 minutes centrifugation at 16,000×g. Precipitation of nucleic acids was accomplished by adding 0.7 volume isopropanol to the supernatant followed by cen- $_{65}$ trifugation for 7 minutes at 12,000×g. After a 70% ethanol wash, the pellet was resuspended in 50 µl TE.

Template Preparation for Sequencing.

CYP82E4v2 and CYP82E5v2 each contain two exons separated by a large intron (1001 bp and 1054 bp, respectively). PCR primers are designed to specifically amplify exon 1 and exon 2 for each gene. Due to the high sequence homology shared among the cDNAs of CYP82E2 gene familv members, two primer pairs correspond to intron sequence in order to obtain amplification products specific to the desired gene (owing to the fact that the intron sequences of this gene family are not nearly as conserved as the exon sequences). The exon 1-specific primers for CYP82E4v2 are 5'-TGGAATTATGCCCATCCTACA-3' (forward) (SEQ ID NO:32) and 5'-CATTAGTGGTTGCACCTGAGG-3' (reverse) (SEQ ID NO:33). CYP82E4v2 exon 2-specific primers are 5'-GATGAGATGTGTGCATACTTG-3' (forward) (SEQ ID NO: 34) and 5'-CCAAATTAGAAAAACTCGTACTG-3' (reverse) (SEQ ID NO:35). For CYP82E5v2, the primers specific for amplifying exon 1 are 5'-ATTGTAGGACTAG-TAACCCTTACAC-3' (forward) (SEQ ID NO:36) and 5'-GAGGCACAAAGAATTCTCATC-3' (reverse) (SEQ ID NO:37); and for CYP82E5v2 exon 2, the primers 5'-GAG-TAGAGGGATTGTTTCCG-3' (forward) (SEQ ID NO:38) and 5'-GTACAATCAAGATAAAACATCTAAGG-3' (reverse) (SEQ ID NO:39) are used. PCR reactions were conducted in 96-well microtiter plates using 1 µl of template in a 25 µl reaction volume containing 10 pmoles each primer, 200 µM dNTP, 1.5 mM MgCl and 1.4 units of high fidelity Tag DNA polymerase (Roche). DNA amplification of CYP82E4v2 sequences was performed using an initial 3 minutes denaturation at 94° C., followed by 30 cycles of denaturation at 94° C. for 30 seconds, primer annealing at 55° C. for 30 seconds, and extension at 72° C. for 1.5 minutes, followed by a final 7 minutes 72° C. extension. Amplification conditions for CYP82E5v2 gene fragments were the same except the annealing temperature was 53° C. PCR reaction products were purified using Millipore's Montage PCRµ96 filter units. Agarose gel electrophoresis was used to estimate the quantity and quality of the cleaned up PCR products prior to sequencing.

Sequence Analysis.

The PCR products were subjected to cycle sequencing using Applied Biosystems Big Dye Version 3.1 in 96-well format sequencing reactions according to the manufacturer's instructions. The primers used for sequencing (3.2 pmoles per reaction) are as follows: 5'-TGGAATTATGCCCATC-CTACA-3' (SEQ ID NO:32) and 5'-CCTATAAAAAG-GAAGTTGCCG-3' (SEQ ID NO:44) (forward and reverse primers for exon 1 of CYP82E4v2); 5'-GATGAGATGTGT-GCATACTTG-3' (SEQ ID NO:34) and 5'-CCAAATTA-GAAAAACTCGTACTG-3' (SEQ ID NO:35) (forward and reverse primers for exon 2 of CYP82E4v2): 5'-ATTGTAG-GACTAGTAACCCTTACAC-3' (SEQ ID NO:36) and 5'-CT-CATCTTTTTTCCATTTATCATTC-3' (SEQ ID NO:45) (forward and reverse primers for exon 1 of CYP82E5v2); and 5'-CAAGGTTCGGCAGATAGTG-3' (SEQ ID NO:46) and 5'-GTACAATCAAGATAAAACATCTAAGG-3' (SEQ ID NO:39) (forward and reverse primers for exon 2 of CYP82E5v2). Sequencing reactions were cleaned up using EdgeBioSystems 96-well plates and analyzed using highthroughput Applied Biosystems 3700 or 3730 capillary sequencers. Sequences were aligned using the Clustal W algorithm as represented in the Vector NTI software package

(Invitrogen). Putative mutations are verified by repeating the sequence analysis on sibling M1 plants grown from the cognate Mo seed lot.

Alkaloid Analysis

Alkaloid analyses are performed by gas chromatography as described previously. Briefly, alkaloid analyses are performed by detaching tobacco leaves and dipping them in a solution of 1% ethephon, then, air-curing in a growth chamber for 7-10 days. Cured leaves are dried at 50° C. for two days and ground to a fine powder. A quantitative determination of the nicotine, nornicotine, anatabine and anabasine content was made using a Perkin Elmer Autosystem XL Gas Chromatograph according to the "LC-Protocol" established 1 at the University of Kentucky (available online at the University of Kentucky's website), hereby incorporated by reference.

EXAMPLE 1

Isolation and Characterization of CYP82E5V2

Nicotine Conversion in N. tomentosiformis and Various Tobacco Genotypes

To assess the alkaloid composition of wild-type plants, concentrations of nicotine and nornicotine were determined in the green and cured leaves of N. tomentosiformis and various tobacco genotypes using gas chromatography. In N. tomentosiformis, nornicotine appears as the predominant 30 alkaloid in both the green and cured leaves (Table 1). Converter Burley cultivar DH-98-325-6 and converter Flue-Cured cultivar SC58C contain low levels of nornicotine (about 3%) in the green leaves, but a large percentage (about 35 95 and 25%, respectively) nicotine content is metabolized into nornicotine once the leaves senesce (see, Table 1). Levels of nornicotine are low in both the green and cured leaves of nonconverter Burley and Flue-Cured cultivars DH98-325-5 and SC58NC, respectively, although nicotine conversion 40 slightly increases after curing.

40

where the primers are complementary to the 5' and 3' termini of the NtabCYP82E3 coding region. Primer design was based on the observation that all members of the closely-related CYP82E2 gene subfamily display a high degree of DNA sequence identity at the regions immediately following the ATG initiation signal and preceding the TAA stop codon (Siminszky et al. (2005) Proc. Natl. Acad. Sci. U.S. A. 102: 14919-14924) providing well conserved primer target sites for the amplification of additional family members. A PCR using CYP82E3-specific primers and permissive annealing temperatures was conducted to amplify putative nicotine demethylase genes from a tobacco green leaf cDNA library. DNA sequence analysis of 20 randomly selected clones generated from the PCR products yielded 19 NtabCYP82E3 cDNA sequences and one cDNA whose DNA sequence is different from all functionally characterized members of the CYP82E2 gene subfamily (Siminszky et al. (2005) Proc. ²⁰ Natl. Acad. Sci. U.S.A. 102: 14919-14924). A homology search against the GenBank database using the BLAST algorithm reveals that the clone representing the unique DNA sequence shares a 99.7% nucleotide and a 99.2% predicted amino acid identity with the tobacco CYP82E5v1 cDNA of 25 unknown function. According to the guidelines of the P450 nomenclature committee, the cDNA was named NtabCYP82E5v2. The predicted amino acid sequence of NtabCYP82E5v2 is 89.2, 91.9 and 91.3% identical to that of NtabCYP82E2, NtabCYP82E3 and NtabCYP82E4v2, respectively.

NtabCYP82E5v2 Encodes for a Functional Nicotine Demethylase

To determine whether NtabCYP82E5v2 encodes nicotine demethylase activity, the N.tabCYP82E5 cDNA is expressed in the WAT11 yeast strain. The WAT11 strain was engineered to provide high efficiency transgene expression and an optimal redox environment for P450-mediated reactions by expressing the Arabidopsis P450 reductase gene, ATR1, under the transcriptional control of a galactose-inducible pro-

TABLE 1

Alkaloid analysis of green and cured leaves of <i>N. tomentosiformis</i> and various tobacco genotypes.							
	% Nicotine ^a % Nornicotine ^a % Conversion ^b						
Plant	Phenotype	Green	Cured ^c	Green	Cured ^c	Green	Cured
N. tomentosiformis	GLC	0.003^d (0.0002)	0.007	0.188	0.299 (0.031)	98.4 (2.4)	97.7 (2.1)
DH98-325-5	NC	1.243 (0.334)	1.332 (0.321)	0.032	0.040	2.6 (0.2)	3.0 (0.3)
DH98-325-6	SLC	1.113 (0.095)	0.082	0.041 (0.004)	1.416 (0.187)	3.5 (0.1)	94.4 (1.6)
SC58NC	NC	1.422	1.487	0.032	0.039 (0.001)	2.2	2.5
SC58C	SLC	1.171	0.501	0.029	0.164	2.4	24.7

^aPercentage of leaf drv weight.

^b[% nornicotine (% nicotine + % nornicotine)⁻¹] x 100

"Leaves are treated with 0.2% ethephon and cured until they turned yellow.

^dMean over the alkaloid content of three plants, except SC58C where one plant was used due to the low frequency

of conversion. "Values in parenthesis represent standard error of the mean.

Abbreviations: GLC, green-leaf converter; NC, nonconverter; SLC, senescing-leaf converter.

Isolation of the NtabCYP82E5v2 cDNA from Tobacco

To identify putative nicotine demethylase genes from tobacco, a PCR-based gene amplification strategy was used

65 moter (Pompon et al. (1996) Methods Enzymol. 272: 51-64). Microsomal fragments isolated from yeast expressing NtabCYP82E5v2 actively catalyze the N-demethylation of

35

45

nicotine. The K_m and v_{max} values for the NtabCYP82E5v2mediated nicotine demethylase reaction are $5.6\pm1.4 \mu$ M and 0.7±0.02 nmol min⁻¹ mg⁻¹ protein (mean±standard error), respectively. In contrast, no nicotine demethylase activity is evident when microsomes isolated from yeast transformed with an empty plasmid are added to the catalytic assay. NtabCYP82E5v2 is Derived from Progenitor N. tomentosiformis

As the allotetraploid genome of tobacco consists of two 10 genetic components, the S genome donated by N. sylvestris and the T genome derived from N. tomentosiformis, which of the two progenitor species contributed NtabCYP82E5v2 to tobacco is unknown. To this end, CYP82E2-specific primers were designed to amplify a genomic fragment of the CYP82E5 genes from tobacco and the NtabCYP82E5v2donating progenitor. PCR amplifications produce a distinct 2243 bp product when genomic DNA isolated from tobacco or *N. tomentosiformis* is used as template, but amplicons are 20 not detected when genomic DNA of N. sylvestris is amplified indicating that NtabCYP82E5v2 was derived from N. tomentosiformis. DNA sequence analysis reveals that the genomic fragments of NtabCYP82E5v2 and NtomCYP82E5 contain a 1054 bp intron flanked by a 604 bp 5' and a 585 bp 3' coding 25 region and differ at 15 positions of which nine and six are located within the coding region and intron, respectively.

Furthermore, the coding regions of ten randomly selected genomic clones isolated from tobacco share 100% nucleotide 30 identity with the NtabCYP82E5v2 cDNA suggesting that NtabCYP82E5v2 is the only CYP82E5 variant located in DH98-325-6 tobacco.

NtabCYP82E5v2 is Expressed at Low Levels in the Green and Yellow Leaves

To investigate the catalytic role of CYP82E5 in planta, the transcriptional profile of CYP82E5v2 in the green and senescing leaves of converter (DH98-325-6) and nonconverter (DH98-325-5) Burley, converter and nonconverter Flue-Cured (SC58) tobacco, and N. tomentosiformis was determined using quantitative real-time (qrt)-PCR analysis.

TABLE 2

Absolute quantification of the CYP82E4v2 and CYP82E5v2 cDNA derived from the green and cured leaves of N. tomentosiformis and different tobacco genotypes by quantitative real-time polymerase chain reaction analysis

	CYP82	2E4v2	CYP82E5v2		- 50
Sample	Green	Cured Pg	Green	Cured	
N. tomentosiformis	0.35c (0.16)	17.94e (2.57)	1.73d (0.72)	0.18c (0.09)	-

SEQ ID NO: 13

42

TABLE 2-continued

Absolute quantification of the CYP82E4v2 and CYP82E5v2
cDNA derived from the green and cured leaves of N. tomentosiformis
and different tobacco genotypes by quantitative real-time
polymerase chain reaction analysis

	CYP82E4v2		CYP82E5v2	
Sample	Green	Cured Pg	Green	Cured
DH98-325-5	0.0004a (0.0001)	0.007b 0.003)	0.08c	0.26c (0.17)
DH98-325-6	0.004b (0.001)	26.09e (3.50)	0.137c (0.003)	0.311c (0.109)
SC58NC	0.0006a (0.0001)	0.005b (0.003)	0.070c (0.012)	0.067c (0.005)
SC58C	0.0006a	1.81d	0.11c	0.083c

*Means are for 1 leaf of 3 independent plants, 3 cDNA measurements per sample (n = 9), except SCS8C where one plant was used (n = 3). *Numbers followed by different letters are significantly different according to Fisher's protected LSD (0.05).

Low levels of NtabCYP82E5v2 express in the green or yellow leaves of all tobacco cultivars (see, Table 2). The expression of NtabCYP82E5v2 significantly increased in the cured versus green leaves of converter and nonconverter Burley cultivars but remained unchanged in Flue-Cured tobacco. In the green leaves of N. tomentosiformis, expression of NtomCYP82E5 was higher than in tobacco, but no difference was noted between the cured leaves of the two species (Table 2). To compare the expression levels of CYP82E5v2 with that of CYP82E4v2, a previously characterized nicotine demethylase gene whose transcription was shown to be sharply upregulated in senescing leaves of N. tomentosiformis and converter tobacco (Gavilano et al. (2007) J. Biol. Chem. 282: 249-256), the transcript accumulation of CYP82E4v2 was compared in the same extracts used for quantifying the CYP82E5v2 mRNA. Transcription of NtabCYP82E5v2 is significantly higher than that of NtabCYP82E4v2 in the green leaves of all tobacco cultivars, N. tomentosiformis and in the cured leaves of nonconverter tobacco plants. However, the trend is reversed in the cured leaves of converter tobacco and N. tomentosiformis (Table 2). Of the two nicotine demethylase genes characterized thus far CYP82E4v2 is the dominant factor in senescence-inducible nornicotine production in N. tomentosiformis and tobacco. In addition, the activity of NtabCYP82E5v2 is the key determinant of nicotine conversion in the green leaves of tobacco.

EXAMPLE 2

Sequence Identity to Cytochrome P450 Gene Family Members

Even though CYP82E5v2 and CYP82E4v2 are both nicotine demethylases, they have less sequence homology to each other than CYP82E5v2 does to CYP82E3, for example.

TABLE 3

Amino acid sequence identity shared between the CYP82E4v2 nicotine demethylase enzyme and the other CYP82E2 proteins that have been assayed for nicotine demethylase activity.							
	CYP82E4v6	CYP82E4v12	58-166	CYP82E3	CYP82E2v1	CYP82E2v2	CYP82E5v2
YP82E4v2	99.6%	99.4%	94.8%	94.2%	92.6%	92.6%	91.1%

CYP82E5v2 and CYP82E4v2 are the only proteins represented that have nicotine demethylase activity.

30

Amino acid sequence identity shared between the CYP82E4v2 nicotine demethylase enzyme and other CYP82E2 proteins.						
CYP82E4v2 CYP82E3 CYP82E2						
CYP82E5v2 SEQ ID NO: 2	91.1%	91.9%	91.3%			

CYP82E5v2 and CYP82E4v2 are the only proteins represented that have nicotine demethylase activity

ТΛ	DT	\mathbf{D}	5	
$\mathbf{I}A$	DL	'nD.	2	

44	
----	--

location using the complementary primers, combined with visual inspection of the corresponding chromatograms distinguishes plants that are truly heterozygous for a mutation in CYP82E4v2, and differentiate these true heterozygotes from artifactual sequence anomalies.

Mutations in CYP82E4v2

High-throughput sequence analysis of 672 independent M₁ plants resulted in the identification of eleven individuals possessing point mutations within the CYP82E4v2 gene (see, Table 6). Six of the eleven plants identified carrying CYP82E4v2 mutations are homozygous for the mutant allele

Expression profiles for the CYP82E2 gene family members that have been assayed for nicotine demethylase activity.							
CYP82E4v2	CYP82E4v6	CYP82E4v12	58-166	CYP82E3	CYP82E2v1	CYP82E2v2	CYP82E5v2
Extremely Low in Green Leaf; Extremely High in Ethylene- Induced Senescent Leaf	Low in Green Leaf; High in Ethylene- Induced Senescent Leaf	Not Reported	Extremely Low In Both Green and Ethylene- Induced Senescent Leaves	Low In Both Green and Ethylene- Induced Senescent Leaves	Extremely Low in Green Leaf; Extremely High in Ethylene- Induced Senescent Leaf	Not Reported	Low in Green Leaf; Very Low in Ethylene- Induced Senescent Leaf

CYP82E5v2 and CYP82E4v2 are the only proteins represented that were positive for nicotine demethylase activity.

EXAMPLE 3

Mutations IN CYP82E4v2 AND CYP82E5v2

Obtaining EMS-Derived Variants of CYP82E4v2

To facilitate the introduction of random mutations into the tobacco genome, thousands of seeds of the strong Converter Burley line DH98-325-6 were treated with the chemical mutagene ethyl methane sulphonate (EMS). Approximately 4000 mutagenized plants (M_0 generation) were grown in the field and allowed to self-pollinate. Several capsules from each individual plant were combined to create discrete M1 seed 40 populations, each corresponding to an individual M_o plant. Genomic DNAs were isolated from young leaf tissue of a single greenhouse-grown M₁ plant from each M₁ seed pool. To identify plants carrying mutations in the CYP82E4v2 gene, specific primers were designed to independently 45 amplify each of the gene's two exons. DNA sequence information was obtained using 96-well high-throughput sequence analysis of the amplification products. The complete analysis of 96 individual plants involved 4 separate 96-well sequencing reactions: forward and reverse sequenc- 50 ing of the exon 1 amplification product, and forward and reverse sequencing of the exon 2 PCR product.

Mutations were identified by conducting a multi-sequence alignment of all 96 sequences for a given run and looking for deviations from the wild-type sequence. Any given mutation 55 changes a tryptophan codon (TGG) into a premature stop that occurred in a parental Mo plant would be expected to be segregating in 1:2:1 ratio in the M₁ generation for the mutant versus wild-type alleles. M₁ plants that are homozygous for an EMS-induced mutation are readily recognized as polymorphisms deviating from the wild type sequence, verified in 60 both directions. M1 plants that are heterozygous for a mutation in CYP82E4v2 would be expected to have the mutant and wild type alleles each representing 50% of the amplification product. Upon sequencing, the location of such a mutation would be expected to be annotated as an "N", since the fluo- 65 rescence reading at that site would be a mixture of two alternative bases. The appearance of an "N" at the same nucleotide

(see, Table 6). All eleven mutations result in changes in the predicted amino acid sequence of the encoded protein.

TABLE 6

	M ₁ L Posses	M_1 Lines of EMS Treated DH98-325-6 Plants Possessing Mutations in the CYP82E4v2 Gene.				
5	Plant Line	Nucleotide Mutation*	Amino Acid Change**	Zygosity of Mutant Allele		
)	DH98-325-6 #101 DH98-325-6 #121 DH98-325-6 #164 DH98-325-6 #371 DH98-325-6 #377 DH98-325-6 #569 DH98-325-6 #506	C1372T G1092T C113T G601A G506A G1375A G886A	P458S K364N P38L E201K R169Q G459R E296K	Homozygous Homozygous Heterozygous Heterozygous Heterozygous Heterozygous		
5	DH98-325-6 #453 DH98-325-6 #453 DH98-325-6 #775 DH98-325-6 #610 DH98-325-6 #761	G880A C1280T G986A G1126A G511A	E290K T427I W329Stop V376M D171N	Heterozygous Heterozygous Homozygous Homozygous Homozygous		

*Nucleotide location is in reference to the ATG initiator codon of the CYP82E4v2 cDNA (see SEQ ID NO: 50). Nucleotide to the left of the number represents the wild type residue, whereas the nucleotide on the right indicates the mutant residue (e.g. C1372T means that the C residue normally located at position 1372 was mutated to a T residue). **Amino acid location is in reference to the initiator methionine of the CYP82E4v2 polypeptide (see SEQ ID NO: 13). Amino acid to the left of the number represents the wild type residue, whereas the amino acid on the right indicates the residue produced by the mutant allele (e.g. P458S means that the Proline residue normally located at position 458 of the protein sequence has been changed to a Serine). "Stop" indicates the introduction of a memature stop codon. premature stop codon.

Plant #775 possesses a mutation at codon 329 which codon (TAG). This mutation renders the encoded protein completely nonfunctional given that the essential heme-binding domain and other highly conserved regions of the enzyme are located downstream of codon 329. Other plants contain mutations affecting highly conserved motifs shared by the vast majority of P450 enzymes include: plant #101 (where a conserved Pro residue immediately adjacent to the hemebinding Cys amino acid is change to a Ser); plant #164 (Pro residue within the highly conserved Pro-rich "hinge" region of the protein changed to a Leu); and plant #569 (Gly residue that's part of the very highly conserved heme-binding motif changed to an Arg).

Alkaloid Analysis of CYP82E4v2 Variants

An alkaloid analysis was conducted on cured leaves of variant individuals to assess the effect of the mutation on the metabolic conversion of nicotine to nornicotine. As shown in Table 7, three individuals (#101, #121 and #775) show conversion rates similar to nonconverter control plants grown from seed lots that had been screened to eliminate converter individuals (TN90 LC plants). The variant CYP82E4v2 enzymes produced in these plants are completely (or nearly completely) nonfunctional. Two plants (#164 and #761) dis- 10 play high conversion rates and have no detrimental effect on enzyme activity. This result for plant #164 is surprising given that it contains a Pro to Leu mutation within one of the residues defining the highly conserved Pro-rich hinge motif. The Val to Met modification at residue 376 partially inhibits 15 enzyme function as the homozygous mutant plant (#610) shows an intermediate conversion phenotype (41.5%).

TABLE 7

Alkaloid Content as Percent Dry Weight of M_1 Plants Homozygous for Mutations in the CYP82E4v2 Gene						
Plant	Nicotine	Nornicotine	Anabasine	Anatabine	% Conversion*	
DH98-325-6 #101	2.17	0.0376	0.00514	0.0384	1.7%	
DH98-325-6 #121	2.5	0.0333	0.00557	0.043	1.3%	
DH98-325-6 #164	0.216	0.84	0.00499	0.0532	79.5%	
DH98-325-6 #610	0.214	0.152	0.00429	0.0121	41.5%	
DH98-325-6 #761	0.214	0.999	0.00429	0.0426	82.4%	
DH98-325-6 #775	0.505	0.0159	0.00417	0.0136	3.1%	
SC58 CtCt (Control) [†]	0.206	0.565	0.00412	0.0197	73.3%	
TN90 LC (Control)	3.11	0.0647	0.00901	0.0487	2.0%	
TN90 LC (Control)	2.46	0.0521	0.00846	0.0417	2.1%	

*[% nornicotine/(% nicotine + % nornicotine)] × 100

A converter plant of flue-cured genotype SC58 is used as a strong converter control; two plants of genotype

TN90 selected from lots screened for low converters (LC) are used as nonconverter controls.

The substitution of the Lys residue at position 364 to an Asn of plant #121 within this plant does not occur in any motif conserved among P450 enzymes, nor within any region predicted by Xu et al. (Physiologia Plantarum 129: 307-319, 40 2007) to be a substrate recognition site. A BLAST alignment against protein sequences deposited in GenBank reveals several P450s that possess an Asn residue at the analagous position (e.g. the CYP82A1 gene of Pea-accession number Q43068). The variant CYP82E4v2 gene in plant #121 is completely inactive. Another anomaly with plant #121 lies in the nature of EMS-induced mutations. EMS is known to induce mutations through the alkylation of G residues, resulting in G to A or C to T transition mutations (Anderson, Methods Cell Biol. 48:31-58, 1995). Very rarely does EMS lead to a G to T transversion mutation such as that found in plant #121 (see, Table 3). All other mutations described for either the CYP82E4v2 or CYP82E5v2 genes are the expected G to A or C to T transition mutations. 55

Mutations in CYP82E5v2

A similar high-throughput sequencing strategy was used to identify mutations in CYP82E5v2, the other confirmed nicotine N-demethylase gene that is found within the tobacco genome. A screen of 768 M_1 plants revealed 11 individuals possessing mutations in CYP82E5v2 (see, Table 8). Three plants (#744, #561 and #340) contain silent nucleotide substitutions that did not alter the predicted protein sequence. Of the eight mutations that lead to changes in the protein $_{65}$ sequence, two are of particular note. Plant #1013 contains a mutation that would lead to a truncated product that should

render the product completely nonfunctional. The G to A mutation at position 1266 changes a Trp codon (TGG) into a premature stop codon (TGA). The predicted CYP82E5v2 protein produced from this truncated reading frame lacks the final 96 amino acids of the enzyme, a region that includes the essential heme-binding domain. Without being limited by mechanism, this truncated protein is probably incapable of catalyzing the oxidative elimination of the N' methyl group of nicotine to form nornicotine. Whatever the mechanism, the #1013 plant has a nonfunctional CYP82E5v2 allele. Plant #680 possesses a mutation that changes a Pro residue at position 449 to a Leu. This Pro is well conserved in plant P450s and lies immediately adjacent to the F-x-x-G-x-R-x-C-x-G motif that defines the heme binding region. The mutation found in plant #680 has a negative impact on enzyme function.

TABLE 8 M1 Lines of EMS Treated DH98-325-6 Plants

Tossessing wutations in the CTT 82E5v2 Gene					
Plant Line	Nucleotide Mutation*	Amino Acid Change**	Zygosity of Mutant Allele		
DH98-325-6 #198	C703T	P235S	Heterozygous		
DH98-325-6 #680	C1346T	P449L	Homozygous		
DH98-325-6 #744	G558A	None	Homozygous		
DH98-325-6 #163	C521T	S174L	Heterozygous		
DH98-325-6 #561	G1458A	None	Heterozygous		
DH98-325-6 #154	C1229T	A410V	Heterozygous		
DH98-325-6 #340	G1248A	None	Heterozygous		
DH98-325-6 #780	G672A	M224I	Heterozygous		
DH98-325-6 #799	C215T	P72L	Heterozygous		
DH98-325-6 #540	C427T	L143F	Heterozygous		
DH98-325-6 #1013	G1266A	W422Stop	Homozygous		

*Nucleotide location is in reference to the ATG initiator codon of the CYP82E5v2 cDNA (see SEQ ID NO: 1). Nucleotide to the left of the number represents the wild type residue, whereas the nucleotide on the right indicates the mutant residue (e.g. C703T means that the C residue normally located at position 703 was mutated to a T residue).

C restude infinite product a position 705 was initiated to a restude).
C restude infinite product a position for was initiated and a restude).
Polypeptide (see SEQ ID NO: 2). Amino acid to the left of the number represents the wild type residue, whereas the amino acid on the right indicates the residue produced by the mutant allele (e.g. P235S means that the Proline residue normally located at position 235 of the protein sequence has been changed to a Serine). "Stop" indicates the introduction of a premature stop eodon. "None" indicates mutations resulting in silent substitutions that did not alter the predicted amino acid sequence.

Plants with Mutations in CYP82E4v2 and CYP82E5v2

Tobacco plant lines are generated that show uniformity and stability within the limits of environmental influence for reduced conversion of nicotine to nornicotine at levels of about 0.5% when a plant with a mutation that inhibits nicotine demethylase activity of CYP82E4v2 is crossed with a plant having a mutation that inhibits nicotine demethylase activity of CYP82E5v2.

An illustrative cross is between a CYP82E4v2 mutant plant ⁵ (of Table 9) and a CYP82E5v2 mutant plant (of Table 9).

TABLE 9

Illustrative Crosses of DH98-325-6 Plants Possessing Mutations in the CYP82E5v2 and CYP82E4v2 Genes			
	CYP82E5v2		
	DH98-325-6 #198		
	DH98-325-6 #680	15	
	DH98-325-6 #163		
	DH98-325-6 #154		
	DH98-325-6 #780		
	DH98-325-6 #799		
	DH98-325-6 #540		
	DH98-325-6 #1013	20	

TABLE 9-continued					
Illustrative Crosses of DH98-325-6 Plants Possessing Mutations in the CYP82E5v2 and CYP82E4v2 Genes					
	CYP82E4v2				
	DH98-325-6 #101 DH98-325-6 #121 DH98-325-6 #775				

All publications and patent applications mentioned in the specification are indicative of the level of those skilled in the art to which this invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain ²⁰ changes and modifications may be practiced within the scope of the appended claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 50

- <210> SEQ ID NO 1
- <211> LENGTH: 1554

<212> TYPE: DNA <213> ORGANISM: Nicotiana tabacum

<400> SEQUENCE: 1

atggtttctc	ccgtagaagc	cattgtagga	ctagtaaccc	ttacacttct	cttctacttc	60
ctatggccca	aaaaatttca	aataccttca	aaaccattac	caccgaaaat	tcccggaggg	120
tggccggtaa	tcggccatct	tttctacttc	gatgatgacg	gcgacgaccg	tccattagct	180
cgaaaactcg	gagacttagc	tgacaaatac	ggcccggttt	tcactttccg	gctaggcctt	240
ccgcttgtgt	tagttgtaag	cagttacgaa	gctgtaaaag	actgcttctc	tacaaatgac	300
gccattttct	ccaatcgtcc	agcttttctt	tacggtgaat	accttggcta	cagtaatgcc	360
atgctatttt	tgacaaaata	cggaccttat	tggcgaaaaa	atagaaaatt	agtcattcag	420
gaagttctct	ctgctagtcg	tctcgaaaaa	ttgaagcacg	tgagatttgg	taaaattcaa	480
acgagcatta	agagtttata	cactcgaatt	gatggaaatt	cgagtacgat	aaatctaact	540
gattggttag	aagaattgaa	ttttggtctg	atcgtgaaaa	tgatcgctgg	gaaaaattat	600
gaatccggta	aaggagatga	acaagtggag	agatttagga	aagcgtttaa	ggattttata	660
attttatcaa	tggagtttgt	gttatgggat	gcttttccaa	ttccattgtt	caaatgggtg	720
gattttcaag	gccatgttaa	ggccatgaaa	aggacattta	aggatataga	ttctgttttt	780
cagaattggt	tagaggaaca	tgtcaagaaa	agagaaaaaa	tggaggttaa	tgcacaaggg	840
aatgaacaag	atttcattga	tgtggtgctt	tcaaaaatga	gtaatgaata	tcttgatgaa	900
ggttactctc	gtgatactgt	cataaaagca	acagtgttta	gtttggtctt	ggatgctgcg	960
gacacagttg	ctcttcacat	gaattgggga	atggcattac	tgataaacaa	tcaacatgcc	1020
ttgaagaaag	cacaagaaga	gatcgataaa	aaagttggta	aggaaagatg	ggtagaagag	1080
agtgatatta	aggatttggt	ctacctccaa	gctattgtta	aagaagtgtt	acgattatat	1140
ccaccaggac	ctttattagt	acctcatgaa	aatgtagagg	attgtgttgt	tagtggatat	1200
cacattccta	aagggactag	actattcgcg	aacgttatga	aattgcagcg	cgatcctaaa	1260
ctctggtcaa atcctgataa gtttgatcca gagagattct tcgctgatga tattgactac	1320					
--	------					
cgtggtcagc actatgagtt tatcccattt ggttctggaa gacgatcttg tccggggatg	1380					
acttatgcat tacaagcgga acacctaaca atagcacatt tgatccaggg tttcaattac	1440					
aaaactccaa atgacgagcc cttggatatg aaggaaggtg caggattaac tatacgtaaa	1500					
gtaaateetg tagaagtgae aattaegget egeetggeae etgagettta ttaa	1554					
<210> SEQ ID NO 2 <211> LENGTH: 517 <212> TYPE: PRT <213> ORGANISM: Nicotiana tabacum						
<400> SEQUENCE: 2						
Met Val Ser Pro Val Glu Ala Ile Val Gly Leu Val Thr Leu151015						
Leu Phe Tyr Phe Leu Trp Pro Lys Lys Phe Gln Ile Pro Ser Lys Pro 20 25 30						
Leu Pro Pro Lys Ile Pro Gly Gly Trp Pro Val Ile Gly His Leu Phe 35 40 45						
Tyr Phe Asp Asp Asp Gly Asp Asp Arg Pro Leu Ala Arg Lys Leu Gly 50 55 60						
Asp Leu Ala Asp Lys Tyr Gly Pro Val Phe Thr Phe Arg Leu Gly Leu 65 70 75 80						
Pro Leu Val Leu Val Val Ser Ser Tyr Glu Ala Val Lys Asp Cys Phe 85 90 95						
Ser Thr Asn Asp Ala Ile Phe Ser Asn Arg Pro Ala Phe Leu Tyr Gly						
Glu Tyr Leu Gly Tyr Ser Asn Ala Met Leu Phe Leu Thr Lys Tyr Gly						
Pro Tyr Trp Arg Lys Asn Arg Lys Leu Val Ile Gln Glu Val Leu Ser						
Ala Ser Arg Leu Glu Lys Leu Lys His Val Arg Phe Gly Lys Ile Gln 145 150 155 160						
Thr Ser Ile Lys Ser Leu Tyr Thr Arg Ile Asp Gly Asn Ser Ser Thr 165 170 175						
Ile Asn Leu Thr Asp Trp Leu Glu Glu Leu Asn Phe Gly Leu Ile Val						
Lys Met Ile Ala Gly Lys Asn Tyr Glu Ser Gly Lys Gly Asp Glu Gln						
Val Glu Arg Phe Arg Lys Ala Phe Lys Asp Phe Ile Ile Leu Ser Met						
Glu Phe Val Leu Trp Asp Ala Phe Pro Ile Pro Leu Phe Lys Trp Val						
Asp Phe Gln Gly His Val Lys Ala Met Lys Arg Thr Phe Lys Asp Ile						
Asp Ser Val Phe Gln Asn Trp Leu Glu Glu His Val Lys Lys Arg Glu						
260 265 270 Lys Met Glu Val Asn Ala Gln Gly Asn Glu Gln Asp Phe Ile Asp Val						
275 280 285 Val Leu Ser Lys Met Ser Asn Glu Tvr Leu Asn Glu Glv Tvr Ser Arg						
290 295 300						
ASP INF VAL HE LYS ALA TNE VAL PNE SEE LEU VAL LEU ASP ALA ALA 305 310 315 320						
Asp Thr Val Ala Leu His Met Asn Trp Gly Met Ala Leu Leu Ile Asn						

325 330	335	
Asn Gln His Ala Leu Lys Lys Ala Gln Glu Glu I 340 345	le Asp Lys Lys Val 350	
Gly Lys Glu Arg Trp Val Glu Glu Ser Asp Ile L 355 360	ys Asp Leu Val Tyr 365	
Leu Gln Ala Ile Val Lys Glu Val Leu Arg Leu T 370 375 3	'yr Pro Pro Gly Pro 80	
Leu Leu Val Pro His Glu Asn Val Glu Asp Cys V 385 390 395	al Val Ser Gly Tyr 400	
His Ile Pro Lys Gly Thr Arg Leu Phe Ala Asn V 405 410	al Met Lys Leu Gln 415	
Arg Asp Pro Lys Leu Trp Ser Asn Pro Asp Lys P. 420 425	he Asp Pro Glu Arg 430	
Phe Phe Ala Asp Asp Ile Asp Tyr Arg Gly Gln H 435 440	lis Tyr Glu Phe Ile 445	
Pro Phe Gly Ser Gly Arg Arg Ser Cys Pro Gly M 450 455 4	let Thr Tyr Ala Leu 60	
Gln Ala Glu His Leu Thr Ile Ala His Leu Ile G 465 470 475	In Gly Phe Asn Tyr 480	
Lys Thr Pro Asn Asp Glu Pro Leu Asp Met Lys G 485 490	lu Gly Ala Gly Leu 495	
Thr Ile Arg Lys Val Asn Pro Val Glu Val Thr I 500 505	le Thr Ala Arg Leu	
Ala Pro Glu Leu Tyr	510	
<210> SEQ ID NO 3 <211> LENGTH: 1054 <212> TYPE: DNA <213> ORGANISM: Nicotiana tabacum		
<400> SEQUENCE: 3		
gtaagttcat tttcattttt cattattcag tctgattttg a	ggaatagac aggttaataa	60
taatttaagt aattagatta tctaaatact aaggatgatt a	tatatagta aaaatgtaga	120
atgataaatg gaaaaaagat gagaattett tgtgeetega e	taatctata tatctttggg	180
agttaaaagt getteaceaa aggggaettt teeteatage t	caagttaga agtttgatta	240
tagatgaaag agtatttatc acttcacgaa ctctgatgat a	aaagtaaat gagatataac	300
cagttataat tgatagaata aaacttcatt actcccattg a	gcataaaaa aaaaagtaaa	360
agggacttct tctcttttt ttagggagaa attctttaat t	gtttgttaa atatagattc	420
atgtttttt tttcttctat ttctaataat aatggttctt g	aatcaggtc gttgactttg	480
tagcagcaat atagtcaaag ctaatatcca tgttatttgg t	tttcgaaca agttatactg	540
aaattatata tacgggtatt aaataataac attattattt a	taggatata cttttttat	600
tgggtaaata ttacaacaac aacaactgac tcagtgaaat t	ttactagtg gggtatgggg	660
agggtagtgt gtatgcagac cttaccccta ccccgaagga g	tagagggat tgtttccgaa	720
agaccctcgg ctcaagaaaa caaaaagaga caatatcagt a	ccaccacag atcatattat	780
taggtaaatg ttattttatt gaattaaaga tgaaatatac a	ggtaaggta taaaacgtgt	840
atttgatttt acactagata aatttgacct cgtacatctc t	aagagaaag ctgaaataaa	900
tgaattttag atttaaaaaa aaaattcatt agtataatga g	atgtgcata cttgacaatt	960
actatactaa atagaacaag gttcggcaga tagtgacact a	acctacttt tgcattgaat	1020

-continued

tatccttttt aattttattc taatttgtct acag

<210> SEQ : <211> LENG	ID NO 4 TH: 2608					
<212> TYPE <213> ORGAI	: DNA NISM: Nicot:	lana tabacum	n			
<400> SEQUI	ENCE: 4					
atggtttctc	ccgtagaagc	cattgtagga	ctagtaaccc	ttacacttct	cttctacttc	60
ctatggccca	aaaaatttca	aataccttca	aaaccattac	caccgaaaat	tcccggaggg	120
tggccggtaa	tcggccatct	tttctacttc	gatgatgacg	gcgacgaccg	tccattagct	180
cgaaaactcg	gagacttagc	tgacaaatac	ggcccggttt	tcactttccg	gctaggcctt	240
ccgcttgtgt	tagttgtaag	cagttacgaa	gctgtaaaag	actgcttctc	tacaaatgac	300
gccattttct	ccaatcgtcc	agcttttctt	tacggtgaat	accttggcta	cagtaatgcc	360
atgctatttt	tgacaaaata	cggaccttat	tggcgaaaaa	atagaaaatt	agtcattcag	420
gaagttetet	ctgctagtcg	tctcgaaaaa	ttgaagcacg	tgagatttgg	taaaattcaa	480
acgagcatta	agagtttata	cactcgaatt	gatggaaatt	cgagtacgat	aaatctaact	540
gattggttag	aagaattgaa	ttttggtctg	atcgtgaaaa	tgatcgctgg	gaaaaattat	600
gaatccggta	aaggagatga	acaagtggag	agatttagga	aagcgtttaa	ggattttata	660
attttatcaa	tggagtttgt	gttatgggat	gcttttccaa	ttccattgtt	caaatgggtg	720
gattttcaag	gccatgttaa	ggccatgaaa	aggacattta	aggatataga	ttctgttttt	780
cagaattggt	tagaggaaca	tgtcaagaaa	agagaaaaaa	tggaggttaa	tgcacaaggg	840
aatgaacaag	atttcattga	tgtggtgctt	tcaaaaatga	gtaatgaata	tcttgatgaa	900
ggttactctc	gtgatactgt	cataaaagca	acagtgtttg	taagttcatt	ttcatttttc	960
attattcagt	ctgattttga	ggaatagaca	ggttaataat	aatttaagta	attagattat	1020
ctaaatacta	aggatgatta	tatatagtaa	aaatgtagaa	tgataaatgg	aaaaaagatg	1080
agaattcttt	gtgcctcgac	taatctatat	atctttggga	gttaaaagtg	cttcaccaaa	1140
ggggactttt	cctcatagct	caagttagaa	gtttgattat	agatgaaaga	gtatttatca	1200
cttcacgaac	tctgatgata	aaagtaaatg	agatataacc	agttataatt	gatagaataa	1260
aacttcatta	ctcccattga	gcataaaaaa	aaaagtaaaa	gggacttctt	ctctttttt	1320
tagggagaaa	ttctttaatt	gtttgttaaa	tatagattca	tgttttttt	ttcttctatt	1380
tctaataata	atggttcttg	aatcaggtcg	ttgactttgt	agcagcaata	tagtcaaagc	1440
taatatccat	gttatttggt	tttcgaacaa	gttatactga	aattatatat	acgggtatta	1500
aataataaca	ttattattta	taggatatac	ttttttatt	gggtaaatat	tacaacaaca	1560
acaactgact	cagtgaaatt	ttactagtgg	ggtatgggga	gggtagtgtg	tatgcagacc	1620
ttacccctac	cccgaaggag	tagagggatt	gtttccgaaa	gaccctcggc	tcaagaaaac	1680
aaaaagagac	aatatcagta	ccaccacaga	tcatattatt	aggtaaatgt	tattttattg	1740
aattaaagat	gaaatataca	ggtaaggtat	aaaacgtgta	tttgatttta	cactagataa	1800
atttgacctc	gtacatctct	aagagaaagc	tgaaataaat	gaattttaga	tttaaaaaaa	1860
aaattcatta	gtataatgag	atgtgcatac	ttgacaatta	ctatactaaa	tagaacaagg	1920
ttcggcagat	agtgacacta	acctactttt	gcattgaatt	atccttttta	attttattct	1980
aatttgtcta	cagagtttgg	tcttggatgc	tgcggacaca	gttgctcttc	acatgaattg	2040

US 9,228,194 B2

56

2100

55

gggaatggca ttactgataa acaatcaaca tgccttgaag aaagcacaag aagagatcga

taaa	aaad	gtt d	ggta	aggaa	aa a	atqq	gtaqa	a aqa	agaqt	tgat	att	aaqa	att t	gqt	ctacct	2160	
ccaa	iqcta	att d	qtta	aagaa	aq to	qtta	cqatt	t ata	atcca	acca	qqa	cetti	tat t	taqta	acctca	a 2220	
tgaa	aato	yta d	gagga	attgi	tg ti	- tgtta	agtgo	g ata	atca	catt	cct	aaag	gga (taga	actatt	2280	
cgcc	jaaco	ytt a	atga	aatto	- gc a	gegeo	gatco	c taa	aacto	ctgq	tca	aatc	ctg a	ataad	gtttqa	a 2340	
tcca	igaga	iga 1	teti	teget	tg a'	 tgata	attga	a cta	accgi	tggt	cag	cact	- atg a	agtti	atccc	2400	
attt	ggtt	ct q	ggaa	gacga	at c	ttgt	ccggg	g gat	zgact	ttat	gca	ttac	aag o	cggaa	acacct	2460	
aaca	ataç	gca (catt	tgat	cc a	gggti	tcaa	a tta	acaaa	aact	cca	aatg	acg a	ageed	ettgga	a 2520	
tate	jaago	jaa q	ggtg	cagga	at t	aacta	ataco	g taa	aagta	aaat	cct	gtag	aag t	gaca	aattac	2580	
ggct	cgco	tg o	gcac	ctga	gc ti	ttati	zaa									2608	
<210 <211 <212 <213)> SI .> LH 2> TY 3> OH	EQ II ENGTH (PE : RGAN]	D NO H: 5: PRT ISM:	5 17 Nico	otia	na ta	abacı	ım									
<400)> SI	EQUEI	NCE :	5													
Met 1	Val	Ser	Pro	Val 5	Glu	Ala	Ile	Val	Gly 10	Leu	Val	Thr	Leu	Thr 15	Leu		
Leu	Phe	Tyr	Phe 20	Leu	Trp	Pro	Гла	Lув 25	Phe	Gln	Ile	Pro	Ser 30	Гла	Pro		
Leu	Pro	Pro 35	ГЛа	Ile	Pro	Gly	Gly 40	Trp	Pro	Val	Ile	Gly 45	His	Leu	Phe		
Tyr	Phe 50	Asp	Asp	Asp	Gly	Asp 55	Asp	Arg	Pro	Leu	Ala 60	Arg	Lys	Leu	Gly		
Asp 65	Leu	Ala	Asp	Lys	Tyr 70	Gly	Pro	Val	Phe	Thr 75	Phe	Arg	Leu	Gly	Leu 80		
Pro	Leu	Val	Leu	Val 85	Val	Ser	Ser	Tyr	Glu 90	Ala	Val	ГЛа	Asp	Суз 95	Phe		
Ser	Thr	Asn	Asp 100	Ala	Ile	Phe	Ser	Asn 105	Arg	Pro	Ala	Phe	Leu 110	Tyr	Gly		
Glu	Tyr	Leu 115	Gly	Tyr	Ser	Asn	Ala 120	Met	Leu	Phe	Leu	Thr 125	Lys	Tyr	Gly		
Pro	Tyr 130	Trp	Arg	Гла	Asn	Arg 135	Lys	Leu	Val	Ile	Gln 140	Glu	Val	Leu	Ser		
Ala 145	Ser	Arg	Leu	Glu	Lys 150	Leu	Lys	His	Val	Arg 155	Phe	Gly	Lys	Ile	Gln 160		
Thr	Ser	Ile	Гла	Ser 165	Leu	Tyr	Thr	Arg	Ile 170	Asp	Gly	Asn	Ser	Ser 175	Thr		
Ile	Asn	Leu	Thr 180	Asp	Trp	Leu	Glu	Glu 185	Leu	Asn	Phe	Gly	Leu 190	Ile	Val		
Lys	Met	Ile 195	Ala	Gly	ГЛа	Asn	Tyr 200	Glu	Ser	Gly	ГЛЗ	Gly 205	Asp	Glu	Gln		
Val	Glu 210	Arg	Phe	Arg	ГЛа	Ala 215	Phe	Гла	Asp	Phe	Ile 220	Ile	Leu	Ser	Met		
Glu 225	Phe	Val	Leu	Trp	Asp 230	Ala	Phe	Pro	Ile	Ser 235	Leu	Phe	Гла	Trp	Val 240		
Aap	Phe	Gln	Gly	His 245	Val	ГЛа	Ala	Met	Lys 250	Arg	Thr	Phe	Lys	Asp 255	Ile		
Asp	Ser	Val	Phe	Gln	Asn	Trp	Leu	Glu	Glu	His	Val	ГЛа	Lys	Arg	Glu		

				_			_					_			_
Lys	Met	Glu 275	Val	Asn	Ala	Gln	Gly 280	Asn	Glu	Gln	Asp	Phe 285	Ile	Asp	Val
Val	Leu 290	Ser	Lys	Met	Ser	Asn 295	Glu	Tyr	Leu	Asp	Glu 300	Gly	Tyr	Ser	Arg
Asp 305	Thr	Val	Ile	ГЛа	Ala 310	Thr	Val	Phe	Ser	Leu 315	Val	Leu	Asp	Ala	Ala 320
Asp	Thr	Val	Ala	Leu 325	His	Met	Asn	Trp	Gly 330	Met	Ala	Leu	Leu	Ile 335	Asn
Asn	Gln	His	Ala 340	Leu	ГЛа	ГЛа	Ala	Gln 345	Glu	Glu	Ile	Asp	Lys 350	Lys	Val
Gly	Lys	Glu 355	Arg	Trp	Val	Glu	Glu 360	Ser	Asp	Ile	Lys	Asp 365	Leu	Val	Tyr
Leu	Gln 370	Ala	Ile	Val	Lys	Glu 375	Val	Leu	Arg	Leu	Tyr 380	Pro	Pro	Gly	Pro
Leu 385	Leu	Val	Pro	His	Glu 390	Asn	Val	Glu	Asp	Cys 395	Val	Val	Ser	Gly	Tyr 400
His	Ile	Pro	ГЛа	Gly 405	Thr	Arg	Leu	Phe	Ala 410	Asn	Val	Met	Lys	Leu 415	Gln
Arg	Asp	Pro	Lys 420	Leu	Trp	Ser	Asn	Pro 425	Asp	Lys	Phe	Asp	Pro 430	Glu	Arg
Phe	Phe	Ala 435	Asp	Asp	Ile	Aap	Tyr 440	Arg	Gly	Gln	His	Tyr 445	Glu	Phe	Ile
Pro	Phe 450	Gly	Ser	Gly	Arg	Arg 455	Ser	Сүз	Pro	Gly	Met 460	Thr	Tyr	Ala	Leu
Gln 465	Ala	Glu	His	Leu	Thr 470	Ile	Ala	His	Leu	Ile 475	Gln	Gly	Phe	Asn	Tyr 480
Гуз	Thr	Pro	Asn	Asp 485	Glu	Pro	Leu	Asp	Met 490	Lys	Glu	Gly	Ala	Gly 495	Leu
Thr	Ile	Arg	Lys 500	Val	Asn	Pro	Val	Glu 505	Val	Thr	Ile	Thr	Ala 510	Arg	Leu
Ala	Pro	Glu 515	Leu	Tyr				505					510		
.01/		70 11		c											
<210 <211 <212)> SH L> LH 2> TY	EQ II ENGTH PE:) NO H: 5: PRT	6 17											
<213	8> 0F	RGAN:	ISM:	Nico	otia	na ta	abacı	m							
<4UU Mot	val	Cor.	Pro	6 Val	Glu	۵la	Tla	Val	Glv	Leu	Val	Thr	Ι.φ11	Thr	Leu
1 Lou	Dho	Der	Dho	5	Tre	Dro	Inc	Ina	10 Dho	Cln	Tlo	Dro	Cor	15	Dro
Leu	Pne	ıyr	20	Leu	Trp	Pro	гуа	цув 25	Pne	GIN	IIe	Pro	30	гуа	Pro
Leu	Pro	Pro 35	ГЛа	Ile	Pro	Gly	Gly 40	Trp	Pro	Val	Ile	Gly 45	His	Leu	Phe
Tyr	Phe 50	Asp	Asp	Asp	Gly	Asp 55	Asp	Arg	Pro	Leu	Ala 60	Arg	Lys	Leu	Gly
Asp 65	Leu	Ala	Asp	ГЛа	Tyr 70	Gly	Pro	Val	Phe	Thr 75	Phe	Arg	Leu	Gly	Leu 80
Pro	Leu	Val	Leu	Val 85	Val	Ser	Ser	Tyr	Glu 90	Ala	Val	ГЛа	Asp	Сүз 95	Phe
Ser	Thr	Asn	Asp 100	Ala	Ile	Phe	Ser	Asn 105	Arg	Pro	Ala	Phe	Leu 110	Tyr	Gly
Glu	Tyr	Leu 115	Gly	Tyr	Ser	Asn	Ala 120	Met	Leu	Phe	Leu	Thr 125	Lys	Tyr	Gly

-continued

Pro Tyr Trp Arg Lys Asn Arg Lys Leu Val Ile Gln Glu Val Leu Ser Ala Ser Arg Leu Glu Lys Leu Lys His Val Arg Phe Gly Lys Ile Gln Thr Ser Ile Lys Ser Leu Tyr Thr Arg Ile Asp Gly Asn Ser Ser Thr Ile Asn Leu Thr Asp Trp Leu Glu Glu Leu Asn Phe Gly Leu Ile Val Lys Met Ile Ala Gly Lys Asn Tyr Glu Ser Gly Lys Gly Asp Glu Gln Val Glu Arg Phe Arg Lys Ala Phe Lys Asp Phe Ile Ile Leu Ser Met 210 215 220 Glu Phe Val Leu Trp Asp Ala Phe Pro Ile Pro Leu Phe Lys Trp Val225230235240 Asp Phe Gln Gly His Val Lys Ala Met Lys Arg Thr Phe Lys Asp Ile 245 250 255 Asp Ser Val Phe Gln Asn Trp Leu Glu Glu His Val Lys Lys Arg Glu Lys Met Glu Val Asn Ala Gln Gly Asn Glu Gln Asp Phe Ile Asp Val Val Leu Ser Lys Met Ser Asn Glu Tyr Leu Asp Glu Gly Tyr Ser Arg Asp Thr Val Ile Lys Ala Thr Val Phe Ser Leu Val Leu Asp Ala Ala Asp Thr Val Ala Leu His Met Asn Trp Gly Met Ala Leu Leu Ile Asn Asn Gln His Ala Leu Lys Lys Ala Gln Glu Glu Ile Asp Lys Lys Val Gly Lys Glu Arg Trp Val Glu Glu Ser Asp Ile Lys Asp Leu Val Tyr Leu Gln Ala Ile Val Lys Glu Val Leu Arg Leu Tyr Pro Pro Gly Pro Leu Leu Val Pro His Glu Asn Val Glu Asp Cys Val Val Ser Gly Tyr His Ile Pro Lys Gly Thr Arg Leu Phe Ala Asn Val Met Lys Leu Gln Arg Asp Pro Lys Leu Trp Ser Asn Pro Asp Lys Phe Asp Pro Glu Arg Phe Phe Ala Asp Asp Ile Asp Tyr Arg Gly Gln His Tyr Glu Phe Ile 435 440 445 Leu Phe Gly Ser Gly Arg Arg Ser Cys Pro Gly Met Thr Tyr Ala Leu Gln Ala Glu His Leu Thr Ile Ala His Leu Ile Gln Gly Phe Asn Tyr Lys Thr Pro Asn Asp Glu Pro Leu Asp Met Lys Glu Gly Ala Gly Leu Thr Ile Arg Lys Val Asn Pro Val Glu Val Thr Ile Thr Ala Arg Leu Ala Pro Glu Leu Tyr

<210> SEQ ID NO 7 <211> LENGTH: 517

<212 <213	2 > T 3 > O	YPE : RGANI	PRT ISM:	Nico	otia	na ta	abacı	ım							
<400)> SI	EQUEI	NCE :	7											
Met 1	Val	Ser	Pro	Val 5	Glu	Ala	Ile	Val	Gly 10	Leu	Val	Thr	Leu	Thr 15	Leu
Leu	Phe	Tyr	Phe 20	Leu	Trp	Pro	Lys	Lуз 25	Phe	Gln	Ile	Pro	Ser 30	Lys	Pro
Leu	Pro	Pro 35	Lys	Ile	Pro	Gly	Gly 40	Trp	Pro	Val	Ile	Gly 45	His	Leu	Phe
Tyr	Phe 50	Asp	Asp	Asp	Gly	Asp 55	Asp	Arg	Pro	Leu	Ala 60	Arg	Lys	Leu	Gly
Asp 65	Leu	Ala	Asp	ГЛЗ	Tyr 70	Gly	Pro	Val	Phe	Thr 75	Phe	Arg	Leu	Gly	Leu 80
Pro	Leu	Val	Leu	Val 85	Val	Ser	Ser	Tyr	Glu 90	Ala	Val	ГЛа	Asp	Суз 95	Phe
Ser	Thr	Asn	Asp 100	Ala	Ile	Phe	Ser	Asn 105	Arg	Pro	Ala	Phe	Leu 110	Tyr	Gly
Glu	Tyr	Leu 115	Gly	Tyr	Ser	Asn	Ala 120	Met	Leu	Phe	Leu	Thr 125	ГЛа	Tyr	Gly
Pro	Tyr 130	Trp	Arg	ГЛЗ	Asn	Arg 135	Lys	Leu	Val	Ile	Gln 140	Glu	Val	Leu	Ser
Ala 145	Ser	Arg	Leu	Glu	Lys 150	Leu	Lys	His	Val	Arg 155	Phe	Gly	Lys	Ile	Gln 160
Thr	Ser	Ile	Lys	Ser 165	Leu	Tyr	Thr	Arg	Ile 170	Asp	Gly	Asn	Leu	Ser 175	Thr
Ile	Asn	Leu	Thr 180	Asp	Trp	Leu	Glu	Glu 185	Leu	Asn	Phe	Gly	Leu 190	Ile	Val
Lys	Met	Ile 195	Ala	Gly	ГЛЗ	Asn	Tyr 200	Glu	Ser	Gly	Lys	Gly 205	Aab	Glu	Gln
Val	Glu 210	Arg	Phe	Arg	ГЛЗ	Ala 215	Phe	Lys	Asp	Phe	Ile 220	Ile	Leu	Ser	Met
Glu 225	Phe	Val	Leu	Trp	Asp 230	Ala	Phe	Pro	Ile	Pro 235	Leu	Phe	Lys	Trp	Val 240
Asp	Phe	Gln	Gly	His 245	Val	Lys	Ala	Met	Lys 250	Arg	Thr	Phe	Lys	Asp 255	Ile
Asp	Ser	Val	Phe 260	Gln	Asn	Trp	Leu	Glu 265	Glu	His	Val	Lys	Lys 270	Arg	Glu
ГЛа	Met	Glu 275	Val	Asn	Ala	Gln	Gly 280	Asn	Glu	Gln	Asp	Phe 285	Ile	Asp	Val
Val	Leu 290	Ser	ГЛа	Met	Ser	Asn 295	Glu	Tyr	Leu	Asp	Glu 300	Gly	Tyr	Ser	Arg
Asp 305	Thr	Val	Ile	ГЛа	Ala 310	Thr	Val	Phe	Ser	Leu 315	Val	Leu	Asb	Ala	Ala 320
Asp	Thr	Val	Ala	Leu 325	His	Met	Asn	Trp	Gly 330	Met	Ala	Leu	Leu	Ile 335	Asn
Asn	Gln	His	Ala 340	Leu	Гла	Lys	Ala	Gln 345	Glu	Glu	Ile	Asp	Lys 350	Lys	Val
Gly	Lys	Glu 355	Arg	Trp	Val	Glu	Glu 360	Ser	Asp	Ile	ГЛа	Asp 365	Leu	Val	Tyr
Leu	Gln 370	Ala	Ile	Val	ГЛа	Glu 375	Val	Leu	Arg	Leu	Tyr 380	Pro	Pro	Gly	Pro
Leu 385	Leu	Val	Pro	His	Glu 390	Asn	Val	Glu	Asp	Сув 395	Val	Val	Ser	Gly	Tyr 400

His Ile Pro Lys Gly Thr Arg Leu Phe Ala Asn Val Met Lys Leu Gln Arg Asp Pro Lys Leu Trp Ser Asn Pro Asp Lys Phe Asp Pro Glu Arg Phe Phe Ala Asp Asp Ile Asp Tyr Arg Gly Gln His Tyr Glu Phe Ile Pro Phe Gly Ser Gly Arg Arg Ser Cys Pro Gly Met Thr Tyr Ala Leu Gln Ala Glu His Leu Thr Ile Ala His Leu Ile Gln Gly Phe Asn Tyr Lys Thr Pro Asn Asp Glu Pro Leu Asp Met Lys Glu Gly Ala Gly Leu Thr Ile Arg Lys Val Asn Pro Val Glu Val Thr Ile Thr Ala Arg Leu Ala Pro Glu Leu Tyr <210> SEO ID NO 8 <211> LENGTH: 517 <212> TYPE: PRT <213> ORGANISM: Nicotiana tabacum <400> SEOUENCE: 8 Met Val Ser Pro Val Glu Ala Ile Val Gly Leu Val Thr Leu Thr Leu Leu Phe Tyr Phe Leu Trp Pro Lys Lys Phe Gln Ile Pro Ser Lys Pro Leu Pro Pro Lys Ile Pro Gly Gly Trp Pro Val Ile Gly His Leu Phe Tyr Phe Asp Asp Asp Gly Asp Asp Arg Pro Leu Ala Arg Lys Leu Gly Asp Leu Ala Asp Lys Tyr Gly Pro Val Phe Thr Phe Arg Leu Gly Leu Pro Leu Val Leu Val Val Ser Ser Tyr Glu Ala Val Lys Asp Cys Phe Ser Thr Asn Asp Ala Ile Phe Ser Asn Arg Pro Ala Phe Leu Tyr Gly Glu Tyr Leu Gly Tyr Ser Asn Ala Met Leu Phe Leu Thr Lys Tyr Gly Pro Tyr Trp Arg Lys Asn Arg Lys Leu Val Ile Gln Glu Val Leu Ser Ala Ser Arg Leu Glu Lys Leu Lys His Val Arg Phe Gly Lys Ile Gln Thr Ser Ile Lys Ser Leu Tyr Thr Arg Ile Asp Gly Asn Ser Ser Thr Ile Asn Leu Thr Asp Trp Leu Glu Glu Leu Asn Phe Gly Leu Ile Val Lys Met Ile Ala Gly Lys Asn Tyr Glu Ser Gly Lys Gly Asp Glu Gln Val Glu Arg Phe Arg Lys Ala Phe Lys Asp Phe Ile Ile Leu Ser Met Glu Phe Val Leu Trp Asp Ala Phe Pro Ile Pro Leu Phe Lys Trp Val

Asp Phe Gln Gly His Val Lys Ala Met Lys Arg Thr Phe Lys Asp Ile

continued

												con	tin	ued	
				245					250					255	
Asj	Ser	Val	Phe 260	Gln	Asn	Trp	Leu	Glu 265	Glu	His	Val	ГЛЗ	Lys 270	Arg	Glu
LY	3 Met	Glu 275	Val	Asn	Ala	Gln	Gly 280	Asn	Glu	Gln	Asp	Phe 285	Ile	Asp	Val
Va	l Leu 290	. Ser	Lys	Met	Ser	Asn 295	Glu	Tyr	Leu	Asp	Glu 300	Gly	Tyr	Ser	Arg
As] 30!	p Thr 5	Val	Ile	Lys	Ala 310	Thr	Val	Phe	Ser	Leu 315	Val	Leu	Asp	Ala	Ala 320
Asj	o Thr	Val	Ala	Leu 325	His	Met	Asn	Trp	Gly 330	Met	Ala	Leu	Leu	Ile 335	Asn
Ası	n Glr	. His	Ala 340	Leu	Lys	Lys	Ala	Gln 345	Glu	Glu	Ile	Asp	Lys 350	Lys	Val
Gl	ү Цуа	Glu 355	Arg	Trp	Val	Glu	Glu 360	Ser	Asp	Ile	ГЛа	Asp 365	Leu	Val	Tyr
Lei	ı Glr 370	. Ala	Ile	Val	ГЛа	Glu 375	Val	Leu	Arg	Leu	Tyr 380	Pro	Pro	Gly	Pro
Le: 38!	ı Leu 5	. Val	Pro	His	Glu 390	Asn	Val	Glu	Asp	Суя 395	Val	Val	Ser	Gly	Tyr 400
Hi	3 Ile	Pro	Lys	Gly 405	Thr	Arg	Leu	Phe	Val 410	Asn	Val	Met	Lys	Leu 415	Gln
Arg	g Asp	Pro	Lys 420	Leu	Trp	Ser	Asn	Pro 425	Asp	Lys	Phe	Asp	Pro 430	Glu	Arg
Phe	e Phe	Ala 435	Asp	Asp	Ile	Asp	Tyr 440	Arg	Gly	Gln	His	Tyr 445	Glu	Phe	Ile
Pro	> Phe 450	Gly	Ser	Gly	Arg	Arg 455	Ser	Сүз	Pro	Gly	Met 460	Thr	Tyr	Ala	Leu
Gl: 465	n Ala 5	Glu	His	Leu	Thr 470	Ile	Ala	His	Leu	Ile 475	Gln	Gly	Phe	Asn	Tyr 480
Ly	s Thr	Pro	Asn	Asp 485	Glu	Pro	Leu	Asp	Met 490	ГЛа	Glu	Gly	Ala	Gly 495	Leu
Th:	r Ile	Arg	Lys 500	Val	Asn	Pro	Val	Glu 505	Val	Thr	Ile	Thr	Ala 510	Arg	Leu
Ala	a Pro	Glu 515	Leu	Tyr											
< 2.	10> 5	EQ II	d No	9											
<23 <23	l1> I l2> I	ENGT YPE :	H: 5 PRT	17											
< 23	L3> C	RGAN	ISM:	Nico	otia	na t	abacı	um							
< 40)0> S	EQUE	NCE:	9	_	_	_					_		_	
Mei 1	: Val	Ser	Pro	Val 5	Glu	Ala	Ile	Val	Gly 10	Leu	Val	Thr	Leu	Thr 15	Leu
Lei	ı Ph∈	Tyr	Phe 20	Leu	Trp	Pro	Lys	Lys 25	Phe	Gln	Ile	Pro	Ser 30	Lys	Pro
Lei	ı Pro	Pro 35	Lys	Ile	Pro	Gly	Gly 40	Trp	Pro	Val	Ile	Gly 45	His	Leu	Phe
Ту	r Phe 50	Asp	Asp	Asp	Gly	Asp 55	Asp	Arg	Pro	Leu	Ala 60	Arg	ГЛа	Leu	Gly
As] 65	p Leu	Ala	Asp	ГЛа	Tyr 70	Gly	Pro	Val	Phe	Thr 75	Phe	Arg	Leu	Gly	Leu 80
Pro	o Leu	. Val	Leu	Val 85	Val	Ser	Ser	Tyr	Glu 90	Ala	Val	ГЛа	Asp	Сүз 95	Phe

-continued

_															
Ser	Thr	Asn	Asp 100	Ala	Ile	Phe	Ser	Asn 105	Arg	Pro	Ala	Phe	Leu 110	Tyr	Gly
Glu	Tyr	Leu 115	Gly	Tyr	Ser	Asn	Ala 120	Met	Leu	Phe	Leu	Thr 125	Lys	Tyr	Gly
Pro	Tyr 130	Trp	Arg	ГЛа	Asn	Arg 135	ГЛа	Leu	Val	Ile	Gln 140	Glu	Val	Leu	Ser
Ala 145	Ser	Arg	Leu	Glu	Lys 150	Leu	ГЛа	His	Val	Arg 155	Phe	Gly	ГЛа	Ile	Gln 160
Thr	Ser	Ile	Lys	Ser 165	Leu	Tyr	Thr	Arg	Ile 170	Asp	Gly	Asn	Ser	Ser 175	Thr
Ile	Asn	Leu	Thr 180	Asp	Trp	Leu	Glu	Glu 185	Leu	Asn	Phe	Gly	Leu 190	Ile	Val
Гла	Met	Ile 195	Ala	Gly	ГЛа	Asn	Tyr 200	Glu	Ser	Gly	ГЛа	Gly 205	Asp	Glu	Gln
Val	Glu 210	Arg	Phe	Arg	Гла	Ala 215	Phe	Lys	Asp	Phe	Ile 220	Ile	Leu	Ser	Ile
Glu 225	Phe	Val	Leu	Trp	Asp 230	Ala	Phe	Pro	Ile	Pro 235	Leu	Phe	ГÀа	Trp	Val 240
Asp	Phe	Gln	Gly	His 245	Val	Lys	Ala	Met	Lys 250	Arg	Thr	Phe	Lys	Asp 255	Ile
Asp	Ser	Val	Phe 260	Gln	Asn	Trp	Leu	Glu 265	Glu	His	Val	Гла	Lys 270	Arg	Glu
Гла	Met	Glu 275	Val	Asn	Ala	Gln	Gly 280	Asn	Glu	Gln	Asp	Phe 285	Ile	Asp	Val
Val	Leu 290	Ser	Lys	Met	Ser	Asn 295	Glu	Tyr	Leu	Asp	Glu 300	Gly	Tyr	Ser	Arg
Asp 305	Thr	Val	Ile	Lys	Ala 310	Thr	Val	Phe	Ser	Leu 315	Val	Leu	Asp	Ala	Ala 320
Asp	Thr	Val	Ala	Leu 325	His	Met	Asn	Trp	Gly 330	Met	Ala	Leu	Leu	Ile 335	Asn
Asn	Gln	His	Ala 340	Leu	ГЛЗ	Lys	Ala	Gln 345	Glu	Glu	Ile	Asp	Lys 350	Lys	Val
Gly	Lys	Glu 355	Arg	Trp	Val	Glu	Glu 360	Ser	Asp	Ile	Lys	Asp 365	Leu	Val	Tyr
Leu	Gln 370	Ala	Ile	Val	Lys	Glu 375	Val	Leu	Arg	Leu	Tyr 380	Pro	Pro	Gly	Pro
Leu 385	Leu	Val	Pro	His	Glu 390	Asn	Val	Glu	Asp	Cys 395	Val	Val	Ser	Gly	Tyr 400
His	Ile	Pro	Lys	Gly 405	Thr	Arg	Leu	Phe	Ala 410	Asn	Val	Met	Lys	Leu 415	Gln
Arg	Asp	Pro	Lys 420	Leu	Trp	Ser	Asn	Pro 425	Asp	Lys	Phe	Asp	Pro 430	Glu	Arg
Phe	Phe	Ala 435	Asp	Asp	Ile	Asp	Tyr 440	Arg	Gly	Gln	His	Tyr 445	Glu	Phe	Ile
Pro	Phe 450	Gly	Ser	Gly	Arg	Arg 455	Ser	Сүз	Pro	Gly	Met 460	Thr	Tyr	Ala	Leu
Gln 465	Ala	Glu	His	Leu	Thr 470	Ile	Ala	His	Leu	Ile 475	Gln	Gly	Phe	Asn	Tyr 480
Lys	Thr	Pro	Asn	Asp 485	Glu	Pro	Leu	Asp	Met 490	Lys	Glu	Gly	Ala	Gly 495	Leu
Thr	Ile	Arg	Lys 500	Val	Asn	Pro	Val	Glu 505	Val	Thr	Ile	Thr	Ala 510	Arg	Leu
	Dress	01	Terr	m e											

<210 <211 <212)> SH .> LH 2> TY	EQ II ENGTH PE:) NO I: 51 PRT	10 .7											
<213	8> OF	RGANI	SM:	Nico	otiar	na ta	abacu	ım							
<400)> SE	QUEN	ICE :	10											
Met 1	Val	Ser	Pro	Val 5	Glu	Ala	Ile	Val	Gly 10	Leu	Val	Thr	Leu	Thr 15	Leu
Leu	Phe	Tyr	Phe 20	Leu	Trp	Pro	Lys	Lys 25	Phe	Gln	Ile	Pro	Ser 30	Lys	Pro
Leu	Pro	Pro 35	ГЛа	Ile	Pro	Gly	Gly 40	Trp	Pro	Val	Ile	Gly 45	His	Leu	Phe
Tyr	Phe 50	Asp	Asp	Asp	Gly	Asp 55	Asp	Arg	Pro	Leu	Ala 60	Arg	Lys	Leu	Gly
Asp 65	Leu	Ala	Asp	Lys	Tyr 70	Gly	Leu	Val	Phe	Thr 75	Phe	Arg	Leu	Gly	Leu 80
Pro	Leu	Val	Leu	Val 85	Val	Ser	Ser	Tyr	Glu 90	Ala	Val	Lys	Asp	Сув 95	Phe
Ser	Thr	Asn	Asp 100	Ala	Ile	Phe	Ser	Asn 105	Arg	Pro	Ala	Phe	Leu 110	Tyr	Gly
Glu	Tyr	Leu 115	Gly	Tyr	Ser	Asn	Ala 120	Met	Leu	Phe	Leu	Thr 125	Lys	Tyr	Gly
Pro	Tyr 130	Trp	Arg	Lys	Asn	Arg 135	Lys	Leu	Val	Ile	Gln 140	Glu	Val	Leu	Ser
Ala 145	Ser	Arg	Leu	Glu	Lys 150	Leu	Lys	His	Val	Arg 155	Phe	Gly	Lys	Ile	Gln 160
Thr	Ser	Ile	Lys	Ser 165	Leu	Tyr	Thr	Arg	Ile 170	Asp	Gly	Asn	Ser	Ser 175	Thr
Ile	Asn	Leu	Thr 180	Asp	Trp	Leu	Glu	Glu 185	Leu	Asn	Phe	Gly	Leu 190	Ile	Val
Lys	Met	Ile 195	Ala	Gly	Lys	Asn	Tyr 200	Glu	Ser	Gly	Lys	Gly 205	Asp	Glu	Gln
Val	Glu 210	Arg	Phe	Arg	ГÀа	Ala 215	Phe	Lys	Asp	Phe	Ile 220	Ile	Leu	Ser	Met
Glu 225	Phe	Val	Leu	Trp	Aap 230	Ala	Phe	Pro	Ile	Pro 235	Leu	Phe	Lys	Trp	Val 240
Asp	Phe	Gln	Gly	His 245	Val	Lys	Ala	Met	Lys 250	Arg	Thr	Phe	ГЛа	Asp 255	Ile
Asp	Ser	Val	Phe 260	Gln	Asn	Trp	Leu	Glu 265	Glu	His	Val	ГЛа	Lys 270	Arg	Glu
Lys	Met	Glu 275	Val	Asn	Ala	Gln	Gly 280	Asn	Glu	Gln	Asp	Phe 285	Ile	Asp	Val
Val	Leu 290	Ser	Lys	Met	Ser	Asn 295	Glu	Tyr	Leu	Asp	Glu 300	Gly	Tyr	Ser	Arg
Asp 305	Thr	Val	Ile	Lys	Ala 310	Thr	Val	Phe	Ser	Leu 315	Val	Leu	Asp	Ala	Ala 320
Asp	Thr	Val	Ala	Leu 325	His	Met	Asn	Trp	Gly 330	Met	Ala	Leu	Leu	Ile 335	Asn
Asn	Gln	His	Ala 340	Leu	Lys	Lys	Ala	Gln 345	Glu	Glu	Ile	Asp	Lys 350	Lya	Val
Gly	LYa	Glu 355	Arg	Trp	Val	Glu	Glu 360	Ser	Asp	Ile	Lys	Asp 365	Leu	Val	Tyr

Leu	Gln	Ala	Ile	Val	LYa	Glu	Val	Leu	Arg	Leu	Tyr	Pro	Pro	Gly	Pro
_	370		_			375		<i>a</i> 1	_	~	380		a	<i>a</i> .	_
Leu 385	Leu	Val	Pro	His	Glu 390	Asn	Val	Glu	Aab	Сув 395	Val	Val	Ser	Gly	Tyr 400
His	Ile	Pro	Lys	Gly 405	Thr	Arg	Leu	Phe	Ala 410	Asn	Val	Met	Lys	Leu 415	Gln
Arg	Asp	Pro	Lys 420	Leu	Trp	Ser	Asn	Pro 425	Asp	ГЛЗ	Phe	Asp	Pro 430	Glu	Arg
Phe	Phe	Ala 435	Asp	Asp	Ile	Asp	Tyr 440	Arg	Gly	Gln	His	Tyr 445	Glu	Phe	Ile
Pro	Phe 450	Gly	Ser	Gly	Arg	Arg 455	Ser	Сүз	Pro	Gly	Met 460	Thr	Tyr	Ala	Leu
Gln 465	Ala	Glu	His	Leu	Thr 470	Ile	Ala	His	Leu	Ile 475	Gln	Gly	Phe	Asn	Tyr 480
Lys	Thr	Pro	Asn	Asp 485	Glu	Pro	Leu	Asp	Met 490	Lys	Glu	Gly	Ala	Gly 495	Leu
Thr	Ile	Arg	Lys 500	Val	Asn	Pro	Val	Glu 505	Val	Thr	Ile	Thr	Ala 510	Arg	Leu
Ala	Pro	Glu 515	Leu	Tyr											
<210 <211)> SH L> LH	EQ II ENGTH) NO 1: 5:	11 17											
<212 <213	2> TY 3> OF	(PE : RGAN	PRT [SM:	Nico	otiar	na ta	abacı	ım							
<400)> SH	EQUEI	ICE :	11											
Met 1	Val	Ser	Pro	Val 5	Glu	Ala	Ile	Val	Gly 10	Leu	Val	Thr	Leu	Thr 15	Leu
Leu	Phe	Tyr	Phe 20	Leu	Trp	Pro	Lys	Lys 25	Phe	Gln	Ile	Pro	Ser 30	Lys	Pro
Leu	Pro	Pro 35	Lys	Ile	Pro	Gly	Gly 40	Trp	Pro	Val	Ile	Gly 45	His	Leu	Phe
Tyr	Phe 50	Asp	Asp	Asp	Gly	Asp 55	Asp	Arg	Pro	Leu	Ala 60	Arg	Lys	Leu	Gly
Asp 65	Leu	Ala	Asp	ГЛа	Tyr 70	Gly	Pro	Val	Phe	Thr 75	Phe	Arg	Leu	Gly	Leu 80
Asp 65 Pro	Leu Leu	Ala Val	Asp Leu	Lys Val 85	Tyr 70 Val	Gly Ser	Pro Ser	Val Tyr	Phe Glu 90	Thr 75 Ala	Phe Val	Arg Lys	Leu Asp	Gly Cys 95	Leu 80 Phe
Asp 65 Pro Ser	Leu Leu Thr	Ala Val Asn	Asp Leu Asp 100	Lys Val 85 Ala	Tyr 70 Val Ile	Gly Ser Phe	Pro Ser Ser	Val Tyr Asn 105	Phe Glu 90 Arg	Thr 75 Ala Pro	Phe Val Ala	Arg Lys Phe	Leu Asp Leu 110	Gly Cys 95 Tyr	Leu 80 Phe Gly
Asp 65 Pro Ser Glu	Leu Leu Thr Tyr	Ala Val Asn Leu 115	Asp Leu Asp 100 Gly	Lys Val 85 Ala Tyr	Tyr 70 Val Ile Ser	Gly Ser Phe Asn	Pro Ser Ser Ala 120	Val Tyr Asn 105 Met	Phe Glu 90 Arg Leu	Thr 75 Ala Pro Phe	Phe Val Ala Leu	Arg Lys Phe Thr 125	Leu Asp Leu 110 Lys	Gly Cya 95 Tyr Tyr	Leu 80 Phe Gly Gly
Asp 65 Pro Ser Glu Pro	Leu Leu Thr Tyr Tyr	Ala Val Asn Leu 115 Trp	Asp Leu Asp 100 Gly Arg	Lys Val 85 Ala Tyr Lys	Tyr 70 Val Ile Ser Asn	Gly Ser Phe Asn Arg 135	Pro Ser Ser Ala 120 Lys	Val Tyr Asn 105 Met Leu	Phe Glu 90 Arg Leu Val	Thr 75 Ala Pro Phe Ile	Phe Val Ala Leu Gln 140	Arg Lys Phe Thr 125 Glu	Leu Asp Leu 110 Lys Val	Gly Cys 95 Tyr Tyr Phe	Leu 80 Phe Gly Gly Ser
Asp 65 Pro Ser Glu Pro	Leu Leu Thr Tyr 130	Ala Val Asn Leu 115 Trp	Asp Leu Asp 100 Gly Arg	Lys Val 85 Ala Tyr Lys	Tyr 70 Val Ile Ser Asn	Gly Ser Phe Asn Arg 135	Pro Ser Ser Ala 120 Lys	Val Tyr Asn 105 Met Leu	Phe Glu 90 Arg Leu Val	Thr 75 Ala Pro Phe Ile	Phe Val Ala Leu Gln 140 Phe	Arg Lys Phe Thr 125 Glu	Leu Asp Leu 110 Lys Val	Gly Cys 95 Tyr Tyr Phe	Leu 80 Phe Gly Gly Ser
Asp 65 Pro Ser Glu Pro Ala 145	Leu Leu Thr Tyr 130 Ser	Ala Val Asn Leu 115 Trp Arg	Asp Leu Asp 100 Gly Arg Leu	Lys Val 85 Ala Tyr Lys Glu	Tyr 70 Val Ile Ser Asn Lys 150	Gly Ser Phe Asn Arg 135 Leu	Pro Ser Ser Ala 120 Lys Lys	Val Tyr Asn 105 Met Leu His	Phe Glu 90 Arg Leu Val Val	Thr 75 Ala Pro Phe Ile Arg 155	Phe Val Ala Leu Gln 140 Phe	Arg Lys Phe Thr 125 Glu Gly	Leu Asp Leu 110 Lys Val Lys	Gly 95 Tyr Tyr Phe Ile	Leu 80 Phe Gly Gly Ser Gln 160
Asp 65 Pro Ser Glu Pro Ala 145 Thr	Leu Leu Thr Tyr 130 Ser Ser	Ala Val Asn Leu 115 Trp Arg Ile	Asp Leu Asp 100 Gly Arg Leu Lys	Lys Val 85 Ala Tyr Lys Glu Ser 165	Tyr 70 Val Ile Ser Asn Lys 150 Leu	Gly Ser Phe Asn Arg 135 Leu Tyr	Pro Ser Ser Ala 120 Lys Lys Thr	Val Tyr Asn 105 Met Leu His Arg	Phe Glu 90 Arg Leu Val Val Ile 170	Thr 75 Ala Pro Phe Ile Arg 155 Asp	Phe Val Ala Leu Gln 140 Phe Gly	Arg Lys Phe Thr 125 Glu Gly Asn	Leu Asp Leu 110 Lys Val Lys Ser	Gly 95 Tyr Tyr Phe Ile Ser 175	Leu 80 Phe Gly Gly Ser Gln 160 Thr
Asp 65 Pro Ser Glu Pro Ala 145 Thr Ile	Leu Leu Thr Tyr 130 Ser Ser Asn	Ala Val Asn Leu 115 Trp Arg Ile Leu	Asp Leu Asp 100 Gly Arg Leu Lys Thr 180	Lys Val 85 Ala Tyr Lys Glu Ser 165 Asp	Tyr 70 Val Ile Ser Asn Lys 150 Leu Trp	Gly Ser Phe Asn Arg 135 Leu Tyr Leu	Pro Ser Ser Ala 120 Lys Lys Thr Glu	Val Tyr Asn 105 Met Leu His Arg Glu 185	Phe Glu 90 Arg Leu Val Val Ile 170 Leu	Thr 75 Ala Pro Phe Ile Arg 155 Asp Asn	Phe Val Ala Leu Gln 140 Phe Gly Phe	Arg Lys Phe Thr 125 Glu Gly Asn Gly	Leu Asp Leu 110 Lys Val Lys Ser Leu 190	Gly 95 Tyr Tyr Phe Ile Ser 175 Ile	Leu 80 Phe Gly Gly Ser Gln 160 Thr Val
Asp 65 Pro Ser Glu Pro Ala 145 Thr Ile Lys	Leu Leu Thr Tyr 130 Ser Ser Asn Met	Ala Val Asn Leu 115 Trp Arg Ile Leu Ile	Asp Leu Asp 100 Gly Arg Leu Lys Thr 180 Ala	Lys Val Ala Tyr Lys Glu Ser 165 Asp Gly	Tyr 70 Val Ile Ser Asn Lys Lys Leu Trp Lys	Gly Ser Phe Asn Arg 135 Leu Tyr Leu Asn	Pro Ser Ser Ala Lys Lys Thr Glu Tyr 200	Val Tyr Asn 105 Met Leu His Arg Glu 185 Glu	Phe Glu 90 Arg Leu Val Val Ile 170 Leu Ser	Thr 75 Ala Pro Phe Ile Arg 155 Asp Asn Gly	Phe Val Ala Leu Gln 140 Phe Gly Phe Lys	Arg Lys Phe Thr 125 Glu Gly Asn Gly Gly 205	Leu Asp Leu Lys Val Lys Ser Leu 190 Asp	Gly Cys 95 Tyr Tyr Phe Ile Ser 175 Ile Glu	Leu 80 Phe Gly Ser Gln 160 Thr Val Gln

-continued

Glu Phe Val Leu Trp Asp Ala Phe Pro Ile Pro Leu Phe Lys Trp Val Asp Phe Gln Gly His Val Lys Ala Met Lys Arg Thr Phe Lys Asp Ile Asp Ser Val Phe Gln Asn Trp Leu Glu Glu His Val Lys Lys Arg Glu Lys Met Glu Val Asn Ala Gln Gly Asn Glu Gln Asp Phe Ile Asp Val Val Leu Ser Lys Met Ser Asn Glu Tyr Leu Asp Glu Gly Tyr Ser Arg Asp Thr Val Ile Lys Ala Thr Val Phe Ser Leu Val Leu Asp Ala Ala Asp Thr Val Ala Leu His Met Asn Trp Gly Met Ala Leu Leu Ile Asn Asn Gln His Ala Leu Lys Lys Ala Gln Glu Glu Ile Asp Lys Lys Val Gly Lys Glu Arg Trp Val Glu Glu Ser Asp Ile Lys Asp Leu Val Tyr Leu Gln Ala Ile Val Lys Glu Val Leu Arg Leu Tyr Pro Pro Gly Pro Leu Leu Val Pro His Glu Asn Val Glu Asp Cys Val Val Ser Gly Tyr His Ile Pro Lys Gly Thr Arg Leu Phe Ala Asn Val Met Lys Leu Gln Arg Asp Pro Lys Leu Trp Ser Asn Pro Asp Lys Phe Asp Pro Glu Arg Phe Phe Ala Asp Asp Ile Asp Tyr Arg Gly Gln His Tyr Glu Phe Ile Pro Phe Gly Ser Gly Arg Arg Ser Cys Pro Gly Met Thr Tyr Ala Leu Gln Ala Glu His Leu Thr Ile Ala His Leu Ile Gln Gly Phe Asn Tyr Lys Thr Pro Asn Asp Glu Pro Leu Asp Met Lys Glu Gly Ala Gly Leu Thr Ile Arg Lys Val Asn Pro Val Glu Val Thr Ile Thr Ala Arg Leu Ala Pro Glu Leu Tyr <210> SEQ ID NO 12 <211> LENGTH: 421 <212> TYPE: PRT <213> ORGANISM: Nicotiana tabacum <400> SEQUENCE: 12 Met Val Ser Pro Val Glu Ala Ile Val Gly Leu Val Thr Leu Thr Leu Leu Phe Tyr Phe Leu Trp Pro Lys Lys Phe Gln Ile Pro Ser Lys Pro Leu Pro Pro Lys Ile Pro Gly Gly Trp Pro Val Ile Gly His Leu Phe Tyr Phe Asp Asp Asp Gly Asp Asp Arg Pro Leu Ala Arg Lys Leu Gly Asp Leu Ala Asp Lys Tyr Gly Pro Val Phe Thr Phe Arg Leu Gly Leu

65	5					70					75					80
Pr	0	Leu	Val	Leu	Val 85	Val	Ser	Ser	Tyr	Glu 90	Ala	Val	ГЛЗ	Asp	Сув 95	Phe
Se	er	Thr	Asn	Asp 100	Ala	Ile	Phe	Ser	Asn 105	Arg	Pro	Ala	Phe	Leu 110	Tyr	Gly
Gl	lu	Tyr	Leu 115	Gly	Tyr	Ser	Asn	Ala 120	Met	Leu	Phe	Leu	Thr 125	Lys	Tyr	Gly
Pr	ro	Tyr 130	Trp	Arg	Lys	Asn	Arg 135	Lys	Leu	Val	Ile	Gln 140	Glu	Val	Leu	Ser
A1 14	La 15	Ser	Arg	Leu	Glu	Lys 150	Leu	ГЛа	His	Val	Arg 155	Phe	Gly	ГÀа	Ile	Gln 160
Th	ır	Ser	Ile	Lys	Ser 165	Leu	Tyr	Thr	Arg	Ile 170	Asp	Gly	Asn	Ser	Ser 175	Thr
Il	Le	Asn	Leu	Thr 180	Asp	Trp	Leu	Glu	Glu 185	Leu	Asn	Phe	Gly	Leu 190	Ile	Val
Γλ	/s	Met	Ile 195	Ala	Gly	Гла	Asn	Tyr 200	Glu	Ser	Gly	ГЛа	Gly 205	Asp	Glu	Gln
Va	al	Glu 210	Arg	Phe	Arg	ГЛа	Ala 215	Phe	Гла	Asp	Phe	Ile 220	Ile	Leu	Ser	Met
G1 22	lu 25	Phe	Val	Leu	Trp	Asp 230	Ala	Phe	Pro	Ile	Pro 235	Leu	Phe	Lys	Trp	Val 240
Αs	ab	Phe	Gln	Gly	His 245	Val	ГЛа	Ala	Met	Lys 250	Arg	Thr	Phe	Lys	Asp 255	Ile
As	ab	Ser	Val	Phe 260	Gln	Asn	Trp	Leu	Glu 265	Glu	His	Val	Lys	Lys 270	Arg	Glu
LУ	/5	Met	Glu 275	Val	Asn	Ala	Gln	Gly 280	Asn	Glu	Gln	Asp	Phe 285	Ile	Asp	Val
Va	al	Leu 290	Ser	Lys	Met	Ser	Asn 295	Glu	Tyr	Leu	Asp	Glu 300	Gly	Tyr	Ser	Arg
Ая 30	sp)5	Thr	Val	Ile	Гүз	Ala 310	Thr	Val	Phe	Ser	Leu 315	Val	Leu	Asp	Ala	Ala 320
As	p	Thr	Val	Ala	Leu 325	His	Met	Asn	Trp	Gly 330	Met	Ala	Leu	Leu	Ile 335	Asn
As	m	Gln	His	Ala 340	Leu	ГЛа	LYa	Ala	Gln 345	Glu	Glu	Ile	Asp	Lys 350	Lys	Val
Gl	Ly	Lys	Glu 355	Arg	Trp	Val	Glu	Glu 360	Ser	Asp	Ile	ГЛа	Asp 365	Leu	Val	Tyr
Le	eu	Gln 370	Ala	Ile	Val	ГЛа	Glu 375	Val	Leu	Arg	Leu	Tyr 380	Pro	Pro	Gly	Pro
Le 38	eu 85	Leu	Val	Pro	His	Glu 390	Asn	Val	Glu	Asp	Сув 395	Val	Val	Ser	Gly	Tyr 400
Hi	ls	Ile	Pro	Lys	Gly 405	Thr	Arg	Leu	Phe	Ala 410	Asn	Val	Met	Lys	Leu 415	Gln
Ar	g	Asp	Pro	Lys 420	Leu											
-				720												
<2 <2 <2	210 211 212	> SH .> LH .> TY	SQ II ENGTH (PE:) NO 1: 51 PRT	13 17											
< 2	213	> OF	RGANI	ISM:	Nico	otian	na ta	abacı	ım							
<4 M∈	et	Leu	Ser	Pro	Ile	Glu	Ala	Ile	Val	Gly	Leu	Val	Thr	Phe	Thr	Phe
1			-	-	5			-	. 3	10				-	15	-

Leu	Phe	Phe	Phe 20	Leu	Trp	Thr	Lys	Lys 25	Ser	Gln	Lys	Pro	Ser 30	Lys	Pro
Leu	Pro	Pro 35	Гла	Ile	Pro	Gly	Gly 40	Trp	Pro	Val	Ile	Gly 45	His	Leu	Phe
His	Phe 50	Asn	Asp	Asp	Gly	Asp 55	Asp	Arg	Pro	Leu	Ala 60	Arg	Lys	Leu	Gly
Asp 65	Leu	Ala	Asp	ГЛа	Tyr 70	Gly	Pro	Val	Phe	Thr 75	Phe	Arg	Leu	Gly	Leu 80
Pro	Leu	Val	Leu	Val 85	Val	Ser	Ser	Tyr	Glu 90	Ala	Val	ГЛа	Asp	Сув 95	Phe
Ser	Thr	Asn	Asp 100	Ala	Ile	Phe	Ser	Asn 105	Arg	Pro	Ala	Phe	Leu 110	Tyr	Gly
Asp	Tyr	Leu 115	Gly	Tyr	Asn	Asn	Ala 120	Met	Leu	Phe	Leu	Ala 125	Asn	Tyr	Gly
Pro	Tyr 130	Trp	Arg	ГЛа	Asn	Arg 135	Гла	Leu	Val	Ile	Gln 140	Glu	Val	Leu	Ser
Ala 145	Ser	Arg	Leu	Glu	Lys 150	Phe	Lys	His	Val	Arg 155	Phe	Ala	Arg	Ile	Gln 160
Ala	Ser	Ile	Lys	Asn 165	Leu	Tyr	Thr	Arg	Ile 170	Asp	Gly	Asn	Ser	Ser 175	Thr
Ile	Asn	Leu	Thr 180	Asp	Trp	Leu	Glu	Glu 185	Leu	Asn	Phe	Gly	Leu 190	Ile	Val
Lys	Met	Ile 195	Ala	Gly	Lys	Asn	Tyr 200	Glu	Ser	Gly	Гүз	Gly 205	Asp	Glu	Gln
Val	Glu 210	Arg	Phe	Lys	Lys	Ala 215	Phe	Lys	Asp	Phe	Met 220	Ile	Leu	Ser	Met
Glu 225	Phe	Val	Leu	Trp	Asp 230	Ala	Phe	Pro	Ile	Pro 235	Leu	Phe	Lys	Trp	Val 240
Asp	Phe	Gln	Gly	His 245	Val	Lys	Ala	Met	Lys 250	Arg	Thr	Phe	Lys	Asp 255	Ile
Asp	Ser	Val	Phe 260	Gln	Asn	Trp	Leu	Glu 265	Glu	His	Ile	Asn	Lys 270	Arg	Glu
LYa	Met	Glu 275	Val	Asn	Ala	Glu	Gly 280	Asn	Glu	Gln	Asp	Phe 285	Ile	Asp	Val
Val	Leu 290	Ser	ГЛа	Met	Ser	Asn 295	Glu	Tyr	Leu	Gly	Glu 300	Gly	Tyr	Ser	Arg
Asp 305	Thr	Val	Ile	Гла	Ala 310	Thr	Val	Phe	Ser	Leu 315	Val	Leu	Asp	Ala	Ala 320
Asp	Thr	Val	Ala	Leu 325	His	Ile	Asn	Trp	Gly 330	Met	Ala	Leu	Leu	Ile 335	Asn
Asn	Gln	Lys	Ala 340	Leu	Thr	ГЛа	Ala	Gln 345	Glu	Glu	Ile	Asp	Thr 350	Lys	Val
Gly	Lys	Asp 355	Arg	Trp	Val	Glu	Glu 360	Ser	Asp	Ile	ГЛа	Asp 365	Leu	Val	Tyr
Leu	Gln 370	Ala	Ile	Val	ГЛа	Glu 375	Val	Leu	Arg	Leu	Tyr 380	Pro	Pro	Gly	Pro
Leu 385	Leu	Val	Pro	His	Glu 390	Asn	Val	Glu	Asp	Суз 395	Val	Val	Ser	Gly	Tyr 400
His	Ile	Pro	Lys	Gly 405	Thr	Arg	Leu	Phe	Ala 410	Asn	Val	Met	Lys	Leu 415	Gln
Arg	Asp	Pro	Lys 420	Leu	Trp	Ser	Asp	Pro 425	Asp	Thr	Phe	Asp	Pro 430	Glu	Arg
Dha	Tle	Ala	Thr	Asp	Tle	Asp	Phe	Ara	Glv	Gln	Tvr	Tvr	Lvs	Tvr	Ile

		435					440					445			
Pro	Phe 450	Gly	Ser	Gly	Arg	Arg 455	Ser	Сүз	Pro	Gly	Met 460	Thr	Tyr	Ala	Leu
Gln 465	Val	Glu	His	Leu	Thr 470	Met	Ala	His	Leu	Ile 475	Gln	Gly	Phe	Asn	Tyr 480
Arg	Thr	Pro	Asn	Asp 485	Glu	Pro	Leu	Asp	Met 490	Lys	Glu	Gly	Ala	Gly 495	Ile
Thr	Ile	Arg	Lys 500	Val	Asn	Pro	Val	Glu 505	Leu	Ile	Ile	Ala	Pro 510	Arg	Leu
Ala	Pro	Glu 515	Leu	Tyr											
<21 <21 <21 <21	0> SI 1> LI 2> T 3> OI	EQ II ENGTI YPE : RGANI	D NO H: 5: PRT ISM:	14 17 Nico	otia	na ta	abacı	um							
<40	0> S1	EQUEI	NCE :	14											
Met 1	Leu	Ser	Pro	Ile 5	Glu	Ala	Ile	Val	Gly 10	Leu	Val	Thr	Phe	Thr 15	Phe
Leu	Phe	Phe	Phe 20	Leu	Trp	Thr	Lys	Lys 25	Ser	Gln	Lys	Pro	Ser 30	Lys	Pro
Leu	Pro	Pro 35	Lys	Ile	Pro	Gly	Gly 40	Trp	Pro	Val	Ile	Gly 45	His	Leu	Phe
His	Phe 50	Asn	Asp	Asp	Gly	Asp 55	Asp	Arg	Pro	Leu	Ala 60	Arg	Lys	Leu	Gly
Asp 65	Leu	Ala	Asp	Lys	Tyr 70	Gly	Pro	Val	Phe	Thr 75	Phe	Arg	Leu	Gly	Leu 80
Pro	Leu	Val	Leu	Val 85	Val	Ser	Ser	Tyr	Glu 90	Ala	Val	ГЛа	Asp	Суз 95	Phe
Ser	Thr	Asn	Asp 100	Ala	Ile	Phe	Ser	Asn 105	Arg	Pro	Ala	Phe	Leu 110	Tyr	Gly
Asp	Tyr	Leu 115	Gly	Tyr	Asn	Asn	Ala 120	Met	Leu	Phe	Leu	Ala 125	Asn	Tyr	Gly
Pro	Tyr 130	Trp	Arg	ГЛа	Asn	Arg 135	ГЛа	Leu	Val	Ile	Gln 140	Glu	Val	Leu	Ser
Ala 145	Ser	Arg	Leu	Glu	Lys 150	Phe	Lys	His	Val	Arg 155	Phe	Ala	Arg	Ile	Gln 160
Ala	Ser	Ile	Lys	Asn 165	Leu	Tyr	Thr	Arg	Ile 170	Asp	Gly	Asn	Ser	Ser 175	Thr
Ile	Asn	Leu	Thr 180	Asp	Trp	Leu	Glu	Glu 185	Leu	Asn	Phe	Gly	Leu 190	Ile	Val
ГЛа	Met	Ile 195	Ala	Gly	ГЛа	Asn	Tyr 200	Glu	Ser	Gly	ГЛа	Gly 205	Asp	Glu	Gln
Val	Glu 210	Arg	Phe	ГЛа	Lys	Ala 215	Phe	Lys	Asp	Phe	Met 220	Ile	Leu	Ser	Met
Glu 225	Phe	Val	Leu	Trp	Asp 230	Ala	Phe	Pro	Ile	Pro 235	Leu	Phe	Lys	Trp	Val 240
Aap	Phe	Gln	Gly	His 245	Val	ГЛа	Ala	Met	Lys 250	Arg	Thr	Phe	Lys	Asp 255	Ile
Asp	Ser	Val	Phe	Gln	Asn	Trp	Leu	Glu	Glu	His	Ile	Asn	Lys	Arg	Glu
Lys	Met	Glu	260 Val	Asn	Ala	Glu	Gly	265 Asn	Glu	Gln	Asp	Phe	270 Ile	Asp	Val
		275					280					285			

Val	Leu 290	Ser	Гла	Met	Ser	Asn 295	Glu	Tyr	Leu	Gly	Glu 300	Gly	Tyr	Ser	Arg
Asp 305	Thr	Val	Ile	Lys	Ala 310	Thr	Val	Phe	Ser	Leu 315	Val	Leu	Asp	Ala	Ala 320
Asp	Thr	Val	Ala	Leu 325	His	Ile	Asn	Trp	Gly 330	Met	Ala	Leu	Leu	Ile 335	Asn
Asn	Gln	Гла	Ala 340	Leu	Thr	ГЛа	Ala	Gln 345	Glu	Glu	Ile	Asp	Thr 350	Lys	Val
Gly	Lys	Asp 355	Arg	Trp	Val	Glu	Glu 360	Ser	Asp	Ile	Lys	Asp 365	Leu	Val	Tyr
Leu	Gln 370	Ala	Ile	Val	ГЛа	Glu 375	Val	Leu	Arg	Leu	Tyr 380	Pro	Pro	Gly	Pro
Leu 385	Leu	Val	Pro	His	Glu 390	Asn	Val	Glu	Asp	Сув 395	Val	Val	Ser	Gly	Tyr 400
His	Ile	Pro	ГЛа	Gly 405	Thr	Arg	Leu	Phe	Ala 410	Asn	Val	Met	ГÀа	Leu 415	Gln
Arg	Asp	Pro	Lys 420	Leu	Trp	Ser	Asp	Pro 425	Asp	Thr	Phe	Asp	Pro 430	Glu	Arg
Phe	Ile	Ala 435	Thr	Asp	Ile	Asp	Phe 440	Arg	Gly	Gln	Tyr	Tyr 445	Гла	Tyr	Ile
Pro	Phe 450	Gly	Ser	Gly	Arg	Arg 455	Ser	Сүз	Ser	Gly	Met 460	Thr	Tyr	Ala	Leu
Gln 465	Val	Glu	His	Leu	Thr 470	Met	Ala	His	Leu	Ile 475	Gln	Gly	Phe	Asn	Tyr 480
Arg	Thr	Pro	Asn	Asp 485	Glu	Pro	Leu	Asp	Met 490	Lys	Glu	Gly	Ala	Gly 495	Ile
Thr	Ile	Arg	Lys 500	Val	Asn	Pro	Val	Glu 505	Leu	Ile	Ile	Ala	Pro 510	Arg	Leu
Ala	Pro	Glu 515	Leu	Tyr											
<210 <213 <213 <213	0> SI 1> LI 2> T 3> OI	EQ II ENGTH ZPE : RGANI	D NO H: 5: PRT ISM:	15 17 Nico	otia	na ta	abacı	ım							
<40	0> SI	EQUEI	NCE :	15											
Met 1	Leu	Ser	Pro	Ile 5	Glu	Ala	Ile	Val	Gly 10	Leu	Val	Thr	Phe	Thr 15	Phe
Leu	Phe	Phe	Phe 20	Leu	Trp	Thr	Lys	Lуз 25	Ser	Gln	ГАЗ	Pro	Ser 30	Lys	Pro
Leu	Pro	Pro 35	ГЛЗ	Ile	Pro	Gly	Gly 40	Trp	Pro	Val	Ile	Gly 45	His	Leu	Phe
His	Phe 50	Asn	Asp	Asp	Gly	Asp 55	Asp	Arg	Pro	Leu	Ala 60	Arg	Гла	Leu	Gly
Asp 65	Leu	Ala	Asp	ГАз	Tyr 70	Gly	Pro	Val	Phe	Thr 75	Phe	Arg	Leu	Gly	Leu 80
Pro	Leu	Val	Leu	Val 85	Val	Ser	Ser	Tyr	Glu 90	Ala	Val	ГЛа	Asp	Сув 95	Phe
Ser	Thr	Asn	Asp 100	Ala	Ile	Phe	Ser	Asn 105	Arg	Pro	Ala	Phe	Leu 110	Tyr	Gly
Asp	Tyr	Leu 115	Gly	Tyr	Asn	Asn	Ala 120	Met	Leu	Phe	Leu	Ala 125	Asn	Tyr	Gly
Pro	Tyr 130	Trp	Arg	ГЛа	Asn	Arg 135	Lys	Leu	Val	Ile	Gln 140	Glu	Val	Leu	Ser

-continued

Ala 145	Ser	Arg	Leu	Glu	Lys 150	Phe	Lys	His	Val	Arg 155	Phe	Ala	Arg	Ile	Gln 160
Ala	Ser	Ile	Lys	Asn 165	Leu	Tyr	Thr	Arg	Ile 170	Asp	Gly	Asn	Ser	Ser 175	Thr
Ile	Asn	Leu	Thr 180	Asp	Trp	Leu	Glu	Glu 185	Leu	Asn	Phe	Gly	Leu 190	Ile	Val
Lys	Met	Ile 195	Ala	Gly	Lys	Asn	Tyr 200	Glu	Ser	Gly	Lys	Gly 205	Asp	Glu	Gln
Val	Glu 210	Arg	Phe	Lys	Lys	Ala 215	Phe	Lys	Asp	Phe	Met 220	Ile	Leu	Ser	Met
Glu 225	Phe	Val	Leu	Trp	Asp 230	Ala	Phe	Pro	Ile	Pro 235	Leu	Phe	ГЛа	Trp	Val 240
Asp	Phe	Gln	Gly	His 245	Val	ГЛа	Ala	Met	Lys 250	Arg	Thr	Phe	ГЛа	Asp 255	Ile
Asp	Ser	Val	Phe 260	Gln	Asn	Trp	Leu	Glu 265	Glu	His	Ile	Asn	Lys 270	Arg	Glu
ГЛа	Met	Glu 275	Val	Asn	Ala	Glu	Gly 280	Asn	Glu	Gln	Asp	Phe 285	Ile	Asp	Val
Val	Leu 290	Ser	Lys	Met	Ser	Asn 295	Glu	Tyr	Leu	Gly	Glu 300	Gly	Tyr	Ser	Arg
Asp 305	Thr	Val	Ile	Lys	Ala 310	Thr	Val	Phe	Ser	Leu 315	Val	Leu	Asp	Ala	Ala 320
Asp	Thr	Val	Ala	Leu 325	His	Ile	Asn	Trp	Gly 330	Met	Ala	Leu	Leu	Ile 335	Asn
Asn	Gln	Lys	Ala 340	Leu	Thr	Lys	Ala	Gln 345	Glu	Glu	Ile	Asp	Thr 350	Lys	Val
Gly	Lys	Asp 355	Arg	Trp	Val	Glu	Glu 360	Ser	Asp	Ile	Asn	Asp 365	Leu	Val	Tyr
Leu	Gln 370	Ala	Ile	Val	Lys	Glu 375	Val	Leu	Arg	Leu	Tyr 380	Pro	Pro	Gly	Pro
Leu 385	Leu	Val	Pro	His	Glu 390	Asn	Val	Glu	Asp	Cys 395	Val	Val	Ser	Gly	Tyr 400
His	Ile	Pro	Lys	Gly 405	Thr	Arg	Leu	Phe	Ala 410	Asn	Val	Met	Lys	Leu 415	Gln
Arg	Asp	Pro	Lys 420	Leu	Trp	Ser	Asp	Pro 425	Asp	Thr	Phe	Asp	Pro 430	Glu	Arg
Phe	Ile	Ala 435	Thr	Asp	Ile	Asb	Phe 440	Arg	Gly	Gln	Tyr	Tyr 445	Lys	Tyr	Ile
Pro	Phe 450	Gly	Ser	Gly	Arg	Arg 455	Ser	Cys	Pro	Gly	Met 460	Thr	Tyr	Ala	Leu
Gln 465	Val	Glu	His	Leu	Thr 470	Met	Ala	His	Leu	Ile 475	Gln	Gly	Phe	Asn	Tyr 480
Arg	Thr	Pro	Asn	Asp 485	Glu	Pro	Leu	Asp	Met 490	Lys	Glu	Gly	Ala	Gly 495	Ile
Thr	Ile	Arg	Lys 500	Val	Asn	Pro	Val	Glu 505	Leu	Ile	Ile	Ala	Pro 510	Arg	Leu
Ala	Pro	Glu 515	Leu	Tyr											
<210)> SI	EQ II	NO	16											

<211> LENGTH: 517
<211> TYPE: PRT
<213> ORGANISM: Nicotiana tabacum

<400)> SE	EQUEL	ICE :	16											
Met 1	Leu	Ser	Pro	Ile 5	Glu	Ala	Ile	Val	Gly 10	Leu	Val	Thr	Phe	Thr 15	Phe
Leu	Phe	Phe	Phe 20	Leu	Trp	Thr	Lys	Lys 25	Ser	Gln	Lys	Pro	Ser 30	Lys	Pro
Leu	Pro	Pro 35	Lys	Ile	Leu	Gly	Gly 40	Trp	Pro	Val	Ile	Gly 45	His	Leu	Phe
His	Phe 50	Asn	Asp	Asp	Gly	Asp 55	Asp	Arg	Pro	Leu	Ala 60	Arg	Lys	Leu	Gly
Asp 65	Leu	Ala	Asp	Lys	Tyr 70	Gly	Pro	Val	Phe	Thr 75	Phe	Arg	Leu	Gly	Leu 80
Pro	Leu	Val	Leu	Val 85	Val	Ser	Ser	Tyr	Glu 90	Ala	Val	LÀa	Asp	Суз 95	Phe
Ser	Thr	Asn	Asp 100	Ala	Ile	Phe	Ser	Asn 105	Arg	Pro	Ala	Phe	Leu 110	Tyr	Gly
Aap	Tyr	Leu 115	Gly	Tyr	Asn	Asn	Ala 120	Met	Leu	Phe	Leu	Ala 125	Asn	Tyr	Gly
Pro	Tyr 130	Trp	Arg	ГÀа	Asn	Arg 135	Lys	Leu	Val	Ile	Gln 140	Glu	Val	Leu	Ser
Ala 145	Ser	Arg	Leu	Glu	Lys 150	Phe	Lys	His	Val	Arg 155	Phe	Ala	Arg	Ile	Gln 160
Ala	Ser	Ile	Lys	Asn 165	Leu	Tyr	Thr	Arg	Ile 170	Asp	Gly	Asn	Ser	Ser 175	Thr
Ile	Asn	Leu	Thr 180	Asp	Trp	Leu	Glu	Glu 185	Leu	Asn	Phe	Gly	Leu 190	Ile	Val
Lys	Met	Ile 195	Ala	Gly	Lys	Asn	Tyr 200	Glu	Ser	Gly	Lys	Gly 205	Asp	Glu	Gln
Val	Glu 210	Arg	Phe	Lys	Lys	Ala 215	Phe	Lys	Asp	Phe	Met 220	Ile	Leu	Ser	Met
Glu 225	Phe	Val	Leu	Trp	Asp 230	Ala	Phe	Pro	Ile	Pro 235	Leu	Phe	Lys	Trp	Val 240
Aap	Phe	Gln	Gly	His 245	Val	LÀa	Ala	Met	Lys 250	Arg	Thr	Phe	Lys	Asp 255	Ile
Aap	Ser	Val	Phe 260	Gln	Asn	Trp	Leu	Glu 265	Glu	His	Ile	Asn	Lys 270	Arg	Glu
ГЛа	Met	Glu 275	Val	Asn	Ala	Glu	Gly 280	Asn	Glu	Gln	Asp	Phe 285	Ile	Asp	Val
Val	Leu 290	Ser	ГЛа	Met	Ser	Asn 295	Glu	Tyr	Leu	Gly	Glu 300	Gly	Tyr	Ser	Arg
Asp 305	Thr	Val	Ile	ГÀа	Ala 310	Thr	Val	Phe	Ser	Leu 315	Val	Leu	Asp	Ala	Ala 320
Asp	Thr	Val	Ala	Leu 325	His	Ile	Asn	Trp	Gly 330	Met	Ala	Leu	Leu	Ile 335	Asn
Asn	Gln	Lys	Ala 340	Leu	Thr	Lys	Ala	Gln 345	Glu	Glu	Ile	Asp	Thr 350	Lys	Val
Gly	Lys	Asp 355	Arg	Trp	Val	Glu	Glu 360	Ser	Asp	Ile	Lys	Asp 365	Leu	Val	Tyr
Leu	Gln 370	Ala	Ile	Val	Lys	Glu 375	Val	Leu	Arg	Leu	Tyr 380	Pro	Pro	Gly	Pro
Leu 385	Leu	Val	Pro	His	Glu 390	Asn	Val	Glu	Asp	Сув 395	Val	Val	Ser	Gly	Tyr 400
His	Ile	Pro	Lys	Gly 405	Thr	Arg	Leu	Phe	Ala 410	Asn	Val	Met	Lys	Leu 415	Gln

-continued

Arg Asp Pro Lys Leu Trp Ser Asp Pro Asp Thr Phe Asp Pro Glu Arg Phe Ile Ala Thr Asp Ile Asp Phe Arg Gly Gln Tyr Tyr Lys Tyr Ile Pro Phe Gly Ser Gly Arg Arg Ser Cys Pro Gly Met Thr Tyr Ala Leu Gln Val Glu His Leu Thr Met Ala His Leu Ile Gln Gly Phe Asn Tyr Arg Thr Pro Asn Asp Glu Pro Leu Asp Met Lys Glu Gly Ala Gly Ile Thr Ile Arg Lys Val Asn Pro Val Glu Leu Ile Ile Ala Pro Arg Leu Ala Pro Glu Leu Tyr <210> SEQ ID NO 17 <211> LENGTH: 517 <212> TYPE: PRT <213> ORGANISM: Nicotiana tabacum <400> SEQUENCE: 17 Met Leu Ser Pro Ile Glu Ala Ile Val Gly Leu Val Thr Phe Thr Phe Leu Phe Phe Leu Trp Thr Lys Lys Ser Gln Lys Pro Ser Lys Pro 20 25 30 Leu Pro Pro Lys Ile Pro Gly Gly Trp Pro Val Ile Gly His Leu Phe His Phe Asn Asp Asp Gly Asp Asp Arg Pro Leu Ala Arg Lys Leu Gly Asp Leu Ala Asp Lys Tyr Gly Pro Val Phe Thr Phe Arg Leu Gly Leu Pro Leu Val Leu Val Val Ser Ser Tyr Glu Ala Val Lys Asp Cys Phe Ser Thr Asn Asp Ala Ile Phe Ser Asn Arg Pro Ala Phe Leu Tyr Gly Asp Tyr Leu Gly Tyr Asn Asn Ala Met Leu Phe Leu Ala Asn Tyr Gly Pro Tyr Trp Arg Lys Asn Arg Lys Leu Val Ile Gln Glu Val Leu Ser Ala Ser Arg Leu Glu Lys Phe Lys His Val Arg Phe Ala Arg Ile Gln Ala Ser Ile Lys Asn Leu Tyr Thr Arg Ile Asp Gly Asn Ser Ser Thr Ile Asn Leu Thr Asp Trp Leu Glu Glu Leu Asn Phe Gly Leu Ile Val Lys Met Ile Ala Gly Lys Asn Tyr Lys Ser Gly Lys Gly Asp Glu Gln Val Glu Arg Phe Lys Lys Ala Phe Lys Asp Phe Met Ile Leu Ser Met Glu Phe Val Leu Trp Asp Ala Phe Pro Ile Pro Leu Phe Lys Trp Val Asp Phe Gln Gly His Val Lys Ala Met Lys Arg Thr Phe Lys Asp Ile

Asp Ser Val Phe Gln Asn Trp Leu Glu Glu His Ile Asn Lys Arg Glu

_														C 111	ucu	
_	_	_	_	260	_	_	_	_	265	_	_	_	_	270	_	_
\mathbf{L}_{j}	уs	Met	Glu 275	Val	Asn	Ala	Glu	Gly 280	Asn	Glu	Gln	Asp	Phe 285	Ile	Asp	Val
v	al	Leu 290	Ser	Lys	Met	Ser	Asn 295	Glu	Tyr	Leu	Gly	Glu 300	Gly	Tyr	Ser	Arg
А: З	sp 05	Thr	Val	Ile	Lys	Ala 310	Thr	Val	Phe	Ser	Leu 315	Val	Leu	Asp	Ala	Ala 320
A	ap	Thr	Val	Ala	Leu 325	His	Ile	Asn	Trp	Gly 330	Met	Ala	Leu	Leu	Ile 335	Asn
A	sn	Gln	Гла	Ala 340	Leu	Thr	Lys	Ala	Gln 345	Glu	Glu	Ile	Asp	Thr 350	Гла	Val
G	ly	Lys	Asp 355	Arg	Trp	Val	Glu	Glu 360	Ser	Asp	Ile	ГЛа	Asp 365	Leu	Val	Tyr
L	eu	Gln 370	Ala	Ile	Val	Lys	Glu 375	Val	Leu	Arg	Leu	Tyr 380	Pro	Pro	Gly	Pro
L- 3:	eu 85	Leu	Val	Pro	His	Glu 390	Asn	Val	Glu	Asp	Суз 395	Val	Val	Ser	Gly	Tyr 400
н	is	Ile	Pro	ГЛа	Gly 405	Thr	Arg	Leu	Phe	Ala 410	Asn	Val	Met	ГЛа	Leu 415	Gln
A:	rg	Asp	Pro	Lys 420	Leu	Trp	Ser	Asp	Pro 425	Asp	Thr	Phe	Asp	Pro 430	Glu	Arg
Pl	he	Ile	Ala 435	Thr	Asp	Ile	Asp	Phe 440	Arg	Gly	Gln	Tyr	Tyr 445	Lys	Tyr	Ile
P	ro	Phe 450	Gly	Ser	Gly	Arg	Arg 455	Ser	Сүз	Pro	Gly	Met 460	Thr	Tyr	Ala	Leu
G 4	ln 65	Val	Glu	His	Leu	Thr 470	Met	Ala	His	Leu	Ile 475	Gln	Gly	Phe	Asn	Tyr 480
A	rg	Thr	Pro	Asn	Asp 485	Glu	Pro	Leu	Asp	Met 490	Lys	Glu	Gly	Ala	Gly 495	Ile
T	hr	Ile	Arg	Lys	Val	Asn	Pro	Val	Glu	Leu	Ile	Ile	Ala	Pro	Arg	Leu
A	la	Pro	Glu	Leu	Tyr				503					210		
			515													
< : < :	210 211)> SH L> LH	EQ II ENGTH) NO H: 5:	18 17											
< <	212 213	2> TY 3> OF	(PE : RGAN	PRT [SM:	Nico	otia	na ta	abacı	ım							
<-	400)> SH	EQUEI	ICE :	18											
M 1	et	Leu	Ser	Pro	Ile 5	Glu	Ala	Ile	Val	Gly 10	Leu	Val	Thr	Phe	Thr 15	Phe
L	eu	Phe	Phe	Phe 20	Leu	Trp	Thr	Lys	Lys 25	Ser	Gln	Lys	Pro	Ser 30	Lys	Pro
L	eu	Pro	Pro 35	Lys	Ile	Pro	Gly	Gly 40	Trp	Pro	Val	Ile	Gly 45	His	Leu	Phe
н	is	Phe 50	Asn	Asp	Asp	Gly	Asp 55	Asp	Arg	Pro	Leu	Ala 60	Arg	Lys	Leu	Gly
A	ab	Leu	Ala	Asp	ГЛа	Tyr	Gly	Pro	Val	Phe	Thr	Phe	Arg	Leu	Gly	Leu
ь: P:	ro	Leu	Val	Leu	Val	70 Val	Ser	Ser	Tyr	Glu	/5 Ala	Val	Lys	Asp	Cys	Phe
S	er	Thr	Asn	Asp	85 Ala	Ile	Phe	Ser	Asn	90 Arg	Pro	Ala	Phe	Leu	95 Tyr	Gly
				100	-				105	5		-		110	-	-

Asp	Tyr	Leu 115	Gly	Tyr	Asn	Asn	Ala 120	Met	Leu	Phe	Leu	Ala 125	Asn	Tyr	Gly
Pro	Tyr 130	Trp	Arg	Гүз	Asn	Arg 135	Lys	Leu	Val	Ile	Gln 140	Glu	Val	Leu	Ser
Ala 145	Ser	Arg	Leu	Glu	Lys 150	Phe	Lys	His	Val	Arg 155	Phe	Ala	Arg	Ile	Gln 160
Ala	Ser	Ile	Lys	Asn 165	Leu	Tyr	Thr	Gln	Ile 170	Asp	Gly	Asn	Ser	Ser 175	Thr
Ile	Asn	Leu	Thr 180	Asp	Trp	Leu	Glu	Glu 185	Leu	Asn	Phe	Gly	Leu 190	Ile	Val
Lys	Met	Ile 195	Ala	Gly	Гла	Asn	Tyr 200	Glu	Ser	Gly	Lys	Gly 205	Asp	Glu	Gln
Val	Glu 210	Arg	Phe	ГЛа	ГЛа	Ala 215	Phe	Lys	Asp	Phe	Met 220	Ile	Leu	Ser	Met
Glu 225	Phe	Val	Leu	Trp	Asp 230	Ala	Phe	Pro	Ile	Pro 235	Leu	Phe	Гла	Trp	Val 240
Asp	Phe	Gln	Gly	His 245	Val	Гла	Ala	Met	Lys 250	Arg	Thr	Phe	Lys	Asp 255	Ile
Asp	Ser	Val	Phe 260	Gln	Asn	Trp	Leu	Glu 265	Glu	His	Ile	Asn	Lys 270	Arg	Glu
Lys	Met	Glu 275	Val	Asn	Ala	Glu	Gly 280	Asn	Glu	Gln	Asp	Phe 285	Ile	Asp	Val
Val	Leu 290	Ser	Lys	Met	Ser	Asn 295	Glu	Tyr	Leu	Gly	Glu 300	Gly	Tyr	Ser	Arg
Asp 305	Thr	Val	Ile	ГАЗ	Ala 310	Thr	Val	Phe	Ser	Leu 315	Val	Leu	Asp	Ala	Ala 320
Asp	Thr	Val	Ala	Leu 325	His	Ile	Asn	Trp	Gly 330	Met	Ala	Leu	Leu	Ile 335	Asn
Asn	Gln	Lys	Ala 340	Leu	Thr	Lys	Ala	Gln 345	Glu	Glu	Ile	Asp	Thr 350	Lys	Val
Gly	ГЛа	Asp 355	Arg	Trp	Val	Glu	Glu 360	Ser	Asp	Ile	ГЛа	Asp 365	Leu	Val	Tyr
Leu	Gln 370	Ala	Ile	Val	Гλа	Glu 375	Val	Leu	Arg	Leu	Tyr 380	Pro	Pro	Gly	Pro
Leu 385	Leu	Val	Pro	His	Glu 390	Asn	Val	Glu	Asp	Сув 395	Val	Val	Ser	Gly	Tyr 400
His	Ile	Pro	Lys	Gly 405	Thr	Arg	Leu	Phe	Ala 410	Asn	Val	Met	Гλа	Leu 415	Gln
Arg	Asp	Pro	Lys 420	Leu	Trp	Ser	Asp	Pro 425	Asp	Thr	Phe	Asp	Pro 430	Glu	Arg
Phe	Ile	Ala 435	Thr	Asp	Ile	Asp	Phe 440	Arg	Gly	Gln	Tyr	Tyr 445	Lys	Tyr	Ile
Pro	Phe 450	Gly	Ser	Gly	Arg	Arg 455	Ser	Сув	Pro	Gly	Met 460	Thr	Tyr	Ala	Leu
Gln 465	Val	Glu	His	Leu	Thr 470	Met	Ala	His	Leu	Ile 475	Gln	Gly	Phe	Asn	Tyr 480
Arg	Thr	Pro	Asn	Asp 485	Glu	Pro	Leu	Asp	Met 490	Lys	Glu	Gly	Ala	Gly 495	Ile
Thr	Ile	Arg	Lys 500	Val	Asn	Pro	Val	Glu 505	Leu	Ile	Ile	Ala	Pro 510	Arg	Leu
Ala	Pro	Glu 515	Leu	Tyr											

<210 <211)> SH L> LH	EQ II ENGTH) NO H: 51	19 17											
<212 <213	2> TY 3> OF	(PE : RGANI	PRT [SM:	Nico	otiar	na ta	abacu	ım							
<400)> SI	EQUEI	ICE :	19											
Met 1	Leu	Ser	Pro	Ile 5	Glu	Ala	Ile	Val	Gly 10	Leu	Val	Thr	Phe	Thr 15	Phe
Leu	Phe	Phe	Phe 20	Leu	Trp	Thr	Lys	Lys 25	Ser	Gln	Lys	Pro	Ser 30	Lys	Pro
Leu	Pro	Pro 35	Lys	Ile	Pro	Gly	Gly 40	Trp	Pro	Val	Ile	Gly 45	His	Leu	Phe
His	Phe 50	Asn	Asp	Asp	Gly	Asp 55	Asp	Arg	Pro	Leu	Ala 60	Arg	Lys	Leu	Gly
Asp 65	Leu	Ala	Asp	Lys	Tyr 70	Gly	Pro	Val	Phe	Thr 75	Phe	Arg	Leu	Gly	Leu 80
Pro	Leu	Val	Leu	Val 85	Val	Ser	Ser	Tyr	Glu 90	Ala	Val	Lys	Asp	Сув 95	Phe
Ser	Thr	Asn	Asp 100	Ala	Ile	Phe	Ser	Asn 105	Arg	Pro	Ala	Phe	Leu 110	Tyr	Gly
Asp	Tyr	Leu 115	Gly	Tyr	Asn	Asn	Ala 120	Met	Leu	Phe	Leu	Ala 125	Asn	Tyr	Gly
Pro	Tyr 130	Trp	Arg	Lys	Asn	Arg 135	Lys	Leu	Val	Ile	Gln 140	Glu	Val	Leu	Ser
Ala 145	Ser	Arg	Leu	Glu	Lys 150	Phe	Lys	His	Val	Arg 155	Phe	Ala	Arg	Ile	Gln 160
Ala	Ser	Ile	Lys	Asn 165	Leu	Tyr	Thr	Arg	Ile 170	Asp	Gly	Asn	Ser	Ser 175	Thr
Ile	Asn	Leu	Thr 180	Asp	Trp	Leu	Glu	Glu 185	Leu	Asn	Phe	Gly	Leu 190	Ile	Val
Lys	Met	Ile 195	Ala	Gly	Lys	Asn	Tyr 200	Glu	Ser	Gly	Lys	Gly 205	Asp	Glu	Gln
Val	Glu 210	Arg	Phe	Lys	Lys	Ala 215	Phe	Lys	Asp	Phe	Met 220	Ile	Leu	Ser	Met
Glu 225	Phe	Val	Leu	Trp	Asp 230	Ala	Phe	Pro	Ile	Pro 235	Leu	Phe	Lys	Trp	Val 240
Asp	Phe	Gln	Gly	His 245	Val	ГЛа	Ala	Met	Lys 250	Arg	Thr	Phe	Lys	Asp 255	Ile
Asp	Ser	Val	Phe 260	Gln	Asn	Trp	Leu	Glu 265	Glu	His	Ile	Asn	Lys 270	Arg	Glu
LÀa	Met	Glu 275	Val	Asn	Ala	Glu	Gly 280	Asn	Glu	Gln	Asp	Phe 285	Ile	Asp	Val
Val	Leu 290	Ser	Lys	Met	Ser	Asn 295	Glu	Tyr	Leu	Gly	Glu 300	Gly	Tyr	Ser	Arg
Asp 305	Thr	Val	Ile	Lys	Ala 310	Thr	Val	Phe	Ser	Leu 315	Val	Leu	Asp	Ala	Ala 320
Asp	Thr	Val	Ala	Leu 325	His	Ile	Asn	Trp	Gly 330	Met	Ala	Leu	Leu	Ile 335	Asn
Asn	Gln	Lys	Ala 340	Leu	Thr	Lys	Ala	Gln 345	Glu	Glu	Ile	Asp	Thr 350	Lys	Val
Gly	Lys	Asp 355	Arg	Trp	Val	Glu	Glu 360	Ser	Asp	Ile	Гла	Asp 365	Leu	Val	Tyr
Leu	Gln 370	Ala	Ile	Val	ГЛа	Glu 375	Val	Leu	Arg	Leu	Tyr 380	Pro	Pro	Gly	Pro

-continued

Leu 385	. Leu	Val	Pro	His	Glu 390	Asn	Val	Glu	Asp	Сув 395	Val	Val	Ser	Gly	Tyr 400
His	Ile	Pro	Lys	Gly 405	Thr	Arg	Leu	Phe	Ala 410	Asn	Val	Met	Lys	Leu 415	Gln
Arg	Asp	Pro	Lys 420	Leu	Trp	Ser	Asp	Pro 425	Asp	Thr	Phe	Asp	Pro 430	Glu	Arg
Phe	Ile	Ala 435	Thr	Asp	Ile	Asp	Phe 440	Arg	Gly	Gln	Tyr	Tyr 445	Lys	Tyr	Ile
Pro	Phe 450	Gly	Ser	Gly	Arg	Arg 455	Ser	Cys	Pro	Arg	Met 460	Thr	Tyr	Ala	Leu
Gln 465	Val	Glu	His	Leu	Thr 470	Met	Ala	His	Leu	Ile 475	Gln	Gly	Phe	Asn	Tyr 480
Arg	Thr	Pro	Asn	Asp 485	Glu	Pro	Leu	Asp	Met 490	Lys	Glu	Gly	Ala	Gly 495	Ile
Thr	Ile	Arg	Lys 500	Val	Asn	Pro	Val	Glu 505	Leu	Ile	Ile	Ala	Pro 510	Arg	Leu
Ala	Pro	Glu 515	Leu	Tyr											
<21 <21 <21 <21	0> SI 1> LI 2> T 3> OI	EQ II ENGTI YPE : RGANI	D NO H: 5: PRT ISM:	20 17 Nico	otia	na ta	abacı	ım							
<40	0> SI	EQUEI	NCE :	20											
Met 1	Leu	Ser	Pro	Ile 5	Glu	Ala	Ile	Val	Gly 10	Leu	Val	Thr	Phe	Thr 15	Phe
Leu	. Phe	Phe	Phe 20	Leu	Trp	Thr	Lys	Lys 25	Ser	Gln	ГЛЗ	Pro	Ser 30	Lys	Pro
Leu	. Pro	Pro 35	ГЛа	Ile	Pro	Gly	Gly 40	Trp	Pro	Val	Ile	Gly 45	His	Leu	Phe
His	Phe 50	Asn	Asp	Asp	Gly	Asp 55	Asp	Arg	Pro	Leu	Ala 60	Arg	Lys	Leu	Gly
Asp 65	Leu	Ala	Asp	ГЛЗ	Tyr 70	Gly	Pro	Val	Phe	Thr 75	Phe	Arg	Leu	Gly	Leu 80
Pro	Leu	Val	Leu	Val 85	Val	Ser	Ser	Tyr	Glu 90	Ala	Val	ГЛа	Asp	Суз 95	Phe
Ser	Thr	Asn	Asp 100	Ala	Ile	Phe	Ser	Asn 105	Arg	Pro	Ala	Phe	Leu 110	Tyr	Gly
Aap	Tyr	Leu 115	Gly	Tyr	Asn	Asn	Ala 120	Met	Leu	Phe	Leu	Ala 125	Asn	Tyr	Gly
Pro	Tyr 130	Trp	Arg	ГЛа	Asn	Arg 135	ГЛа	Leu	Val	Ile	Gln 140	Glu	Val	Leu	Ser
Ala 145	Ser	Arg	Leu	Glu	Lys 150	Phe	Lys	His	Val	Arg 155	Phe	Ala	Arg	Ile	Gln 160
Ala	Ser	Ile	Lys	Asn 165	Leu	Tyr	Thr	Arg	Ile 170	Asp	Gly	Asn	Ser	Ser 175	Thr
Ile	Asn	Leu	Thr 180	Asp	Trp	Leu	Glu	Glu 185	Leu	Asn	Phe	Gly	Leu 190	Ile	Val
Lys	Met	Ile 195	Ala	Gly	Гла	Asn	Tyr 200	Glu	Ser	Gly	Lys	Gly 205	Asp	Glu	Gln
Val	Glu 210	Arg	Phe	ГЛа	ГЛа	Ala 215	Phe	Lys	Asp	Phe	Met 220	Ile	Leu	Ser	Met
Glu 225	Phe	Val	Leu	Trp	Asp 230	Ala	Phe	Pro	Ile	Pro 235	Leu	Phe	ГЛа	Trp	Val 240

Asp	Phe	Gln	Gly	His 245	Val	Lys	Ala	Met	Lys 250	Arg	Thr	Phe	Lys	Asp 255	Ile
Asp	Ser	Val	Phe 260	Gln	Asn	Trp	Leu	Glu 265	Glu	His	Ile	Asn	Lys 270	Arg	Glu
Lys	Met	Glu 275	Val	Asn	Ala	Glu	Gly 280	Asn	Glu	Gln	Asp	Phe 285	Ile	Asp	Val
Val	Leu 290	Ser	Lys	Met	Ser	Asn 295	Lys	Tyr	Leu	Gly	Glu 300	Gly	Tyr	Ser	Arg
Asp 305	Thr	Val	Ile	Гла	Ala 310	Thr	Val	Phe	Ser	Leu 315	Val	Leu	Asp	Ala	Ala 320
Asp	Thr	Val	Ala	Leu 325	His	Ile	Asn	Trp	Gly 330	Met	Ala	Leu	Leu	Ile 335	Asn
Asn	Gln	Гла	Ala 340	Leu	Thr	Lys	Ala	Gln 345	Glu	Glu	Ile	Asp	Thr 350	Lys	Val
Gly	Lys	Asp 355	Arg	Trp	Val	Glu	Glu 360	Ser	Asp	Ile	Гла	Asp 365	Leu	Val	Tyr
Leu	Gln 370	Ala	Ile	Val	Lys	Glu 375	Val	Leu	Arg	Leu	Tyr 380	Pro	Pro	Gly	Pro
Leu 385	Leu	Val	Pro	His	Glu 390	Asn	Val	Glu	Asp	Сув 395	Val	Val	Ser	Gly	Tyr 400
His	Ile	Pro	ГЛа	Gly 405	Thr	Arg	Leu	Phe	Ala 410	Asn	Val	Met	Гла	Leu 415	Gln
Arg	Asp	Pro	Lys 420	Leu	Trp	Ser	Asp	Pro 425	Asp	Thr	Phe	Asp	Pro 430	Glu	Arg
Phe	Ile	Ala 435	Thr	Asp	Ile	Asp	Phe 440	Arg	Gly	Gln	Tyr	Tyr 445	Lys	Tyr	Ile
Pro	Phe 450	Gly	Ser	Gly	Arg	Arg 455	Ser	Сүз	Pro	Gly	Met 460	Thr	Tyr	Ala	Leu
Gln 465	Val	Glu	His	Leu	Thr 470	Met	Ala	His	Leu	Ile 475	Gln	Gly	Phe	Asn	Tyr 480
Arg	Thr	Pro	Asn	Asp 485	Glu	Pro	Leu	Asp	Met 490	Lys	Glu	Gly	Ala	Gly 495	Ile
Thr	Ile	Arg	Lys 500	Val	Asn	Pro	Val	Glu 505	Leu	Ile	Ile	Ala	Pro 510	Arg	Leu
Ala	Pro	Glu 515	Leu	Tyr											
<210 <211 <212 <213	0> SH L> LH 2> TY 3> OH	EQ II ENGTH ZPE : RGANI) NO H: 5: PRT [SM:	21 17 Nico	otiar	na ta	abacı	ım							
< 400)> SH	EQUEI	ICE :	21											
Met 1	Leu	Ser	Pro	Ile 5	Glu	Ala	Ile	Val	Gly 10	Leu	Val	Thr	Phe	Thr 15	Phe
Leu	Phe	Phe	Phe 20	Leu	Trp	Thr	Lys	Lys 25	Ser	Gln	Гла	Pro	Ser 30	Lys	Pro
Leu	Pro	Pro 35	Lys	Ile	Pro	Gly	Gly 40	Trp	Pro	Val	Ile	Gly 45	His	Leu	Phe
His	Phe 50	Asn	Asp	Asp	Gly	Asp 55	Asp	Arg	Pro	Leu	Ala 60	Arg	Lys	Leu	Gly
Asp 65	Leu	Ala	Asp	Lys	Tyr 70	Gly	Pro	Val	Phe	Thr 75	Phe	Arg	Leu	Gly	Leu 80
Pro	Leu	Val	Leu	Val	Val	Ser	Ser	Tyr	Glu	Ala	Val	Lys	Asp	Cys	Phe

					85					90					95	
S	er	Thr	Asn	Asp 100	Ala	Ile	Phe	Ser	Asn 105	Arg	Pro	Ala	Phe	Leu 110	Tyr	Gly
A	ab	Tyr	Leu 115	Gly	Tyr	Asn	Asn	Ala 120	Met	Leu	Phe	Leu	Ala 125	Asn	Tyr	Gly
P	ro	Tyr 130	Trp	Arg	ГАЗ	Asn	Arg 135	Lys	Leu	Val	Ile	Gln 140	Glu	Val	Leu	Ser
A 1-	la 45	Ser	Arg	Leu	Glu	Lys 150	Phe	Lys	His	Val	Arg 155	Phe	Ala	Arg	Ile	Gln 160
A	la	Ser	Ile	Гла	Asn 165	Leu	Tyr	Thr	Arg	Ile 170	Asp	Gly	Asn	Ser	Ser 175	Thr
I	le	Asn	Leu	Thr 180	Asp	Trp	Leu	Glu	Glu 185	Leu	Asn	Phe	Gly	Leu 190	Ile	Val
$\mathbf{L}_{\mathbf{j}}$	γя	Met	Ile 195	Ala	Gly	ГЛа	Asn	Tyr 200	Glu	Ser	Gly	ГЛа	Gly 205	Asp	Glu	Gln
v	al	Glu 210	Arg	Phe	ГЛа	ГЛа	Ala 215	Phe	Lys	Asp	Phe	Met 220	Ile	Leu	Ser	Met
G 2	lu 25	Phe	Val	Leu	Trp	Asp 230	Ala	Phe	Pro	Ile	Pro 235	Leu	Phe	ГЛа	Trp	Val 240
A	ab	Phe	Gln	Gly	His 245	Val	Гла	Ala	Met	Lув 250	Arg	Thr	Phe	Lys	Asp 255	Ile
A	ab	Ser	Val	Phe 260	Gln	Asn	Trp	Leu	Glu 265	Glu	His	Ile	Asn	Lys 270	Arg	Glu
Ŀ	үs	Met	Glu 275	Val	Asn	Ala	Glu	Gly 280	Asn	Glu	Gln	Asp	Phe 285	Ile	Asp	Val
v	al	Leu 290	Ser	Lys	Met	Ser	Asn 295	Glu	Tyr	Leu	Gly	Glu 300	Gly	Tyr	Ser	Arg
А: З	ap 05	Thr	Val	Ile	ГЛа	Ala 310	Thr	Val	Phe	Ser	Leu 315	Val	Leu	Asp	Ala	Ala 320
A	ab	Thr	Val	Ala	Leu 325	His	Ile	Asn	Trp	Gly 330	Met	Ala	Leu	Leu	Ile 335	Asn
A	sn	Gln	Lys	Ala 340	Leu	Thr	Lys	Ala	Gln 345	Glu	Glu	Ile	Asp	Thr 350	Lys	Val
G	ly	Lys	Asp 355	Arg	Trp	Val	Glu	Glu 360	Ser	Asp	Ile	Lys	Asp 365	Leu	Val	Tyr
L	eu	Gln 370	Ala	Ile	Val	Гла	Glu 375	Val	Leu	Arg	Leu	Tyr 380	Pro	Pro	Gly	Pro
L- 3	eu 85	Leu	Val	Pro	His	Glu 390	Asn	Val	Glu	Asp	Сув 395	Val	Val	Ser	Gly	Tyr 400
H	is	Ile	Pro	Lys	Gly 405	Thr	Arg	Leu	Phe	Ala 410	Asn	Val	Met	Lys	Leu 415	Gln
A	rg	Asp	Pro	Lys 420	Leu	Trp	Ser	Asp	Pro 425	Asp	Ile	Phe	Asp	Pro 430	Glu	Arg
Pl	he	Ile	Ala 435	Thr	Asp	Ile	Asp	Phe 440	Arg	Gly	Gln	Tyr	Tyr 445	Lys	Tyr	Ile
P	ro	Phe 450	Gly	Ser	Gly	Arg	Arg 455	Ser	Сув	Pro	Gly	Met 460	Thr	Tyr	Ala	Leu
G 4	ln 65	Val	Glu	His	Leu	Thr 470	Met	Ala	His	Leu	Ile 475	Gln	Gly	Phe	Asn	Tyr 480
A	rg	Thr	Pro	Asn	Asp 485	Glu	Pro	Leu	Asp	Met 490	Гуз	Glu	Gly	Ala	Gly 495	Ile
T	hr	Ile	Arg	Lys	Val	Asn	Pro	Val	Glu 505	Leu	Ile	Ile	Ala	Pro	Arg	Leu

Ala Pro Glu Leu Tyr 515

<210)> SE > LE	EQ II ENGTH	0 NO 1: 32	22 28											
<212	2> TY	PE:	PRT	.0											
<213	8> OF	RGANI	SM:	Nico	otiar	na t <i>a</i>	ibacu	ım							
<400)> SE	EQUEN	ICE :	22											
Met 1	Leu	Ser	Pro	Ile 5	Glu	Ala	Ile	Val	Gly 10	Leu	Val	Thr	Phe	Thr 15	Phe
Leu	Phe	Phe	Phe 20	Leu	Trp	Thr	Lys	Lys 25	Ser	Gln	Lys	Pro	Ser 30	Lys	Pro
Leu	Pro	Pro 35	Lys	Ile	Pro	Gly	Gly 40	Trp	Pro	Val	Ile	Gly 45	His	Leu	Phe
His	Phe 50	Asn	Asp	Asp	Gly	Asp 55	Asp	Arg	Pro	Leu	Ala 60	Arg	Lys	Leu	Gly
Asp 65	Leu	Ala	Asp	Lys	Tyr 70	Gly	Pro	Val	Phe	Thr 75	Phe	Arg	Leu	Gly	Leu 80
Pro	Leu	Val	Leu	Val 85	Val	Ser	Ser	Tyr	Glu 90	Ala	Val	Lys	Asp	Суз 95	Phe
Ser	Thr	Asn	Asp 100	Ala	Ile	Phe	Ser	Asn 105	Arg	Pro	Ala	Phe	Leu 110	Tyr	Gly
Asp	Tyr	Leu 115	Gly	Tyr	Asn	Asn	Ala 120	Met	Leu	Phe	Leu	Ala 125	Asn	Tyr	Gly
Pro	Tyr 130	Trp	Arg	Lys	Asn	Arg 135	Lys	Leu	Val	Ile	Gln 140	Glu	Val	Leu	Ser
Ala 145	Ser	Arg	Leu	Glu	Lys 150	Phe	Lys	His	Val	Arg 155	Phe	Ala	Arg	Ile	Gln 160
Ala	Ser	Ile	Lys	Asn 165	Leu	Tyr	Thr	Arg	Ile 170	Asp	Gly	Asn	Ser	Ser 175	Thr
Ile	Asn	Leu	Thr 180	Asp	Trp	Leu	Glu	Glu 185	Leu	Asn	Phe	Gly	Leu 190	Ile	Val
Lys	Met	Ile 195	Ala	Gly	Lys	Asn	Tyr 200	Glu	Ser	Gly	Lys	Gly 205	Asp	Glu	Gln
Val	Glu 210	Arg	Phe	Lys	Lys	Ala 215	Phe	Lys	Asp	Phe	Met 220	Ile	Leu	Ser	Met
Glu 225	Phe	Val	Leu	Trp	Asp 230	Ala	Phe	Pro	Ile	Pro 235	Leu	Phe	Lys	Trp	Val 240
Asp	Phe	Gln	Gly	His 245	Val	Гла	Ala	Met	Lys 250	Arg	Thr	Phe	Lys	Asp 255	Ile
Asp	Ser	Val	Phe 260	Gln	Asn	Trp	Leu	Glu 265	Glu	His	Ile	Asn	Lys 270	Arg	Glu
Lys	Met	Glu 275	Val	Asn	Ala	Glu	Gly 280	Asn	Glu	Gln	Asp	Phe 285	Ile	Asp	Val
Val	Leu 290	Ser	Lys	Met	Ser	Asn 295	Glu	Tyr	Leu	Gly	Glu 300	Gly	Tyr	Ser	Arg
Asp 305	Thr	Val	Ile	Lys	Ala 310	Thr	Val	Phe	Ser	Leu 315	Val	Leu	Asp	Ala	Ala 320
Asp	Thr	Val	Ala	Leu 325	His	Ile	Asn								

<210> SEQ ID NO 23 <211> LENGTH: 517 <212> TYPE: PRT

<213> ORGANISM:	Nicotiana taba	cum	
<400> SEQUENCE:	23		
Met Leu Ser Pro	Ile Glu Ala Il	e Val Gly Leu Val.	Thr Phe Thr Phe
1	5	10	15
Leu Phe Phe Phe	Leu Trp Thr Ly	rs Lys Ser Gln Lys	Pro Ser Lys Pro
20		25	30
Leu Pro Pro Lys	Ile Pro Gly Gl	y Trp Pro Val Ile	Gly His Leu Phe
35	40		45
His Phe Asn Asp	Asp Gly Asp As	p Arg Pro Leu Ala	Arg Lys Leu Gly
50	55	60	
Asp Leu Ala Asp	Lys Tyr Gly Pr	o Val Phe Thr Phe	Arg Leu Gly Leu
65	70	75	80
Pro Leu Val Leu	Val Val Ser Se	er Tyr Glu Ala Val	Lys Asp Cys Phe
	85	90	95
Ser Thr Asn Asp	Ala Ile Phe Se	er Asn Arg Pro Ala	Phe Leu Tyr Gly
100		105	110
Asp Tyr Leu Gly 115	iyr Asn Asn Al 12	a met Leu Phe Leu	Ala Ash Tyr Giy 125 Glu Val Leu Ser
130 Ala Ser Arg Leu	Glu Lys Phe Ly	s Leu val fie Gin 140 zs His Val Arg Phe	Ala Arg Ile Gin
145	150	r Arq Ile Asp Glv	160
Ala Ser Ile Lvs	Asn Leu Tvr Th		Asn Ser Ser Thr
Ile Asn Leu Thr	165	170	175
	Asp Trp Leu Gl	.u Glu Leu Asn Phe	Gly Leu Ile Val
180	Gly Lys Asn Ty	185	190
Lys Met Ile Ala		Yr Glu Ser Gly Lys	Gly Asp Glu Gln
195	20	00	205
Val Glu Arg Phe	Lys Lys Ala Ph	Ne Lys Asp Phe Met	Ile Leu Ser Met
210	215	220	
Glu Phe Val Leu	Trp Asp Ala Ph	e Pro Ile Pro Leu	Phe Lys Trp Val
225	230	235	240
Asp Pne Gin Giy	Als Val Lys Al 245 Gln Len Tro Le	a Met Lys Arg Inr 250 w Clu Clu Hig Ile	255
Lvs Met Glu Val	Asn Ala Glu Gl	265 V Asn Glu Gln Asp	270 Phe Ile Asp Val
275 Val Leu Ser Lvs	Met Ser Asn Gl	u Tyr Leu Glv Glu	285 Gly Tyr Ser Arg
290 Asp Thr Val Tle	295 Lvs Ala Thr Va	1 Phe Ser Leu Val	Leu Asp Ala Ala
305 Asp Thr Val Ala	310	315	Jeu Leu Ile Asp
Ach Cin Lee Die	325	330	335
Ash Gin Lys Ala 340	Deu Inr Lys Al	345	Asp Inr Lys Val 350
GIY LYS Asp Arg	Trp Val Glu Gl	u Ser Asp Ile Lys	Asp Leu Val Tyr
355	36	0	365
Leu Gln Ala Ile	Val Lys Glu Me	t Leu Arg Leu Tyr	Pro Pro Gly Pro
370	375	380	
Leu Leu Val Pro	His Glu Asn Va	il Glu Asp Cys Val	Val Ser Gly Tyr
385	390	395	400

His	Ile	Pro	Lys	Gly 405	Thr	Arg	Leu	Phe	Ala 410	Asn	Val	Met	Lys	Leu 415	Gln
Arg	Asp	Pro	Lys 420	Leu	Trp	Ser	Asp	Pro 425	Asp	Thr	Phe	Asp	Pro 430	Glu	Arg
Phe	Ile	Ala 435	Thr	Asp	Ile	Asp	Phe 440	Arg	Gly	Gln	Tyr	Tyr 445	Lys	Tyr	Ile
Prc	Phe 450	Gly	Ser	Gly	Arg	Arg 455	Ser	Сүз	Pro	Gly	Met 460	Thr	Tyr	Ala	Leu
Gln 465	Val	Glu	His	Leu	Thr 470	Met	Ala	His	Leu	Ile 475	Gln	Gly	Phe	Asn	Tyr 480
Arg	Thr	Pro	Asn	Asp 485	Glu	Pro	Leu	Asp	Met 490	Lys	Glu	Gly	Ala	Gly 495	Ile
Thr	Ile	Arg	Lys 500	Val	Asn	Pro	Val	Glu 505	Leu	Ile	Ile	Ala	Pro 510	Arg	Leu
Ala	Pro	Glu 515	Leu	Tyr											
<21 <21	0> SI 1> Li	EQ II ENGTI	О NO H: 5:	24 17											
<21 <21	2> T 3> 01	YPE : RGANI	PRT ISM:	Nico	otian	na ta	abacı	ım							
< 40	0> SI	EQUEI	NCE :	24											
Met 1	Leu	Ser	Pro	Ile 5	Glu	Ala	Ile	Val	Gly 10	Leu	Val	Thr	Phe	Thr 15	Phe
Leu	. Phe	Phe	Phe 20	Leu	Trp	Thr	Lys	Lys 25	Ser	Gln	ГЛЗ	Pro	Ser 30	Lys	Pro
Leu	Pro	Pro 35	Lys	Ile	Pro	Gly	Gly 40	Trp	Pro	Val	Ile	Gly 45	His	Leu	Phe
His	Phe 50	Asn	Asp	Asp	Gly	Asp 55	Aab	Arg	Pro	Leu	Ala 60	Arg	Lys	Leu	Gly
Asp 65	Leu	Ala	Asp	ГЛа	Tyr 70	Gly	Pro	Val	Phe	Thr 75	Phe	Arg	Leu	Gly	Leu 80
Prc	Leu	Val	Leu	Val 85	Val	Ser	Ser	Tyr	Glu 90	Ala	Val	Гла	Asp	Суз 95	Phe
Ser	Thr	Asn	Asp 100	Ala	Ile	Phe	Ser	Asn 105	Arg	Pro	Ala	Phe	Leu 110	Tyr	Gly
Asp	Tyr	Leu 115	Gly	Tyr	Asn	Asn	Ala 120	Met	Leu	Phe	Leu	Ala 125	Asn	Tyr	Gly
Prc	Tyr 130	Trp	Arg	ГЛа	Asn	Arg 135	Lys	Leu	Val	Ile	Gln 140	Glu	Val	Leu	Ser
Ala 145	Ser	Arg	Leu	Glu	Lys 150	Phe	Гла	His	Val	Arg 155	Phe	Ala	Arg	Ile	Gln 160
Ala	Ser	Ile	Lys	Asn 165	Leu	Tyr	Thr	Arg	Ile 170	Asn	Gly	Asn	Ser	Ser 175	Thr
Ile	Asn	Leu	Thr 180	Asp	Trp	Leu	Glu	Glu 185	Leu	Asn	Phe	Gly	Leu 190	Ile	Val
Lys	Met	Ile 195	Ala	Gly	Гүз	Asn	Tyr 200	Glu	Ser	Gly	Гүз	Gly 205	Asp	Glu	Gln
Val	Glu 210	Arg	Phe	Гла	Гла	Ala 215	Phe	Lys	Asp	Phe	Met 220	Ile	Leu	Ser	Met
Glu 225	Phe	Val	Leu	Trp	Asp 230	Ala	Phe	Pro	Ile	Pro 235	Leu	Phe	Lys	Trp	Val 240
Asp	Phe	Gln	Gly	His 245	Val	ГЛа	Ala	Met	Lys 250	Arg	Thr	Phe	Lys	Asp 255	Ile

Asp	Ser	Val	Phe 260	Gln	Asn	Trp	Leu	Glu 265	Glu	His	Ile	Asn	Lys 270	Arg	Glu
Lys	Met	Glu 275	Val	Asn	Ala	Glu	Gly 280	Asn	Glu	Gln	Asp	Phe 285	Ile	Asp	Val
Val	Leu 290	Ser	Lys	Met	Ser	Asn 295	Glu	Tyr	Leu	Gly	Glu 300	Gly	Tyr	Ser	Arg
Asp 305	Thr	Val	Ile	Гла	Ala 310	Thr	Val	Phe	Ser	Leu 315	Val	Leu	Asp	Ala	Ala 320
Asp	Thr	Val	Ala	Leu 325	His	Ile	Asn	Trp	Gly 330	Met	Ala	Leu	Leu	Ile 335	Asn
Asn	Gln	Lys	Ala 340	Leu	Thr	Lys	Ala	Gln 345	Glu	Glu	Ile	Asp	Thr 350	Lys	Val
Gly	Lys	Asp 355	Arg	Trp	Val	Glu	Glu 360	Ser	Asp	Ile	ГЛЗ	Asp 365	Leu	Val	Tyr
Leu	Gln 370	Ala	Ile	Val	ГÀа	Glu 375	Val	Leu	Arg	Leu	Tyr 380	Pro	Pro	Gly	Pro
Leu 385	Leu	Val	Pro	His	Glu 390	Asn	Val	Glu	Asp	Сув 395	Val	Val	Ser	Gly	Tyr 400
His	Ile	Pro	Гла	Gly 405	Thr	Arg	Leu	Phe	Ala 410	Asn	Val	Met	Lys	Leu 415	Gln
Arg	Asp	Pro	Lys 420	Leu	Trp	Ser	Asp	Pro 425	Asp	Thr	Phe	Asp	Pro 430	Glu	Arg
Phe	Ile	Ala 435	Thr	Asp	Ile	Asp	Phe 440	Arg	Gly	Gln	Tyr	Tyr 445	Lys	Tyr	Ile
Pro	Phe 450	Gly	Ser	Gly	Arg	Arg 455	Ser	Сүз	Pro	Gly	Met 460	Thr	Tyr	Ala	Leu
Gln 465	Val	Glu	His	Leu	Thr 470	Met	Ala	His	Leu	Ile 475	Gln	Gly	Phe	Asn	Tyr 480
Arg	Thr	Pro	Asn	Asp 485	Glu	Pro	Leu	Asp	Met 490	Lys	Glu	Gly	Ala	Gly 495	Ile
Thr	Ile	Arg	Lys 500	Val	Asn	Pro	Val	Glu 505	Leu	Ile	Ile	Ala	Pro 510	Arg	Leu
Ala	Pro	Glu 515	Leu	Tyr											
<210 <211 <211 <211	0> SH L> LH 2> TY 3> OH	EQ II ENGTH ZPE : RGANI) NO H: 5: PRT ISM:	25 17 Nico	otiar	na ta	abacı	ım							
<40	D> SI	EQUEI	ICE :	25											
Met 1	Leu	Ser	Pro	Ile 5	Glu	Ala	Ile	Val	Gly 10	Leu	Val	Thr	Phe	Thr 15	Phe
Leu	Phe	Phe	Phe 20	Leu	Trp	Thr	Lys	Lув 25	Ser	Gln	ГЛа	Pro	Ser 30	Lys	Pro
Leu	Pro	Pro 35	Lys	Ile	Pro	Gly	Gly 40	Trp	Pro	Val	Ile	Gly 45	His	Leu	Phe
His	Phe 50	Asn	Asp	Asp	Gly	Asp 55	Asp	Arg	Pro	Leu	Ala 60	Arg	Lys	Leu	Gly
Asp 65	Leu	Ala	Asp	Гүз	Tyr 70	Gly	Pro	Val	Phe	Thr 75	Phe	Arg	Leu	Gly	Leu 80
Pro	Leu	Val	Leu	Val 85	Val	Ser	Ser	Tyr	Glu 90	Ala	Val	Гла	Asp	Сув 95	Phe
Ser	Thr	Asn	Asp	Ala	Ile	Phe	Ser	Asn	Arg	Pro	Ala	Phe	Leu	Tyr	Gly

			100					105					110		
Asp	Tyr	Leu 115	Gly	Tyr	Asn	Asn	Ala 120	Met	Leu	Phe	Leu	Ala 125	Asn	Tyr	Gly
Pro	Tyr 130	Trp	Arg	ГЛа	Asn	Arg 135	Lys	Leu	Val	Ile	Gln 140	Glu	Val	Leu	Ser
Ala 145	Ser	Arg	Leu	Glu	Lys 150	Phe	Lys	His	Val	Arg 155	Phe	Ala	Arg	Ile	Gln 160
Ala	Ser	Ile	Lys	Asn 165	Leu	Tyr	Thr	Arg	Ile 170	Asp	Gly	Asn	Ser	Ser 175	Thr
Ile	Asn	Leu	Thr 180	Asp	Trp	Leu	Glu	Glu 185	Leu	Asn	Phe	Gly	Leu 190	Ile	Val
ГЛа	Met	Ile 195	Ala	Gly	ГЛЗ	Asn	Tyr 200	Glu	Ser	Gly	ГЛа	Gly 205	Asp	Glu	Gln
Val	Glu 210	Arg	Phe	Lys	Lys	Ala 215	Phe	Lys	Asp	Phe	Met 220	Ile	Leu	Ser	Met
Glu 225	Phe	Val	Leu	Trp	Asp 230	Ala	Phe	Pro	Ile	Pro 235	Leu	Phe	Lys	Trp	Val 240
Asp	Phe	Gln	Gly	His 245	Val	Lys	Ala	Met	Lys 250	Arg	Thr	Phe	Lys	Asp 255	Ile
Asp	Ser	Val	Phe 260	Gln	Asn	Trp	Leu	Glu 265	Glu	His	Ile	Asn	Lys 270	Arg	Glu
Lys	Met	Glu 275	Val	Asn	Ala	Glu	Gly 280	Asn	Glu	Gln	Asp	Phe 285	Ile	Asp	Val
Val	Leu 290	Ser	ГЛЗ	Met	Ser	Asn 295	Glu	Tyr	Leu	Gly	Glu 300	Gly	Tyr	Ser	Arg
Asp 305	Thr	Val	Ile	ГЛа	Ala 310	Thr	Val	Phe	Ser	Leu 315	Val	Leu	Asp	Ala	Ala 320
Asp	Thr	Val	Ala	Leu 325	His	Ile	Asn	Trp	Gly 330	Met	Ala	Leu	Leu	Ile 335	Asn
Asn	Gln	Lys	Ala 340	Leu	Thr	Lys	Ala	Gln 345	Glu	Glu	Ile	Asp	Thr 350	Lys	Val
Gly	Lys	Asp 355	Arg	Trp	Val	Glu	Glu 360	Ser	Asp	Ile	Lys	Asp 365	Leu	Val	Tyr
Leu	Gln 370	Ala	Ile	Val	Lys	Glu 375	Val	Leu	Arg	Leu	Tyr 380	Pro	Pro	Gly	Pro
Leu 385	Leu	Val	Pro	His	Glu 390	Asn	Val	Glu	Asp	Суз 395	Val	Val	Ser	Gly	Tyr 400
His	Ile	Pro	Lys	Gly 405	Thr	Arg	Leu	Phe	Ala 410	Asn	Val	Met	Lys	Leu 415	Gln
Arg	Asp	Pro	Lys 420	Leu	Trp	Ser	Asp	Pro 425	Asp	Thr	Phe	Asp	Pro 430	Glu	Arg
Phe	Ile	Ala 435	Thr	Asp	Ile	Asp	Phe 440	Arg	Gly	Gln	Tyr	Tyr 445	Гла	Tyr	Ile
Pro	Phe 450	Gly	Ser	Gly	Arg	Arg 455	Ser	Сүз	Pro	Gly	Met 460	Thr	Tyr	Ala	Leu
Gln 465	Val	Glu	His	Leu	Thr 470	Met	Ala	His	Leu	Ile 475	Gln	Gly	Phe	Asn	Tyr 480
Arg	Thr	Pro	Asn	Asp 485	Glu	Pro	Leu	Asp	Met 490	Гла	Glu	Gly	Ala	Gly 495	Ile
Thr	Ile	Arg	Lys 500	Val	Asn	Pro	Val	Glu 505	Leu	Ile	Ile	Ala	Pro 510	Arg	Leu
Ala	Pro	Glu 515	Leu	Tyr											

<210)> SH	EQ II) NO	26												
<211> LENGTH: 517 <212> TYPE: PRT																
<213	8> OF	RGANI	SM:	Nicotiana tabacum												
<400)> SE	EQUE	ICE :	26												
Met 1	Leu	Ser	Pro	Ile 5	Glu	Ala	Ile	Val	Gly 10	Leu	Val	Thr	Phe	Thr 15	Phe	
Leu	Phe	Phe	Phe 20	Leu	Trp	Thr	Lys	Lys 25	Ser	Gln	Lys	Pro	Ser 30	Lys	Pro	
Leu	Pro	Pro 35	Lys	Ile	Pro	Gly	Gly 40	Trp	Pro	Val	Ile	Gly 45	His	Leu	Phe	
His	Phe 50	Asn	Asp	Asp	Gly	Asp 55	Asp	Arg	Pro	Leu	Ala 60	Arg	Lys	Leu	Gly	
Asp 65	Leu	Ala	Asp	ГЛа	Tyr 70	Gly	Pro	Val	Phe	Thr 75	Phe	Arg	Leu	Gly	Leu 80	
Pro	Leu	Val	Leu	Val 85	Val	Ser	Ser	Tyr	Glu 90	Ala	Val	Lys	Asp	Сув 95	Phe	
Ser	Thr	Asn	Asp 100	Ala	Ile	Phe	Ser	Asn 105	Arg	Pro	Ala	Phe	Leu 110	Tyr	Gly	
Asp	Tyr	Leu 115	Gly	Tyr	Asn	Asn	Ala 120	Met	Leu	Phe	Leu	Ala 125	Asn	Tyr	Gly	
Pro	Tyr 130	Trp	Arg	Lys	Asn	Arg 135	Lys	Leu	Val	Ile	Gln 140	Glu	Val	Leu	Ser	
Ala 145	Ser	Arg	Leu	Glu	Lys 150	Phe	Lys	His	Val	Arg 155	Phe	Ala	Arg	Ile	Gln 160	
Ala	Ser	Ile	Lys	Asn 165	Leu	Tyr	Thr	Arg	Ile 170	Asp	Gly	Asn	Ser	Ser 175	Thr	
Ile	Asn	Leu	Thr 180	Asp	Trp	Leu	Glu	Glu 185	Leu	Asn	Phe	Gly	Leu 190	Ile	Val	
ГЛЗ	Met	Ile 195	Ala	Gly	Lys	Asn	Tyr 200	Glu	Ser	Gly	Lys	Gly 205	Asp	Glu	Gln	
Val	Glu 210	Arg	Phe	Lys	Lys	Ala 215	Phe	Lys	Asp	Phe	Met 220	Ile	Leu	Ser	Met	
Glu 225	Phe	Val	Leu	Trp	Asp 230	Ala	Phe	Pro	Ile	Pro 235	Leu	Phe	Lys	Trp	Val 240	
Asp	Phe	Gln	Gly	His 245	Val	Lys	Ala	Met	Lys 250	Arg	Thr	Phe	Lys	Asp 255	Ile	
Asp	Ser	Val	Phe 260	Gln	Asn	Trp	Leu	Glu 265	Glu	His	Ile	Asn	Lys 270	Arg	Glu	
ГЛЗ	Met	Glu 275	Val	Asn	Ala	Glu	Gly 280	Asn	Glu	Gln	Asp	Phe 285	Ile	Asp	Val	
Val	Leu 290	Ser	Гла	Met	Ser	Asn 295	Glu	Tyr	Leu	Gly	Glu 300	Gly	Tyr	Ser	Arg	
Asp 305	Thr	Val	Ile	Lys	Ala 310	Thr	Val	Phe	Ser	Leu 315	Val	Leu	Asp	Ala	Ala 320	
Asp	Thr	Val	Ala	Leu 325	His	Ile	Asn	Trp	Gly 330	Met	Ala	Leu	Leu	Ile 335	Asn	
Asn	Gln	Lys	Ala 340	Leu	Thr	Lys	Ala	Gln 345	Glu	Glu	Ile	Asp	Thr 350	Lys	Val	
Gly	Lys	Asp 355	Arg	Trp	Val	Glu	Glu 360	Ser	Asp	Ile	Lys	Asp 365	Leu	Val	Tyr	
_				_			_								_	

Leu Gln Ala Ile Val Lys Glu Val Leu Arg Leu Tyr Pro Pro Gly Pro

-continued

	370					375					380				
Leu 385	Leu	Val	Pro	His	Glu 390	Asn	Val	Glu	Asp	Суз 395	Val	Val	Ser	Gly	Tyr 400
His	Ile	Pro	Гла	Gly 405	Thr	Arg	Leu	Phe	Ala 410	Asn	Val	Met	Lys	Leu 415	Leu
Arg	Asp	Pro	Lys 420	Leu	Trp	Pro	Asp	Pro 425	Asp	Thr	Phe	Asp	Pro 430	Glu	Arg
Phe	Ile	Ala 435	Thr	Asp	Ile	Asp	Phe 440	Arg	Gly	Gln	Tyr	Tyr 445	Lys	Tyr	Ile
Pro	Phe 450	Gly	Ser	Gly	Arg	Arg 455	Ser	Суз	Pro	Gly	Met 460	Thr	Tyr	Ala	Leu
Gln 465	Val	Glu	His	Leu	Thr 470	Met	Ala	His	Leu	Ile 475	Gln	Gly	Phe	Asn	Tyr 480
Arg	Thr	Pro	Asn	Asp 485	Glu	Pro	Leu	Asp	Met 490	Lys	Glu	Gly	Ala	Gly 495	Ile
Thr	Ile	Arg	Lys 500	Val	Asn	Pro	Val	Glu 505	Leu	Ile	Ile	Ala	Pro 510	Arg	Leu
Ala	Pro	Glu 515	Leu	Tyr											
<210 <211 <212 <213)> SH .> LH ?> TY ?> OF	EQ II ENGTH ZPE : RGANI	D NO H: 5: PRT LSM:	27 17 Nico	otiar	na ta	abacı	ım							
<400)> SH	EQUEN	ICE :	27											
Met 1	Leu	Ser	Pro	Ile 5	Glu	Ala	Ile	Val	Gly 10	Leu	Val	Thr	Phe	Thr 15	Phe
Leu	Phe	Phe	Phe 20	Leu	Trp	Thr	Lys	Lуз 25	Ser	Gln	ГЛа	Pro	Ser 30	Lys	Pro
Leu	Pro	Pro 35	Lys	Ile	Pro	Gly	Gly 40	Trp	Pro	Val	Ile	Gly 45	His	Leu	Phe
His	Phe 50	Asn	Asp	Asp	Gly	Asp 55	Asp	Arg	Pro	Leu	Ala 60	Arg	Lys	Leu	Gly
Asp 65	Leu	Ala	Asp	Lys	Tyr 70	Gly	Pro	Val	Phe	Thr 75	Phe	Arg	Leu	Gly	Leu 80
Pro	Leu	Val	Leu	Val 85	Val	Ser	Ser	Tyr	Glu 90	Ala	Val	Lys	Asp	Суз 95	Phe
Ser	Thr	Asn	Asp 100	Ala	Ile	Phe	Ser	Asn 105	Arg	Pro	Ala	Phe	Leu 110	Tyr	Gly
Aap	Tyr	Leu 115	Gly	Tyr	Asn	Asn	Ala 120	Met	Leu	Phe	Leu	Ala 125	Asn	Tyr	Gly
Pro	Tyr 130	Trp	Arg	ГЛа	Asn	Arg 135	LÀa	Leu	Val	Ile	Gln 140	Glu	Val	Leu	Ser
Ala 145	Ser	Arg	Leu	Glu	Lys 150	Phe	Lys	His	Val	Arg 155	Phe	Ala	Arg	Ile	Gln 160
Ala	Ser	Ile	Lys	Asn 165	Leu	Tyr	Thr	Arg	Ile 170	Asp	Gly	Asn	Ser	Ser 175	Thr
Ile	Asn	Leu	Thr 180	Asp	Trp	Leu	Glu	Glu 185	Leu	Asn	Phe	Gly	Leu 190	Ile	Val
Lys	Met	Ile 195	Ala	Gly	ГЛа	Asn	Tyr 200	Glu	Ser	Gly	Гла	Gly 205	Asp	Glu	Gln
Val	Glu 210	Arg	Phe	ГЛЗ	ГЛа	Ala 215	Phe	ГЛа	Asp	Phe	Met 220	Ile	Leu	Ser	Met

Glu 225	Phe	Val	Leu	Trp	Asp 230	Ala	Phe	Pro	Ile	Pro 235	Leu	Phe	Lys	Trp	Val 240
Asp	Phe	Gln	Gly	His 245	Val	ГЛа	Ala	Met	Lув 250	Arg	Thr	Phe	Lys	Asp 255	Ile
Asp	Ser	Val	Phe 260	Gln	Asn	Trp	Leu	Glu 265	Glu	His	Ile	Asn	Lys 270	Arg	Glu
ГЛЗ	Met	Glu 275	Val	Asn	Ala	Glu	Gly 280	Asn	Glu	Gln	Asp	Phe 285	Ile	Asp	Val
Val	Leu 290	Ser	Lys	Met	Ser	Asn 295	Glu	Tyr	Leu	Gly	Glu 300	Gly	Tyr	Ser	Arg
Asp 305	Thr	Val	Ile	Glu	Ala 310	Thr	Val	Phe	Ser	Leu 315	Val	Leu	Asp	Ala	Ala 320
Asp	Thr	Val	Ala	Leu 325	His	Ile	Asn	Trp	Gly 330	Met	Ala	Leu	Leu	Ile 335	Asn
Asn	Gln	Lys	Ala 340	Leu	Thr	ГЛа	Ala	Gln 345	Glu	Glu	Ile	Asp	Thr 350	Lys	Val
Сүз	Lys	Asp 355	Arg	Trp	Val	Glu	Glu 360	Ser	Asp	Ile	ГЛа	Asp 365	Leu	Val	Tyr
Leu	Gln 370	Ala	Ile	Val	ГЛа	Glu 375	Val	Leu	Arg	Leu	Tyr 380	Pro	Pro	Gly	Pro
Leu 385	Leu	Val	Pro	His	Glu 390	Asn	Val	Glu	Asp	Сув 395	Val	Val	Ser	Gly	Tyr 400
His	Ile	Pro	Гла	Gly 405	Thr	Arg	Leu	Phe	Ala 410	Asn	Val	Met	Lys	Leu 415	Gln
Arg	Asp	Pro	Lys 420	Leu	Trp	Ser	Asp	Pro 425	Asp	Thr	Phe	Asp	Pro 430	Glu	Arg
Phe	Ile	Ala 435	Thr	Asp	Ile	Asp	Phe 440	Arg	Gly	Gln	Tyr	Tyr 445	Lys	Tyr	Ile
Pro	Phe 450	Gly	Pro	Gly	Arg	Arg 455	Ser	Суз	Pro	Gly	Met 460	Thr	Tyr	Ala	Leu
Gln 465	Val	Glu	His	Leu	Thr 470	Met	Ala	His	Leu	Ile 475	Gln	Gly	Phe	Asn	Tyr 480
Arg	Thr	Pro	Asn	Asp 485	Glu	Pro	Leu	Asp	Met 490	Lys	Glu	Gly	Ala	Gly 495	Ile
Thr	Ile	Arg	Lys 500	Val	Asn	Pro	Val	Glu 505	Leu	Ile	Ile	Ala	Pro 510	Arg	Leu
Ala	Pro	Glu 515	Leu	Tyr											
<210 <211)> SH L> LH	EQ II ENGTH) NO H: 5:	28 17											
<212 <213	2> TY 3> OF	(PE : RGAN	PRT [SM:	Nico	otia	na ta	abacı	ım							
<400)> SI	EQUEI	ICE :	28											
Met 1	Val	Phe	Pro	Ile 5	Glu	Ala	Ile	Val	Gly 10	Leu	Val	Thr	Phe	Thr 15	Phe
Leu	Phe	Tyr	Phe 20	Leu	Trp	Thr	Гла	Lув 25	Ser	Gln	ГЛа	Pro	Ser 30	Гла	Pro
Leu	Pro	Pro 35	Гла	Ile	Pro	Gly	Gly 40	Trp	Pro	Val	Ile	Gly 45	His	Leu	Phe
His	Phe 50	Asn	Asp	Asp	Gly	Asn 55	Asp	Arg	Pro	Leu	Ala 60	Arg	Lys	Leu	Gly
Asp 65	Leu	Ala	Asp	Гла	Tyr 70	Gly	Pro	Val	Phe	Thr 75	Phe	Arg	Leu	Gly	Leu 80

Pro Leu Val Leu Val Val Ser Ser Tyr Glu Ala Ile Lys Asp Cys Phe Ser Thr Asn Asp Ala Ile Phe Ser Asn Arg Pro Ala Phe Leu Tyr Gly Glu Tyr Leu Gly Tyr Asn Asn Ala Met Leu Phe Leu Ala Asn Tyr Gly Pro Tyr Trp Arg Lys Asn Arg Lys Leu Val Ile Gln Glu Val Leu Ser Ala Ser Arg Leu Lys Lys Phe Lys His Val Arg Phe Ala Arg Ile Gln Thr Ser Ile Lys Asn Leu Tyr Thr Arg Ile Asp Arg Asn Ser Ser Thr Ile Asn Leu Thr Asp Trp Leu Glu Glu Leu Asn Phe Gly Leu Ile Val Lys Met Ile Ala Gly Lys Asn Tyr Glu Ser Gly Lys Gly Asp Glu Gln Val Glu Arg Phe Lys Lys Ala Phe Lys Asp Phe Met Ile Ile Ser Met Glu Phe Val Leu Trp Asp Ala Phe Pro Ile Pro Leu Phe Lys Trp Val Asp Phe Gln Gly His Val Lys Ala Met Lys Arg Thr Phe Lys Asp Ile Asp Ser Val Phe Gln Asn Trp Leu Glu Glu His Ile Asn Lys Arg Glu Lys Met Glu Val Asn Ala Glu Gly Asn Glu Gln Asp Phe Ile Asp Val Val Leu Ser Lys Met Ser Asn Glu Tyr Leu Gly Glu Gly Tyr Ser Arg Asp Thr Val Ile Lys Ala Thr Val Phe Ser Leu Val Leu Asp Ala Ala Asp Thr Val Ala Leu His Ile Asn Trp Gly Met Ala Leu Leu Ile Asn Asn Gln Asn Ala Leu Lys Lys Ala Gln Glu Glu Ile Asp Thr Ile Val Gly Lys Asp Arg Trp Val Glu Glu Ser Asp Ile Lys Asp Leu Val Tyr Leu Gln Ala Ile Val Lys Glu Val Leu Arg Leu Tyr Pro Pro Gly Pro Leu Leu Val Pro His Glu Asn Ile Glu Asp Cys Val Val Ser Gly Tyr Tyr Ile Ser Lys Gly Thr Arg Leu Phe Ala Asn Val Met Lys Leu Gln Arg Asp Pro Lys Leu Trp Pro Asn Pro Asp Asn Phe Asp Pro Glu Arg Phe Val Ala Ala Gly Ile Asp Phe Arg Gly Gln His Tyr Glu Tyr Ile Pro Phe Gly Ser Gly Arg Arg Ser Cys Pro Gly Met Thr Tyr Ala Leu Gln Val Glu His Leu Thr Met Ala His Leu Ile Gln Gly Phe Asn Tyr Ser Thr Pro Asn Asp Glu Pro Leu Asp Met Lys Glu Gly Ala Gly Ile

Thr	Ile	Arg	Lys 500	Val	Asn	Pro	Val	Glu 505	Val	Ile	Ile	Met	Pro 510	Arg	Leu
Ala	Pro	Glu 515	Leu	Tyr											
<21 <21 <21 <21	0> SH 1> LH 2> TY 3> OH	EQ II ENGTI (PE : RGAN]	D NO H: 5: PRT ISM:	29 18 Nico	otia	na ta	abacı	ım							
<40	0> SI	EQUEI	NCE :	29											
Met 1	Val	Phe	Pro	Val 5	Glu	Ala	Ile	Val	Gly 10	Leu	Val	Thr	Phe	Thr 15	Phe
Leu	Phe	Tyr	Phe 20	Leu	Trp	Thr	Lys	Lys 25	Ser	Gln	Lys	Pro	Ser 30	Lys	Pro
Leu	Pro	Pro 35	ГЛа	Ile	Pro	Gly	Gly 40	Trp	Pro	Val	Ile	Gly 45	His	Leu	Phe
Tyr	Phe 50	Asp	Asp	Asp	Gly	Asp 55	Asp	Arg	Pro	Leu	Ala 60	Arg	Lys	Leu	Gly
Asp 65	Leu	Ala	Asp	ГЛа	Tyr 70	Gly	Pro	Val	Phe	Thr 75	Phe	Arg	Leu	Gly	Leu 80
Pro	Leu	Val	Leu	Val 85	Val	Ser	Ser	Tyr	Glu 90	Ala	Ile	ГÀа	Asp	Сув 95	Phe
Ser	Thr	Asn	Asp 100	Ala	Ile	Phe	Ser	Asn 105	Arg	Pro	Ala	Phe	Leu 110	Tyr	Gly
Glu	Tyr	Leu 115	Gly	Tyr	ГЛа	Asn	Ala 120	Met	Leu	Phe	Leu	Ala 125	Asn	Tyr	Gly
Ser	Tyr 130	Trp	Arg	Гла	Asn	Arg 135	Lys	Leu	Ile	Ile	Gln 140	Glu	Val	Leu	Ser
Ala 145	Ser	Arg	Leu	Glu	Lys 150	Phe	Lys	His	Val	Arg 155	Phe	Ala	Arg	Ile	Gln 160
Thr	Ser	Ile	Lys	Asn 165	Leu	Tyr	Thr	Arg	Ile 170	Asp	Gly	Asn	Ser	Ser 175	Thr
Ile	Asn	Leu	Thr 180	Asp	Trp	Leu	Glu	Glu 185	Leu	Asn	Phe	Gly	Leu 190	Ile	Val
ГЛа	Met	Ile 195	Ala	Gly	ГЛа	Asn	Tyr 200	Glu	Ser	Gly	ГЛа	Gly 205	Asp	Glu	Gln
Val	Glu 210	Arg	Phe	ГЛа	ГЛа	Ala 215	Phe	ГЛа	Asp	Phe	Met 220	Ile	Leu	Ser	Met
Glu 225	Phe	Val	Leu	Trp	Asp 230	Ala	Phe	Pro	Ile	Pro 235	Leu	Phe	Lys	Trp	Val 240
Asp	Phe	Gln	Gly	His 245	Val	ГЛа	Ala	Met	Lys 250	Arg	Thr	Phe	ГЛа	Asp 255	Ile
Asp	Ser	Val	Phe 260	Gln	Asn	Trp	Leu	Glu 265	Glu	His	Ile	Гла	Lys 270	Arg	Glu
Гла	Ile	Met 275	Glu	Val	Gly	Thr	Glu 280	Gly	Asn	Glu	Gln	Asp 285	Phe	Ile	Asp
Val	Val 290	Leu	Ser	ГЛЗ	Met	Ser 295	Asn	Glu	Tyr	Leu	Gly 300	Glu	Gly	Tyr	Ser
Arg 305	Asp	Thr	Val	Ile	Lys 310	Ala	Thr	Val	Phe	Ser 315	Leu	Val	Leu	Asp	Ala 320
Ala	Asp	Thr	Val	Ala 325	Leu	His	Ile	Asn	Сув 330	Gly	Met	Ala	Leu	Leu 335	Ile
Asn	Asn	Gln	Asn 340	Ala	Leu	Lys	Lys	Ala 345	Gln	Glu	Glu	Ile	Asp 350	Thr	Гла
-continued

Val	Gly	Lys 355	Asp	Arg	Trp	Val	Glu 360	Glu	Ser	Asp	Ile	Lys 365	Asp	Leu	Val
Tyr	Leu 370	Gln	Ala	Ile	Val	Lys 375	Glu	Val	Leu	Arg	Leu 380	Tyr	Pro	Pro	Gly
Pro 385	Leu	Leu	Val	Pro	His 390	Glu	Asn	Val	Glu	Asp 395	Суз	Val	Val	Ser	Gly 400
Tyr	His	Ile	Pro	Lys 405	Gly	Thr	Arg	Leu	Phe 410	Ala	Asn	Val	Met	Lys 415	Leu
Gln	Arg	Asp	Pro 420	Гүз	Leu	Trp	Ser	Asn 425	Pro	Asp	Гүз	Phe	Asn 430	Pro	Glu
Arg	Phe	Ile 435	Ala	Arg	Asp	Ile	Asp 440	Phe	His	Gly	Gln	His 445	Tyr	Glu	Tyr
Ile	Pro 450	Phe	Gly	Ser	Gly	Arg 455	Arg	Ser	Суз	Pro	Gly 460	Met	Thr	Tyr	Ala
Leu 465	Gln	Val	Glu	His	Leu 470	Thr	Met	Ala	His	Leu 475	Ile	Gln	Gly	Phe	Asn 480
Tyr	Arg	Thr	Pro	Thr 485	Asp	Glu	Pro	Leu	Asp 490	Met	Lys	Glu	Gly	Ala 495	Gly
Ile	Thr	Ile	Arg 500	Lys	Val	Asn	Pro	Val 505	Lys	Val	Ile	Ile	Thr 510	Pro	Arg
Leu	Ala	Pro 515	Glu	Leu	Tyr										
<210 <211 <212 <213	0> SH L> LH 2> TY 3> OH	EQ II ENGTH (PE : RGANI) NO H: 5: PRT [SM:	30 17 Nico	otiar	na ta	abacı	ım							
-100) - CT		ICF.	20											
<400)> SH	EQUEI	ICE :	30	_	_		_	_		_				
<400 Met 1)> SH Val	EQUE1 Phe	ICE: Pro	30 Ile 5	Glu	Ala	Phe	Val	Gly 10	Leu	Val	Thr	Phe	Thr 15	Phe
<400 Met 1 Leu	D> SH Val Leu	EQUE1 Phe Tyr	NCE: Pro Phe 20	30 Ile 5 Leu	Glu Trp	Ala Thr	Phe Lys	Val Lys 25	Gly 10 Ser	Leu Gln	Val Lys	Thr Leu	Phe Pro 30	Thr 15 Lys	Phe Pro
<400 Met 1 Leu Leu	D> SH Val Leu Pro	Phe Tyr Pro 35	NCE: Pro Phe 20 Lys	30 Ile 5 Leu Ile	Glu Trp Pro	Ala Thr Gly	Phe Lys Gly 40	Val Lys 25 Trp	Gly 10 Ser Pro	Leu Gln Val	Val Lys Ile	Thr Leu Gly 45	Phe Pro 30 His	Thr 15 Lys Leu	Phe Pro Phe
<400 Met 1 Leu Leu)> SH Val Leu Pro Phe 50	Phe Tyr Pro 35 Asn	ICE : Pro Phe 20 Lys Asn	30 Ile 5 Leu Ile Asp	Glu Trp Pro Gly	Ala Thr Gly Asp 55	Phe Lys Gly 40 Asp	Val Lys 25 Trp Arg	Gly 10 Ser Pro Pro	Leu Gln Val Leu	Val Lys Ile Ala 60	Thr Leu Gly 45 Arg	Phe Pro 30 His Lys	Thr 15 Lys Leu Leu	Phe Pro Phe Gly
<400 Met 1 Leu His Asp 65)> SI Val Leu Pro Phe 50 Leu	Phe Tyr Pro 35 Asn Ala	NCE: Pro Phe 20 Lys Asn Asp	30 Ile Leu Ile Asp Lys	Glu Trp Pro Gly Tyr 70	Ala Thr Gly Asp 55 Gly	Phe Lys Gly 40 Asp Pro	Val Lys 25 Trp Arg Val	Gly 10 Ser Pro Pro Phe	Leu Gln Val Leu Thr 75	Val Lys Ile Ala 60 Phe	Thr Leu Gly 45 Arg Arg	Phe Pro 30 His Lys Leu	Thr 15 Lys Leu Leu Gly	Phe Pro Phe Gly Leu 80
<400 Met 1 Leu His Asp 65 Pro)> SE Val Leu Pro Phe 50 Leu Leu	QUE Phe Tyr Pro 35 Asn Ala Val	NCE: Pro Phe 20 Lys Asn Asp Leu	30 Ile Leu Ile Asp Lys Val 85	Glu Trp Pro Gly Tyr 70 Val	Ala Thr Gly Asp 55 Gly Ser	Phe Lys Gly 40 Asp Pro Ser	Val Lys 25 Trp Arg Val Tyr	Gly 10 Ser Pro Pro Phe Glu 90	Leu Gln Val Leu Thr 75 Ala	Val Lys Ile Ala 60 Phe Ile	Thr Leu Gly 45 Arg Arg Lys	Phe Pro 30 His Lys Leu Asp	Thr 15 Lys Leu Leu Gly Cys 95	Phe Pro Phe Gly Leu 80 Phe
<400 Met 1 Leu Leu His Asp 65 Pro Ser	<pre>>> SI Val Leu Pro Phe 50 Leu Leu Thr</pre>	CQUET Phe Tyr Pro 35 Asn Ala Val Asn	NCE: Pro Phe 20 Lys Asn Asp Leu Asp 100	30 Ile Leu Ile Asp Lys Val 85 Ala	Glu Trp Pro Gly Tyr 70 Val Ile	Ala Thr Gly Asp 55 Gly Ser Phe	Phe Lys Gly 40 Asp Pro Ser Ser	Val Lys 25 Trp Arg Val Tyr Tyr Asn 105	Gly 10 Ser Pro Pro Phe Glu 90 Arg	Leu Gln Val Leu Thr 75 Ala Pro	Val Lys Ile Ala 60 Phe Ile Ala	Thr Leu Gly 45 Arg Arg Lys Phe	Phe Pro 30 His Lys Leu Asp Leu 110	Thr 15 Lys Leu Leu Gly S 95 Tyr	Phe Pro Phe Gly Leu 80 Phe Gly
<400 Met 1 Leu Leu His 65 Pro Ser Glu	<pre>>> SI Val Leu Pro Phe 50 Leu Leu Thr Tyr</pre>	QUEN Phe Tyr Pro 35 Asn Ala Val Asn Leu 115	NCE: Pro Phe 20 Lys Asn Asp Leu Asp 100 Gly	30 Ile Leu Ile Asp Lys Val 85 Ala Tyr	Glu Trp Pro Gly Tyr 70 Val Ile Asn	Ala Thr Gly Asp 55 Gly Ser Phe Asn	Phe Lys Gly 40 Asp Pro Ser Ser Thr 120	Val Lys 25 Trp Arg Val Tyr Asn 105 Met	Gly 10 Ser Pro Pro Phe Glu 90 Arg Leu	Leu Gln Val Leu Thr 75 Ala Pro Phe	Val Lys Ile Ala 60 Phe Ile Ala Leu	Thr Leu Gly 45 Arg Lys Phe Ala 125	Phe Pro 30 His Lys Leu Leu 110 Asn	Thr 15 Lys Leu Gly Cys 95 Tyr Tyr	Phe Pro Phe Gly Leu 80 Phe Gly Gly
<400 Met 1 Leu His Asp 65 Pro Ser Glu Pro	<pre>>> SI Val Leu Pro Phe 50 Leu Leu Thr Tyr Tyr 130</pre>	QUUEN Phe Tyr Pro 35 Asn Ala Val Asn Leu 115 Trp	NCE: Pro Phe 20 Lys Asn Asp Leu Asp 100 Gly Arg	30 Ile Leu Ile Asp Lys Val 85 Ala Tyr Lys	Glu Trp Pro Gly Tyr 70 Val Ile Asn	Ala Thr Gly Asp 55 Gly Ser Phe Asn Arg 135	Phe Lys Gly 40 Asp Pro Ser Ser Thr 120 Lys	Val Lys 25 Trp Arg Val Tyr Asn 105 Met Leu	Gly 10 Ser Pro Pro Phe Glu Arg Leu Val	Leu Gln Val Leu Thr 75 Ala Pro Phe Ile	Val Lys Ile Ala 60 Phe Ile Ala Leu Gln 140	Thr Leu Gly 45 Arg Lys Phe Ala 125 Glu	Phe Pro 30 His Lys Leu Leu 110 Asn Val	Thr 15 Lys Leu Gly Cys 95 Tyr Tyr Leu	Phe Pro Phe Gly Leu 80 Phe Gly Gly Ser
<400 Met 1 Leu His Asp 65 Pro Ser Glu Pro Ala 145	<pre>>> SI Val Leu Pro Phe 50 Leu Leu Thr Tyr Tyr Tyr 130 Ser</pre>	QUEN Phe Tyr Pro 35 Asn Ala Val Asn Leu 115 Trp Arg	NCE: Pro Phe 20 Lys Asn Asp Leu Asp 100 Gly Arg Leu	30 Ile Leu Ile Asp Lys Val S5 Ala Tyr Lys Glu	Glu Trp Pro Gly Tyr 70 Val Ile Asn Asn Lys 150	Ala Thr Gly Asp 55 Gly Ser Phe Asn Arg 135 Phe	Phe Lys Gly 40 Asp Pro Ser Ser Ser Thr 120 Lys	Val Lys 25 Trp Arg Val Tyr Asn 105 Met Leu Gln	Gly 10 Ser Pro Pro Phe Glu 90 Arg Leu Val	Leu Gln Val Leu Thr 75 Ala Pro Phe Ile Arg 155	Val Lys Ile Ala 60 Phe Ile Ala Leu Gln 140 Phe	Thr Leu Gly 45 Arg Lys Phe Ala 125 Glu Thr	Phe Pro 30 His Lys Leu Asp Leu 110 Asn Val Arg	Thr 15 Lys Leu Gly Cys 95 Tyr Tyr Leu Ile	Phe Pro Gly Leu 80 Phe Gly Gly Ser Gln 160
<400 Met 1 Leu His Asp 65 Pro Ser Glu Pro Ala 145 Thr	<pre>>> SI Val Leu Pro Phe 50 Leu Leu Thr Tyr Tyr 130 Ser Ser</pre>	QUEN Phe Tyr Pro 35 Asn Ala Val Leu 115 Trp Arg Ile	NCE: Pro Phe 20 Lys Asn Asp Leu Asp 100 Gly Arg Leu Lys	30 Jle Leu Ile Lys Val Tyr Lys Glu Asn 165	Glu Trp Pro Gly Tyr 70 Val Ile Asn Lys 150 Leu	Ala Thr Gly Asp 55 Gly Ser Phe Asn Arg 135 Phe Tyr	Phe Lys Gly 40 Asp Pro Ser Ser Thr 120 Lys Lys Thr	Val Lys 25 Trp Arg Val Tyr Asn 105 Met Leu Gln Arg	Gly Ser Pro Pro Glu Glu Arg Leu Val Val Ile 170	Leu Gln Val Leu Thr 75 Ala Pro Phe Ile Arg 155 Asn	Val Lys Ile Ala 60 Phe Ile Ala Leu Gln 140 Phe Gly	Thr Leu Gly 45 Arg Lys Phe Ala 125 Glu Thr Asn	Phe Pro 30 His Lys Leu Leu 110 Asn Val Arg Ser	Thr 15 Lys Leu Gly Cys 7yr Tyr Leu Ile Ser 175	Phe Pro Phe Gly Leu 80 Phe Gly Gly Gly Ser Gln 160 Thr
<400 Met 1 Leu His Asp 65 Pro Ser Glu Pro Ala 145 Thr Ile	<pre>>> SI Val Leu Pro Phe 50 Leu Leu Thr Tyr Tyr 130 Ser Ser Asn</pre>	QUEP Phe Tyr Pro 35 Asn Ala Val Asn Leu 115 Trp Arg Ile Leu	NCE: Pro Phe 20 Lys Asn Asp Leu Asp 100 Gly Arg Leu Lys Thr 180	30 Jle Leu Lys Val Xyal Ala Tyr Lys Glu Asn 165 Asp	Glu Trp Pro Gly Tyr 70 Val Ile Asn Lys 150 Leu Trp	Ala Thr Gly Asp 55 Gly Ser Phe Asn Arg 135 Phe Tyr Leu	Phe Lys Gly 40 Asp Pro Ser Ser Thr 120 Lys Lys Thr Glu	Val Lys 25 Arg Val Tyr Asn 105 Met Leu Gln Arg Glu 185	Gly 10 Ser Pro Pro Glu 90 Arg Leu Val Val Ile 170 Leu	Leu Gln Val Leu Thr 75 Ala Pro Phe Ile Arg 155 Asn Asn	Val Lys Ile Ala 60 Phe Ile Ala Leu Gln 140 Phe Gly Phe	Thr Leu Gly 45 Arg Lys Phe Ala 125 Glu Thr Asn Gly	Phe So His Lys Leu Asp Leu 110 Asn Val Arg Ser Leu 190	Thr 15 Lys Leu Gly 95 Tyr Tyr Leu Ile Ser 175 Ile	Phe Pro Gly Leu 80 Phe Gly Gly Ser Gln 160 Thr Val

Val	Glu 210	Arg	Phe	Lys	Asn	Ala 215	Phe	Lys	Asp	Phe	Met 220	Val	Leu	Ser	Met
Glu 225	Phe	Val	Leu	Trp	Asp 230	Ala	Phe	Pro	Ile	Pro 235	Leu	Phe	Lys	Trp	Val 240
Asp	Phe	Gln	Gly	His 245	Ile	Гла	Ala	Met	Lys 250	Arg	Thr	Phe	Lys	Asp 255	Ile
Asp	Ser	Val	Phe 260	Gln	Asn	Trp	Leu	Glu 265	Glu	His	Ile	Asn	Lys 270	Arg	Glu
Lys	Met	Glu 275	Val	Gly	Ala	Glu	Gly 280	Asn	Glu	Gln	Asp	Phe 285	Ile	Asp	Val
Val	Leu 290	Ser	Lys	Leu	Ser	Lys 295	Glu	Tyr	Leu	Asp	Glu 300	Gly	Tyr	Ser	Arg
Asp 305	Thr	Val	Ile	Lys	Ala 310	Thr	Val	Phe	Ser	Leu 315	Val	Leu	Asp	Ala	Ala 320
Asp	Thr	Val	Ala	Leu 325	His	Ile	Asn	Trp	Gly 330	Met	Thr	Leu	Leu	Ile 335	Asn
Asn	Gln	Asn	Ala 340	Leu	Met	ГЛЗ	Ala	Gln 345	Glu	Glu	Ile	Asp	Thr 350	Lys	Val
Gly	Lys	Asp 355	Arg	Trp	Val	Glu	Glu 360	Ser	Asp	Ile	Гла	Asp 365	Leu	Val	Tyr
Leu	Gln 370	Ala	Ile	Val	Lys	Lys 375	Val	Leu	Arg	Leu	Tyr 380	Pro	Pro	Gly	Pro
Leu 385	Leu	Val	Pro	His	Glu 390	Asn	Val	Lys	Asp	Суз 395	Val	Val	Ser	Gly	Tyr 400
His	Ile	Pro	Lys	Gly 405	Thr	Arg	Leu	Phe	Ala 410	Asn	Val	Met	Lys	Leu 415	Gln
Arg	Asp	Pro	Lys 420	Leu	Leu	Ser	Asn	Pro 425	Asp	Lys	Phe	Asp	Pro 430	Glu	Arg
Phe	Ile	Ala 435	Gly	Asp	Ile	Aap	Phe 440	Arg	Gly	His	His	Tyr 445	Glu	Phe	Ile
Pro	Phe 450	Gly	Ser	Gly	Arg	Arg 455	Ser	Cys	Pro	Gly	Met 460	Thr	Tyr	Ala	Leu
Gln 465	Val	Glu	His	Leu	Thr 470	Met	Ala	His	Leu	Ile 475	Gln	Gly	Phe	Asn	Tyr 480
ГЛа	Thr	Pro	Asn	Asp 485	Glu	Ala	Leu	Asp	Met 490	Lys	Glu	Gly	Ala	Gly 495	Ile
Thr	Ile	Arg	Lys	Val	Asn	Pro	Val	Glu 505	Leu	Ile	Ile	Thr	Pro	Arg	Leu
Ala	Pro	Glu	Leu	Tyr				505					510		
.01		515		21											
<210 <211 <212)> SE L> LE 2> TY	SQ 11 ENGTH (PE :) NO 1: 51 PRT	31 L7											
<213	3> OF)> SE	RGANI	ISM:	Nico 31	otian	na ta	abacı	ım							
Met	Leu	Ser	Pro	Ile 5	Glu	Ala	Phe	Val	Gly	Leu	Val	Thr	Phe	Thr	Phe
- Leu	Leu	Tyr	Phe	Leu	Trp	Thr	Lys	Lys	Ser	Gln	Lys	Leu	Pro	гэ	Pro
Leu	Pro	Pro	20 Lys	Ile	Pro	Gly	Gly	25 Trp	Pro	Val	Ile	Gly	30 His	Leu	Phe
		35			'	-1	40	- T .	'			45	'		

His	Phe 50	Asn	Asn	Asp	Gly	Asp 55	Asp	Arg	Pro	Leu	Ala 60	Arg	Lys	Leu	Gly
Asp 65	Leu	Ala	Asp	ГЛа	Tyr 70	Gly	Pro	Val	Phe	Thr 75	Phe	Arg	Leu	Gly	Leu 80
Pro	Leu	Val	Leu	Val 85	Val	Ser	Ser	Tyr	Glu 90	Ala	Ile	ГЛЗ	Asp	Cys 95	Phe
Ser	Thr	Asn	Asp 100	Ala	Ile	Phe	Ser	Asn 105	Arg	Pro	Ala	Leu	Leu 110	Tyr	Gly
Glu	Tyr	Leu 115	Gly	Tyr	Asn	Asn	Thr 120	Met	Leu	Phe	Leu	Ala 125	Asn	Tyr	Gly
Pro	Tyr 130	Trp	Arg	ГЛЗ	Asn	Arg 135	Lys	Leu	Val	Ile	Gln 140	Glu	Val	Leu	Ser
Ala 145	Ser	Arg	Leu	Glu	Lys 150	Phe	Lys	Gln	Val	Arg 155	Phe	Thr	Arg	Ile	Gln 160
Thr	Ser	Ile	Lys	Asn 165	Leu	Tyr	Thr	Arg	Ile 170	Asn	Gly	Asn	Ser	Ser 175	Thr
Ile	Asn	Leu	Thr 180	Asp	Trp	Leu	Glu	Glu 185	Leu	Asn	Phe	Gly	Leu 190	Ile	Val
Lys	Met	Ile 195	Ala	Gly	ГЛа	Asn	Tyr 200	Glu	Ser	Gly	Гла	Gly 205	Aab	Glu	Gln
Val	Glu 210	Arg	Phe	Гла	Asn	Ala 215	Phe	Lys	Asp	Phe	Met 220	Val	Leu	Ser	Met
Glu 225	Phe	Val	Leu	Trp	Asp 230	Ala	Phe	Pro	Ile	Pro 235	Leu	Phe	Lys	Trp	Val 240
Asp	Phe	Gln	Gly	His 245	Ile	Lys	Ala	Met	Lys 250	Arg	Thr	Phe	Lys	Asp 255	Ile
Asp	Ser	Val	Phe 260	Gln	Asn	Trp	Leu	Glu 265	Glu	His	Ile	Asn	Lys 270	Arg	Glu
Lys	Ile	Glu 275	Val	Gly	Ala	Glu	Gly 280	Asn	Glu	Gln	Asp	Phe 285	Ile	Asp	Val
Val	Leu 290	Ser	Lys	Leu	Ser	Lys 295	Glu	Tyr	Leu	Asp	Glu 300	Gly	Tyr	Ser	Arg
Asp 305	Thr	Val	Ile	ГЛа	Ala 310	Thr	Val	Phe	Ser	Leu 315	Val	Leu	Asb	Ala	Ala 320
Asp	Thr	Val	Ala	Leu 325	His	Ile	Asn	Trp	Gly 330	Met	Thr	Leu	Leu	Ile 335	Asn
Asn	Gln	Asn	Ala 340	Leu	Met	Lys	Ala	Gln 345	Glu	Glu	Ile	Asp	Thr 350	Lys	Val
Gly	Lys	Asp 355	Arg	Trp	Val	Glu	Glu 360	Ser	Asp	Ile	Гла	Asp 365	Leu	Val	Tyr
Leu	Gln 370	Ala	Ile	Val	ГЛа	Lys 375	Val	Leu	Arg	Leu	Tyr 380	Pro	Pro	Gly	Pro
Leu 385	Leu	Val	Pro	His	Glu 390	Asn	Val	ГЛа	Asp	Сув 395	Val	Val	Ser	Gly	Tyr 400
His	Ile	Pro	Lys	Gly 405	Thr	Arg	Leu	Phe	Ala 410	Asn	Val	Met	Lys	Leu 415	Gln
Arg	Asp	Pro	Lys 420	Leu	Leu	Ser	Asn	Pro 425	Asp	Lys	Phe	Asp	Pro 430	Glu	Arg
Phe	Ile	Ala 435	Gly	Asp	Ile	Asp	Phe 440	Arg	Gly	His	His	Tyr 445	Glu	Phe	Ile
Pro	Phe 450	Gly	Ser	Gly	Arg	Arg 455	Ser	Суз	Pro	Gly	Met 460	Thr	Tyr	Ala	Leu
Gln	Val	Glu	His	Leu	Thr	Met	Ala	His	Leu	Ile	Gln	Gly	Phe	Asn	Tyr

465 470 475 480 Lys Thr Pro Asn Asp Glu Ala Leu Asp Met Lys Glu Gly Ala Gly Ile 485 490 495 Thr Ile Arg Lys Val Asn Pro Val Glu Leu Ile Ile Thr Pro Arg Leu 500 505 510 Ala Pro Glu Leu Tyr 515 <210> SEQ ID NO 32 <211> LENGTH: 21 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer <400> SEQUENCE: 32 tqqaattatq cccatcctac a 21 <210> SEQ ID NO 33 <211> LENGTH: 21 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer <400> SEQUENCE: 33 cattagtggt tgcacctgag g 21 <210> SEQ ID NO 34 <211> LENGTH: 21 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer <400> SEQUENCE: 34 gatgagatgt gtgcatactt g 21 <210> SEQ ID NO 35 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer <400> SEQUENCE: 35 23 ccaaattaga aaaactcgta ctg <210> SEQ ID NO 36 <211> LENGTH: 25 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer <400> SEQUENCE: 36 attgtaggac tagtaaccct tacac 25 <210> SEQ ID NO 37 <211> LENGTH: 21

-continued

130

<212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer <400> SEQUENCE: 37 gaggcacaaa gaattctcat c 21 <210> SEQ ID NO 38 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer <400> SEQUENCE: 38 20 gagtagaggg attgtttccg <210> SEQ ID NO 39 <211> LENGTH: 26 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer <400> SEQUENCE: 39 26 gtacaatcaa gataaaacat ctaagg <210> SEQ ID NO 40 <211> LENGTH: 21 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer <400> SEQUENCE: 40 cggtgaatac cttggctaca g 21 <210> SEQ ID NO 41 <211> LENGTH: 25 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer <400> SEQUENCE: 41 25 aattgtcact tctacaggat ttact <210> SEQ ID NO 42 <211> LENGTH: 18 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer <400> SEQUENCE: 42 atggtttttc cggtagaa 18 <210> SEQ ID NO 43 <211> LENGTH: 18 <212> TYPE: DNA

<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Description of Artificial Sequence:	Synthetic
primer	
ADD SECTIONCE, AD	
CHOON SEQUENCE: 45	
ttaataaagg tgaggtgg	19
ccaacaaage ccaygege	10
<210> SEO ID NO 44	
<211> LENGTH: 23	
<212> TYPE: DNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Description of Artificial Sequence:	Synthetic
primer	
<400> SEQUENCE: 44	
acgrgateet aaaeterggt erg	23
210. CEO ID NO 4E	
<210> SEQ 15 NO 43	
<212> TYPE: DNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<pre><223> OTHER INFORMATION: Description of Artificial Sequence:</pre>	Synthetic
primer	-1
-	
<400> SEQUENCE: 45	
gcctgcacct tccttcatg	19
<210> SEQ ID NO 46	
<211> LENGTH: 23	
<212> TYPE: DNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Description of Artificial Sequence:	Synthetic
primer	
ADDS SECUENCE: 46	
(100) BEgolmen. 10	
gggatgaett atgeattaea age	23
3334034000 40304044 430	20
<210> SEQ ID NO 47	
<211> LENGTH: 25	
<212> TYPE: DNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Description of Artificial Sequence:	Synthetic
primer	
<400> SEQUENCE: 47	
	25
aallyleaet tetaeaggat ttaet	25
<210> SEO ID NO 48	
<211> LENGTH: 20	
<212> TYPE: DNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Description of Artificial Sequence:	Synthetic
primer	
•	
<400> SEQUENCE: 48	
-	
qqtqtccaca qacttcqtqq	20
<210> SEQ ID NO 49	
<211> LENGTH: 20	
<212> TYPE: DNA	
<213> ORGANISM: Artificial Sequence	
-	

<220> FEATURE: <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer														
<400> SEQUENCE: 49														
gactecteae ageageacea	20													
<pre>11> SEQ ID NO 50 11> LENGTH: 1733 212> TYPE: DNA 213> ORGANISM: Nicotiana sp. 220> FEATURE: 221> NAME/KEY: CDS 222> LOCATION: (1)(1551)</pre>														
<400> SEQUENCE: 3														
atg ctt tct ccc ata gaa gcc att gta gga cta gta acc ttc aca ttt Met Leu Ser Pro Ile Glu Ala Ile Val Gly Leu Val Thr Phe Thr Phe 1 5 10 15	48													
ctc ttc ttc cta tgg aca aaa aaa tct caa aaa cct tca aaa ccc Leu Phe Phe Leu Trp Thr Lys Lys Ser Gln Lys Pro Ser Lys Pro 20 25 30	96													
tta cca ccg aaa atc ccc gga gga tgg ccg gta atc ggc cat ctt ttc Leu Pro Pro Lys Ile Pro Gly Gly Trp Pro Val Ile Gly His Leu Phe 35 40 45	44													
cac ttc aat gac gac ggc gac gac cgt cca tta gct cga aaa ctc gga 19 His Phe Asn Asp Asp Gly Asp Asp Arg Pro Leu Ala Arg Lys Leu Gly 50 55 60	92													
gac tta gct gac aaa tac ggc ccc gtt ttc act ttt cgg cta ggc ctt24Asp Leu Ala Asp Lys Tyr Gly Pro Val Phe Thr Phe Arg Leu Gly Leu65707580	40													
ccc ctt gtc tta gtt gta agc agt tac gaa gct gta aaa gac tgt ttc 23 Pro Leu Val Leu Val Val Ser Ser Tyr Glu Ala Val Lys Asp Cys Phe 85 90 95	88													
tct aca aat gac gcc att ttt tcc aat cgt cca gct ttt ctt tac ggc 33 Ser Thr Asn Asp Ala Ile Phe Ser Asn Arg Pro Ala Phe Leu Tyr Gly 100 105 110	36													
gat tac ctt ggc tac aat aat gcc atg cta ttt ttg gcc aat tac gga 34 Asp Tyr Leu Gly Tyr Asn Asn Ala Met Leu Phe Leu Ala Asn Tyr Gly 115 120 125	84													
cct tac tgg cga aaa aat cga aaa tta gtt att cag gaa gtt ctc tcc 43 Pro Tyr Trp Arg Lys Asn Arg Lys Leu Val Ile Gln Glu Val Leu Ser 130 135 140	32													
gct agt cgt ctc gaa aaa ttc aaa cac gtg aga ttt gca aga att caa 43 Ala Ser Arg Leu Glu Lys Phe Lys His Val Arg Phe Ala Arg Ile Gln 145 150 155 160	80													
gcg agc att aag aat tta tat act cga att gat gga aat tcg agt acg Ala Ser Ile Lys Asn Leu Tyr Thr Arg Ile Asp Gly Asn Ser Ser Thr 165 170 175	28													
ata aat tta act gat tgg tta gaa gaa ttg aat ttt ggt ctg atc gtg 5 Ile Asn Leu Thr Asp Trp Leu Glu Glu Leu Asn Phe Gly Leu Ile Val 180 185 190	76													
aag atg atc gct gga aaa aat tat gaa tcc ggt aaa gga gat gaa caa 6: Lys Met Ile Ala Gly Lys Asn Tyr Glu Ser Gly Lys Gly Asp Glu Gln 195 200 205	24													
gtg gag aga ttt aag aaa gcg ttt aag gat ttt atg att tta tca atg Val Glu Arg Phe Lys Lys Ala Phe Lys Asp Phe Met Ile Leu Ser Met 210 215 220	72													
gag ttt gtg tta tgg gat gca ttt cca att cca tta ttt aaa tgg gtg72Glu Phe Val Leu Trp Asp Ala Phe Pro Ile Pro Leu Phe Lys Trp Val225230235240	20													

-continued

gat Asp	ttt Phe	caa Gln	glà aaa	cat His 245	gtt Val	aag Lys	gct Ala	atg Met	aaa Lys 250	agg Arg	act Thr	ttt Phe	aaa Lys	gat Asp 255	ata Ile	768		
gat Asp	tct Ser	gtt Val	ttt Phe 260	cag Gln	aat Asn	tgg Trp	tta Leu	gag Glu 265	gaa Glu	cat His	att Ile	aat Asn	aaa Lys 270	aga Arg	gaa Glu	816		
aaa Lys	atg Met	gag Glu 275	gtt Val	aat Asn	gca Ala	gaa Glu	999 Gly 280	aat Asn	gaa Glu	caa Gln	gat Asp	ttc Phe 285	att Ile	gat Asp	gtg Val	864		
gtg Val	ctt Leu 290	tca Ser	aaa Lys	atg Met	agt Ser	aat Asn 295	gaa Glu	tat Tyr	ctt Leu	ggt Gly	gaa Glu 300	ggt Gly	tac Tyr	tct Ser	cgt Arg	912		
gat Asp 305	act Thr	gtc Val	att Ile	aaa Lys	gca Ala 310	acg Thr	gtg Val	ttt Phe	agt Ser	ttg Leu 315	gtc Val	ttg Leu	gat Asp	gca Ala	gca Ala 320	960		
gac Asp	aca Thr	gtt Val	gct Ala	ctt Leu 325	cac His	ata Ile	aat Asn	tgg Trp	gga Gly 330	atg Met	gca Ala	tta Leu	ttg Leu	ata Ile 335	aac Asn	1008		
aat Asn	caa Gln	aag Lys	gcc Ala 340	ttg Leu	acg Thr	aaa Lys	gca Ala	caa Gln 345	gaa Glu	gag Glu	ata Ile	gac Asp	aca Thr 350	aaa Lys	gtt Val	1056		
ggt Gly	aag Lys	gac Asp 355	aga Arg	tgg Trp	gta Val	gaa Glu	gag Glu 360	agt Ser	gat Asp	att Ile	aag Lys	gat Asp 365	ttg Leu	gta Val	tac Tyr	1104		
ctc Leu	caa Gln 370	gct Ala	att Ile	gtt Val	aaa Lys	gaa Glu 375	gtg Val	tta Leu	cga Arg	tta Leu	tat Tyr 380	cca Pro	cca Pro	gga Gly	cct Pro	1152		
ttg Leu 385	tta Leu	gta Val	cca Pro	cac His	gaa Glu 390	aat Asn	gta Val	gaa Glu	gat Asp	tgt Cys 395	gtt Val	gtt Val	agt Ser	gga Gly	tat Tyr 400	1200		
cac His	att Ile	cct Pro	aaa Lys	999 Gly 405	aca Thr	aga Arg	tta Leu	ttc Phe	gca Ala 410	aac Asn	gtc Val	atg Met	aaa Lys	ctg Leu 415	caa Gln	1248		
cgt Arg	gat Asp	cct Pro	aaa Lys 420	ctc Leu	tgg Trp	tct Ser	gat Asp	cct Pro 425	gat Asp	act Thr	ttc Phe	gat Asp	cca Pro 430	gag Glu	aga Arg	1296		
ttc Phe	att Ile	gct Ala 435	act Thr	gat Asp	att Ile	gac Asp	ttt Phe 440	cgt Arg	ggt Gly	cag Gln	tac Tyr	tat Tyr 445	aag Lys	tat Tyr	atc Ile	1344		
ccg Pro	ttt Phe 450	ggt Gly	tct Ser	gga Gly	aga Arg	cga Arg 455	tct Ser	tgt Cys	cca Pro	glà dàð	atg Met 460	act Thr	tat Tyr	gca Ala	ttg Leu	1392		
caa Gln 465	gtg Val	gaa Glu	cac His	tta Leu	aca Thr 470	atg Met	gca Ala	cat His	ttg Leu	atc Ile 475	caa Gln	ggt Gly	ttc Phe	aat Asn	tac Tyr 480	1440		
aga Arg	act Thr	cca Pro	aat Asn	gac Asp 485	gag Glu	ccc Pro	ttg Leu	gat Asp	atg Met 490	aag Lys	gaa Glu	ggt Gly	gca Ala	ggc Gly 495	ata Ile	1488		
act Thr	ata Ile	cgt Arg	aag Lys 500	gta Val	aat Asn	cct Pro	gtg Val	gaa Glu 505	ctg Leu	ata Ile	ata Ile	gcg Ala	cct Pro 510	cgc Arg	ctg Leu	1536		
gca Ala	cct Pro	gag Glu 515	ctt Leu	tat Tyr	taa	aacc	taa	gatc	ttte	at ci	ttgg [.]	ttga	t ca	ttgta	ataa	1591		
tac	tccta	aaa 1	tgqa	tatt	ca ti	ttac	cttti	t at	caati	taat	tgt	cagt	acg .	agtti	tttcta	1651		
att	tgata	aca 1	ttta	taat	aa ta	aaqt	aaaa	a ata	aatto	qtac	taa	tata	taa .	aqqti	ttqtaq	1711		
aad	ataa	ta :	acto	atta	ta a	a St								55	5 5	1733		
uay	ncad		accy	Lucy		-												

35

1. A tobacco product comprising cured tobacco material made from a tobacco plant comprising a first mutation within the coding region of a wild type allele encoding a nicotine demethylase having the amino acid sequence of SEQ ID NO: 2, wherein said first mutation reduces the activity of said nicotine demethylase.

137

2. The tobacco product according to claim **1**, wherein said tobacco plant further comprises a second mutation within the coding region of a wild type allele encoding a nicotine demethylase having the amino acid sequence of SEQ ID NO:13, wherein said second mutation reduces the activity of said nicotine demethylase.

3. The tobacco product according to claim **1**, wherein said 15 tobacco material is made by a curing process selected from the group consisting of flue curing, air curing, and fire curing.

4. The tobacco product according to claim **1**, wherein said tobacco product is selected from the group consisting of a cigar, a cigarette, pipe tobacco, a cigarillo, a non-ventilated or $_{20}$ vented recess filter cigarette, a dissolving strip, a gum, a tablet, snuff, and chewing tobacco.

5. The tobacco product according to claim **1**, wherein said cured tobacco material comprises a lower level of nornicotine or N'-nitrosonornicotine compared to cured tobacco material ²⁵ made from a tobacco plant comprising said wild type allele.

6. The tobacco product according to claim **1**, wherein the conversion of nicotine to nornicotine in said cured tobacco material is selected from the group consisting of less than 0.3%, less than 0.5%, less than 0.7%, between 0.1%-0.5%, between 0.1%-0.4%, between 0.1%-0.7%, and between 0.1%-1.0%.

7. The tobacco product according to claim 1, wherein said tobacco plant has a non-converter phenotype and reduced nicotine demethylase activity from said first locus.

8. The tobacco product according to claim **1**, wherein said tobacco plant is a hybrid.

9. The tobacco product according to claim **1**, wherein said tobacco plant is selected from the group consisting of a Burley type, a dark type, a flue-cured type, and an Oriental type.

10. The tobacco product according to claim 1, wherein said first mutation results in a substitution at an amino acid residue in a position selected from the group consisting of residues 235, 449, 174, 410, 224, 72, 143, and 422, wherein said amino acid position numbering is according to SEQ ID NO:2.

11. The tobacco product according to claim 1, wherein said first mutation results in a Tryptophan to stop codon substitution at amino acid position 422, wherein said amino acid position numbering is according to SEQ ID NO:2.

12. The tobacco product according to claim **2**, wherein said first or second mutation is selected from the group consisting of a point mutation, a deletion, an insertion, and an inversion.

13. The tobacco product according to claim 2, wherein said second mutation results in a Tryptophan to stop codon substitution at amino acid position 329, wherein said amino acid position numbering is according to SEQ ID NO:13.

14. The tobacco product according to claim 2, wherein said tobacco plant is homozygous for said first or second mutation, or both.

15. Cured tobacco material made from a tobacco plant comprising a first mutation within the coding region of a wild type allele encoding a nicotine demethylase having the amino acid sequence of SEQ ID NO:2, wherein said first mutation reduces the activity of said nicotine demethylase.

16. The cured tobacco material according to claim **15**, wherein said tobacco plant further comprises a second mutation within the coding region of a wild type allele encoding a nicotine demethylase having the amino acid sequence of SEQ ID NO: 13, wherein said second mutation reduces the activity of said nicotine demethylase.

17. The tobacco product according to claim **11**, wherein said second mutation results in a Tryptophan to stop codon substitution at amino acid position 329, wherein said amino acid position numbering is according to SEQ ID NO:13.

* * * * *