

7-6-2014

Effects of Plant Virus and Its Insect Vector on *Encarsia Formosa*, a Biocontrol Agent of Whiteflies

Xiaoyuan Liu

Chinese Academy of Agricultural Sciences, China

Wensheng Xiang

Northeast Agricultural University, China

Xiaoguo Jiao

Chinese Academy of Agricultural Sciences, China

Youjun Zhang

Chinese Academy of Agricultural Sciences, China

Wen Xie

Chinese Academy of Agricultural Sciences, China

See next page for additional authors

Right click to open a feedback form in a new tab to let us know how this document benefits you.

Follow this and additional works at: https://uknowledge.uky.edu/entomology_facpub

 Part of the [Entomology Commons](#)

Repository Citation

Liu, Xiaoyuan; Xiang, Wensheng; Jiao, Xiaoguo; Zhang, Youjun; Xie, Wen; Wu, Qingjun; Zhou, Xuguo; and Wang, Shaoli, "Effects of Plant Virus and Its Insect Vector on *Encarsia Formosa*, a Biocontrol Agent of Whiteflies" (2014). *Entomology Faculty Publications*. 54. https://uknowledge.uky.edu/entomology_facpub/54

This Article is brought to you for free and open access by the Entomology at UKnowledge. It has been accepted for inclusion in Entomology Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Authors

Xiaoyuan Liu, Wensheng Xiang, Xiaoguo Jiao, Youjun Zhang, Wen Xie, Qingjun Wu, Xuguo Zhou, and Shaoli Wang

Effects of Plant Virus and Its Insect Vector on *Encarsia Formosa*, a Biocontrol Agent of Whiteflies**Notes/Citation Information**

Published in *Scientific Reports*, v. 4, 5926.

This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder in order to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>

Digital Object Identifier (DOI)

<http://dx.doi.org/10.1038/srep05926>



OPEN

SUBJECT AREAS:

AGRI-ECOLOGY

INVASIVE SPECIES

BEHAVIOURAL ECOLOGY

Received

8 January 2014

Accepted

13 June 2014

Published

6 August 2014

Correspondence and requests for materials should be addressed to S.W. (wangshaoli@caas.cn) or X.Z. (xuguozhou@uky.edu)

Effects of plant virus and its insect vector on *Encarsia formosa*, a biocontrol agent of whiteflies

Xiaoyuan Liu^{1,2}, Wensheng Xiang², Xiaoguo Jiao¹, Youjun Zhang¹, Wen Xie¹, Qingjun Wu¹, Xuguo Zhou³ & Shaoli Wang¹

¹Department of Plant Protection, Institute of Vegetables and Flowers, Chinese Academy of Agricultural Sciences, Beijing, 100081, P. R. China, ²Department of Life Science, Northeast Agricultural University, Harbin 150030, P. R. China, ³Department of Entomology, University of Kentucky, Lexington, KY 40546-0091, U.S.A.

In this study, we investigated the tritrophic interactions among a persistently transmitted plant virus, *Tomato yellow leaf curl virus* (TYLCV), its insect vector, the sweetpotato whitefly *Bemisia tabaci*, and a parasitoid, *Encarsia formosa* Gahan, one of the most extensively used biological control agents. As an emerging invasive pest worldwide, the two most damaging whiteflies are *B. tabaci* B and Q cryptic species. On healthy tomato plants, parasitoid-induced mortality was significantly higher in *B. tabaci* B than in Q. In contrast, similar mortality levels of B and Q were observed on TYLCV-infected plants. A higher rate of parasitism was consistently observed in B, independent of the TYLCV infection. Similarly, the life history traits of *E. formosa* were influenced by both TYLCV and the two cryptic species of *B. tabaci*. Specifically, *E. formosa* parasitizing B had a greater adult longevity and shorter developmental time on healthy plants, whereas the parasitoids developing from Q has a greater adult longevity on TYLCV-infected plants. The emergence rate of *E. formosa* was unaffected by either *B. tabaci* cryptic species or the virus. These results suggest that the vector-borne pathogen can manipulate the host suitability of a parasitoid and hence the parasitoid-host interactions.

The sweetpotato whitefly, *Bemisia tabaci* (Gennadius) (Homoptera: Aleyrodidae), attacks many economically important crops, including vegetables, cotton, and ornamentals. Recent studies suggest that *B. tabaci* is a species complex containing at least 24 morphologically indistinguishable cryptic species^{1–3}. The two most invasive and widespread groups of *B. tabaci* are the Middle East-Asia Minor I species (referred to as B hereafter) and the Mediterranean species (referred to as Q hereafter)^{1–3}. The *B. tabaci* B was first recorded in China in the early 1990s, and it subsequently spread throughout much of the country, but has been recently displaced by Q in many areas^{4,5}.

Although *B. tabaci* can damage crops directly by feeding, it causes greater damage by transmitting over 100 begomoviruses that can have catastrophic impacts on plant growth and survival³. *Tomato yellow leaf curl virus* (TYLCV) is a single-stranded DNA plant virus that originated in the Middle East⁶. TYLCV is transmitted by *B. tabaci*, the only known vector, in a circulative, persistent manner⁷. TYLCV was first detected in Shanghai, China in 2006⁸ and then quickly spread from the Southern to Northern parts of China⁹. Although both B and Q can vector TYLCV horizontally, Q exhibited a much higher efficiency than B; as a result, Q appears to be a more competent viral vector than B^{9–11}. The spread of TYLCV coincided with the rapid displacement of the *B. tabaci* B by Q in China⁹. Because both cryptic species have developed resistance to various synthetic insecticides, especially in Q^{12,13}, biological control is one of the best alternatives available for the sustainable management of *B. tabaci* populations.

Encarsia formosa Gahan (Hymenoptera: Aphelinidae) is a solitary endoparasitoid that has been used commercially for the biocontrol of whiteflies, including the sweetpotato whitefly *B. tabaci* and the greenhouse whitefly *Trialeurodes vaporariorum*, worldwide¹⁴. As one of the most extensively used natural enemies, *E. formosa* parasitizes at least fifteen species of whiteflies in eight genera¹⁴. Specially, adult females parasitize and kill whitefly nymph, although the developmental time of *E. formosa* depends on the developmental stage of *B. tabaci* nymph¹⁵.

Bemisia tabaci Q has now become the primary whitefly cryptic species in China^{4,5}, and as noted, the rapid spread of Q has been accompanied by the recent outbreaks of TYLCV⁹. Information regarding the tritrophic interactions among TYLCV, a persistently transmitted plant virus, *B. tabaci*, the only known insect vector, and *E.*




Figure 1 | Biocontrol properties of *E. formosa* on the sweetpotato whitefly *B. tabaci*. The 3rd instar nymphs of *B. tabaci* B and Q cryptic species were reared on healthy (NV) and TYLCV-infected tomato plants (V), respectively, before submitting them to a hymenopterous parasitoid, *E. formosa*. The percent mortality (A) and the parasitism rate (B) of *B. tabaci* were documented. Values (means \pm SE) are from three biological replications. Within each panel, different letters indicate significant differences among treatments ($P < 0.05$).

formosa, the primary biocontrol agent for whiteflies, are grossly lacking. In this study, we are specifically investigating the impacts of this virus and its insect vector on the performance and life history traits of *E. formosa*. A better understanding of such interactions will, in turn, offer valuable insight into the long-term sustainable control of whiteflies, with or without the occurrence of epidemic plant virus.

Results

Biocontrol properties of *E. formosa* on the sweetpotato whitefly *B. tabaci*. The mortality rate of *E. formosa* was significantly affected by the *B. tabaci* cryptic species ($F_{3, 236} = 34.759$, $P < 0.0001$), virus infection status ($F_{3, 236} = 54.068$, $P < 0.0001$), and the interaction between the two ($F_{3, 236} = 23.432$, $P < 0.0001$). The mortality rate was significantly lower in Q reared on healthy plants than in B reared

on healthy plants or in either B or Q reared on TYLCV-infected plants (Figure 1A, Figure S1).

The parasitism rate was significantly affected by the *B. tabaci* cryptic species ($F_{3, 236} = 190.939$, $P < 0.0001$), virus infection status ($F_{3, 236} = 15.896$, $P < 0.0001$), and the interaction between the two ($F_{3, 236} = 27.190$, $P < 0.0001$). The parasitism rate was significantly lower in Q reared on healthy plants than in B reared on healthy plants or in either B or Q reared on TYLCV-infected plants. In addition, the parasitism rate was lower in Q reared on TYLCV-infected plants than in B reared on healthy or TYLCV-infected plants (Figure 1B, Figure S1).

The impacts of TYLCV and *B. tabaci* cryptic species on the life history traits of *E. formosa*. The developmental time of *E. formosa* was significantly affected by the virus infection status ($F_{3, 236} =$



10.960, $P = 0.001$) and the interaction between *B. tabaci* cryptic species and virus infection status ($F_{3, 236} = 11.954$, $P = 0.001$), but was not significantly affected by *B. tabaci* cryptic species ($F_{3, 236} = 3.640$, $P = 0.058$). Developmental time was shorter in B reared on healthy plants than on TYLCV-infected plants or Q reared on either healthy plants or TYLCV-infected plants. But it was not significantly different among B reared on TYLCV-infected plants or Q reared on either healthy plants or TYLCV-infected plants (Figure 2A, Figure S2).

The emergence rate of *E. formosa*, i.e., the percentage of pupae from which adult wasps emerged, was not significantly affected by *B. tabaci* cryptic species ($F_{3, 236} = 0.723$, $P = 0.396$), virus infection status ($F_{3, 236} = 0.238$, $P = 0.626$), or the interaction between the two ($F_{3, 236} = 2.239$, $P = 0.136$) (Figure 2B, Figure S2).

The longevity of *E. formosa* was significantly affected by the *B. tabaci* cryptic species ($F_{3, 236} = 6.086$, $P = 0.014$), virus infection status ($F_{3, 236} = 80.644$, $P < 0.0001$), and the interaction between the two ($F_{3, 236} = 294.839$, $P < 0.0001$). Longevity was significantly shorter in Q reared on healthy plants than on TYLCV-infected plants or B reared on either healthy plants or TYLCV-infected plants. In addition, the longevity of *E. formosa* parasitizing *B. tabaci* Q from the TYLCV-infected tomato was significantly higher than those wasps parasitized whiteflies from the TYLCV-free tomato. In case of *B. tabaci* B, however, the trend was reversed (Figure 2C, Figure S2).

Discussion

The influence of insect vector on the performance and fitness of *E. formosa*. Although parasitism of *B. tabaci* by *E. formosa* has been studied previously^{16–18}, none of them has been focused on the potential impact of cryptic species. As the most invasive and destructive *B. tabaci* cryptic species, B and Q vary in their mating and feeding behavior and prefer different host plants^{10,19,20}. On healthy tomato plants, *E. formosa* performed better (higher mortality and parasitism rates) on *B. tabaci* B than on Q (Figure 3). This is consistent with the results from a congeneric parasitoid, *E. sophia*²¹. In regard to fitness, the developmental times of *E. formosa* on the 3rd instar nymphs of *B. tabaci* were 14.16 and 14.52 days, respectively for B and Q. A similar trend was documented in *E. sophia*, in which the developmental time of the parasitoid on *B. tabaci* B and Q were not significantly different²¹. Similarly, the emergence rate of *E. formosa* was not significantly affected by the two cryptic species. As for adult longevity, however, *E. formosa* lived longer in B than in Q. The combined results from parasitoid's performance and fitness indicate that *B. tabaci* B is a more suitable host for the *Encarsia* spp. than Q, although the underlying mechanisms are largely unknown. The *E. formosa* colony was obtained initially from the United Kingdom and was introduced into the vegetable-growing areas in China to control whiteflies in the 1980s. *E. formosa* has been acclimated into the field conditions in China for over 30 years, and experienced the invasions of *B. tabaci* B and Q in the 1990s and 2000s, respectively²². By 2011, Q had displaced B and become the dominant cryptic *B. tabaci* species in most areas in China^{5,23}. The present study suggests that parasitic *E. formosa* is more effective against *B. tabaci* B than against Q, and this difference in their suitability for the parasitoid could partially contribute to the competitive displacement of *B. tabaci* B by Q in China.

The impact of virus infection of the host plant on the performance and fitness of *E. formosa*. Virus induced plant responses can influence the behavior and physiology of insect vectors, and, sometimes, lead to behavioral changes in the vectors that favor viral transmission^{24,25}. Although tritrophic interactions among vector-borne pathogens, its vector, and the parasitoid (*E. formosa*) add another layer of complexity, it is germane to speculate that vector-borne pathogen can exert its impact on a parasitoid through the vector. Multiple lines of evidence have shown that TYLCV can manipulate

the behavior (e.g. host preference) of its exclusive vector, *B. tabaci* species complex, for the benefit of viral transmission^{9,10,26,27}. TYLCV-infected whiteflies fed more readily than uninfected individuals and spent more time salivating, which is essential for the spread of a persistently transmitted plant virus¹⁰. Field surveys indicate an apparent mutualistic relationship between the outbreaks of TYLCV and the introduction and rapid range expansion of *B. tabaci* Q⁵. Plant defensive compounds, specifically terpenoids, play a key role in mediating vector–pathogen mutualistic relationships. Gas chromatography–mass spectrometry (GCMS) analysis showed that healthy plants released significantly more terpenoids than TYLCV-infected plants²⁶. The higher level of terpenoids, a suite of defensive compounds from plants, however, compromised the fitness level of *B. tabaci*. Suppression of terpenoid synthesis pathway via gene silencing, in return, enhanced the fitness level of *B. tabaci*²⁷. That means TYLCV can influence host-derived cues in ways that have positive effects on vector attraction^{10,26}. Among the two cryptic species, *B. tabaci* Q preferentially settled and oviposited on TYLCV-infected plants rather than on healthy plants, and had a better overall fitness, including a greater fecundity, survival rate, and female body size, and a shorter developmental time^{9,28}.

In this study, a differential response was observed toward virus infection among the two *B. tabaci* cryptic species. Specifically, for B, the virus has no apparent effects on either the performance or the fitness of *E. formosa*. Increased mortality and parasitism rates, as well as enhanced life history traits, however, were documented in *E. formosa* parasitizing Q from TYLCV-infected tomatoes. Improved parasitism or predation by the virus-infected host is not without precedent. Belliure *et al.* noticed an increase in the developmental time of thrips on infected host plants, which led a reduced risk of predation by predatory mites²⁹. Infected with *Tomato spotted wilt virus*, thrips larvae grew too big to be preyed on by the predatory mites. However, when the prey size is not the limiting factor, plant virus could not influence the predation risk (a predatory beetle can consume all sizes of thrips larvae)²⁹. The seven-spotted ladybug, *Coccinella septempunctata*, consumed more aphids when the aphids were infected with *Rhopalosiphum padi virus*³⁰. Also, a higher rate of parasitism and the greater overall population suppression by the parasitoid *Aphidius colemani* were observed when their aphid hosts were infected with *Cereal yellow dwarf virus*³¹.

Conclusion. A persistently transmitted virus, like TYLCV, can enhance the quality of host plant to attract insect vector and increase the viral transmission³². The quality of the insect vector, the host for parasitoids, can be improved accordingly. Higher quality of the host, consequently, can impact the performance of parasitoids and can, potentially, shift their host preference (Figure 3). Previous study showed that TYLCV-infected plants improved the growth, development, and fitness of Q but decreased the fitness of B⁹. In this study, when infected, the fitness level of *E. formosa* decreased in B, whereas both fitness and performance of the parasitic wasp were significantly improved in Q.

The present work showed the improved parasitism of *E. formosa* on the host of *B. tabaci* Q when the virus factor was added. The combined results suggest that the parasitoids may contribute to the suppression of TYLCV in the field by disproportionately influencing the vectors feeding on the infected plants. In addition, for the concurrent control of *B. tabaci* B and Q in the field, our results are encouraging in the context of integrated pest management, because they suggest that the current biological practice with parasitoids may contribute indirectly to the management of TYLCV by impacting on more competent virus-carrying vectors, *B. tabaci* Q, and consequently reducing the proportion of vectors in the population that are infectious.




Figure 2 | The impact of TYLCV and *B. tabaci* cryptic species on the life history traits of *E. formosa*. To evaluate the potential impacts of plant virus and insect vector on a primary whitefly parasitoid, life history traits, including the developmental time (A), emergence rate (B), and longevity (C), of *E. formosa* were recorded. The 3rd instar nymphs of *B. tabaci* B and Q cryptic species were reared on healthy (NV) and TYLCV-infected tomato plants (V), respectively. Values (means \pm SE) are from three biological replications. Within each panel, different letters indicate significant differences among treatments ($P < 0.05$).




Figure 3 | Patterns of parasite behavior in relation to plant pathogen and insect vectors. In this schematic, the parasitoid is an endoparasitoid, *E. formosa*, the pathogen is a plant virus, *Tomato yellow leaf curl virus* (infected plant is discoloured in yellow), and the vector is the sweetpotato whitefly *B. tabaci* B and Q cryptic species. Both insect vector and plant virus have a significant influence on the performance and fitness of *E. formosa*. Parasitism was performed better on *B. tabaci* B than on Q on healthy plants. Whereas, virus induced plant responses influences the behavior and physiology of the insect vector, and consequently exerts its impact on the parasitoids indirectly. For *B. tabaci* Q, TYLCV had a positive effect on both performance and fitness of *E. formosa*. In contrast, a negative impact on the life history traits of *E. formosa* was observed when B was fed on the infected tomato plants. Please note the number of *B. tabaci* nymphs infected with *E. formosa* is purely hypothetical, implicating either positive (more) or negative (less) effects.

Methods

Insect and parasitoid cultures. *Bemisia tabaci* B population was originally collected from an infested cabbage field in Beijing, China in 2004³³, and the Q population was collected from a poinsettia plant in Beijing, China in 2009. Their identities were confirmed by random amplified polymorphic DNA polymerase chain reaction (RAPD-PCR)³⁴. Both *B. tabaci* populations were maintained on tomato plants in a growth chamber at $27 \pm 1^\circ\text{C}$, $70 \pm 5\%$ RH, and a 16/8 L/D photoperiod.

Encarsia formosa culture was a gift from the Beijing Academy of Agriculture and Forestry Sciences. The parasitic wasp had been maintained on *B. tabaci* B on tobacco plants for more than five generations. Upon arrival in our laboratory, the original culture was divided into two groups; one culture was maintained on B and the other was maintained on Q.

Plant materials. Tomato (*L. esculentum* Mill, No. 9 Zhongza) was used as the host plant for *B. tabaci* B and Q. The tomato plants were grown in the greenhouse and covered with 100-mesh insect-proof screen to exclude whiteflies and other pests.

TYLCV-infected tomato plants were obtained by *Agrobacterium tumefaciens*-mediated inoculation using a cloned TYLCV genome (GenBank accession ID: AM282874). The plants were inoculated at the 3 to 4 true-leaf stage. TYLCV infection of the inoculated tomato plants was confirmed using a set of TYLCV primers C473 and V61³⁵. *Bemisia tabaci* B and Q populations were maintained on healthy and TYLCV-infected tomato plants, respectively, when host plants reached 6 to 7 true-leaf stage.

Effects of *B. tabaci* cryptic species and plant virus on the parasitism of *E. formosa*.

The parasitoid-induced *B. tabaci* mortality, the percentage of *B. tabaci* parasitized, the percentage of *E. formosa* pupae from which adult wasps emerged, *E. formosa* longevity, and *E. formosa* developmental time were determined using the clip-cages method. 3rd-instar nymphs were used in this study because the development of *E. formosa* is more synchronized at this life stage^{17,36}. Twenty newly emerged *B. tabaci* adults (B or Q) were placed on the abaxial surface of a selected host leaf of a tomato plant (infected or not infected with TYLCV); the adults and leaf were contained in a leaf clip-cage (one clip cage per plant). After 24 h, the adults were removed, and the eggs were monitored and counted daily until they developed into 3rd-instar nymphs. The number of 3rd-instar nymphs was then adjusted to 30 per clip cage. One newly emerged *E. formosa* female was then introduced into each leaf clip-cage, and each of the four treatments (two cryptic species and two virus conditions) was represented by 20 replicate clip-cages. The plants with the cages were maintained in growth chambers at $27 \pm 1^\circ\text{C}$, $70 \pm 5\%$ RH, and a 14/10 L/D photoperiod. After 24 h, the adult *E. formosa* was removed from each clip-cage. When parasitized, each *B. tabaci* nymph was turned into one *E. formosa* pupa under the above mentioned

experimental conditions. The endpoint measurements, including parasitism rate, mortality, and emergence rate were determined for each cohort of 30 *B. tabaci* nymphs in their respective clip-cage after 6–10 days assay period.

A second experiment was carried out to determine the impact of *B. tabaci* cryptic species and plant virus on *E. formosa* longevity. This experiment used the same conditions and similar materials as described in the previous paragraph. One newly emerged *E. formosa* female adult was introduced into a clip-cage on a tomato leaf (infected or not infected with TYLCV) infested with 30 3rd-instar nymphs of *B. tabaci* (B or Q). Each of the four treatments (two cryptic species and two virus conditions) was represented by 20 replicate clip-cages. Each day until the wasp died, the wasp in each clip-cage was transferred to a new leaf with 30 3rd-instar nymphs. The longevity of each *E. formosa* adult was recorded. Both experiments have been independently replicated three times.

Data analysis. Data were first checked for normality and transformed when necessary to meet the assumption of normal distribution. The data for parasitism rate, *B. tabaci* mortality rate, and *E. formosa* emergence rate were arcsine-square root transformed for analyses. The generalized linear model (GLM, two-way ANOVA) was used to test the effects of *B. tabaci* cryptic species, virus infection status of the tomato plant, and the interaction of cryptic species and virus infection status on the following parameters: *B. tabaci* mortality rate, the percentage of *B. tabaci* parasitized by *E. formosa*, the emergence rate of *E. formosa* adults from pupae, the longevity of *E. formosa* adults, and *E. formosa* developmental time. SPSS (version 13.0; SPSS Inc., Chicago, IL, USA) was used for statistical analyses (one-way ANOVA, Tukey's test).

- Dinsdale, A., Cook, L., Riginos, C., Buckley, Y. M. & De Barro, P. Refined global analysis of *Bemisia tabaci* (Hemiptera: Sternorrhyncha: Aleyrodoidea: Aleyrodidae) mitochondrial cytochrome oxidase I to identify species level genetic boundaries. *Ann. Entomol. Soc. Am.* **103**, 196–208 (2010).
- Xu, J., De Barro, P. J. & Liu, S. S. Reproductive incompatibility among genetic groups of *Bemisia tabaci* supports the proposition that the whitefly is a cryptic species complex. *Bull. Entomol. Res.* **100**, 359–366 (2010).
- De Barro, P. J., Liu, S. S., Boykin, L. M. & Dinsdale, A. B. *Bemisia tabaci*: A statement of species status. *Annu. Rev. Entomol.* **1**, 1–19 (2011).
- Chu, D., Wan, F. H., Zhang, Y. J. & Brown, J. K. Change in the biotype composition of *Bemisia tabaci* in Shandong Province of China from 2005 to 2008. *Environ. Entomol.* **39**, 1028–1036 (2010).
- Pan, H. P. *et al.* Further spread of and domination by *Bemisia tabaci* biotype Q on field crops in China. *J. Econ. Entomol.* **104**, 978–985 (2011).
- Varma, A. & Malathi, V. G. Emerging geminivirus problems: a serious threat to crop production. *Ann. Appl. Biol.* **142**, 145–164 (2003).



7. Brown, J. K. & Czosnek, H. Whitefly transmission of plant viruses. *Adv. Bot. Res.* **36**, 65–76 (2002).
8. Wu, J. B., Dai, F. M. & Zhou, X. P. First report of *Tomato yellow leaf curl virus* in China. *Plant Disease* **90**, 1359–1359 (2006).
9. Pan, H. P. *et al.* Rapid spread of *Tomato yellow leaf curl virus* in China is aided differentially by two invasive whiteflies. *PLoS ONE* **7**, e34817 (2012).
10. Liu, B. M. *et al.* Multiple forms of vector manipulation by a plant-infecting virus: *Bemisia tabaci* and *tomato yellow leaf curl virus*. *J. Virol.* **87**, 4929–4937 (2013).
11. Jiang, Y. X., de Blas, C., Barrios, L. & Fereres, A. Correlation between whitefly (Homoptera: Aleyrodidae) feeding behavior and transmission of *tomato yellow leaf curl virus*. *Ann. Entomol. Soc. Am.* **93**, 573–579 (2000).
12. Luo, C. *et al.* Insecticide resistance in *Bemisia tabaci* biotype Q (Hemiptera: Aleyrodidae) from China. *Crop Prot.* **29**, 429–434 (2010).
13. Li, X. C. *et al.* Baseline susceptibilities of B- and Q-biotype *Bemisia tabaci* to anthranilic diamides in Arizona. *Pest Manag. Sci.* **68**, 83–91 (2012).
14. Hoddle, M. S., Van Driesche, R. G. & Sanderson, J. P. Biology and use of the whitefly parasitoid *Encarsia formosa*. *Annu. Rev. Entomol.* **43**, 645–69 (1998).
15. Zhang, S. Z., Wan, F. H., Zhang, F. & Hua, B. Z. Parasitic suitability of two strains of *Encarsia formosa* on *Bemisia tabaci*. *Chinese J. Biol. Control* **19**, 149–153 (2003).
16. Qiu, Y. T., Van Lenteren, J. C., Drost, Y. C. & Posthuma–Doodeman, J. A. M. Life-history parameters of *Encarsia formosa*, *Eretmocerus eremicus* and *E. mundus*, aphelinid parasitoids of *Bemisia argentifoli* (Hemiptera: Aleyrodidae). *Eur. J. Entomol.* **101**, 83–94 (2004).
17. Takahashi, K. M., Berti Filho, E. & Lourenção, A. L. Biology of *Bemisia tabaci* (Genn.) B-biotype and parasitism by *Encarsia formosa* (Gahan) on collard, soybean and tomato plants. *Sci. Agr.* **65**, 639–642 (2008).
18. Grille, G., Lorenzo, M. E., Burla, J. P., Franco, J. & Basso, C. Parasitoid Niches of *Encarsia formosa* and *Encarsia Lycopersici* (Hymenoptera: Aphelinidae) Exploiting *Trialeurodes vaporariorum* (Hemiptera: Aleyrodidae). *Fla. Entomol.* **95**, 1024–1030 (2012).
19. Elbaz, M., Lahav, N., & Morin, S. Evidence for pre-zygotic reproductive barrier between the B and Q biotypes of *Bemisia tabaci* (Hemiptera: Aleyrodidae). *B. Entomol. Res.* **100**, 581–590 (2010).
20. Tsueda, H., & Tsuchida, K. Reproductive differences between Q and B whiteflies, *Bemisia tabaci*, on three host plants and negative interactions in mixed cohorts. *Entomol. Exp. Appl.* **141**, 197–207 (2011).
21. Wang, J. H., Luo, C. & Liu, T. X. Effects of *Bemisia tabaci* (Hemiptera: Aleyrodidae) biotype on host selection and development of *Encarsia Sophia* (Hymenoptera: Aphelinidae). *Acta. Entomol. Sin.* **54**, 687–693 (2011).
22. Chu, D. *et al.* The introduction of the exotic biotype Q of *Bemisia tabaci* from the Mediterranean region into China on ornamental crops. *Fla. Entomol.* **89**, 168–174 (2006).
23. Wang, Z. Y., Yan, H. F., Yang, Y. H. & Wu, Y. D. Biotype and insecticide resistance status of the whitefly *Bemisia tabaci* from China. *Pest Manag. Sci.* **66**, 1360–1366 (2010).
24. Zang, L. S. & Liu, T. X. Host-feeding of three parasitoid species on *Bemisia tabaci* biotype B and implications for whitefly biological control. *Entomol. Exp. Appl.* **127**, 55–63 (2008).
25. Stafford, C. A., Walker, G. P. & Ullman, D. E. Infection with a plant virus modifies vector feeding behavior. *P. Natl. Acad. Sci. USA.* **108**, 9350–9355 (2011).
26. Fang, Y. *et al.* *Tomato yellow leaf curl virus* alters the host preferences of its vector *Bemisia tabaci*. *Sci. Rep.* **3**, 2876 (2013).
27. Luan, J. B. *et al.* Suppression of terpenoid synthesis in plants by a virus promotes its mutualism with vectors. *Eco. Lett.* **16**, 390–398 (2013).
28. Chen, G. *et al.* Virus infection of a weed increases vector attraction to and vector fitness on the weed. *Sci. Rep.* **3**, 2253 (2013).
29. Belliure, B., Janssen, A. & Sabelis, M. W. Herbivore benefits from vectoring plant virus through reduction of period of vulnerability to predation. *Oecologia.* **156**, 797–806 (2008).
30. Ban, L., Ahmed, E., Ninkovic, V., Delp, G. & Glinwood, R. Infection with an insect virus affects olfactory behaviour and interactions with host plant and natural enemies in an aphid. *Entomol. Exp. Appl.* **127**, 108–117 (2008).
31. de Oliveira, C. F., Long, E. Y. & Finke, D. L. A negative effect of a pathogen on its vector? A plant pathogen increases the vulnerability of its vector to attack by natural enemies. *Oecologia* **174**, 1169–1177 (2014).
32. Mauck, K., Bosque-Pérez, N. A., Eigenbrode, S. D., DeMoraes, C. M. & Mescher, M. C. Transmission mechanisms shape pathogen effects on host-vector interactions: evidence from plant viruses. *Funct. Ecol.* **26**, 1162–1175 (2012).
33. Xie, W. *et al.* Induction effects of host plants on insecticide susceptibility and detoxification enzymes of *Bemisia tabaci* (Hemiptera: Aleyrodidae). *Pest Manag. Sci.* **67**, 87–93 (2011).
34. Wu, X. X., Hu, D. X., Li, Z. X. & Shen, Z. M. Using RAPD-PCR to distinguish biotypes of *Bemisia tabaci* (Homoptera: Aleyrodidae) in China. *Entomol. Sin.* **9**, 1–8 (2002).
35. Ghanim, M., Morin, S., Zeidan, M. & Czosnek, H. Evidence for transovarial transmission of *Tomato Yellow Leaf Curl Virus* by its vector, the whitefly *Bemisia tabaci*. *Virology* **240**, 295–303 (1998).
36. Hu, J. S., Gelman, D. B. & Blackburn, B. B. Age-specific interaction between the parasitoid, *Encarsia formosa* and its host, the silverleaf whitefly, *Bemisia tabaci* (Strain B). *J. Insect Sci.* **3**, 28 (2003).

Acknowledgments

Special thanks go to John J. Obrycki (University of Kentucky) for his comments and English editing. This research was supported by the 973 Program (2013CB127602), the National Natural Science Foundation of China (31171857), the China Agriculture Research System (CARS-26-10), and the Beijing Key Laboratory for Pest Control and Sustainable Control. The granting agencies had no role in study design, data collection and analysis, decision to publish, or manuscript preparation.

Author contributions

S.L.W., Y.J.Z., X.G.Z. and X.Y.L. designed the experiment. X.Y.L. performed the experiment. Q.J.W., S.L.W., W.X. and W.S.X. contributed reagents/materials. X.Y.L., S.L.W., X.G.J. and X.G.Z. wrote the paper. All authors reviewed the manuscript.

Additional information

Supplementary information accompanies this paper at <http://www.nature.com/scientificreports>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Liu, X. *et al.* Effects of plant virus and its insect vector on *Encarsia formosa*, a biocontrol agent of whiteflies. *Sci. Rep.* **4**, 5926; DOI:10.1038/srep05926 (2014).



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder in order to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>