



9-17-2012

Serotonin and Synaptic Transmission at Invertebrate Neuromuscular Junctions

Wen-Hui Wu
University of Kentucky

Robin L. Cooper
University of Kentucky, rlcoop1@uky.edu

Right click to open a feedback form in a new tab to let us know how this document benefits you.

Follow this and additional works at: https://uknowledge.uky.edu/biology_facpub

 Part of the [Biology Commons](#)

Repository Citation

Wu, Wen-Hui and Cooper, Robin L., "Serotonin and Synaptic Transmission at Invertebrate Neuromuscular Junctions" (2012). *Biology Faculty Publications*. 60.

https://uknowledge.uky.edu/biology_facpub/60

This Article is brought to you for free and open access by the Biology at UKnowledge. It has been accepted for inclusion in Biology Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Serotonin and Synaptic Transmission at Invertebrate Neuromuscular Junctions

Notes/Citation Information

Published in *Experimental Neurobiology*, v. 21, no. 3, p. 101-112.

Copyright © Experimental Neurobiology 2012.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Digital Object Identifier (DOI)

<http://dx.doi.org/10.5607/en.2012.21.3.101>

Serotonin and Synaptic Transmission at Invertebrate Neuromuscular Junctions

Wen-Hui Wu and Robin L. Cooper*

Department of Biology & Center for Muscle Biology, University of Kentucky, Lexington, KY 40506-0225, USA

The serotonergic system in vertebrates and invertebrates has been a focus for over 50 years and will likely continue in the future. Recently, genomic analysis and discovery of alternative splicing and differential expression in tissues have increased the knowledge of serotonin (5-HT) receptor types. Comparative studies can provide useful insights to the wide variety of mechanistic actions of 5-HT responsible for behaviors regulated or modified by 5-HT. To determine cellular responses and influences on neural systems as well as the efferent control of behaviors by the motor units, preparations amenable to detailed studies of synapses are beneficial as working models. The invertebrate neuromuscular junctions (NMJs) offer some unique advantages for such investigations; action of 5-HT at crustacean NMJs has been widely studied, and leech and *Aplysia* continue to be key organisms. However, there are few studies in insects likely due to the focus in modulation within the CNS and lack of evidence of substantial action of 5-HT at the *Drosophila* NMJs. There are only a few reports in gastropods and annelids as well as other invertebrates. In this review we highlight some of the key findings of 5-HT actions and receptor types associated at NMJs in a variety of invertebrate preparations in hopes that future studies will build on this knowledge base.

Key words: synapse, reserve pool, readily releasable pool

GENERAL BACKGROUND OF 5-HYDROXYTRYPTAMINE AND RECEPTORS

5-Hydroxytryptamine (5-HT, serotonin) is a common biogenic amine found in both vertebrates and invertebrates as well as in plants [1, 2]. The precursor to 5-HT, tryptophan, is likely important in the early evolution of life and perhaps the early presence of tryptophan is a reason for 5-HT to be potentially the first neurotransmitter noted with the development of a nervous system [2]. 5-HT acts as both a neurotransmitter and neurohormone and as a potent modulator of neurons and various

tissues in many animal species [3]. Generally 5-HT actions are elicited by transmembrane G protein coupled receptors (GPCRs), which then activate or inhibit different intracellular second messenger cascades. 5-HT receptors from some organisms have been classified based on sequence or pharmacology [4, 5]; for example in the vertebrates, 7 families (5-HT₁₋₇), 14 subtypes have been identified, whereas in *Drosophila* four 5-HT receptors named 5-HT_{1Adro} 5-HT_{1Bdro} 5-HT_{2dro} 5-HT_{7dro} [5-10] have been classified. 5-HT receptors appear to be present on invertebrate presynaptic nerve terminals and on muscle membranes; receptors of a cricket (*Gryllus domestica*) mandibular muscle have a similar pharmacological profile as a 5-HT₂-like receptor subtype [11]. Profiling the 5-HT receptor subtypes directly on skeletal muscle within invertebrates is an area of research that is lacking. The 5-HT₄ and 5-HT₇ receptors are shown to have alternate splice variants which increase the number of receptor subtypes and may

Received June 2, 2012, Accepted June 19, 2012

*To whom correspondence should be addressed.
TEL: 1-859-257-5950, FAX: 1-859-257-1717
e-mail: RLCOOP1@email.uky.edu

alter the selectivity to pharmacological agents [12]. In addition, 5-HT₂ receptors can have different RNA-edited isoforms [13, 14].

With the use of the genetically modifiable model *D. melanogaster*, a number of studies have examined over-expression and under-expression of receptors subtypes on the effects of development, behavior and physiology as well as the general actions of 5-HT in *D. melanogaster* [7, 15-19]. Based on physiological and pharmacological studies in crustaceans there may be a larger number of 5-HT receptors present than in *D. melanogaster* [20-26]. Two receptors types have been cloned and characterized in crustaceans [9, 27, 28] and in a pond snail [29]. A 5-HT receptor 5-HT(apAC1) has been cloned, sequenced and characterized in *Aplysia* sensory neurons [30]. 5-HT receptors are being cloned in a variety of invertebrates and surely more will be forthcoming with the rapid development in genomic sequencing abilities. There are a plethora of reports on the effects of 5-HT for sensory and central neurons as well as on behaviors in invertebrates which are worthy of multiple reviews. However, for this brief review we focus on the physiological effects of 5-HT at the skeletal neuromuscular junctions in some of the key model invertebrates. The invertebrate neuromuscular junctions (NMJs) are very diverse across species and within species in structure and function [31-37]. The recent majority of reports on structure and function of NMJs are of *D. melanogaster* due to the genetic approaches and manipulations being utilized [38-41].

WHY FOCUS ON NMJS?

The synaptic communication between neurons and target cells depends on the specialized anatomy and physiology of the synapses [42]. The regulation and modulation of neurotransmitter release is the basis of chemical synaptic transmission. For nervous systems to function properly, the efficacy of synapses are finely regulated and adjustable to respond to changing circumstance and requirement. Too high or too low synaptic input both result in inappropriate communication of target cells. Both pre- and postsynaptic factors can influence the synaptic strength. The amount of neurotransmitter released and the sensitivity of the postsynaptic membrane both are important for measuring synaptic strength. Each step in the process of synaptic transmission can be the target of many factors that lead to alteration of synaptic strength. For example, the phosphorylation state of SNARE proteins that are involved in vesicle docking, or the density of active zones where transmitter is released, can influence the number of quantal units released per impulse (presynaptic mechanism). Postsynaptically, the number of active receptors, the postsynaptic input resistance, the area and the ultrastructure of

subsynaptic reticulum, all can alter the effectiveness of quantum release and thus influence synaptic strength.

The ease in accessibility to the synaptic sites at NMJs allows one to record intracellular or very close to synapses by extracellular recordings (focal macropatch over a varicosity) in order to minimize cable properties in signal decrement [43-45]. Such signal loss occurs with recordings in a neuron cell body to measure synaptic function in the dendritic trees. The localized recording over a NMJ allows one the ability to measure properties of single and multiple vesicular quanta for very precise quantal analysis (occurrences, size and shape) to index synaptic function [46-49]. In addition, invertebrate NMJs are relatively stable for hours in a minimal saline at room temperature as compared to mammalian NMJs. Since most muscles in invertebrates are innervated by relatively few motor neurons, for the most part, they are identifiable anatomically and physiologically from preparation to preparation [50, 51]. Since the fine structure and detailed quantal analysis is feasible for many invertebrate NMJs, the acute and chronic actions of modulators on structure and function can be examined for their mechanistic actions [52-54].

INSECTS

Given such a diverse group of animals within the class Insecta, it would not be surprising to find a wide range of anatomic and physiologic profiles in the innervation of skeletal NMJs. For example, the innervation of the genital chamber of the female cricket, *Acheta domestica*, shows 5-HT-immunoreactive nerve terminals that contact the muscle fibers which likely releases 5-HT in a type of volume transmission over the muscle as there are no defined synapses [55]. Such 5-HT containing nerve endings are also present in earthworm skeletal muscles [56]. However, no serotonin is associated with the oviducts or the innervation to the oviducts in the locust [57]. Earlier studies did not elucidate if the effect of 5-HT was directly on the presynaptic terminal or on the muscle but reported overall changes in force of muscle contraction. In a locust leg muscle, 5-HT produces an overall decrease in force development [58] but the mechanism of action still needs to be determined. It is suggested that in some of the earlier studies with insects, the high concentrations of 5-HT used may indeed block synaptic transmission by impeding the postsynaptic receptors [36, 58].

Despite the intense investigations in synaptic structure and plasticity in *D. melanogaster* related to genetic and mutational manipulations, there are few reports on the modulation of synaptic efficacy by peptides or modulators at the skeletal NMJ [54, 59-64]. As for the influence of 5-HT at the NMJ, the scantiness

of studies is likely due to the mild effects observed by using 5-HT itself as well as pharmacological agonists/antagonists of 5-HT receptors. However, application of 5-HT to the intact larval CNS does enhance the drive of motor neurons (MN) [17]. The most commonly studied *Drosophila* neuromuscular junctions are those in the most prominent ventral longitudinal abdominal muscle fiber muscles 6 and 7 [65], which have the simplest innervation pattern among the *Drosophila* body wall muscles. Both electrophysiological and morphological studies imply that each of these two muscles is innervated by only 2 axons [66, 67]. Application of 5-HT to these NMJs appears to slightly depress synaptic strength [68,69]. We are not aware of any attempt to investigate actions of 5-HT on adult skeletal NMJs. However, with the recent advent of designer receptors exclusively activated by designer drugs (DREAD) in motor neurons allows one to examine mechanisms of activating second messenger cascades as if receptors for modulators existed on presynaptic nerve terminals or on the muscles themselves [70, 71].

CRUSTACEANS

The NMJs in crustaceans offer many advantages for addressing mechanism of action in modulation of synaptic efficacy at NMJs, but crustaceans do fall short in being able to genetically modify the properties for investigations. Potentially approaches with RNAi might be practical to address more species-specific manipulations in synaptic function in a variety of crustaceans [72-75]. The same physiological and anatomical advantages of the *Drosophila* NMJs apply for the crustaceans, but in addition, the wide range in known diversity in synapses within crustaceans makes them attractable for comparative studies in commonalities of mechanisms in low- and high-output synapses or ones that facilitate or depress rapidly [31, 32, 76]. The parallels to vertebrate central synaptic physiology of phenomenon described at crustacean NMJs are likely one reason of continual interest to a wide variety of researchers investigating synaptic transmission. In addition, the historical contribution of crustaceans in synaptic physiology is unsurpassed [77-80]. The ability to combine direct structure and function in defined labeled synapses offers the ability to unravel synaptic structural complexity with function [31, 32, 43-45, 81].

It was demonstrated as early as 1954 that 5-HT enhances synaptic transmission at the crustacean NMJs [82, 83] and that the effect was likely a presynaptic enhancement of mean quantal content came afterwards [84]. The 5-HT that modulates most crustacean skeletal NMJs does so through the exposure of hemolymph. 5-HT is released from nerve endings in thoracic roots and from the pericardial organs into the hemolymph [85].

Thus, 5-HT is accessible to all the exposed NMJs. The excitatory as well as inhibitory NMJs are enhanced in transmission by 5-HT [86, 87]. The quantal effects are explained by increased probability of vesicular fusion during evoked transmission likely caused by an increase in the number of vesicular vesicles being docked and possibility their sensitivity of fusing due to enhanced Ca^{2+} sensitivity or presence of free Ca^{2+} within the terminals [88]. However, several studies have shown that a presynaptic rise in free Ca^{2+} is not substantial enough to account as a primary mechanism of 5-HT's action [86, 89-91]. Since there is a steep rise in sensitivity to Ca^{2+} for enhancing synaptic efficacy at crustacean NMJs [92] a low release from internal stores may account well enough for part of the effect [20]. This notion of an internal release of Ca^{2+} is also supported by experiments conducted by Glusman and Kravitz [91] in which they showed that a calcium-free bath, along with EGTA and high $MgCl_2$, 5-HT could still cause spontaneous release of transmitter for lobster NMJs. The enhanced spontaneous and evoked fusion events relates to an increase in 'n' (number of sites) and 'p' (probability of release) to explain the enhanced 'm' (mean quantal content; $m=np$) after exposure to 5-HT [69, 93, 94]. An interesting observation, but not yet explained mechanistically, is that 5-HT produced an effect with low or zero extracellular calcium at a crayfish NMJ but 5-HT's effect depended on extracellular sodium concentration [89].

Low- and high-output NMJs in crayfish and crab show differential responses to 5-HT [95-97]. This could be due to the larger reserve pool of vesicles in tonic (low-output) terminals than the phasic (high-output) terminals and the fact that higher-output synapses in crustaceans have more complex synapses containing more active zones in close apposition on synapses than lower output synapses [45, 98-100].

NMJs investigated in lobster and crab revealed similar findings to those of the crayfish. 5-HT also enhances both excitatory and inhibitor NMJs that have been examined in *Homarus americanus* (lobster) [101, 102]. 5-HT also promotes the force of nerve-evoked contractions of the gastric mill muscle of the crab, *Cancer borealis* [103].

The differential responses and cellular mechanism of 5-HT's action at crustacean NMJs is likely accounted for by the density and receptor subtypes on the presynaptic terminals. Vertebrate 5-HT₂-like receptors were physiologically identified for *Procambarus clarkii* at NMJs [21-24, 69]. Since this subtype of receptor has been sequenced in a crab and crayfish [28] these may be the subtypes present at the NMJs; however the blockers for the vertebrate 5-HT₂-like receptors could not block the entire 5-HT enhancement of synaptic enhancement [24]. Also 5-HT₂ agonists did not mimic the responses fully at the crayfish NMJ [24], so

potential affinity in binding 5-HT and pharmacological agents differ in crustaceans to vertebrate subtype receptor analogs. The pharmacology of monoamines in the cardiac ganglion of lobsters also does not mimic vertebrate classifications [104]. Care needs to be taken in assuming the pharmacology of mammalian 5-HT receptors applies to invertebrates [22].

Given there is at least some pharmacological and sequence similarity to vertebrate 5-HT₂ receptor subtype present in crayfish and that injection of an IP₃ analog (adenophostin-A) in the presynaptic motor nerve terminals enhances release [20], a potential mechanism is that 5-HT receptors on the presynaptic membrane mediate activation of G coupled receptors which leads to activation of phospholipase C (PLC) which in turn produces 1,4,5-trisphosphate (IP₃) and diacylglycerol (DAG) [105]. The production of IP₃ can directly result in Ca²⁺ release from internal stores (i.e., ER) through IP₃ receptors on the ER [106].

Since vertebrate 5-HT₂ receptor family activates phospholipase C (PLC) [9] a similar receptor activated cascade is possible at the crayfish NMJs. Such mechanisms are established in other systems [105, 107, 108] and given that caffeine and ryanodine actions are in concurrence with IP₃ receptors potentially on the ER in crayfish presynaptic motor nerve terminals [20] we have to consider this mechanism as a likely possibility. The rise in Ca²⁺, even a slight rise, could activate calmodulin and in turn activate CaM-Kinase (CaM-K), which can lead to phosphorylation of proteins such as synapsin. The possibility is that vesicles would then be able to leave the tethers to the cytoskeleton and dock to the presynaptic membrane, which is also a phosphorylation step [109-111]. The increased docked vesicles could be subjected to the calcium influx and release from internal stores [112]. This would account for the increase in the occurrence of spontaneous quantal events and enhanced evoked responses with 5-HT exposure. In the invertebrate *Aplysia*, it was shown that exposure of neurons to 5-HT results in phosphorylation of synapsins [113]. cAMP was also suggested to be involved in 5-HT action [30, 114-116]. cAMP has been shown to activate Protein Kinase A (PKA) which then can lead to phosphorylation of transcriptional factors such as CREB. Such action can regulate synthesis of proteins used in synaptic transmission [117-119]. It has also been suggested that the cAMP and calmodulin pathways may work together and promote transcription [120]. When phosphatases are inhibited at the crayfish NMJ the effect of 5-HT is enhanced, thus demonstrating the significance of phosphorylation [121] which is known to occur with exposure to 5-HT at crustacean NMJs [122].

In a recent study addressing the potential mechanisms of 5-HT, as well as stimulation of the motor nerve terminal, in recruiting vesicles from a reserve pool (RP) to a readily releasable pool (RRP)

within the presynaptic nerve terminals of crayfish NMJs, we developed a model to account for the observations and previous reports. In a current study, we inhibited the packaging of glutamate by blocking the vesicular glutamate transporter (VGlut) with the drug bafilomycin A1 (BA) [123-125]. In this way, the rapidly recycling vesicles within the RRP will be empty with repetitive stimulation. However, if the RP is spared from being recruited by low frequency stimulation and if they are already packaged with transmitter, prior to exposure to BA, then 5-HT should be able to recruit these RP vesicles to the RRP and synaptic transmission restored temporarily. This is exactly what was observed indicating that the RP and RRP can be physiologically differentiated into distinct functional groups and that 5-HT recruits the RP into action [126, 127]. To deplete or use up the packaged RRP vesicles, continuous stimulation was provided since the opener NMJ preparation is low-output and fatigue resistant. A high frequency of 40 Hz was used for comparative purposes to 20 Hz continuous stimulation. As expected, preparations stimulated at 40 Hz depressed faster than the ones stimulated at 20 Hz and there was a reduced effect for the 40 Hz stimulated preparations to exposure of 5-HT. This suggests that a higher stimulation frequency is able to recruit some of the RP to the RRP. This is illustrated in a model (Fig. 1). To address if PLC is an intermediate step within the cascade of events activated by 5-HT mediated responses, we used a PLC inhibitor (U73122) and an inactive analog (U73343) to serve as a negative control [128]. We found that the treatment of U73122 caused a significant decrease of 5-HT effect on synaptic transmission. This result confirmed the involvement of PLC signaling cascade in inducing the enhancement of synaptic transmission by 5-HT at a different physiological condition. There are observations in other preparations that indicate the presence of two distinct vesicle pools: RRP and RP. In the cat superior sympathetic ganglion, Prado et al. [129] separated the two pools by electrically stimulating the nerve to deplete the RRP of acetylcholine, and then recruit RP vesicles by tityustoxin. Using FM 1-43 dye, the two pools have been identified in a temperature-sensitive mutant *Drosophila* line, *shibire*, and later in WT [130, 131]. However in our study with the crayfish NMJ, a novel approach with bafilomycin A1 was used together with continuous stimulation to deplete the RRP, and then 5-HT was applied to recruit RP vesicles and the recruitment involves a PLC signaling cascade. A mechanistic illustration is detailed in Fig. 1.

There does not appear to be a substantial direct effect on crustacean skeletal muscle to account for an increase in EPSP or IPSP amplitude due to an increase in input resistance of the fibers [82, 114, 132, 133]. A small increase in input resistance, by exposure to 5-HT, accounts for a slight increase in the EPSP




Fig. 1. Schematic illustration of 5-HT in recruiting vesicles from a reserve pool. (A) Two vesicle recycling pathways have been proposed. In a resting synapse, vesicles in RP slowly join in to the RRP (1), and then recycle back to RP either through or bypass endosome (2). This is called slow recycling loop. However, in an active synapse, in addition to the slow recycling loop, vesicles in RRP recycle quickly within the RRP (3) which is named quick recycling loop. Recycling vesicles are refilled with glutamate to be able to participate in the coming synaptic activities. (B) In an active synapse treated with Bafilomycin A1, vesicles in RRP can be used up in time with stimulation because recycling vesicles can no longer be refilled. Synaptic depression occurs sooner than the one without Bafilomycin A1 treatment. (C) If 5-HT is added after depression, 5-HT possibly activates PLC signaling cascade and recruits RP vesicles to revitalize the synaptic transmission in a fast manner. (D) In time, synaptic depression occurs again because most RRP and RP vesicles are empty. Yellow colored vesicles represent partially full of glutamate. (E) Even when RRP vesicles are not depleted by Bafilomycin A1, 5-HT can also recruit RP vesicles into RRP via one possible mechanism (PLC). (F) It is also possible that 5-HT can activate silence synapse most likely in low-output terminals. (G) The PLC activation of 5-HT effect is confirmed with PLC non-selective inhibitor. RP, reserved pool; RRP, readily releasable pool; SY, synapse; GluT, glutamate transporter; BA, Bafilomycin A1.

amplitude for superficial flexor muscle fibers of crayfish [94]. More substantial alteration in input resistance can occur in crustacean neurons due to 5-HT exposure [134] so there could be some effect on the presynaptic motor nerve terminals.

In comparison to the smooth muscle in the intestine of vertebrates, the muscles of the crayfish hindgut are striated with gap junctions and generate intrinsic pacemaker activity [135, 136]. Application of 5-HT [137] and octopamine [138] to GI tract increases the frequency and strength of contractions. 5-HT and dopamine are highly concentrated in CNS and GI tract and they are directly responsible for the peristalsis and muscle contraction [137, 139].

ANNELIDS

The leech has served as a model organism in neurobiology for many years [140] but few studies have directly focused attention at NMJs in the leech and even fewer on the effects of 5-HT in synaptic transmission at NMJs. However, studies have examined the effect of 5-HT on the drive of motor neurons and innervation patterns [141-147]. 5-HT exposure has a relaxing effect on skeletal muscle in the leech [148] and enhances muscle force and work production during locomotion and feeding [149]. This is physiological relevant since Retzius neurons do directly innervate skeletal muscle in the leech and these cells do release 5-HT [144, 150-152]. In the earthworm and polychaete (*Sabellastarte magnifica*) muscle contraction is reduced by 5-HT [153, 154] which lead to the idea that 5-HT might be acting as inhibitory transmitter in these preparations [155].

GASTROPODS

A few studies with gastropods have been approached for the direct effect of 5-HT at the NMJ. 5-HT produces facilitation for an evoked response in buccal muscle within *Aplysia* [156]. The presynaptic actions of 5-HT is to enhance transmitter release [157]. Like for some of the actions in annelids, 5-HT can also produce muscle relaxation and reduce force in *Aplysia* [158]. Such effects on muscle contraction and force maybe dependent on 5-HT concentration and the species studied, since in *Aplysia brasiliiana* 5-HT increases a Ca^{2+} influx that promotes muscle contraction used for swimming [159].

OTHER INVERTEBRATES

In a sea urchin (*Parechinus*), 5-HT apparently had no effect at the NMJ [160]. However in a sea cucumber (*Apostichopus*

japonicas), 5-HT inhibited evoked contractions induced by acetylcholine and there appears to be 5-HT innervation directly to muscles of the body wall [161].

SUMMARY

Although headway has been made in describing the various actions of 5-HT at NMJs in invertebrates, the cellular mechanisms of these actions are still lacking. Additional pharmacological and molecular profiling in a variety of invertebrate preparations will increase our knowledge of both the uniqueness and similarities among the invertebrates. As history has taught us in physiology, and in particular neurobiology, what is learned in invertebrate preparations paves the way to new views and mechanistic cellular understanding of complex processes within the vertebrates.

ACKNOWLEDGEMENTS

We thank Mr. Josh Titlow, Ms. Rachel Holsinger and Mr. Zana Majeed for reviewing this manuscript and their constructive suggestions. Supported by personal funds of authors.

REFERENCES

1. Azmitia EC (2001) Modern views on an ancient chemical: serotonin effects on cell proliferation, maturation, and apoptosis. *Brain Res Bull* 56:413-424.
2. Azmitia EC (2007) Serotonin and brain: evolution, neuroplasticity, and homeostasis. *Int Rev Neurobiol* 77:31-56.
3. Bunin MA, Wightman RM (1999) Paracrine neurotransmission in the CNS: involvement of 5-HT. *Trends Neurosci* 22:377-382.
4. Monastirioti M (1999) Biogenic amine systems in the fruit fly *Drosophila melanogaster*. *Microsc Res Tech* 45:106-121.
5. Tierney AJ (2001) Structure and function of invertebrate 5-HT receptors: a review. *Comp Biochem Physiol A Mol Integr Physiol* 128:791-804.
6. Blenau W, Baumann A (2001) Molecular and pharmacological properties of insect biogenic amine receptors: lessons from *Drosophila melanogaster* and *Apis mellifera*. *Arch Insect Biochem Physiol* 48:13-38.
7. Colas JF, Launay JM, Maroteaux L (1999) Maternal and zygotic control of serotonin biosynthesis are both necessary for *Drosophila* germband extension. *Mech Dev* 87:67-76.
8. Saudou F, Boschert U, Amlaiky N, Plassat JL, Hen R (1992) A family of *Drosophila* serotonin receptors with distinct intracellular signalling properties and expression patterns. *EMBO J* 11:7-17.
9. Saudou F, Hen R (1994) 5-Hydroxytryptamine receptor subtypes in vertebrates and invertebrates. *Neurochem Int* 25:503-532.
10. Witz P, Amlaiky N, Plassat JL, Maroteaux L, Borrelli E, Hen R (1990) Cloning and characterization of a *Drosophila* serotonin receptor that activates adenylate cyclase. *Proc Natl Acad Sci U S A* 87:8940-8944.
11. Baines RA, Downer RG (1991) Pharmacological characterization of a 5-hydroxytryptamine-sensitive receptor/adenylate cyclase complex in the mandibular closer muscles of the cricket, *Gryllus domestica*. *Arch Insect Biochem Physiol* 16:153-163.
12. Hoyer D, Hannon JP, Martin GR (2002) Molecular, pharmacological and functional diversity of 5-HT receptors. *Pharmacol Biochem Behav* 71:533-554.
13. Burns CM, Chu H, Rueter SM, Hutchinson LK, Canton H, Sanders-Bush E, Emeson RB (1997) Regulation of serotonin-2C receptor G-protein coupling by RNA editing. *Nature* 387:303-308.
14. Niswender CM, Sanders-Bush E, Emeson RB (1998) Identification and characterization of RNA editing events within the 5-HT_{2C} receptor. *Ann N Y Acad Sci* 861:38-48.
15. Colas JF, Launay JM, Kellermann O, Rosay P, Maroteaux L (1995) *Drosophila* 5-HT₂ serotonin receptor: coexpression with fushi-tarazu during segmentation. *Proc Natl Acad Sci U S A* 92:5441-5445.
16. Colas JF, Launay JM, Vonesch JL, Hickel P, Maroteaux L (1999) Serotonin synchronises convergent extension of ectoderm with morphogenetic gastrulation movements in *Drosophila*. *Mech Dev* 87:77-91.
17. Dasari S, Cooper RL (2004) Modulation of sensory-CNS-motor circuits by serotonin, octopamine, and dopamine in semi-intact *Drosophila* larva. *Neurosci Res* 48:221-227.
18. Dasari S, Cooper RL (2006) Direct influence of serotonin on the larval heart of *Drosophila melanogaster*. *J Comp Physiol B* 176:349-357.
19. Dasari S, Wang L, Harrison DA, Cooper RL (2009) Reduced and misexpression of 5-HT₂ receptors alters development, behavior and CNS activity in *Drosophila melanogaster*. *Int J Zool Res* 5:101-114.
20. Dropic AJ, Brailoiu E, Cooper RL (2005) Presynaptic mechanism of action induced by 5-HT in nerve terminals: possible involvement of ryanodine and IP₃ sensitive Ca²⁺ stores. *Comp Biochem Physiol A Mol Integr Physiol* 142:355-361.
21. Sosa MA, Spitzer N, Edwards DH, Baro DJ (2004) A

- crustacean serotonin receptor: cloning and distribution in the thoracic ganglia of crayfish and freshwater prawn. *J Comp Neurol* 473:526-537.
22. Sparks GM, Brailoiu E, Brailoiu GC, Dun NJ, Tabor J, Cooper RL (2003) Effects of m-CPP in altering neuronal function: blocking depolarization in invertebrate motor and sensory neurons but exciting rat dorsal horn neurons. *Brain Res* 969:14-26.
 23. Sparks GM, Dasari S, Cooper RL (2004) Actions of MDMA at glutamatergic neuromuscular junctions. *Neurosci Res* 48:431-438.
 24. Tabor JN, Cooper RL (2002) Physiologically identified 5-HT₂-like receptors at the crayfish neuromuscular junction. *Brain Res* 932:91-98.
 25. Tierney AJ, Mangiamele LA (2001) Effects of serotonin and serotonin analogs on posture and agonistic behavior in crayfish. *J Comp Physiol A* 187:757-767.
 26. Tierney AJ, Greenlaw MA, Dams-O'Connor K, Aig SD, Perna AM (2004) Behavioral effects of serotonin and serotonin agonists in two crayfish species, *Procambarus clarkii* and *Orconectes rusticus*. *Comp Biochem Physiol A Mol Integr Physiol* 139:495-502.
 27. Clark MC, Dever TE, Dever JJ, Xu P, Rehder V, Sosa MA, Baro DJ (2004) Arthropod 5-HT₂ receptors: a neurohormonal receptor in decapod crustaceans that displays agonist independent activity resulting from an evolutionary alteration to the DRY motif. *J Neurosci* 24:3421-3435.
 28. Spitzer N, Edwards DH, Baro DJ (2008) Conservation of structure, signaling and pharmacology between two serotonin receptor subtypes from decapod crustaceans, *Panulirus interruptus* and *Procambarus clarkii*. *J Exp Biol* 211:92-105.
 29. Mapara S, Parries S, Quarrington C, Ahn KC, Gallin WJ, Goldberg JI (2008) Identification, molecular structure and expression of two cloned serotonin receptors from the pond snail, *Helisoma trivolvis*. *J Exp Biol* 211:900-910.
 30. Lee YS, Choi SL, Lee SH, Kim H, Park H, Lee N, Lee SH, Chae YS, Jang DJ, Kandel ER, Kaang BK (2009) Identification of a serotonin receptor coupled to adenylyl cyclase involved in learning-related heterosynaptic facilitation in *Aplysia*. *Proc Natl Acad Sci U S A* 106:14634-14639.
 31. Atwood HL, Cooper RL (1995) Functional and structural parallels in crustacean and *Drosophila* neuromuscular systems. *Am Zool* 35:556-565.
 32. Atwood HL, Cooper RL (1996) Synaptic diversity and differentiation: crustacean neuromuscular junctions. *Invert Neurosci* 1:291-307.
 33. Collins CA, DiAntonio A (2007) Synaptic development: insights from *Drosophila*. *Curr Opin Neurobiol* 17:35-42.
 34. Govind CK, Walrond JP (1989) Structural plasticity at crustacean neuromuscular synapses. *J Neurobiol* 20:409-421.
 35. Westfall IA (1996) Ultrastructure of synapses in the first-evolved nervous systems. *J Neurocytol* 25:735-746.
 36. Gerschenfeld HM (1973) Chemical transmission in invertebrate central nervous systems and neuromuscular junctions. *Physiol Rev* 53:1-119.
 37. Whim MD, Church PJ, Lloyd PE (1993) Functional roles of peptide cotransmitters at neuromuscular synapses in *Aplysia*. *Mol Neurobiol* 7:335-347.
 38. Lee JY, Bhatt D, Bhatt D, Chung WY, Cooper RL (2009) Furthering pharmacological and physiological assessment of the glutamatergic receptors at the *Drosophila* neuromuscular junction. *Comp Biochem Physiol C Toxicol Pharmacol* 150:546-557.
 39. Li H, Cooper RL (2001) Effects of the ecdysoneless mutant on synaptic efficacy and structure at the neuromuscular junction in *Drosophila* larvae during normal and prolonged development. *Neuroscience* 106:193-200.
 40. Li H, Harrison D, Jones G, Jones D, Cooper RL (2001) Alterations in development, behavior, and physiology in *Drosophila* larva that have reduced ecdysone production. *J Neurophysiol* 85:98-104.
 41. Li H, Peng X, Cooper RL (2002) Development of *Drosophila* larval neuromuscular junctions: maintaining synaptic strength. *Neuroscience* 115:505-513.
 42. Sherrington CS (1906) *The integrative action of the nervous system*. Scribner, New York.
 43. Cooper RL, Marin L, Atwood HL (1995) Synaptic differentiation of a single motor neuron: conjoint definition of transmitter release, presynaptic calcium signals, and ultrastructure. *J Neurosci* 15:4209-4222.
 44. Cooper RL, Harrington CC, Marin L, Atwood HL (1996) Quantal release at visualized terminals of a crayfish motor axon: intraterminal and regional differences. *J Comp Neurol* 375:583-600.
 45. Cooper RL, Winslow JL, Govind CK, Atwood HL (1996) Synaptic structural complexity as a factor enhancing probability of calcium-mediated transmitter release. *J Neurophysiol* 75:2451-2466.
 46. Cooper RL, Stewart BA, Wojtowicz JM, Wang S, Atwood HL (1995) Quantal measurement and analysis methods compared for crayfish and *Drosophila* neuromuscular junctions, and rat hippocampus. *J Neurosci Methods* 61:67-78.

47. Viele K, Stromberg AJ, Cooper RL (2003) Estimating the number of release sites and probability of firing within the nerve terminal by statistical analysis of synaptic charge. *Synapse* 47:15-25.
48. Viele K, Lancaster M, Cooper RL (2006) Self-modeling structure of evoked postsynaptic potentials. *Synapse* 60:32-44.
49. Lancaster M, Viele K, Johnstone AF, Cooper RL (2007) Automated classification of evoked quantal events. *J Neurosci Methods* 159:325-336.
50. Atwood HL, Cooper RL (1996) Assessing ultrastructure of crustacean and insect neuromuscular junctions. *J Neurosci Methods* 69:51-58.
51. Worden MK (1998) Modulation of vertebrate and invertebrate neuromuscular junctions. *Curr Opin Neurobiol* 8:740-745.
52. Cooper RL, Ruffner ME (1998) Depression of synaptic efficacy at intermolt in crayfish neuromuscular junctions by 20-Hydroxyecdysone, a molting hormone. *J Neurophysiol* 79:1931-1941.
53. Logsdon S, Johnstone AF, Viele K, Cooper RL (2006) Regulation of synaptic vesicles pools within motor nerve terminals during short-term facilitation and neuromodulation. *J Appl Physiol* 100:662-671.
54. Ruffner ME, Cromarty SI, Cooper RL (1999) Depression of synaptic efficacy in high- and low-output *Drosophila* neuromuscular junctions by the molting hormone (20-HE). *J Neurophysiol* 81:788-794.
55. Elekes K, Hustert R (1988) The efferent innervation of the genital chamber by an identified serotonergic neuron in the female cricket *Acheta domestica*. *Cell Tissue Res* 252:449-457.
56. Nishihara H (1967) The fine structure of the earthworm body wall muscle. *Acta Anat Nippon* 42:38-39.
57. Lange AB (2004) A neurohormonal role for serotonin in the control of locust oviducts. *Arch Insect Biochem Physiol* 56:179-190.
58. Hill RB, Usherwood PN (1961) The action of 5-hydroxytryptamine and related compounds on neuromuscular transmission in the locust *Schistocerca gregaria*. *J Physiol* 157:393-401.
59. Badre NH, Martin ME, Cooper RL (2005) The physiological and behavioral effects of carbon dioxide on *Drosophila melanogaster* larvae. *Comp Biochem Physiol A Mol Integr Physiol* 140:363-376.
60. Chen X, Ganetzky B (2012) A neuropeptide signaling pathway regulates synaptic growth in *Drosophila*. *J Cell Biol* 196:529-543.
61. Cooper RL, Neckameyer WS (1999) Dopaminergic modulation of motor neuron activity and neuromuscular function in *Drosophila melanogaster*. *Comp Biochem Physiol B Biochem Mol Biol* 122:199-210.
62. Dunn TW, Mercier AJ (2005) Synaptic modulation by a *Drosophila* neuropeptide is motor neuron-specific and requires CaMKII activity. *Peptides* 26:269-276.
63. Middleton CA, Nongthomba U, Parry K, Sweeney ST, Sparrow JC, Elliott CJ (2006) Neuromuscular organization and aminergic modulation of contractions in the *Drosophila* ovary. *BMC Biol* 4:17.
64. Nagaya Y, Kutsukake M, Chigusa SI, Komatsu A (2002) A trace amine, tyramine, functions as a neuromodulator in *Drosophila melanogaster*. *Neurosci Lett* 329:324-328.
65. Crossley CA (1978) The morphology and development of the *Drosophila* muscular system. In: *The genetics and biology of drosophila* (Ashburner M, Novitski E, eds), pp 499-599. Academic Press, New York.
66. Jan LY, Jan YN (1976) Properties of the larval neuromuscular junction in *Drosophila melanogaster*. *J Physiol* 262:189-214.
67. Sink H, Whittington PM (1991) Location and connectivity of abdominal motoneurons in the embryo and larva of *Drosophila melanogaster*. *J Neurobiol* 22:298-311.
68. Dasari S, Viele K, Turner AC, Cooper RL (2007) Influence of PCPA and MDMA (ecstasy) on physiology, development and behavior in *Drosophila melanogaster*. *Eur J Neurosci* 26:424-438.
69. Sparks GM, Cooper RL (2004) 5-HT offsets homeostasis of synaptic transmission during short-term facilitation. *J Appl Physiol* 96:1681-1690.
70. Nichols CD, Becnel J, Johnson O, Majeed ZR, Tran V, Yu B, Roth BL, Cooper RL (2012) DREADD receptor control of behavior, signalling, and physiology in the model organism *Drosophila melanogaster*. Thorlabs (Photonics) meeting abstract. Optogenetics and Pharmacogenetics in Neuronal Function and Dysfunction. New Orleans, LA, USA. 11-12 October 2012.
71. Majeed ZR, Cooper RL, Nichols CD (2012) The influence of DREAD receptors activation in the CNS of *Drosophila melanogaster*. Annual Meeting of Society for Neuroscience, 2012 Oct 13-17. New Orleans, LA, USA.
72. Estrada B, Gisselbrecht SS, Michelson AM (2007) The transmembrane protein Perdido interacts with Grip and integrins to mediate myotube projection and attachment in the *Drosophila* embryo. *Development* 134:4469-4478.
73. Kato Y, Shiga Y, Kobayashi K, Tokishita S, Yamagata H, Iguchi

- T, Watanabe H (2011) Development of an RNA interference method in the cladoceran crustacean *Daphnia magna*. *Dev Genes Evol* 220:337-345.
74. Estrada MP, Lugo JM, Acosta J, Carpio Y, Borroto I, Morera Y, González O, Rodríguez T, Ramos L, Huberman A (2007) Effects of RNA interference on gene functions of aquatic organisms. *Biotechnol Appl* 24:178-182.
75. Pekhletski R, Cooper RL, Atwood HL, Hampson DR (1996) Expression profiling of mRNA obtained from single identified crustacean motor neurons: determination of specificity of hybridization. *Invert Neurosci* 1:341-349.
76. Wu WH, Cooper RL (2010) Physiological recordings of high and low output NMJs on the Crayfish leg extensor muscle. *J Vis Exp* (45):e2319. Available from: <http://www.jove.com/index/details.stp?id=2319>.
77. Atwood HL (1976) Organization and synaptic physiology of crustacean neuromuscular systems. *Prog Neurobiol* 7:291-391.
78. Atwood HL (1967) Variation in physiological properties of crustacean motor synapses. *Nature* 215:57-58.
79. Cooper AS, Cooper RL (2009) Historical view and physiology demonstration at the NMJ of the crayfish opener muscle. *J Vis Exp* (33):e1595. Available from: <http://www.jove.com/index/details.stp?id=1595>.
80. Wiese K (2002) *The crustacean nervous system*. Springer, Berlin.
81. Johnstone AF, Viele K, Cooper RL (2011) Structure/function assessment of synapses at motor nerve terminals. *Synapse* 65:287-299.
82. Florey E, Florey E (1954) Über die mögliche Bedeutung von Enteramin (5-oxytryptamin) als nervöser Aktivitätsstoff bei cephalopoden und dekapoden Crustacean. *Z Naturforsch* 9B:58-68.
83. Fischer L, Florey E (1983) Modulation of synaptic transmission and excitation-contraction coupling in the opener muscle of the crayfish, *Astacus leptodactylus*, by 5-hydroxytryptamine and octopamine. *J Exp Biol* 102:187-198.
84. Dudel J (1965) Facilitatory effects of 5-hydroxy-tryptamine on the crayfish neuromuscular junction. *Naunyn Schmiedebergs Arch Pharmacol* 249:515-528.
85. Beltz BS, Kravitz EA (1983) Mapping of serotonin-like immunoreactivity in the lobster nervous system. *J Neurosci* 3:585-602.
86. Vyshedskiy A, Delaney KR, Lin JW (1998) Neuromodulators enhance transmitter release by two separate mechanisms at the inhibitor of crayfish opener muscle. *J Neurosci* 18:5160-5169.
87. Wang C, Zucker RS (1998) Regulation of synaptic vesicle recycling by calcium and serotonin. *Neuron* 21:155-167.
88. Dudel J (1988) Modulation of quantal synaptic release by serotonin and forskolin in crayfish motor nerve terminals. In: *Modulation of synaptic transmission and plasticity in nervous systems*, vol. 19 (Hertting G, Spatz HC, eds), pp 259-270. Springer, Berlin.
89. Dixon D, Atwood HL (1985) Crayfish motor nerve terminal's response to serotonin examined by intracellular microelectrode. *J Neurobiol* 16:409-424.
90. Delaney K, Tank DW, Zucker RS (1991) Presynaptic calcium and serotonin-mediated enhancement of transmitter release at crayfish neuromuscular junction. *J Neurosci* 11:2631-2643.
91. Glusman S, Kravitz EA (1982) The action of serotonin on excitatory nerve terminals in lobster nerve-muscle preparations. *J Physiol* 325:223-241.
92. Dudel J (1981) The effect of reduced calcium on quantal unit current and release at the crayfish neuromuscular junction. *Pflugers Arch* 391:35-40.
93. Southard RC, Haggard J, Crider ME, Whiteheart SW, Cooper RL (2000) Influence of serotonin on the kinetics of vesicular release. *Brain Res* 871:16-28.
94. Strawn JR, Neckameyer WS, Cooper RL (2000) The effects of 5-HT on sensory, central and motor neurons driving the abdominal superficial flexor muscles in the crayfish. *Comp Biochem Physiol B Biochem Mol Biol* 127:533-550. (See Erratum 128:377-378, 2001).
95. Cooper RL, Dönmezer A, Shearer J (2003) Intrinsic differences in sensitivity to 5-HT between high- and low-output terminals innervating the same target. *Neurosci Res* 45:163-172.
96. Djokaj S, Cooper RL, Rathmayer W (2001) Presynaptic effects of octopamine, serotonin, and cocktails of the two modulators on neuromuscular transmission in crustaceans. *J Comp Physiol A* 187:145-154.
97. Johnstone AF, Kellie SS, Cooper RL (2008) Presynaptic depression in phasic motor nerve terminals and influence of 5-HT on vesicle dynamics. *Open Neurosci J* 2:16-23.
98. Bradacs H, Cooper R, Msghina M, Atwood H (1997) Differential physiology and morphology of phasic and tonic motor axons in a crayfish limb extensor muscle. *J Exp Biol* 200:677-691.
99. King MJ, Atwood HL, Govind CK (1996) Structural features of crayfish phasic and tonic neuromuscular terminals. *J Comp Neurol* 372:618-626.
100. Millar AG, Zucker RS, Ellis-Davies GC, Charlton MP,

- Atwood HL (2005) Calcium sensitivity of neurotransmitter release differs at phasic and tonic synapses. *J Neurosci* 25:3113-3125.
101. Hamilton JL, Edwards CR, Holt SR, Worden MK (2007) Temperature dependent modulation of lobster neuromuscular properties by serotonin. *J Exp Biol* 210:1025-1035.
 102. Harris-Warrick RM, Kravitz EA (1984) Cellular mechanisms for modulation of posture by octopamine and serotonin in the lobster. *J Neurosci* 4:1976-1993.
 103. Jorge-Rivera JC, Sen K, Birmingham JT, Abbott LE, Marder E (1998) Temporal dynamics of convergent modulation at a crustacean neuromuscular junction. *J Neurophysiol* 80:2559-2570.
 104. Berlind A (2001) Monoamine pharmacology of the lobster cardiac ganglion. *Comp Biochem Physiol C Toxicol Pharmacol* 128:377-390.
 105. Berridge MJ, Irvine RF (1989) Inositol phosphates and cell signalling. *Nature* 341:197-205.
 106. Hisatsune C, Nakamura K, Kuroda Y, Nakamura T, Mikoshiba K (2005) Amplification of Ca^{2+} signaling by diacylglycerol-mediated inositol 1,4,5-trisphosphate production. *J Biol Chem* 280:11723-11730.
 107. Mattson MP, LaFerla FM, Chan SL, Leissring MA, Shepel PN, Geiger JD (2000) Calcium signaling in the ER: its role in neuronal plasticity and neurodegenerative disorders. *Trends Neurosci* 23:222-229.
 108. Petersen OH, Cancela JM (1999) New Ca^{2+} -releasing messengers: are they important in the nervous system? *Trends Neurosci* 22:488-495.
 109. Chi P, Greengard P, Ryan TA (2003) Synaptic vesicle mobilization is regulated by distinct synapsin I phosphorylation pathways at different frequencies. *Neuron* 38:69-78.
 110. He P, Southard RC, Chen D, Whiteheart SW, Cooper RL (1999) Role of alpha-SNAP in promoting efficient neurotransmission at the crayfish neuromuscular junction. *J Neurophysiol* 82:3406-3416.
 111. Tolar LA, Pallanck L (1998) NSF function in neurotransmitter release involves rearrangement of the SNARE complex downstream of synaptic vesicle docking. *J Neurosci* 18:10250-10256.
 112. Yang SN, Tang YG, Zucker RS (1999) Selective induction of LTP and LTD by postsynaptic $[Ca^{2+}]_i$ elevation. *J Neurophysiol* 81:781-787.
 113. Fiumara F, Giovedi S, Menegon A, Milanese C, Merlo D, Montarolo PG, Valtorta F, Benfenati F, Ghirardi M (2004) Phosphorylation by cAMP-dependent protein kinase is essential for synapsin-induced enhancement of neurotransmitter release in invertebrate neurons. *J Cell Sci* 117:5145-5154.
 114. Battelle BA, Kravitz EA (1978) Targets of octopamine action in the lobster: cyclic nucleotide changes and physiological effects in hemolymph, heart and exoskeletal muscle. *J Pharmacol Exp Ther* 205:438-448.
 115. Dixon D, Atwood HL (1989) Conjoint action of phosphatidylinositol and adenylate cyclase systems in serotonin-induced facilitation at the crayfish neuromuscular junction. *J Neurophysiol* 62:1251-1259.
 116. Goy MF, Kravitz EA (1989) Cyclic AMP only partially mediates the actions of serotonin at lobster neuromuscular junctions. *J Neurosci* 9:369-379.
 117. Geppert M, Goda Y, Stevens CF, Sudhof TC (1997) The small GTP-binding protein Rab3A regulates a late step in synaptic vesicle fusion. *Nature* 387:810-814.
 118. Bolshakov VY, Golan H, Kandel ER, Siegelbaum SA (1997) Recruitment of new sites of synaptic transmission during the cAMP-dependent late phase of LTP at CA3-CA1 synapses in the hippocampus. *Neuron* 19:635-651.
 119. Yao J, Qi J, Chen G (2006) Actin-dependent activation of presynaptic silent synapses contributes to long-term synaptic plasticity in developing hippocampal neurons. *J Neurosci* 26:8137-8147.
 120. Dash PK, Karl KA, Colicos MA, Prywes R, Kandel ER (1991) cAMP response element-binding protein is activated by Ca^{2+} /calmodulin- as well as cAMP-dependent protein kinase. *Proc Natl Acad Sci U S A* 88:5061-5065.
 121. Swain JE, Robitaille R, Dass GR, Charlton MP (1991) Phosphatases modulate transmission and serotonin facilitation at synapses: studies with the inhibitor okadaic acid. *J Neurobiol* 22:855-864.
 122. Goy MF, Schwarz TL, Kravitz EA (1984) Serotonin-induced protein phosphorylation in a lobster neuromuscular preparation. *J Neurosci* 4:611-626.
 123. Bowman EJ, Graham LA, Stevens TH, Bowman BJ (2004) The bafilomycin/concanamycin binding site in subunit c of the V-ATPases from *Neurospora crassa* and *Saccharomyces cerevisiae*. *J Biol Chem* 279:33131-33138.
 124. Cavalier P, Attwell D (2007) Neurotransmitter depletion by bafilomycin is promoted by vesicle turnover. *Neurosci Lett* 412:95-100.
 125. Juge N, Gray JA, Omote H, Miyaji T, Inoue T, Hara C, Uneyama H, Edwards RH, Nicoll RA, Moriyama Y (2010) Metabolic control of vesicular glutamate transport and release. *Neuron* 68:99-112.

126. Wu WH, Cooper RL (2012) The regulation and packaging of synaptic vesicles as related to recruitment within glutamatergic synapses (in review, doi: <http://dx.doi.org/10.1016/j.neuroscience.2012.08.037>).
127. Wu WH, Cooper RL (2012) The regulation and packaging of synaptic vesicles related to recruitment within glutamatergic synapses. Annual Meeting of Society for Neuroscience, 2012 Oct 13-17. New Orleans, LA, USA.
128. Bleasdale JE, Thakur NR, Gremban RS, Bundy GL, Fitzpatrick FA, Smith RJ, Bunting S (1990) Selective inhibition of receptor-coupled phospholipase C-dependent processes in human platelets and polymorphonuclear neutrophils. *J Pharmacol Exp Ther* 255:756-768.
129. Prado MA, Gomez MV, Collier B (1992) Mobilization of the readily releasable pool of acetylcholine from a sympathetic ganglion by tityustoxin in the presence of vesamicol. *J Neurochem* 59:544-552.
130. Betz WJ, Mao F, Bewick GS (1992) Activity-dependent fluorescent staining and destaining of living vertebrate motor nerve terminals. *J Neurosci* 12:363-375.
131. Kuromi H, Kidokoro Y (2000) Tetanic stimulation recruits vesicles from reserve pool via a cAMP-mediated process in *Drosophila* synapses. *Neuron* 27:133-143.
132. Grundfest H, Reuben JP (1961) Neuromuscular synaptic activity in lobster. In: *Nervous inhibition* (Florey E, ed), pp 92-104. Pergamon, New York.
133. Kravitz EA, Glusman S, Harris-Warrick RM, Livingstone MS, Schwarz T, Goy MF (1980) Amines and a peptide as neurohormones in lobsters: actions on neuromuscular preparations and preliminary behavioural studies. *J Exp Biol* 89:159-175.
134. Cooper RL, Ward E, Braxton R, Li H, Warren WM (2003) The effects of serotonin and ecdysone on primary sensory neurons in crayfish. *Microsc Res Tech* 60:336-345.
135. Brenner TL (1999) The physiology of crayfish intestinal striated muscle: histology, histochemistry, and excitation-contraction coupling, MSc thesis. University of Calgary, Calgary, AB, Canada.
136. To TH, Brenner TL, Cavey MJ, Wilkens JL (2004) Histological organization of the intestine in the crayfish *Procambarus clarkii*. *Acta Zool* 85:119-130.
137. Musolf BE (2007) Serotonergic modulation of the Crayfish hindgut: effects on hindgut contractility and regulation of serotonin on hindgut. Biology Dissertations. Paper 32. Available from: http://digitalarchive.gsu.edu/biology_diss/32.
138. Orchard I, Lange AB (1985) Evidence for octopaminergic modulation of an insect visceral muscle. *J Neurobiol* 16:171-181.
139. Cooper AS, Leksrisawat B, Gilberts AB, Mercier AJ, Cooper RL (2011) Physiological experimentation with the crayfish hindgut: a student laboratory exercise. *J Vis Exp* (47):e2324. Available from: <http://www.jove.com/details.php?id=2324>.
140. Nicholls JG, Stent GS, Muller KJ (1981) *Neurobiology of the leech*. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
141. Brodfuehrer PD, Debski EA, O'Gara BA, Friesen WO (1995) Neuronal control of leech swimming. *J Neurobiol* 27:403-418.
142. Stuart AE (1970) Physiological and morphological properties of motoneurons in the central nervous system of the leech. *J Physiol* 209:627-646.
143. Ort CA, Kristan WB, Stent GS (1974) Neuronal control of swimming in the medicinal leech. *J Comp Physiol* 94:121-154.
144. Kuffler DP (1978) Neuromuscular transmission in longitudinal muscle of the leech *Hirudo medicinalis*. *J Comp Physiol* 124:333-338.
145. Cline HT (1983) 3H-GABA uptake selectively labels identifiable neurons in the leech central nervous system. *J Comp Neurol* 215:351-358.
146. O'Gara BA, Illuzzi FA, Chung M, Portnoy AD, Fraga K, Frieman VB (1999) Serotonin induces four pharmacologically separable contractile responses in the pharynx of the leech *Hirudo medicinalis*. *Gen Pharmacol* 32:669-681.
147. Sawada M, Coggeshall RE (1976) A central inhibitory action of 5-hydroxytryptamine in the leech. *J Neurobiol* 7:477-482.
148. Schain RJ (1961) Effects of 5-hydroxytryptamine on the dorsal muscle of the leech (*Hirudo medicinalis*). *Br J Pharmacol Chemother* 16:257-261.
149. Gerry SP, Ellerby DJ (2011) Serotonin modulates muscle function in the medicinal leech *Hirudo verbana*. *Biol Lett* 7:885-888.
150. Mason A, Kristan WB (1982) Neuronal excitation, inhibition and modulation of leech longitudinal muscle. *J Comp Physiol* 146:527-536.
151. Mason A, Sunderland AJ, Leake LD (1979) Effects of leech Retzius cells on body wall muscles. *Comp Biochem Physiol C* 63C:359-361.
152. Yaksta-Sauerland BA, Coggeshall RE (1973) Neuromuscular junctions in the leech. *J Comp Neurol* 151:85-99.
153. Hidaka T, Kuriyama H, Yamamoto T (1969) The mechanical properties of the longitudinal muscle in the earthworm. *J Exp Biol* 50:431-443.

154. Del Carmen Alvarez M, Del Castillo J, Sánchez V (1969) Pharmacological responses of the dorsal longitudinal muscle of *Sabellastarte magnifica*. *Comp Biochem Physiol* 29:931-942.
155. Díaz-Miranda L, de Motta GE, García-Arrarás JE (1992) Monoamines and neuropeptides as transmitters in the sedentary polychaete *Sabellastarte magnifica*: actions on the longitudinal muscle of the body wall. *J Exp Zool* 263:54-67.
156. Fox LE, Lloyd PE (2002) Mechanisms involved in persistent facilitation of neuromuscular synapses in *Aplysia*. *J Neurophysiol* 87:2018-2030.
157. Lotshaw DP, Lloyd PE (1990) Peptidergic and serotonergic facilitation of a neuromuscular synapse in *Aplysia*. *Brain Res* 526:81-94.
158. Evans CG, Vilim FS, Harish O, Kupfermann I, Weiss KR, Cropper EC (1999) Modulation of radula opener muscles in *Aplysia*. *J Neurophysiol* 82:1339-1351.
159. Laurienti PJ, Blankenship JE (1997) Serotonergic modulation of a voltage-gated calcium current in parapodial swim muscle from *Aplysia brasiliana*. *J Neurophysiol* 77:1496-1502.
160. Bolt RE, Ewer DW (1963) Studies on the myoneural physiology of Echinodermata. V. the lantern retractor muscle of *Parechinus*: responses to drugs. *J Exp Biol* 40:727-733.
161. Inoue M, Tamori M, Motokawa T (2002) Innervation of holothurian body wall muscle: inhibitory effects and localization of 5-HT. *Zoolog Sci* 19:1217-1222.