



9-25-2008

# Co-up-regulation of Three P450 Genes in Response to Permethrin Exposure in Permethrin Resistant House Flies, *Musca domestica*

Fang Zhu

*University of Kentucky, fangzhudy@uky.edu*

Ting Li

*Auburn University Main Campus*

Lee Zhang

*Auburn University Main Campus*

Nannan Liu

*Auburn University Main Campus*

**Right click to open a feedback form in a new tab to let us know how this document benefits you.**

Follow this and additional works at: [https://uknowledge.uky.edu/entomology\\_facpub](https://uknowledge.uky.edu/entomology_facpub)

 Part of the [Entomology Commons](#)

## Repository Citation

Zhu, Fang; Li, Ting; Zhang, Lee; and Liu, Nannan, "Co-up-regulation of Three P450 Genes in Response to Permethrin Exposure in Permethrin Resistant House Flies, *Musca domestica*" (2008). *Entomology Faculty Publications*. 63.

[https://uknowledge.uky.edu/entomology\\_facpub/63](https://uknowledge.uky.edu/entomology_facpub/63)

This Article is brought to you for free and open access by the Entomology at UKnowledge. It has been accepted for inclusion in Entomology Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

---

**Co-up-regulation of Three P450 Genes in Response to Permethrin Exposure in Permethrin Resistant House Flies, *Musca domestica***

**Notes/Citation Information**

Published in *BMC Physiology*, v. 8, no. 18, p. 1-13.

© 2008 Zhu et al. This is an Open Access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Digital Object Identifier (DOI)**

<http://dx.doi.org/10.1186/1472-6793-8-18>

Research article

Open Access

## Co-up-regulation of three P450 genes in response to permethrin exposure in permethrin resistant house flies, *Musca domestica*

Fang Zhu<sup>1,2</sup>, Ting Li<sup>1</sup>, Lee Zhang<sup>3</sup> and Nannan Liu<sup>\*1</sup>

Address: <sup>1</sup>Department of Entomology and Plant Pathology, Auburn University, Auburn, AL 36849, USA, <sup>2</sup>Department of Entomology, University of Kentucky, Lexington, KY 40546, USA and <sup>3</sup>Genomics and Sequencing Laboratory, Auburn University, Auburn, AL 36849, USA

Email: Fang Zhu - fangzhudy@uky.edu; Ting Li - tzl0001@auburn.edu; Lee Zhang - zhangl1@auburn.edu; Nannan Liu\* - liunann@auburn.edu  
\* Corresponding author

Published: 25 September 2008

Received: 10 June 2008

BMC Physiology 2008, 8:18 doi:10.1186/1472-6793-8-18

Accepted: 25 September 2008

This article is available from: <http://www.biomedcentral.com/1472-6793/8/18>

© 2008 Zhu et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Background:** Insects may use various biochemical pathways to enable them to tolerate the lethal action of insecticides. For example, increased cytochrome P450 detoxification is known to play an important role in many insect species. Both constitutively increased expression (overexpression) and induction of P450s are thought to be responsible for increased levels of detoxification of insecticides. However, unlike constitutively overexpressed P450 genes, whose expression association with insecticide resistance has been extensively studied, the induction of P450s is less well characterized in insecticide resistance. The current study focuses on the characterization of individual P450 genes that are induced in response to permethrin treatment in permethrin resistant house flies.

**Results:** The expression of 3 P450 genes, *CYP4D4v2*, *CYP4G2*, and *CYP6A38*, was co-up-regulated by permethrin treatment in permethrin resistant ALHF house flies in a time and dose-dependent manner. Comparison of the deduced protein sequences of these three P450s from resistant ALHF and susceptible aabys and CS house flies revealed identical protein sequences. Genetic linkage analysis located *CYP4D4v2* and *CYP6A38* on autosome 5, corresponding to the linkage of P450-mediated resistance in ALHF, whereas *CYP4G2* was located on autosome 3, where the major insecticide resistance factor(s) for ALHF had been mapped but no P450 genes reported prior to this study.

**Conclusion:** Our study provides the first direct evidence that multiple P450 genes are co-up-regulated in permethrin resistant house flies through the induction mechanism, which increases overall expression levels of P450 genes in resistant house flies. Taken together with the significant induction of *CYP4D4v2*, *CYP4G2*, and *CYP6A38* expression by permethrin only in permethrin resistant house flies and the correlation of the linkage of the genes with resistance and/or P450-mediated resistance in resistant ALHF house flies, this study sheds new light on the functional importance of P450 genes in response to insecticide treatment, detoxification of insecticides, the adaptation of insects to their environment, and the evolution of insecticide resistance.

## Background

Cytochrome P450s constitute the largest gene superfamily and are found in the organs and tissues of many organisms, including mammals, fish, plants, arthropods, fungi, and bacteria. Cytochrome P450s have long been of particular interest because they are critical for the detoxification and/or activation of xenobiotics such as drugs, pesticides, plant toxins, chemical carcinogens and mutagens; and for metabolizing endogenous compounds such as hormones, fatty acids, and steroids. Basal and up-regulation of P450 gene expression can significantly affect disposition of xenobiotics or endogenous compounds in the tissues of organisms and thus alter their pharmacological/toxicological effects [1]. Insect cytochrome P450s are known to play an important role in detoxifying insecticides [2,3] and plant toxins [4,5], resulting in the development of resistance to insecticides [3,6-10] and facilitating the adaptation of insects to their plant hosts [11,12]. A significant characteristic of insect P450s that is associated with enhanced metabolic detoxification of insecticides in insects is the constitutively increased levels of P450 proteins and P450 activity that result from constitutively transcriptional overexpression of P450 genes in insecticide resistant insects [3,6-9,13,14]. Another feature of insect P450 genes is that the expression of some P450 genes can be induced by exogenous and endogenous compounds [3], a phenomenon known as induction. It has been suggested that the induction of P450s and their activities in insects is involved in the adaptation of insects to their environment and the development of insecticide resistance [15,16].

While all insects probably possess some capacity to detoxify insecticides and xenobiotics, the degree to which they can metabolize and detoxify these toxic chemicals is of considerable importance to their survival in a chemically unfriendly environment [16] and to the development of resistance. The constitutively increased expression and induction of P450s are thought to be responsible for increased levels of detoxification of insecticides, but unlike the constitutively overexpressed P450 genes, whose association with insecticide resistance has been extensively studied, the induction of P450s by xenobiotics, especially phenobarbital, is less well linked to insecticide resistance, although it is well documented in insects [17-22]. It has been proposed that many chemical inducers act as substrates for P450s and that the induction or modulation of P450s by the substrates will, in turn, reduce the effects of the substrates by enhancing substrate metabolism [23]. It has been suggested that the modulation of gene expression reflects a compromise between the insect's need to both conserve energy and adjust to a rapidly changing environment by enhancing the activity of the detoxification system only when a chemical stimulus occurs [24].


The house fly strain ALHF exhibits high levels of resistance to pyrethroids [25]. Previous research using piperonyl butoxide (PBO), an inhibitor of cytochrome P450s, suggested that P450-mediated detoxification may be one of the major mechanisms involved in the development of pyrethroid resistance in ALHF [25]. Genetic linkage analysis points to the localization of PBO-suppressible-P450-mediated resistance on autosomes 1, 2, and 5 of ALHF [26] and factors on autosome 5 are known to play a major role in P450-mediated resistance. Two P450 cDNAs, *CYP6A36* and *CYP6A5v2*, have recently been identified as being constitutively overexpressed in ALHF and mapped on autosome 5 [27,28], which is correlated with the linkage of resistance in ALHF. These studies suggest the importance of constitutive overexpression of these two genes in increasing metabolic detoxification of permethrin and in the evolution of permethrin resistance in ALHF. To test whether induction of P450s is also involved in the development of permethrin resistance in ALHF, the current study is focused on the characterization of individual P450 genes from house flies that are induced in response to permethrin challenge in ALHF. Three novel P450 genes, *CYP4D4v2*, *CYP4G2*, and *CYP6A38*, were isolated whose expression was induced by permethrin treatment in ALHF. Genetic linkage studies were then conducted in order to identify a further possible causal link between the expression of these genes in response to insecticide treatment and the development of insecticide resistance in ALHF.

## Results

### Identification of P450 genes in response to permethrin challenge

In order to identify the P450 genes raised in response to the insecticide treatment, we used the PCR technique with degenerated PCR primers [29] to amplify P450 cDNAs from house flies. We initially isolated a total of 19 P450 cDNA fragments from ALHF house flies using three primer pairs, C2/Flyh1, C2/Flyc1, and HemeR1/CYP6AD1 (Table 1). Northern blot analysis was conducted using the 19 P450 cDNA fragments as probes to compare expression levels of these cDNAs in both permethrin treated and untreated susceptible CS and resistant ALHF house flies. Three of the 19 P450 cDNAs were found to be significantly induced in the permethrin treated ALHF house flies after 24 hours treatment (Fig. 1) at the pilot dose of LD<sub>50</sub> that caused ~50% mortality of each house fly strain.

The full length of the three putative P450 cDNAs were isolated using 5' and/or 3' RACE with the primers (Table 1) designed based on the 5' and/or 3' end sequences of the putative P450 cDNA fragments. The sequences of the 5' and/or 3' RACE amplified cDNAs overlapped with their corresponding putative P450 cDNA fragments. An entire



**Figure 1**  
**Northern blot analysis of differentially expressed patterns of *CYP4G2*, *CYP4D4v2*, and *CYP6A38* between permethrin treated and untreated susceptible CS and resistant ALHF house flies.** mRNAs were isolated from the whole bodies of 20 surviving house flies 24 h after permethrin treatment with 2  $\mu\text{g}/\text{fly}$ . Blots were hybridized with the cDNA probes derived from three P450 gene fragments. The ethidium bromide stain of 18S ribosomal RNA in agarose gel is shown at the bottom.

cDNA fragment for each of the putative P450 genes was subsequently amplified for both ALHF and aabys house flies by PCR using the primer pair (Table 1) synthesized based on the respective 5' and 3' end sequences of each gene. The sequences were named *CYP4D4v2* [accession number: EF615001, <http://www.ncbi.nlm.nih.gov/entrez/viewer.fcgi?val=EF615001.1>], *CYP4G2* [accession number: EF615002, <http://www.ncbi.nlm.nih.gov/entrez/viewer.fcgi?val=EF615002.1>], and *CYP6A38* [accession number: EF615003, <http://www.ncbi.nlm.nih.gov/entrez/viewer.fcgi?val=EF615003.1>] by the P450 nomenclature committee (Dr. D. Nelson, personal communication). The cDNA sequences of *CYP4D4v2*, *CYP4G2*, and *CYP6A38* have open reading frames of 1515, 1647, and 1500 nucleotides encoding proteins of 505, 549, and 500 residues, respectively. Comparison of the deduced protein sequences of *CYP4D4v2*, *CYP4G2*, and *CYP6A38* between ALHF and aabys revealed identical protein sequences, although several nucleotide polymorphisms were found in the coding regions of these three genes between the two strains (Figs. 2, 3, 4).

#### **Response of P450 genes to permethrin challenge in resistant and susceptible house flies**

Based on this pilot experiment, we focused our study on *CYP4D4v2*, *CYP4G2*, and *CYP6A38* in order to further characterize their overexpression in response to permethrin challenge in resistant and susceptible house flies using the quantitative real-time PCR (qRT-PCR) method, as described below under Materials and Methods. To minimize the possibility that these P450 genes have nothing to do with resistance but arose solely because of a strain-strain difference, the study used ALHF and two susceptible

house fly strains, CS and aabys. To examine the effect of permethrin on induction of the three target P450s, we measured the expression of the genes in house flies challenged with permethrin at a corresponding dose range ( $\text{LD}_{10}$ ,  $\text{LD}_{50}$ , and  $\text{LD}_{90}$ ) for various durations. Although no induction was detected in either the susceptible CS flies of the aabys for the dose range and time intervals tested (data not shown), our results showed that permethrin induced all three P450 genes in ALHF with varying levels in a time (24 h)- and dose ( $\text{LD}_{50}$ -10  $\mu\text{g}/\text{fly}$ )-dependent manner (Fig. 5). Based on these data, a dose of 10  $\mu\text{g}/\text{fly}$  and a time interval of 24 h were chosen for the further induction studies.

We detected no significant induction in the expression of the three P450 genes in susceptible CS and aabys house flies that had either been treated with acetone alone or with permethrin solution in acetone compared with untreated house flies (Fig. 6). Similarly, no significant induction was obtained in acetone treated ALHF house flies compared with their untreated counterparts (Fig. 6). However, these three genes were induced at a variety of levels in permethrin treated ALHF house flies compared with untreated or acetone treated ALHF flies; a marked induction of *CYP4D4v2* and *CYP6A38* mRNA ( $\sim 4$ -fold) were detected in permethrin treated ALHF house flies (Fig. 6B, C), whereas a low level ( $\sim 1.5$ -fold) of induction for *CYP4G2* was detected in the permethrin treated ALHF house flies (Fig. 6A). The significant induction of the three target P450 genes only in ALHF house flies suggests their importance in response to permethrin treatment in the resistant ALHF house flies. No significant differences were observed in the basal expression of *CYP4D4v2*, *CYP4G2*,

**Table 1: The primers used for cloning P450 genes, qRT-PCR reactions, and SNP determinations**

Gene	Primer Name	Function	Primer Sequence	Position (nt)
CYP4G2	AP450HF20F	Full length cloning	5' ATGGATAGTGCAAACAATTCAACTGCG 3'	1 to 27
	SNP450HF20F	SNaP determination	5' AAATGGCCATTTGGTGGCCCAT 3'	174 to 195
	RTP450HF20F	qRT-PCR	5' CGAGGAGGATGATGAAATAAGCAAGC 3'	837 to 862
	RTP450HF20R	qRT-PCR	5' TTGGACATGGCCATCATGGCATCT 3'	957 to 980
	P450HF20R	5'-RACE	5' GCAGGAAGTTGTCACCAAGATG 3'	1140 to 1162
	P450HF20F	3'-RACE	5' CGTGCATCGCAATCCCCAATAC 3'	1341 to 1362
	AP450HF20R	Full length cloning	5' TTACATAGCTTTCATGGCTTCGGGTC 3'	1625 to 1650
CYP4D4v2	AP450HF5F	Full length cloning	5' ATGTTATTGAATTCCTGGTGGGTC 3'	1 to 25
	SNP450HF5F	SNaP determination	5' ACATGACACCACGACAAGTGG 3'	948 to 968
	RTP450HF5F	qRT-PCR	5' AGGATAAGGAGAAACCGGTGACC 3'	1052 to 1074
	RTP450HF5R	qRT-PCR	5' CAATTGTCGGCACCGATGGATAC 3'	1156 to 1134
	P450HF5R	5'-RACE	5' AGCAACTCAAAATGGCGTACC 3'	1430 to 1410
	AP450HF5R	Full length cloning	5' TAACTACTGCGAACTCTCAAACCC 3'	1521 to 1497
CYP6A38	P450HF17P-F	5' flanking region cloning	5' TGGTCTTCTAGGGGAGAAGACTACCTGC 3'	-676 to -649
	SNP450HF17PF	SNaP determination	5' TTCAGGATTGCTGGGTAGCT 3'	-69 to -50
	P450HF16F-3	Full length cloning	5' CATTATGGAGACTTCGGGAGTTTTG 3'	-4 to 21
	P450HF17P	5' flanking region cloning	5' CGTAGGTTCTCATGGGGTATACCCAGC 3'	120 to 93
	RTP450HF17F	qRT-PCR	5' CCCTGATGGGCAACATGAATGGAT 3'	122 to 145
	RTP450HF17R	qRT-PCR	5' TAGTTGTTTGTCCAGCAGCACCAC 3'	270 to 247
	P450HF17R	5' RACE	5' CGCTGTACTTCAATAGATTTCTGC 3'	650 to 626
	P450HF17F	3' RACE	5' TGAGGGTGATACAAAACCAAGC 3'	1017 to 1038
	AP450HF17R	Full length cloning	5' GAGATAATCTCCCACCCCTAAATCG 3'	1520 to 1496
Common	FlyhI		5' GGICCIAGIAACTGCATIGG 3'	
	FlycI		5' GGAAGTNGACACNTTYATGTT 3'	
	CYP6ADI		5' GTNATHGGHHNBTYGCHTTYGG 3'	
	HemeRI		5' CCIATGCAGTTTICTIGGICC 3'	
	Oligo (dt)		5' TAATACGACTCACTATAGGGAGATTTTTTTTTTTTTTTTTT 3'	
	C2		5' TAATACGACTCACTATAGGGAGA 3'	
	API (RACE)		5' CCATCCTAATACGACTCACTATAGGGC 3'	
	API (GenomeWalking)		5' GTAATACGACTCACTATAGGGC 3'	
	ActinSI		5' AGGCGAATCGCGAGAAGATG 3'	
	ActinASI		5' TCAGATCACGACCAGCCAGATC 3'	

and CYP6A38 between ALHF and both or one of the susceptible CS and aabys strains without permethrin treatment (Fig. 6). These results suggest that, unlike some P450s in which constitutive expression may play important role in insecticide resistance [30-32], CYP4D4v2, CYP6A38, and/or CYP4G2, may be uniquely featured in ALHF in response to the insecticide exposure through induction of their expression, which, in turn, enhances their capacity to detoxify the insecticide and leads to enhanced insecticide resistance.

**Chromosomal linkage and allele determination of CYP4D4v2, CYP4G2, and CYP6A38**

We previously demonstrated that permethrin resistance in ALHF house flies was linked to autosomes 1, 2, 3, and 5, with major factors on autosomes 3 and 5 [26]. We also demonstrated that the P450-mediated resistance in ALHF was predominantly linked to autosome 5 [26]. To deter-

mine whether there is a causal link between the P450 genes and insecticide resistance, we examined the genetic linkage of CYP4D4v2, CYP4G2, and CYP6A38 with 5 back-cross (BC<sub>1</sub>) house fly lines derived from crosses of ALHF and a susceptible morphological marker strain, aabys, by allele specific single nucleotide polymorphism (SNP) determination. Sequence comparisons of the three genes between ALHF and aabys revealed several nucleotide polymorphisms in the coding regions of CYP4D4v2 and CYP4G2, while no nucleotide polymorphisms were identified in the coding region of CYP6A38. We therefore cloned a ~700 bp 5' flanking region of CYP6A38 in order to genetically map the CYP6A38 gene. Comparison of the nucleotide sequence of the 5' flanking region of CYP6A38 uncovered several nucleotide polymorphisms between ALHF and aabys (data not shown), so the nucleotide polymorphisms, C to T, C to T, and G to T, in CYP4D4v2, CYP4G2, and CYP6A38, respectively, in ALHF relative to

ALHF	ATGTTATTGAATTCCTGGCGGGTCTCTTGACCCTTCTGCTCGTTGTCGACACGCTGTACAAGACGGCGCATGTTATGATTGCAAAGGCTGGGATACGGGGACCCACACCGGTCGCCG	120
aabys	ATGTTATTGAATTCCTGGCGGGTCTCTTGACCCTTCTGCTCGTTGTCGACACGCTGTACAAGACGGCGCATGTTATGATTGCAAAGGCTGGGATACGGGGACCCACACCGGTCGCCG	120
	M L F E F L A G L L T L L L V V D T L Y K R R R H V M I A K A G I R G P T P L P	40
ALHF	CTTTTGGAAACATGTTGCATTTTGTGGGCAATACAACGGCCAATGCCTATGACCTTTTCGATAAATGACGGCCAAATATGGCAAAGTGGTGGCCATGTGGATGGGTAATCATTTGTGTG	240
aabys	CTTTTGGAAACATGTTGCATTTTGTGGGCAATACAACGGCCAATGCCTATGACCTTTTCGATAAATGACGGCCAAATATGGCAAAGTGGTGGCCATGTGGATGGGTAATCATTTGTGTG	240
	L F G N M L H F V G N T T A N A Y D L F D K L T A K Y G K V V R M W M G N H C V	80
ALHF	TTGTTTGTGCGCGATGCCAAATCTTCGAGGACATCTTCAAAGTCAACAGCTGATCACCACCAAAATCTCTATGATTTGCTAGTGGTTGGCTGGCCAGGGTCTGTTGTTGAGCACC	360
aabys	TTGTTTGTGCGCGATGCCAAATCTTCGAGGACATCTTCAAAGTCAACAGCTGATCACCACCAAAATCTCTATGATTTGCTAGTGGTTGGCTGGCCAGGGTCTGTTGTTGAGCACC	360
	L F V R D A K F F E D I F K S Q Q L I T K N N L Y D L L S G W L G Q G L L S T	120
ALHF	GGACCAAAATGGTTTTCGCGACGTAAGATAATCACTCCAACCTTTCATTTAAATTTGGAAGAATTTGGAGATCTTCGATCAGCAGAGTCATTAATGGTGAAGAGTTTAAATGGAG	480
aabys	GGACCAAAATGGTTTTCGCGACGTAAGATAATCACTCCAACCTTTCATTTAAATTTGGAAGAATTTGGAGATCTTCGATCAGCAGAGTCATTAATGGTGAAGAGTTTAAATGGAG	480
	G P K W F S R R K I I T P T F H F K I L E E F V E I F D Q Q S H V M V K S L M E	160
ALHF	AAGCCGATGGCAAATGTCGTCATATGTTCCCAAAGTTTGTCTCATGGCTTGGATATTATAACGGAGACGGCAATGGCGTGAAGTCCATGCCAGGAACATCCAGAATTCCC	600
aabys	AAGCCGATGGCAAATGTCGTCATATGTTCCCAAAGTTTGTCTCATGGCTTGGATATTATAACGGAGACGGCAATGGCGTGAAGTCCATGCCAGGAACATCCAGAATTCCC	600
	K A D G K T V V H M F P K V C L M A L D I I T E T A M G V K V H A Q E H P E F P	200
ALHF	TATGCCAAGGCGATGACGGATACCTCCAACATTATGTCGGAACCGTTTGTGAAACCTTTCGATCGTTTCGATGGCTATTCCGGATGTTTACAAATCAAAAATACCAACAGCTGCAAAAAT	720
aabys	TATGCCAAGGCGATGACGGATACCTCCAACATTATGTCGGAACCGTTTGTGAAACCTTTCGATCGTTTCGATGGCTATTCCGGATGTTTACAAATCAAAAATACCAACAGCTGCAAAAAT	720
	Y A K A V T D T S N I M S D R F V K P F D R F D G Y F R I F H N Q K Y H Q L Q N	240
ALHF	AATATCAAATATGACGACTTACGGATAAGGTCATCAACGAACGACGTGAGGCTTGCAAAAGTCCATGACAGTGGAACTATCAGGCTGGCAGTAGCGTGCATGAAATGGGCATT	840
aabys	AATATCAAATATGACGACTTACGGATAAGGTCATCAACGAACGACGTGAGGCTTGCAAAAGTCCATGACAGTGGAACTATCAGGCTGGCAGTAGCGTGCATGAAATGGGCATT	840
	N I K I M H D F T D K V I N E R R E A L Q K S I D S G T Y Q A G S S V D E M G I	280
ALHF	AAGAAGCAATGGCTTCCCTGATGTGCTGTGCAATCGACGGTGGTGGAAAACCTTGCAGAAATAGGATATTCGTGAAGAGGTGGATACTTTTATGTTTCAAGGACATGACACCAG	960
aabys	AAGAAGCAATGGCTTCCCTGATGTGCTGTGCAATCGACGGTGGTGGAAAACCTTGCAGAAATAGGATATTCGTGAAGAGGTGGATACTTTTATGTTTCAAGGACATGACACCAG	960
	K K R M A F L D V L L Q S T V G G K P L T N K D I R E E V D T F M F E G H D T T	320
ALHF	ACAAGTGCATTGCTTACTCTCTACTGCTGTCGCGGCATCCTGACGTACAGCAAAAAGTACTCGAGGAGATTCACAGCGTATTGTTGAGGATAAGGAGAAAACCGGTGACCATGAAA	1080
aabys	ACAAGTGCATTGCTTACTCTCTACTGCTGTCGCGGCATCCTGACGTACAGCAAAAAGTACTCGAGGAGATTCACAGCGTATTGTTGAGGATAAGGAGAAAACCGGTGACCATGAAA	1080
	T S G I C F T L Y L L S R H P D V Q Q K V L E E I H S V I G E D K E K P V T M K	360
ALHF	GATTCGAGAGCTTAAATATTTGGATTGTTATCAAGGAAAGTCAGCGTTTGTATCCATCGGTCGACAAATGGTTCGTGTTACCGAACAGGATGTTGTTATAAATGGTGTACCATA	1200
aabys	GATTCGAGAGCTTAAATATTTGGATTGTTATCAAGGAAAGTCAGCGTTTGTATCCATCGGTCGACAAATGGTTCGTGTTACCGAACAGGATGTTGTTATAAATGGTGTACCATA	1200
	D L L Q E A G C V L K Y L D C V I K E S Q R L Y P S V P T I G R V T E Q D V T T A A A T I N G V T I	400
ALHF	CCGGCCAAACCAACATCACCCCTGCTCATGTATGCCCCATGAAGGATCCCGATTATTCCCAAGCCAGAAGAATTTTGGCCGACGTTTCATGAACCCAGGATAAAAATAATCCC	1320
aabys	CCGGCCAAACCAACATCACCCCTGCTCATGTATGCCCCATGAAGGATCCCGATTATTCCCAAGCCAGAAGAATTTTGGCCGACGTTTCATGAACCCAGGATAAAAATAATCCC	1320
	P A N T N I T L L M Y A A M K D P D Y F P K P E E F L P E R F M N T E D K I N P	440
ALHF	TTTGCCTATGTTCCGTTTCAGTGTGACCCCGTAACGTCATAGGTGAGAAATTTGCCATGGCTGAAATGAAAAGTACCGTCAGTAAAATGGTACGCCATTTTGGTGTGCTCCCCCTGGGA	1440
aabys	TTTGCCTATGTTCCGTTTCAGTGTGACCCCGTAACGTCATAGGTGAGAAATTTGCCATGGCTGAAATGAAAAGTACCGTCAGTAAAATGGTACGCCATTTTGGTGTGCTCCCCCTGGGA	1440
	F A Y V P P F S A G P R N C I G Q K F A M A E M K S T V S K M V R H F E L L P L G	480
ALHF	GAGGAGGTGAAACCTGTCATGAATATGGTATTACGTTCCGAGACTGATGCCAAATGGGTTTGGAGTTTCGCAAGTAG	1518
aabys	GAGGAGGTGAAACCTGTCATGAATATGGTATTACGTTCCGAGACTGATGCCAAATGGGTTTGGAGTTTCGCAAGTAG	1518
	E E V K P V M N M V L R S E T D A Q L G L R V R K * 506	

**Figure 2**  
**Alignment of the nucleotide and deduced amino acid sequences of CYP4D4v2 in ALHF and aabys house flies.**  
 The nucleotide polymorphisms between ALHF and aabys are highlighted and underlined.

aabys (Fig. 7), were used to determine the linkage of P450 genes relative to the recessive morphological markers in the aabys strain. The SNP determination reactions were conducted for each of the genes using a specific primer (Fig. 7) designed according to the sequences immediately upstream of the nucleotide polymorphism in order to distinguish the single nucleotide polymorphism for the P450 allele in each house fly strain or line. Our results showed that the BC<sub>1</sub> lines with the genotypes of *ac/ac*, *+/ar*, *+/bwb*, *+/ye*, *+/sw* (A2345), *+/ac*, *ar/ar*, *+/bwb*, *+/ye*, *+/sw* (A1345), *+/ac*, *+/ar*, *bwb/bwb*, *+/ye*, *+/sw* (A1245), and *+/ac*, *+/ar*, *+/bwb*, *ye/ye*, *+/sw* (A1235) were heterozygous for *CYP4D4v2* and *CYP6A38*, whereas the BC<sub>1</sub> line with the genotype of *+/ac*, *+/ar*, *+/bwb*, *+/ye*, *sw/sw* (A1234) was homozygous for both the *CYP4D4v2* and *CYP6A38* alleles from aabys (Table 2). These results strongly indicate that both *CYP4D4v2* and *CYP6A38* are located on autosome 5 in house flies. The BC<sub>1</sub> lines of A2345, A1345, A1234, and

A1235 were heterozygous for *CYP4G2*, whereas the A1245 line was homozygous for the *CYP4G2* allele from aabys (Table 2), indicating that *CYP4G2* is located on autosome 3 in house flies.

**Discussion**

The primary goal of this study was to investigate whether insecticide resistant insects may be uniquely resistant to insecticides due to their ability to mount an adequate cellular response when challenged with insecticides by up-regulating the production of P450s, which, in turn, may significantly diminish toxicological effects of the insecticides to these insects [1]. It also appears that the induction of gene expression may reflect a good compromise between energy saving (i.e., enhancing the activity of the detoxification system only when a chemical stimulus occurs) and adjustment to a rapidly changing environment [33]. Multiple P450 genes that are induced in insects

ALHF	ATGGATAGTCAAACAATCAACTGCGGGCCCGCAACTGATTGAACCAAAATTTGGACAGCCCTTCTGGGTATTGCAGTGGTTGTGAGTTTGTACGAAATATGGCTAAGGAATACCAGG	120
aabys	ATGGATAGTCAAACAATCAACTGCGGGCCCGCAACTGATTGAACCAAAATTTGGACAGCCCTTCTGGGTATTGCAGTGGTTGTGAGTTTGTACGAAATATGGCTAAGGAATACCAGG	120
	M D S A N N S T A G P A T V L N P I W T A L L G I A V V V S L Y E I W L R N T R	40
ALHF	AAATATAAATTAACGGCAAATATGCCAAACCCACCTATGCTGCCACTCATCGGAATGGCCATTGGTGGCCCATGTGACAAATGCCGAAATCCTTGCCCGTGGCATTGGCTATATGCAA	240
aabys	AAATATAAATTAACGGCAAATATGCCAAACCCACCTATGCTGCCACTCATCGGAATGGCCATTGGTGGCCCATGTGACAAATGCCGAAATCCTTGCCCGTGGCATTGGCTATATGCAA	240
	K Y K L T A N M P N P F M L P L I G N G H L V A H L T N A E I L A R G I G Y M Q	80
ALHF	ACCTATGGGGTGTATGCGTGGGCTTTTGGGTCCCATGTTGGTGTGTTCTCTCGTGAACGCTCCGATATTGAATTGATTCTCAGTACCCACACCCATTGGAGAAGTCAATTGAATAT	360
aabys	ACCTATGGGGTGTATGCGTGGGCTTTTGGGTCCCATGTTGGTGTGTTCTCTCGTGAACGCTCCGATATTGAATTGATTCTCAGTACCCACACCCATTGGAGAAGTCAATTGAATAT	360
	T Y G G A M R G F L G P M L V V F L W N A P D I E L I L S T H T H L E K S I E Y	120
ALHF	CGTTTCTCAAACCTGGTTCGGTGTGGTCTACTGATTCGAATGGTACCATTGGCAACATCATCGCAAATGATTGCTCCAACATCCATCAAAGCATTTTGAAGAGTTTCGTGCCA	480
aabys	CGTTTCTCAAACCTGGTTCGGTGTGGTCTACTGATTCGAATGGTACCATTGGCAACATCATCGCAAATGATTGCTCCAACATCCATCAAAGCATTTTGAAGAGTTTCGTGCCA	480
	R F F K P W F G D G L L I S N G H H W Q H H R K M I A P T F H Q S I L K S F V P	160
ALHF	GCCTTTGTACAGCATTCCAAGAAGTGGTGGAAACGATGGCCAAGGAATGGGCAAGAAATCGATGTCATGACTATATGTCACAGACCACCTGTGGAAATTTTGCTCTCCACTGCCATG	600
aabys	GCCTTTGTACAGCATTCCAAGAAGTGGTGGAAACGATGGCCAAGGAATGGGCAAGAAATCGATGTCATGACTATATGTCACAGACCACCTGTGGAAATTTTGCTCTCCACTGCCATG	600
	A F V Q H S K K V E R M A K E L G K E F D V H D Y M S Q T T V E I L S T A T A M	200
ALHF	GGTGTAAAGAGTGGCCAGAGGAAACAAGAGTTTGAATATGCCAAGGCTGTGGTGGACATGTCGACATCATCACAAGCGTCAATTGAAATCTTCTATCGCATGGATGCCCTTAC	720
aabys	GGTGTAAAGAGTGGCCAGAGGAAACAAGAGTTTGAATATGCCAAGGCTGTGGTGGACATGTCGACATCATCACAAGCGTCAATTGAAATCTTCTATCGCATGGATGCCCTTAC	720
	G V K K V P E D N K S L E Y A K A V V D M C D I I H K R Q L K F F Y R M D A L Y	240
ALHF	AATTTGAGCAGCATGAGCGAGAAGGCAAGAAGATGATGGACATTTATTTGGGTATGACCCGTAAGTGGTACGGAACGTCACAGAAATTTCAATGCCGAATCGCGTGCCTTGTGCGAG	840
aabys	AATTTGAGCAGCATGAGCGAGAAGGCAAGAAGATGATGGACATTTATTTGGGTATGACCCGTAAGTGGTACGGAACGTCACAGAAATTTCAATGCCGAATCGCGTGCCTTGTGCGAG	840
	N L S S M S E K G K K M M D I I L G M T R K V V T E R Q Q N F N A E S R A I V E	280
ALHF	GAGGATGATGAAATAAGCAAGCAGAAGCAACAGGCCAAGAAGAAGGAGGGTTTGGCGTATGATTGGATGACATTGATGAAATGATGGGTGCCAAGAACGTTTGGCTCTGCTAGAT	960
aabys	GAGGATGATGAAATAAGCAAGCAGAAGCAACAGGCCAAGAAGAAGGAGGGTTTGGCGTATGATTGGATGACATTGATGAAATGATGGGTGCCAAGAACGTTTGGCTCTGCTAGAT	960
	E D D E I S K Q K Q Q A K K K E G L R D D L D D I D E N D V G A K K R L A L L D	320
ALHF	GCCATGATGGCCATGTCCAAGAACCCCGATGTTGAGTGGACCGACAAGGATGTCATGATGAAGTCAACACCATTATGTTGAGGGCCATGATACCACCTCAGCTGGTCCAGTTTTGTC	1080
aabys	GCCATGATGGCCATGTCCAAGAACCCCGATGTTGAGTGGACCGACAAGGATGTCATGATGAAGTCAACACCATTATGTTGAGGGCCATGATACCACCTCAGCTGGTCCAGTTTTGTC	1080
	A M M A M S K N P D V E T G D K A D V M D E V N T I M F E G H D T T A G S S F V	360
ALHF	CTCTGCATGTTGGGCATCTCAAGGATAATCCAAGAGAAGGCTTGGCCGAACAAAAGGCCATCTTTGGTGACAACCTCCTGCGCGACTGTACCTTTGCCGATACCATGGAATGAAATAC	1200
aabys	CTCTGCATGTTGGGCATCTCAAGGATAATCCAAGAGAAGGCTTGGCCGAACAAAAGGCCATCTTTGGTGACAACCTCCTGCGCGACTGTACCTTTGCCGATACCATGGAATGAAATAC	1200
	L C M L G I Y K D I Q E K V L A E Q K A I F G D N F L R D C T F A D A T M E M K Y	400
ALHF	CTGGAACGTGTGATTATGAAACTTTGGCTTTGTATCCACCAGTACCCCTTATTGCCGCTGCTGAGTTCGATGATAAAATGGCTTCCGGCCCTACACAATCCCAAGGGCACAAAC	1320
aabys	CTGGAACGTGTGATTATGAAACTTTGGCTTTGTATCCACCAGTACCCCTTATTGCCGCTGCTGAGTTCGATGATAAAATGGCTTCCGGCCCTACACAATCCCAAGGGCACAAAC	1320
	L E R V I M E T L R L Y P P V P L I A R R A E F D V K L A S G P Y T I P K G T T	440
ALHF	GTGGTATTGCCAAATTTGCCGTGCATCGCAATCCCAATACTTCCCAATCCCGAGAATTCGATCCTGACAATTTCTACCCGAACGATGGCCAAATCGTCATTACTACAGTTTCATT	1440
aabys	GTGGTATTGCCAAATTTGCCGTGCATCGCAATCCCAATACTTCCCAATCCCGAGAATTCGATCCTGACAATTTCTACCCGAACGATGGCCAAATCGTCATTACTACAGTTTCATT	1440
	V V I A Q F A V H R N P Q Y F P N P E K F D P D N F L P E R M A N R H Y Y S F I	480
ALHF	CCCTTCAGTCCCGCCCGCAGAAGTTGTGCGGTCCGAAATATGCCATGCTGAAATGAAGTCTGCTCAGCATTATTCGTAACATATTCGTGAGAGCAACCAACAGGAGAAGGAC	1560
aabys	CCCTTCAGTCCCGCCCGCAGAAGTTGTGCGGTCCGAAATATGCCATGCTGAAATGAAGTCTGCTCAGCATTATTCGTAACATATTCGTGAGAGCAACCAACAGGAGAAGGAC	1560
	P F T A G P R S C V G R K Y A M L K L G V L G L S T I R N Y S V Q S N Q Q E K D	520
ALHF	TTCAAGTGCAGGCCGATATTATCCTAAAGATTGAAATGGTTTCAACATCATGTTGAACCGACGACCCGAAGCCATGAAAGCTATGTAA	1650
aabys	TTTAAAGTGCAGGCCGATATTATCCTAAAGATTGAAATGGTTTCAACATCATGTTGAACCGACGACCCGAAGCCATGAAAGCTATGTAA	1650
	F K L Q A D I I L K I E N G F N I M L N R R P E A M K A M * 550	

**Figure 3**  
**Alignment of the nucleotide and deduced amino acid sequences of CYP4G2 in ALHF and aabys house flies. The nucleotide polymorphisms between ALHF and aabys are highlighted and underlined.**

in response to host plant allelochemicals or secondary products have been extensively studied [3,34-39] and are fairly well documented in terms of their function in the adaptation of insects in "animal-plant warfare" [40] and in the co-evolution of insects and plants [41]. In contrast, P450 gene induction in response to insecticide resistance is less well understood. The current study, therefore, focused on characterization the P450 genes induced in response to a challenge with insecticides in resistant house flies. We restricted this response to permethrin treatment because it is the insecticide that the house flies are resistant to. We found that resistant house flies exposed to permethrin responded by up-regulating a set of P450 genes compared to the levels found in flies that had not been challenged with permethrin, whereas sus-

ceptible house flies showed no significant response. This finding indicates that exposure to permethrin induced a response in the expression of multiple P450 genes at different levels in a resistance-specific manner. Similar results have also been reported in *Drosophila melanogaster* [42], where the expression of *CYP6g1* and *CYP12d1* were induced in the DDT resistant strains post-exposure to DDT. Our study strongly indicated a clear dose- and time-dependent manner of induction. The lack of induction of P450 gene expression as a result of permethrin treatment at lower doses (LD<sub>10</sub>) in ALHF house flies found in this study is probably due to their rapid metabolism of permethrin at lower doses, which thus never reaches the threshold dose needed for induction [43]. The low or nonexistent levels of induction at higher (LD<sub>90</sub>) doses



ALHF	ATGGAGAC <u>TCGGGAGTTTTGATATATCTCATT</u> TTGGTCATTGCCTC <u>GGCTGTGGTTTATTGCTGAGAAGGAATCTAAGCTATTGGAAGAGGCTGGGTATACCCCATGAGGAACCTACG</u>	120
aabys	ATGGAGAC <u>TCGGGAGTTTTGATATATCTCATT</u> TTGGTCATTGCCTC <u>GGCTGTGGTTTATTGCTGAGAAGGAATCTAAGCTATTGGAAGAGGCTGGGTATACCCCATGAGGAACCTACG</u>	120
	M E T S G V L I Y L I L V I A S A V V Y L L R R N L S Y W K R L G I P H E E P T	40
ALHF	GCCCTGATGGGCAACATGAATGGATTGAGGACCCAGCGGTGTATGGGTGACATTTGGCCGAATAITATAGGAAATTTAAGGGCTGCGGACCCTTTGGCGGAATTTCAATGGTCAGAGA	240
aabys	GCCCTGATGGGCAACATGAATGGATTGAGGACCCAGCGGTGTATGGGTGACATTTGGCCGAATAITATAGGAAATTTAAGGGCTGCGGACCCTTTGGCGGAATTTCAATGGTCAGAGA	240
	A L M G N M N G L R T Q R C M G D I L A E Y Y R K F K G C G P F A G I F N G Q R	80
ALHF	CCCGGGTGGTGTCTGGCAAAACAATAAACAACAGGTCCTGATCAAGGATTTACACAAATTCACCGATCGTGGTCTGTATTACAATGAGAAATCGGATCCCTTTGACAGGACATCTG	360
aabys	CCCGGGTGGTGTCTGGCAAAACAATAAACAACAGGTCCTGATCAAGGATTTACACAAATTCACCGATCGTGGTCTGTATTACAATGAGAAATCGGATCCCTTTGACAGGACATCTG	260
	P G V V L L D K Q L I K Q V L I K D F T Q F T D R G L Y Y N E K S D P L T G H L	120
ALHF	TTCTTTGGTGGATGGCATAAATGGCGGAGCCTGCGCAACAAGCTCTCGCCACCTTTACCTCGGGCAAAATGAAATATATGTATCCGACAGTGTGGAAAGTGGCCGAGCGTTTTATGGAA	480
aabys	TTCTTTGGTGGATGGCATAAATGGCGGAGCCTGCGCAACAAGCTCTCGCCACCTTTACCTCGGGCAAAATGAAATATATGTATCCGACAGTGTGGAAAGTGGCCGAGCGTTTTATGGAA	480
	F F V D G D K W R S L R N K L S P T F T S G K M K Y M Y P T V L E V A E R F M E	160
ALHF	GTTTTGGCGAAAAGTCCCGGAGAATCCATTGGATGTACGGATATGCTGGGACGTTTACCCTGGATGTGATTGCCAGTGTGGCTTTTGGCATTGAATGCAATAGCCTGAGGAAT	600
aabys	GTTTTGGCGAAAAGTCCCGGAGAATCCATTGGATGTACGGATATGCTGGGACGTTTACCCTGGATGTGATTGCCAGTGTGGCTTTTGGCATTGAATGCAATAGCCTGAGGAAT	600
	V L A Q K V R E N S I V D V R D M L G R F T V D V I A S V A F G I E C N S L R N	200
ALHF	GCTGAGATAAATTTCTGCAGCTTGGCAGGAAATCCATTGAAGTACAGCGCCATAATGCCCTGATCATGACATTGATTGATGGGTTTCCCGCCTATCCAGAATGCTGGGCATGCGCACT	720
aabys	GCTGAGATAAATTTCTGCAGCTTGGCAGGAAATCCATTGAAGTACAGCGCCATAATGCCCTGATCATGACATTGATTGATGGGTTTCCCGCCTATCCAGAATGCTGGGCATGCGCACT	720
	A E D K F L Q L G R K S I E V Q R H N A L I M T L I D G F P R L S R M L G M R T	240
ALHF	TTGCCCCAGGATGCCATGAGTTTTTTCATGCAGACCATCCGGGAAACGGTGGAAATATCGTGAGAGGAATAATATCCAAAGAAGTGATTCCTGAATATTCTGATTGAATGAAAAATACT	840
aabys	TTGCCCCAGGATGCCATGAGTTTTTTCATGCAGACCATCCGGGAAACGGTGGAAATATCGTGAGAGGAATAATATCCAAAGAAGTGATTCCTGAATATTCTGATTGAATGAAAAATACT	840
	L P Q D V H E F F M Q T I R E T V E Y R E R N N I Q R S D F L N I L I E L K N T	280
ALHF	GTGGATGACAAGTCAGGCTGGGAGGATGGAAGTGGCTGCCAGGATTTGTTTTCTTTTGGCTGGCTTGGAGACCTCCTCGTCGACCATGGCCATGGCCTTGTATGAA	960
aabys	GTGGATGACAAGTCAGGCTGGGAGGATGGAAGTGGCTGCCAGGATTTGTTTTCTTTTGGCTGGCTTGGAGACCTCCTCGTCGACCATGGCCATGGCCTTGTATGAA	960
	V D D K S G L G G M E L E E L A A Q V F V F F L A G F E T S S S T M A Y A L Y E	320
ALHF	TTGCCCCAAAATCAGCAGATACAGGAGAGATTAAGGGAGGAGATCAATGAGGCATTTGAGGGTGATACAAAACCAAGCTATGAGACCATAATGAATTTGAGTTACCTGGATCAGGTGATA	1080
aabys	TTGCCCCAAAATCAGCAGATACAGGAGAGATTAAGGGAGGAGATCAATGAGGCATTTGAGGGTGATACAAAACCAAGCTATGAGACCATAATGAATTTGAGTTACCTGGATCAGGTGATA	1080
	L A Q N Q Q I E R L R E E I N E A F E G G T D T K P S Y E T I M N L S Y L D Q V I	360
ALHF	TCGGAATCTACGCAATA <u>CCCATTTTACCCTTCTTAAATCGCCAGGCCTTAAATGACTATGTGGTGGCTGGACCCCAAAATTCGAATACCCAAGGGAACCCCATCTTCATACCC</u>	1200
aabys	TCGGAATCTACGCAATA <u>CCCATTTTACCCTTCTTAAATCGCCAGGCCTTAAATGACTATGTGGTGGCTGGACCCCAAAATTCGAATACCCAAGGGAACCCCATCTTCATACCC</u>	1200
	S E T L R K Y P I L P F L N R Q A L N D Y V V P G H P K F R I P K G T I F I P	400
ALHF	GTCATGGGCATACAAATGACCCAGAATCTATCCCAACCCGATGAATTTGATCCGAAACGTTTTTACCAGATATGGTGAAGCAACGTTGATCCATCGAATGGATGCCCTTTGGAGAT	1320
aabys	GTCATGGGCATACAAATGACCCAGAATCTATCCCAACCCGATGAATTTGATCCGAAACGTTTTTACCAGATATGGTGAAGCAACGTTGATCCATCGAATGGATGCCCTTTGGAGAT	1320
	V M G I Q H D P E F Y P Q P D E F D P E R F S P D M V K Q R D S I E W M P F G D	440
ALHF	GGGCCCCGAACTGTATGGCGCCGATTTGCCAAAATGCAAAACCCGCTTGGCTTGGCATGTGTTTTGAAATATTTTCGGGTTGATGTTTGGCCAAAGACCCCATGCACTTGACATTT	1440
aabys	GGGCCCCGAACTGTATGGCGCCGATTTGCCAAAATGCAAAACCCGCTTGGCTTGGCATGTGTTTTGAAATATTTTCGGGTTGATGTTTGGCCAAAGACCCCATGCACTTGACATTT	1440
	G P R N C I G A R F A K M Q T R L G L A C V L K Y F R V V V C P K T P L H L T F	480
ALHF	CAGGCAAGCCTTTGGTGTGACACCACGGCATAATGTCTACCTGAAATTTGAGGCGATTTAG	1503
aabys	CAGGCAAGCCTTTGGTGTGACACCACGGCATAATGTCTACCTGAAATTTGAGGCGATTTAG	1503
	Q A K P L V L T P R H N V Y L K L E A I * 501	


**Figure 4**  
**Alignment of the nucleotide and deduced amino acid sequences of CYP6A38 in ALHF and aabys house flies. The nucleotide polymorphisms between ALHF and aabys are highlighted and underlined.**

may indicate a dysfunction of the induction process in insects that have been highly poisoned. A lack of induction of P450s has also been reported in *Drosophila melanogaster* when the insects were challenged with insecticides at concentrations that exceeded LC<sub>99</sub> [44].

It has been proposed that induction and/or constitutive overexpression of P450s is linked to the adaptation of insects to their environment [15,16]. Further, in many cases increased levels of P450 gene expression have resulted in increased levels of both total P450s and the activities of those P450s, strongly suggesting this as a major cause of insecticide resistance [3,7,10,31,42]. We recently identified two P450 genes that were constitutively overexpressed in ALHF house flies [27,28]. Our current study identified a further three P450 genes that were co-up-regulated in response to permethrin exposure in ALHF house flies. Taken together, these findings suggest both

constitutive overexpression and induction mechanisms participate in increasing P450-mediated metabolic detoxification of permethrin in resistant ALHF house and imply the role of these genes in the evolution of insecticide resistance. It has been proposed that there could be similar regulatory mechanisms governing P450 constitutive overexpression and induction, and that both contribute to the development of insecticide resistance [15]. Accordingly, we hypothesize that both induction and constitutive overexpression of P450 genes in the resistant ALHF share an altered regulatory system, which differs from that in the susceptible strains and regulate P450 gene expression in resistant house flies. Further study of the regulation of both constitutively overexpressed [27,28] and permethrin induced P450 genes will allow us to test this hypothesis.

Early studies in our laboratory on permethrin resistance in ALHF house flies led to the identification of permethrin

**Figure 5**

**Time and dose-dependent induction of the expression of three genes was analyzed by qRT-PCR as described in the materials and methods.** A. The duration of the gene expression following permethrin treatment at the dose of LD<sub>50</sub> (10 µg/fly). B. The expression of the genes 24 h after permethrin treatment with a dose range of LD<sub>10</sub> (5 µg/fly), LD<sub>50</sub> (10 µg/fly), and LD<sub>90</sub> (20 µg/fly). The relative level of gene expression shown in Y axis is the ratio of the gene expression in each treatment in comparison with that in acetone treated flies. The results are shown as the mean ± S.E.


resistance that could be largely suppressed by PBO, an inhibitor of cytochrome P450s [25]. Furthermore, genetic linkage studies had associated permethrin resistance in ALHF to the autosomes 3 and 5, while PBO-suppressible resistance (or P450-mediated resistance) had been mainly tied to autosome 5, with minor factors linked to autosomes 1 and 2 [26]. The genetic linkage between an over-expressed P450 gene or protein and insecticide resistance appears to be an important step in establishing a causal link between a P450 gene and its role in resistance [3,6,32,45-47]. We therefore went on to examine the linkage of the three P450 genes with 5 house fly BC<sub>1</sub> lines derived from crosses of ALHF and a susceptible morphological marker strain, aabys, using allele specific PCR determination. The results revealed that both *CYP4D4v2* and *CYP6A38* were located on autosome 5, whereas *CYP4G2* was located on autosome 3, of the house flies. Given that *CYP4D4v2* and *CYP6A38* are highly induced in ALHF and specifically located on autosome 5, on which P450-mediated resistance in ALHF has been mapped, it seems likely that the induction of *CYP4D4v2* and *CYP6A38* plays an important role in the development of insecticide resistance in ALHF house flies. Compared to the induction of the other two P450 genes, the relatively low level (~1.5-fold) of *CYP4G2* induction obtained in the ALHF house flies may suggest its relatively minor role in resistance.

*CYP4G2* has been linked on autosome 3. To our knowledge, this is the first report of a P450 gene located on auto-

some 3 of house flies. An earlier study indicated that although factors on autosome 3 were very important in the overall level of permethrin resistance in ALHF house flies [26], the resistance in ALHF governed by them was not suppressed by PBO [26]. These conflicting results may suggest that an ~1.5-fold level of induction of a P450 gene is too small to be detected by a synergism study. Alternatively, since PBO appears not to be a perfect inhibitor for some of the P450s responsible for resistance [3,48,49], the product of *CYP4G2* may not be sensitive to the inhibition of PBO and the corresponding level of resistance due to the metabolism of the gene may therefore not be suppressed by PBO.

### Conclusion

This study provides direct evidence that multiple P450 genes, *CYP4D4v2*, *CYP4G2*, and *CYP6A38*, are up-regulated in insecticide resistant house flies through the induction mechanism. Taken together with the induction of *CYP4D4v2*, *CYP4G2*, and *CYP6A38* only in resistant house flies and the correlation linking the genes with the development of resistance and/or P450-mediated resistance in ALHF, this study suggests the functional importance of these three P450 genes in the increased detoxification of insecticides in ALHF. Our previous studies [27,28] have also indicated that two P450 genes are constitutively overexpressed in ALHF house flies. Taken together, these studies indicate that both P450 induction and constitutive overexpression may be co-responsible for detoxification of insecticides, evolutionary insecticide



**Figure 6**

**Induction of *CYP4G2*, *CYP4D4v2*, and *CYP6A38* in CS and ALHF house flies following treatment with permethrin.** The expression of three genes was analyzed by qRT-PCR as described in the materials and methods. A: Relative expression of *CYP4G2* in permethrin treated and untreated susceptible CS, aabys and resistant ALHF house flies. B: Relative expression of *CYP4D4v2* in permethrin treated and untreated CS, aabys and ALHF house flies. C: Relative expression of *CYP6A38* in permethrin treated and untreated CS, aabys and ALHF house flies. The relative level of gene expression shown in Y axis is the ratio of the gene expression in each strain or each treatment in comparison with that in untreated CS flies. The results are shown as the mean  $\pm$  S.E. There was no significant difference ( $P \leq 0.05$ ) in the levels of P450 gene expression among the samples with the same alphabetic letter (i.e., a, b, or c).

selection, and the ability of insects to adapt to changing environments.

## Methods

### House fly strains

Three house fly strains were used in this study. ALHF is a wild-type strain collected from a poultry farm in Alabama in 1998, selected with permethrin for 6 generations, achieving a 6,600-fold resistance, and maintained under biannual selection with permethrin [25,26]. CS is a wild type insecticide-susceptible strain. aabys is an insecticide-susceptible strain with recessive morphological markers *ali-curve* (*ac*), *aristapedia* (*ar*), *brown body* (*bwb*), *yellow eyes* (*ye*), and *snipped wings* (*sw*) on autosomes 1, 2, 3, 4, and 5, respectively. Both CS and aabys were obtained from Dr. J. G. Scott (Cornell University).

### Permethrin challenge experiments

Two-day old resistant ALHF and susceptible CS house flies were treated with permethrin by topical application [25,26] with 0.5  $\mu$ l permethrin solution (in acetone) and acetone alone dropped on the thoracic notum. Preliminary dose range, time course, and P450 gene induction assays were performed with a corresponding dose range of LD<sub>10</sub>, LD<sub>50</sub>, and LD<sub>90</sub> (2 ng/fly, 10 ng/fly, and 20 ng/fly for susceptible CS and aabys; 5  $\mu$ g/fly, 10  $\mu$ g/fly, and 20  $\mu$ g/fly

fly for ALHF). Based on the results, in which the induction of P450s reached its peak at 24 h after ALHF were treated with permethrin at the dose of LD<sub>50</sub>, the dose of 20  $\mu$ g/fly that resulted in 50–60% mortality for resistant ALHF were chosen as the challenge dose. The surviving flies were collected for RNA extraction after exposure to permethrin. The control flies that had not been exposed to permethrin treatment and acetone treated flies were collected on the same day as the permethrin treated flies. The experiments were repeated three times.

### RNA extraction, cDNA preparation, and the putative P450 gene isolation

Total RNAs were extracted from the house flies using the acidic guanidine thiocyanate-phenol-chloroform method [7]. mRNA was isolated with oligotex-dT suspension as described by the manufacturer (QIAGEN). The first strand cDNA was synthesized with SuperScript II reverse transcriptase (Stratagene) and an antisense 5'-anchored oligo(dT) primer (Table 1) [50]. The PCR products were amplified using three primer pairs of C2/Flyh1, C2/Flyc1 and HemeR1/CYP6AD1 (Table 1). CYP6AD1 was designed based on a conserved amino acid region found in the house fly, *Drosophila*, and mosquito P450 sequences after alignment of these insect P450 genes; Flyh1 was designed based on the heme binding consensus sequence



**Figure 7**  
**Graphic representation of CYP4G2, CYP4D4v2, and CYP6A38, showing locations and sequences of SNP-specific primers for SNP determinations and genetic linkage analyses of three P450 genes. A: CYP4G2. B: CYP4D4v2. C: CYP6A38.**

[51]; HemeR1 was generated based on the complementary sequence of Flyh1; and Flyc1 was synthesized based on a conserved 13 amino acid region found in rat, human,

and insect P450 sequences [51]. The PCR products were cloned into PCR™ 2.1 Original TA cloning vector (Invitrogen) and sequenced. Cloning and sequence analyses of

**Table 2: The nucleotide(s) at the polymorphism site of each P450 gene in different house fly strains and lines generated by the crosses and resistant ALHF and susceptible aabys strains.**

House Fly Strain		P450 Gene		
		CYP4G2	CYP4D4v2	CYP6A38
ALHF		C	C	G
aabys		T	T	T
BC <sub>1</sub> Lines* (ALHF ♀ x aabys ♂)	A2345	C/T	T/C	G/T
	A1345	C/T	T/C	G/T
	A1245	T	T/C	G/T
	A1235	C/T	T/C	G/T
	A1234	C/T	T	T
BC <sub>1</sub> Lines* (ALHF ♂ x aabys ♀)	A2345	C/T	T/C	G/T
	A1345	C/T	T/C	G/T
	A1245	T	T/C	G/T
	A1235	C/T	T/C	G/T
	A1234	C/T	T	T

\* These lines were named according to the autosomes bearing wild-type markers from ALHF. For example, the A1234 strain had wild-type markers on autosomes 1, 2, 3, and 4 from ALHF and the recessive mutant marker on autosome 5 from aabys (see details in Methods).

P450 gene fragments were repeated at least three times with different preparations of mRNAs. Three TA clones from each replication were sequenced.

#### **Rapid amplification of cDNA ends (RACE) of the putative P450 gene fragments**

RACE was carried out using the Marathon™ cDNA Amplification Kit (Clontech) as described by the manufacturer and Liu and Zhang [51]. The first strand cDNAs were synthesized with AMV reverse transcriptase using ALHF mRNAs as templates. The double strand cDNA was synthesized following the protocol described by the manufacturer (Clontech). Adaptors were ligated to both ends of each double strand cDNA using T4 DNA ligase as described by the manufacturer. The 5' and/or 3' ends of the P450 cDNA fragments were amplified by PCR using adapter primer AP1 and gene specific primers generated based on the 5' and/or 3' end sequences of the putative P450 cDNA fragments. The full lengths of putative P450 cDNAs were subsequently generated by reverse transcription PCR (RT-PCR) using specific primer pairs of AP450HF5F/AP450HF5R, AP450HF20F/AP450HF20R and P450HF16F-3/AP450HF17R (Table 1) synthesized based on the 5' and 3' end sequences of the putative P450 genes. Cloning and sequence analyses of the P450 cDNAs were repeated at least three times and three TA clones from each replication were verified by sequencing.

#### **Cloning and sequencing of the 5' flanking region of CYP6A38 from ALHF and aabys**

House fly genomic DNAs were digested with 5 different restriction enzymes using the Universal GenomeWalker™ Kit (Clontech) and generated 5 adaptor-ligated ALHF genomic DNA libraries as described by the manufacturer. The adaptor ligated DNA fragments in the GenomeWalker libraries were amplified by PCR with Advantage *Tth* polymerase (Clontech), the antisense primer, P450HF17P (Table 1), based on the 5' coding region of the *CYP6A38*, and a sense primer, AP1, based on the sequence of the adaptor. The PCR products were cloned into the TA cloning vector (Invitrogen) and sequenced. Cloning and sequence analyses of PCR products were repeated at least three times each with three TA clones from each replication. The 5' flanking region of *CYP6A38* in aabys was subsequently generated by PCR from the genomic DNA using a primer pair, P450HF17P-F/P450HF17P (Table 1) designed according to the 5' flanking region of *CYP6A38* in ALHF.

#### **Northern blot analysis**

Northern blot analyses were performed according to Sambrook et al. [52]. Twenty micrograms of total RNA from each sample were fractionated on 1% formaldehyde denaturing agarose gel and transferred to Nytran membranes (Schleicher and Schuell) [52]. The P450 cDNAs were

labeled with [ $\alpha$ -<sup>32</sup>P] dCTP using a Primer-It II Random Primer Labeling kit (Stratagene) and hybridized with RNA blots using QuickHyb solution (Stratagene). The amount of RNA loaded in each lane was standardized by comparing the density of the 18S ribosomal RNA band on agarose gel under UV light before transfer [53]. All Northern blot analyses were repeated three times with different preparations of RNA samples.

#### **Quantitative Real-time PCR (qRT-PCR)**

Total RNA samples (0.5  $\mu$ g/sample) were reverse-transcribed using SuperScript II reverse transcriptase (Stratagene) in a total volume of 20  $\mu$ l. The quantity of cDNAs was measured using a spectrophotometer prior to qRT-PCR. qRT-PCR was performed with the SYBR Green master mix Kit and ABI 7500 Real Time PCR system (Applied Biosystems). Each qRT-PCR reaction (25  $\mu$ l final volume) contained 1 $\times$  SYBR Green master mix, 1  $\mu$ l of cDNA, and a gene specific primer pair, RTP450HF5F/RTP450HF5R, RTP450HF20F/RTP450HF20R, or RTP450HF17F/RTP450HF17R (Table 1), at a final concentration of 3–5  $\mu$ M. A 'no-template' negative control and all samples were performed in triplicate. The reaction cycle consisted of a melting step of 50°C for 2 min then 95°C for 10 min, followed by 40 cycles of 95°C for 15 sec and 60°C for 1 min. Specificity of the PCR reactions was assessed by a melting curve analysis for each PCR reaction using Dissociation Curves software [54]. Relative expression levels for specific genes were calculated by the  $2^{-\Delta\Delta C(t)}$  method using SDS RQ software [55]. The  $\beta$ -actin gene, an endogenous control, was used to normalize expression of target genes [56,57]. The preliminary assay had shown that the  $\beta$ -actin gene remained constant in different tissues and in both permethrin treated and untreated house flies and could therefore be used for internal normalization in qRT-PCR assays. Each experiment was repeated three times with different preparations of RNA samples. The statistical significance of the gene expressions was calculated using a Student's *t*-test for all 2-sample comparisons and a one-way analysis of variance (ANOVA) for multiple sample comparisons (SAS v9.1 software); a value of  $P \leq 0.05$  was considered statistically significant.

#### **Genetic linkage analysis of cytochrome P450 genes**

To determine the genetic linkage of P450 genes, a genetic cross experiment was conducted [26,58]. Briefly, reciprocal crosses between ALHF and aabys were conducted and  $F_1$  males were back-crossed to aabys females. Five lines were saved from the back-cross generation ( $BC_1$ ) with the genotypes of: *ac/ac*, *+/ar*, *+/bwb*, *+/ye*, *+/sw* (A2345); *+/ac*, *ar/ar*, *+/bwb*, *+/ye*, *+/sw* (A1345); *+/ac*, *+/ar*, *bwb/bwb*, *+/ye*, *+/sw* (A1245); *+/ac*, *+/ar*, *+/bwb*, *ye/ye*, *+/sw* (A1235); and *+/ac*, *+/ar*, *+/bwb*, *+/ye*, *sw/sw* (A1234). Since crossing over does not or very rarely occurs in male flies [59], the presence of a mutant phenotype indicated that the respective

autosome with a mutant-type marker was derived from the aabys females. The genotype of each line was homozygous for the recessive mutant allele from aabys and heterozygous for the dominant wild-type alleles from ALHF. These lines were named according to the autosomes bearing wild-type markers from ALHF. For example, the A1234 strain had wild-type markers on autosomes 1, 2, 3, and 4 from ALHF and the recessive mutant marker on autosome 5 from aabys.

The P450 specific alleles in ALHF and aabys and five house fly BC<sub>1</sub> lines were genetically mapped by SNP determination using an ABI Prism SNaPshot Multiplex Kit and analyzed on the ABI Prism® 3100 Genetic Analyzer using Genemapper software according to the manufacturer's instructions (A&B Applied Biosystems). Briefly, the cDNA fragments, which covered the SNP sites of P450 cDNAs, were generated by PCR. The SNP determination reactions were conducted for *CYP4D4v2*, *CYP4G2*, and *CYP6A38* using allele specific primers, SNP450HF5F, SNP450HF20F, and SNP450HF17F, respectively (Table 1, Fig. 7), designed according to the sequences immediately upstream of the nucleotide polymorphism to distinguish the single nucleotide polymorphism for the P450 allele in each house fly strain or line. Three replications of the SNP determination were carried out with different preparations of the PCR templates. To confirm that the PCR products used for the SNP determination were in fact the P450 gene fragments, the PCR products were sequenced at least once each.

### Abbreviations

ANOVA: a one-way analysis of variance; BC<sub>1</sub>: back-cross generation 1; PBO: piperonyl butoxide; qRT-PCR: quantitative real-time PCR; RACE: rapid amplification of cDNA ends; RT-PCR: Reverse transcription PCR; SNP: single nucleotide polymorphism.

### Authors' contributions

FZ was involved in coordinating and carrying out the experiments described in this report and wrote the manuscript. TL was involved in the dose range and time course studies. LZ was involved in the review of data, supervising implementation of the overall process, sequencing, and editing the manuscript. NL was involved in the overall design of study, review of data, and editing the manuscript. All authors read and approved the final manuscript.

### Acknowledgements

This study was supported by U.S. Department of Agriculture National Research Initiative (USDA-NRI) Competitive Grants Program Grants 2001-35302-10956 to N.L. and L.Z., and Hatch Project ALA08-029. We thank Dr. Nelson for naming the P450 genes, Dr. J. G. Scott for supplying the CS and aabys flies, and Jan Szechi for editorial assistance.

### References

- Pavek P, Dvorak Z: **Xenobiotic-induced transcriptional regulation of xenobiotic metabolizing enzymes of the cytochrome P450 superfamily in human extrahepatic tissues.** *Curr Drug Metab* 2008, **9**:129-143.
- Scott JG: **Cytochromes P450 and insecticide resistance.** *Insect Biochem Mol Biol* 1999, **29**:757-777.
- Feyereisen R: **Insect cytochrome P450.** In *Comprehensive Molecular Insect Science* Edited by: Gilbert LI, Iatrou K, Gill S. Oxford: Elsevier; 2005:1-77.
- Berenbaum MR: **Coumarins.** In *Herbivores: Their Interaction with Secondary Plant Metabolites* Edited by: Rosenthal GA, Berenbaum MR. New York: Academic Press; 1991:221-249.
- Schuler M: **The role of cytochrome P450 monooxygenases in plant-insect interactions.** *Plant Physiol* 1996, **112**:1411-1419.
- Carino FA, Koener JF, Plapp FW Jr, Feyereisen R: **Constitutive overexpression of the cytochrome P450 gene *CYP6A1* in a house fly strain with metabolic resistance to insecticides.** *Insect Biochem Mol Biol* 1994, **24**:411-418.
- Liu N, Scott JG: **Phenobarbital induction of *CYP6D1* is due to a trans acting factor on autosome 2 in house flies, *Musca domestica*.** *Insect Mol Biol* 1997, **6**:77-81.
- Liu N, Scott JG: **Increased transcription of *CYP6D1* causes cytochrome P450-mediated insecticide resistance in house fly.** *Insect Biochem Mol Biol* 1998, **28**:531-535.
- Kasai S, Weerasinghe IS, Shono T, Yamakawa M: **Molecular cloning, nucleotide sequence, and gene expression of a cytochrome P450 (*CYP6F1*) from the pyrethroid-resistant mosquito, *Culex quinquefasciatus* Say.** *Insect Biochem Mol Biol* 2000, **30**:163-171.
- Brattsten LB, Lee YH, Keeley LL: **Insecticide resistance: challenge to pest management and basic research.** *Science* 1986, **231**:1255-1260.
- Li X, Berenbaum MR, Schuler MA: **Cytochrome P450 and actin genes expressed in *Helicoverpa zea* and *Helicoverpa armigera*: paralogy/orthology identification, gene conversion and evolution.** *Insect Biochem Mol Biol* 2002, **32**:311-320.
- Wen Z, Pan L, Berenbaum MB, Schuler MA: **Metabolism of linear and angular furanocoumarins by *Papilio polyxenes* *CYP6B1* co-expressed with NADPH cytochrome P450 reductase.** *Insect Biochem Mol Biol* 2003, **33**:937-947.
- Daborn PJ, Yen JL, Bogwitz MR, Le Goff G, Feil E, Jeffers S, Tijet N, Perry T, Heckel D, Batterham P, Feyereisen R, Wilson TG, French-Constant RH: **A single P450 allele associated with insecticide resistance in *Drosophila*.** *Science* 2002, **297**:2253-2256.
- Daborn PJ, Lumb C, Boey A, Wong W, French-Constant RH, Batterham P: **Evaluating the insecticide resistance potential of eight *Drosophila melanogaster* cytochrome P450 genes by transgenic over-expression.** *Insect Biochem Mol Biol* 2007, **37**:512-519.
- Terriere LC: **Enzyme induction, gene amplification, and insect resistance to insecticides.** In *Pest Resistance to Pesticides* Edited by: Georgiopoulos GP, Saito T. New York: Plenum Press; 1983:265-297.
- Terriere LC: **Induction of detoxication enzymes in insects.** *Ann Rev Entomol* 1984, **29**:71-88.
- Ranasinghe C, Hobbs AA: **Induction of cytochrome P450 *CYP6B7* and cytochrome b5 mRNAs from *Helicoverpa armigera* (Hubner) by pyrethroid insecticides in organ culture.** *Insect Mol Biol* 1999, **8**:432-447.
- Scharf ME, Parimi S, Meinke LJ, Chandler LD, Siegfried BD: **Expression and induction of three family 4 cytochrome P450 (*CYP4*) genes identified from insecticide-resistant and susceptible western corn rootworms, *Diabrotica virgifera virgifera*.** *Insect Mol Biol* 2001, **10**:139-146.
- Zhu YC, Snodgrass GL: **Cytochrome P450 *CYP6X1* cDNAs and mRNA expression levels in three strains of the tarnished plant bug *Lygus lineolaris* (Heteroptera: Miridae) having different susceptibilities to pyrethroid insecticide.** *Insect Mol Biol* 2003, **12**:39-49.
- Scott JG, Sridhar P, Liu N: **Adult specific expression and induction of cytochrome P450lpr in house flies.** *Arch Insect Biochem Physiol* 1996, **31**:313-323.
- Bautista M, Tanaka T, Miyata T: **Identification of permethrin-inducible cytochrome P450s from the diamondback moth, *Plutella xylostella* (L.) and the possibility of involvement in permethrin resistance.** *Pestic Biochem Physiol* 2007, **87**:85-93.

22. Le Goff G, Hilliou F, Siegfried BD, Boundy S, Wajnberg E, Sofer L, Audant P, French-Constant RH, Feyereisen R: **Xenobiotic response in *Drosophila melanogaster*: sex dependent of P450 and GST gene induction.** *Insect Biochem Mol Biol* 2006, **36**:674-682.
23. Okey AB: **Enzyme induction in the cytochrome P-450 system.** *Pharmacol Ther* 1990, **45**:241-298.
24. Depardieu F, Podglajen I, Leclercq R, Collatz E, Courvalin P: **Modes and modulations of antibiotic resistance gene expression.** *Clin Microbiol Rev* 2007, **20**:79-114.
25. Liu N, Yue X: **Insecticide resistance and cross-resistance in the house fly (Diptera: Muscidae).** *J Econ Entomol* 2000, **93**:1269-1275.
26. Liu N, Yue X: **Genetics of pyrethroid resistance in a strain (ALHF) of house flies (Diptera: Muscidae).** *Pestic Biochem Physiol* 2001, **70**:151-158.
27. Zhu F, Feng J, Zhang L, Liu N: **Characterization of two novel cytochrome P450 genes in insecticide resistant house flies.** *Insect Mol Biol* 2008, **20**:1365-1583.
28. Zhu F, Liu N: **Differential expression of CYP6A5 and CYP6A5v2 in pyrethroid-resistant house flies, *Musca domestica*.** *Arch Insect Biochem Physiol* 2008, **34**:147-161.
29. Snyder MJ, Scott JA, Andersen JF, Feyereisen R: **Sampling P450 diversity by cloning polymerase chain reaction products obtained with degenerated primers.** *Methods Enzymol* 1996, **272**:304-312.
30. Kasai S, Scott JG: **Expression and regulation of CYP6D3 in the house fly, *Musca domestica* (L.).** *Insect Biochem Mol Biol* 2001, **32**:1-8.
31. Carino FA, Koener JF, Plapp FW Jr, Feyereisen R: **Expression of the cytochrome P450 gene CYP6A1 in the housefly, *Musca domestica*.** In *Molecular Mechanisms of Insecticide Resistance: Diversity Among Insects. ACS Symposium series 505* Edited by: Mullin CA, Scott JG. Washington, DC: American Chemical Society; 1992:31-40.
32. Liu N, Scott JG: **Genetic analysis of factors controlling high-level expression of cytochrome P450, CYP6D1, cytochrome b5, P450 reductase, and monooxygenase activities in LPR house flies, *Musca domestica*.** *Biochem Genet* 1996, **34**:133-147.
33. Brattsten LB: **Biochemical defense mechanisms in herbivores against plant allelochemicals.** In *Herbivores: Their Interaction with Secondary Plant Metabolites* Edited by: Rosenthal GA, Janzen DH. San Diego, CA: Academic Press; 1979:199-270.
34. Cohen MB, Berenbaum MR, Schuler MA: **A host-inducible cytochrome P-450 from a host-specific caterpillar: molecular cloning and evolution.** *Proc Natl Acad Sci USA* 1992, **89**:10920-10924.
35. Ma R, Cohen MB, Berenbaum MR, Schuler MA: **Black swallowtail (*Papilio polyxenes*) alleles encode cytochrome P450s that selectively metabolize linear furanocoumarins.** *Arch Biochem Biophys* 1994, **310**:332-340.
36. Snyder MJ, Stevens JL, Andersen JF, Feyereisen R: **Expression of cytochrome P450 genes of the CYP4 family in midgut and fat body of the tobacco hornworm, *Manduca sexta*.** *Arch Biochem Biophys* 1995, **321**:13-20.
37. Stevens JL, Snyder MJ, Koener JF, Feyereisen R: **Inducible P450s of the CYP9 family from larval *Manduca sexta* midgut.** *Insect Biochem Mol Biol* 2000, **30**:559-568.
38. Li X, Berenbaum MR, Schuler MA: **Molecular cloning and expression of CYP6B8: a xanthotoxin-inducible cytochrome P450 cDNA from *Helicoverpa zea*.** *Insect Biochem Mol Biol* 2000, **30**:75-84.
39. Li W, Berenbaum MR, Schuler MA: **Molecular analysis of multiple CYP6B genes from polyphagous *Papilio* species.** *Insect Biochem Mol Biol* 2001, **31**:999-1011.
40. Gonzalez FJ, Nebert DW: **Evolution of the P450 gene superfamily: animal-plant "warfare", molecular drive, and human genetic differences in drug oxidation.** *Trends Genet* 1990, **6**:182-186.
41. Li X, Baudry J, Berenbaum MR, Schuler MA: **Structural and functional evolution of insect CYP6B proteins: from specialist to generalist P450.** *Proc Natl Acad Sci USA* 2004, **101**:2939-2944.
42. Festucci-Buselli RA, Carvalho-Dias AS, de Oliveira-Andrade M, Caixeta-Nunes C, Li H-M, Stuart JJ, Muir W, Scharf ME, Pittendrigh BR: **Expression of *Cyp6g1* and *Cyp12d1* in DDT resistant and susceptible strains of *Drosophila melanogaster*.** *Insect Mol Biol* 2005, **14**:69-77.
43. Islam Z, Gray JS, Pestka JJ: **p38 mitogen-activated protein kinase mediates IL-8 induction by the ribotoxin deoxyvalenol in human monocytes.** *Toxicol Appl Pharmacol* 2006, **213**:235-244.
44. Willoughby L, Chung H, Lumb C, Robin C, Batterham P, Daborn PJ: **A comparison of *Drosophila melanogaster* detoxification gene induction responses for six insecticides, caffeine and Phenobarbital.** *Insect Biochem Mol Biol* 2006, **36**:934-942.
45. Rose RL, Goh D, Thompson DM, Verma KD, Heckel DG, Gahan LJ, Roe RM, Hodgson E: **Cytochrome P450 (CYP) 9A1 in *Heliothis virescens*: the first member of a new CYP family.** *Insect Biochem Mol Biol* 1997, **27**:605-615.
46. Guzov VM, Unnithan GC, Chernogolov AA, Feyereisen R: **CYP12A1, a mitochondrial cytochrome P450 from the house fly.** *Arch Biochem Biophys* 1998, **359**:231-240.
47. Maitra S, Dombrowski SM, Basu M, Raustol O, Waters LC, Ganguly R: **Factors on the third chromosome affect the level of *Cyp6a2* and *Cyp6a8* expression in *Drosophila melanogaster*.** *Gene* 2000, **248**:147-156.
48. Brown TM, Bryson PK, Payne GT: **Synergism by propynyl aryl ethers in permethrin-resistant tobacco budworm larvae.** *Pestic Sci* 1996, **46**:323-331.
49. Zhang L, Harada K, Shono T: **Genetic analysis of pyriproxifen resistance in the housefly, *Musca domestica* (L.).** *Appl Ent Zool* 1997, **32**:217-226.
50. Tomita T, Scott JG: **cDNA and deduced protein sequence of *Cyp6d1*: the putative gene for a cytochrome P450 responsible for pyrethroid resistance in house fly.** *Insect Biochem Mol Biol* 1995, **25**:275-283.
51. Liu N, Zhang L: **Identification of two cytochrome P450 genes and their 5' flanking regions from the house fly, *Musca domestica* (L.).** *Insect Biochem Mol Biol* 2002, **32**:755-764.
52. Sambrook J, Fritsch EF, Maniatis T: *Molecular Cloning. A Laboratory Manual* Cold Spring Harbor: Cold Spring Harbor Laboratory Press; 1989.
53. Spiess AN, Ivell R: **Normalization of RNA hybridization signals by means of SYBR Green II-stained 28S or 18S ribosomal RNA and phosphor imager.** *BioTechniques* 1998, **26**:46-50.
54. Wittwer CT, Herrmann MG, Moss AA, Rasmussen RP: **Continuous fluorescence monitoring of rapid cycle DNA amplification.** *BioTechniques* 1997, **22**:130-131.
55. Livak KJ, Schmittgen TD: **Analysis of relative gene expression data using real-time quantitative PCR and the 2<sup>-ΔΔCT</sup> method.** *Methods* 2001, **25**:402-408.
56. Aerts JL, Gonzales MI, Topalian SL: **Selection of appropriate control genes to assess expression of tumor antigens using real-time RT-PCR.** *BioTechniques* 2004, **36**:84-86.
57. Dorak MT: **Real-Time PCR.** In *Real-Time PCR (Advanced Methods Series)* Edited by: Dorak MT. Oxford: Taylor & Francis; 2006:1-333.
58. Liu N, Scott JG: **Genetics of resistance to pyrethroid insecticides in the house fly, *Musca domestica*.** *Pestic Biochem Physiol* 1995, **52**:116-124.
59. Gao J, Scott JG: **Role of the transcriptional repressor *mdGfi-1* in CYP6D1v1-mediated insecticide resistance in the house fly, *Musca domestica*** 2006, **36**:387-395.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

