



Messina (*Melilotus siculus*)—A New Pasture Legume for Saltland

Amanda L. Bonython

Cooperative Research Centre for Future Farm Industries, Australia

Andrew D. Craig

Cooperative Research Centre for Future Farm Industries, Australia

Ross A. Ballard

Cooperative Research Centre for Future Farm Industries, Australia

Nigel Charman

Cooperative Research Centre for Future Farm Industries, Australia

Philip G. H. Nichols

Cooperative Research Centre for Future Farm Industries, Australia

See next page for additional authors

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/22/1/6>

The 22nd International Grassland Congress (Revitalising Grasslands to Sustain Our Communities) took place in Sydney, Australia from September 15 through September 19, 2013.

Proceedings Editors: David L. Michalk, Geoffrey D. Millar, Warwick B. Badgery, and Kim M.

Broadfoot

Publisher: New South Wales Department of Primary Industry, Kite St., Orange New South Wales, Australia

Presenter Information

Amanda L. Bonython, Andrew D. Craig, Ross A. Ballard, Nigel Charman, Philip G. H. Nichols, Clinton Revell, and Natasha L. Teakle

Messina (*Melilotus siculus*) – a new pasture legume for saltland

Amanda L Bonython^{AB}, Andrew D Craig^{AB}, Ross A Ballard^{AB}, Nigel Charman^{AB}, Philip GH Nichols^{ACD}, Clinton Revell^{AC}, and Natasha L Teakle^{DE}

^A Cooperative Research Centre for Future Farm Industries, Crawley, Western Australia

^B South Australian Research and Development Institute, Adelaide, South Australia

^C Department of Agriculture and Food Western Australia, South Perth, Western Australia

^D School of Plant Biology, The University of Western Australia, Crawley, Western Australia

^E Centre for Ecohydrology, The University of Western Australia, Crawley, Western Australia

Contact email: amanda.bonython@sa.gov.au

Keywords: Annual fodder legume, dryland salinity, *Melilotus messanensis*, salinity tolerance, Sicilian sweet clover, waterlogging tolerance.

Introduction

Messina (*Melilotus siculus* ((Turra) Vitman ex B.D. Jacks)) is a new annual pasture legume for saltland in temperate Australia and regions of the world that experience Mediterranean climates. Messina has greater tolerance to the combined stresses of salinity and waterlogging than existing commercial pasture legumes. Coupled with desirable agronomic traits these characteristics give messina the capacity to rehabilitate saltland and increase productivity on land where existing legumes fail. This paper reviews the agronomic performance of messina in relation to top soil salinity levels.

Methods

In June 2010, 21 low coumarin messina accessions were sown at three trial sites in the upper South East of South Australia (36.14°S, 140.03°E). All sites had alkaline soils (clay loams to loamy sands) and two incurred inundation over winter. The sites experienced a typical Mediterranean climate. Two elite messina accessions have been identified and their performance has been compared with the current commercial saltland legume options; *Trifolium michelianum* Savi (cv. Frontier), *Medicago polymorpha* L. (cv. Scimitar) and *Melilotus albus* Medik. (cv. Jota). The average number of days from sowing to first flower, seasonal dry matter (DM kg/ha) and seed production (kg/ha) were assessed. Seasonal top soil (0-10 cm) salinity (EC_e, dS/m) was also assessed (data not presented).

Results

The average number of days from sowing to first flower was 118 and 119 for messina A and B respectively, compared to Frontier (117 days) and Scimitar (111 days). Jota was not assessed.

Messina dry matter production in spring 2010 was comparable to Frontier, but significantly greater than Scimitar (Table 1). In winter 2011 messina again performed as well as Frontier, but poorer than Scimitar. In both the winter and spring of 2012 messina dry matter production was significantly greater than the three controls. Overall Jota performed poorly.

Mean seed availability for the 2010, 2011 and 2012 harvests was significantly higher for messina (messina A 2084 kg/ha and messina B 2362 kg/ha) than Frontier (537 kg/ha), Scimitar (1299 kg/ha) and Jota (589 kg/ha).

Discussion

The maturity of messina is comparable to Frontier and Scimitar, indicating it will be adapted to similar rainfall environments (350-500 mm pa). Similar results have been recorded in Western Australia, the major target zone for messina in Australia (P. Nichols unpublished data). Importantly, messina has performed well on moderately acidic saline soils in WA (Nichols *et al.* 2008).

Messina seedlings can germinate at high salinity levels, in contrast to those of Frontier (low tolerance) and

Table 1. Multi-site ANOVA of seasonal dry matter production (kg/ha) at three trial sites.

	Spring 2010A	Winter 2011A	Spring 2011	Winter 2012	Spring 2012
Messina A	3277	1283	5520	612	2826
Messina B	3228	1548	4067	581	2790
Frontier	3988	1856	7420	269	924
Scimitar	1704	2716	5725	223	1200
Jota	320	184	1575	21	553
LSD (P=0.05)	788	870	1578	231	1191

A: Sites 1 and 2 assessed only.

Scimitar (moderate tolerance) (Nichols *et al.* 2009; Rogers *et al.* 2011; Teakle *et al.* 2012). Autumn top soil salinity levels at the three trial sites increased from an average of 4 dS/m (low salinity) in 2011 to 10 dS/m (high salinity) in 2012. Messina outperformed Frontier and Scimitar at these higher salinity levels, indicating that its greatest contribution to increasing pasture productivity will occur in highly saline environments where existing pasture legumes fail. The poor performance of Jota during these trials can be attributed to its low waterlogging tolerance.

Messina is a prolific seed producer. High second year seed reserves are typically the result of substantial hard-seed carryover from year one (Nichols *et al.* 2009). This feature appears important for annual legume persistence on saltland, where environmental uncertainty increases the requirement for a long-term hard-seed residual (Cohen 1967; Venable and Brown 1988).

Conclusion

The greatest contribution of messina to improving pasture productivity appears to be on highly saline soils where existing legumes fail. Prior to its commercial release, some outstanding information is required to ensure the successful uptake of messina. This includes determining its area of adaptation, undertaking studies to confirm its value and safety for grazing animals, and determining management strategies for optimum pasture and animal performance.

Acknowledgements

This research was funded through the Future Farm Industries

CRC and supported by the South Australian of Australian and Food Western Australia. We thank the landholders for their cooperation and technical staff for their assistance.

References

- Cohen D (1967) Optimising reproduction in a randomly varying environment when a correlation may exist between the conditions at the time a choice has to be made and the subsequent outcome. *Journal of Theoretical Biology* **16**, 1-14.
- Teakle NL, Bowman S, Barrett-Lennard EG, Real D, Colmer TD (2012) Comparisons of annual pasture legumes in growth, ion regulation and root porosity demonstrate that *Melilotus siculus* has exceptional tolerance to combinations of salinity and waterlogging. *Environmental and Experimental Botany* **77**, 175-184.
- Rogers ME, Colmer TD, Nichols PGH, Hughes SJ, Frost K, Cornwall D, Chandra S, Miller SM, Craig AD (2011) Salinity and waterlogging tolerance amongst accessions of messina (*Melilotus siculus*). *Crop and Pasture Science* **62**, 225-235.
- Nichols PGH, Craig AD, Rogers ME, Albertsen TO, Miller S, McClements DR, Hughes SJ, D'Antuono MFD, Dear BS (2008) Production and persistence of annual legumes at five saline sites in southern Australia. *Australian Journal of Experimental Agriculture* **48**, 518-535.
- Nichols PGH, Malik AI, Stockdale M, Colmer TD (2009) Salt tolerance and avoidance mechanisms at germination of annual pasture legumes and their importance for adaptation to saline environments. *Plant and Soil* **315**, 241-255.
- Venable DL, Brown JS (1988) The selective interactions of dispersal dormancy and seed size as adaptations for reducing risk in variable environments. *American Naturalist* **131**, 360-384.