



1-1-2018

ICTV Virus Taxonomy Profile: *Partitiviridae*

Eeva J. Vainio

Natural Resources Institute Finland, Finland

Sotaro Chiba

Nagoya University, Japan

Said A. Ghabrial

University of Kentucky, saghab00@uky.edu

Edgar Maiss

Leibniz University Hannover, Germany

Marilyn Roossinck

Pennsylvania State University

See next page for additional authors

[Click here to let us know how access to this document benefits you.](#)

Follow this and additional works at: https://uknowledge.uky.edu/plantpath_facpub

 Part of the [Genetics and Genomics Commons](#), [Virology Commons](#), and the [Viruses Commons](#)

Repository Citation

Vainio, Eeva J.; Chiba, Sotaro; Ghabrial, Said A.; Maiss, Edgar; Roossinck, Marilyn; Sabanadzovic, Sead; Suzuki, Nobuhiro; Xie, Jiatao; Nibert, Max; and ICTV Report Consortium, "ICTV Virus Taxonomy Profile: *Partitiviridae*" (2018). *Plant Pathology Faculty Publications*. 83.

https://uknowledge.uky.edu/plantpath_facpub/83

This Article is brought to you for free and open access by the Plant Pathology at UKnowledge. It has been accepted for inclusion in Plant Pathology Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Authors

Eeva J. Vainio, Sotaro Chiba, Said A. Ghabrial, Edgar Maiss, Marilyn Roossinck, Sead Sabanadzovic, Nobuhiro Suzuki, Jiatao Xie, Max Nibert, and ICTV Report Consortium

ICTV Virus Taxonomy Profile: *Partitiviridae*

Notes/Citation Information

Published in *Journal of General Virology*, v. 99, issue 1, p. 17-18.

This is an open access article published by the Microbiology Society under the [Creative Commons Attribution License](#).

Digital Object Identifier (DOI)

<https://doi.org/10.1099/jgv.0.000985>



ICTV Virus Taxonomy Profile: *Partitiviridae*

Eeva J. Vainio,^{1,*} Sotaro Chiba,² Said A. Ghabrial,³ Edgar Maiss,⁴ Marilyn Roossinck,⁵ Sead Sabanadzovic,⁶ Nobuhiro Suzuki,⁷ Jiatao Xie,⁸ Max Nibert⁹ and ICTV Report Consortium

Abstract

The *Partitiviridae* is a family of small, isometric, non-enveloped viruses with bisegmented double-stranded (ds) RNA genomes of 3–4.8 kbp. The two genome segments are individually encapsidated. The family has five genera, with characteristic hosts for members of each genus: either plants or fungi for genera *Alphapartitivirus* and *Betapartitivirus*, fungi for genus *Gammapartitivirus*, plants for genus *Deltapartitivirus* and protozoa for genus *Cryspovirus*. Partitiviruses are transmitted intracellularly via seeds (plants), oocysts (protozoa) or hyphal anastomosis, cell division and sporogenesis (fungi); there are no known natural vectors. This is a summary of the International Committee on Taxonomy of Viruses (ICTV) Report on the taxonomy of the *Partitiviridae*, which is available at www.ictv.global/report/partitiviridae.

Table 1. Characteristics of the family *Partitiviridae*

Typical member:	<i>Atkinsonella hypoxylon virus</i>, 2H (RNA1, L39125; RNA2, L39126), species <i>Atkinsonella hypoxylon virus</i>, genus <i>Betapartitivirus</i>
Genome	3–4.8 kbp of linear bisegmented dsRNA
Virion	Isometric, non-enveloped, 25–43 nm in diameter; dsRNA1 and dsRNA2 are separately encapsidated
Replication	Cytoplasmic. Genomic RNA acts as a template for mRNA synthesis within the virus particle; transcription occurs by a semiconservative mechanism
Translation	From monocistronic positive-sense transcripts of both genomic dsRNAs
Host range	Plants, fungi and protozoa
Taxonomy	Five genera, including >40 species, and 15 species unassigned to a genus

VIRION

Virus particles are isometric, non-enveloped, and 25–43 nm in diameter (Table 1, Fig. 1a, b). Each capsid is composed of 120 copies of a single protein arranged as 60 dimers with $T=1$ icosahedral symmetry [1]. Dimeric surface protrusions are frequently observed on viral capsids. One or two molecules of RNA-dependent RNA polymerase (RdRP) are packaged inside each particle [2].

REPLICATION

Each dsRNA is monocistronic. The RdRP is believed to function as both a transcriptase and a replicase and catalyzes *in vitro* end-to-end transcription of each dsRNA to produce mRNA by a semi-conservative mechanism. Virions accumulate in the cytoplasm.

GENOME

Members of all five genera possess two essential genome segments, dsRNA1 and dsRNA2, each containing one large ORF on the positive-strand RNA molecule (Fig. 2). The smaller of the two dsRNA genome segments usually encodes the coat protein (CP) and the larger usually encodes the virion-associated RNA polymerase. The linear dsRNA segments are separately encapsidated. Additional dsRNA segments (satellite or defective) may also be present.

TAXONOMY

Alphapartitivirus

Members of the genus *Alphapartitivirus* infect either plants, or ascomycetous or basidiomycetous fungi. The two

Received 21 November 2017; Accepted 23 November 2017

Author affiliations: ¹Natural Resources Institute Finland (Luke), Helsinki 00790, Finland; ²Asian Satellite Campuses Institute, Nagoya University, Nagoya 464-0861, Japan; ³Department of Plant Pathology, University of Kentucky, Lexington, KY 40546, USA; ⁴Institute of Horticultural Production Systems, Leibniz University Hannover, Hannover 30419, Germany; ⁵Center for Infectious Disease Dynamics, Pennsylvania State University, University Park, PA 16802, USA; ⁶Department of Biochemistry, Molecular Biology, Entomology and Plant Pathology, Mississippi State University, MS 39762, USA; ⁷Institute of Plant Science and Resources, Okayama University, Chuo 2-20-1, Kurashiki, 710-0046, Japan; ⁸College of Plant Science and Technology, Huazhong Agricultural University, Wuhan 430070, Hubei Province, PR China; ⁹Department of Microbiology and Immunobiology, Harvard Medical School, Boston, MA 02115, USA.

*Correspondence: Eeva J. Vainio, eeva.vainio@luke.fi

Keywords: *Partitiviridae*; ICTV; taxonomy; *Alphapartitivirus*; *Betapartitivirus*; *Deltapartitivirus*; *Gammapartitivirus*; *Cryspovirus*.

Abbreviations: CP, coat protein; RdRP, RNA-dependent RNA polymerase.




Fig. 1. (a) Transmission electron micrograph of negatively-stained purified particles of *Penicillium stoloniferum* virus S, a representative member of the genus *Gammapartitivirus*. (b) Cryo-EM reconstructions of *Penicillium stoloniferum* virus S at 0.45 nm resolution, and rendered with radial colour mapping.

essential dsRNA genome segments are individually about 1.9–2.0 kbp (dsRNA1) and 1.7–1.9 kbp (dsRNA2), typically containing a poly(A) tract near the plus-strand 3′-terminus. There is a single major CP with predicted *Mr* of 51–57 kDa. Plant alphapartitiviruses cause persistent infections, whereas some fungal alphapartitiviruses cause host effects, such as hypovirulence or a reduced growth rate [3, 4].

Betapartitivirus

Members of the genus *Betapartitivirus* infect either plants, or ascomycetous or basidiomycetous fungi. The two essential dsRNA genome segments are about 2.2–2.4 kbp (dsRNA1) and 2.1–2.4 kbp (dsRNA2), typically containing a poly(A) tract near the plus-strand 3′-terminus. There is a single major CP with predicted *Mr* of 71–77 kDa. Plant betapartitiviruses cause persistent infections [5, 6]. Some fungal betapartitiviruses cause reduced host virulence and changes in colony morphology [7].

Gammapartitivirus

All known members of the genus *Gammapartitivirus* infect ascomycetous fungi. The two essential dsRNA segments are about 1.6–1.8 kbp (dsRNA1) and 1.4–1.6 kbp (dsRNA2). There is a single major CP with predicted *Mr* of 44–47 kDa. Most gammapartitiviruses seem to induce latent infections. *Aspergillus fumigatus* partitivirus 1, a related, unclassified virus, has been associated with host effects.




Fig. 2. *Atkinsonella hypoxylon* virus [10], an isolate of the type species of the genus *Betapartitivirus*, has a bipartite genome consisting of dsRNA1 and dsRNA2.

Deltapartitivirus

All known members of the genus *Deltapartitivirus* induce persistent infections in plants [8]. They are transmitted by ovule and pollen to the seed embryo. The two essential dsRNA segments are individually 1.6–1.7 kbp (dsRNA1) and 1.4–1.6 kbp (dsRNA2). There is a single major CP with predicted *Mr* of 38–49 kDa.

Cryspovirus

Members of the genus *Cryspovirus* infect apicomplexan protozoa of the genus *Cryptosporidium* [9]. The viral genome comprises two dsRNA segments, which are individually 1.5 and 1.8 kbp. There is a single major CP with predicted *Mr* of 37 kDa. Virions are disseminated within *Cryptosporidium* oocysts. Infections of the *Cryptosporidium* host cells appear to be latent.

RESOURCES

Full ICTV Online (10th) Report:
www.ictv.global/report/partitiviridae.

Funding information

Production of this summary, the online chapter, and associated resources was funded by a grant from the Wellcome Trust (WT108418AIA).

Acknowledgements

Members of the ICTV Report Consortium are Elliot J. Lefkowitz, Andrew J. Davison, Stuart G. Siddell, Sead Sabanadzovic, Donald B. Smith, Richard J. Orton and Peter Simmonds.

Conflicts of interest

The authors declare that there are no conflicts of interest.

References

1. Nibert ML, Tang J, Xie J, Collier AM, Ghabrial SA *et al*. 3D structures of fungal partitiviruses. *Adv Virus Res* 2013;86:59–85.
2. Nibert ML, Ghabrial SA, Maiss E, Lesker T, Vainio EJ *et al*. Taxonomic reorganization of family *Partitiviridae* and other recent progress in partitivirus research. *Virus Res* 2014;188:128–141.
3. Chiba S, Lin YH, Kondo H, Kanematsu S, Suzuki N. Effects of defective interfering RNA on symptom induction by, and replication of, a novel partitivirus from a phytopathogenic fungus, *Rosellinia necatrix*. *J Virol* 2013;87:2330–2341.
4. Vainio EJ, Hantula J. Taxonomy, biogeography and importance of *Heterobasidium* viruses. *Virus Res* 2016;219:2–10.
5. Lesker T, Rabenstein F, Maiss E. Molecular characterization of five betacryptoviruses infecting four clover species and dill. *Arch Virol* 2013;158:1943–1952.
6. Roossinck MJ. Lifestyles of plant viruses. *Philos Trans R Soc Lond B Biol Sci* 2010;365:1899–1905.
7. Xiao X, Cheng J, Tang J, Fu Y, Jiang D *et al*. A novel partitivirus that confers hypovirulence on plant pathogenic fungi. *J Virol* 2014;88:10120–10133.
8. Sabanadzovic S, Valverde RA. Properties and detection of two cryptoviruses from pepper (*Capsicum annuum*). *Virus Genes* 2011;43:307–312.
9. Nibert ML, Woods KM, Upton SJ, Ghabrial SA. *Cryspovirus*: a new genus of protozoan viruses in the family *Partitiviridae*. *Arch Virol* 2009;154:1959–1965.
10. Oh CS, Hillman BI. Genome organization of a partitivirus from the filamentous ascomycete *Atkinsonella hypoxylon*. *J Gen Virol* 1995;76:1461–1470.