



University of Kentucky
UKnowledge

Toxicology and Cancer Biology Faculty
Publications

Toxicology and Cancer Biology

12-2020

Epigenetic Regulation of Prostate Cancer

Ruixin Wang

University of Kentucky, rwa284@uky.edu

Xiaoqi Liu

University of Kentucky, Xiaoqi.Liu@uky.edu

Follow this and additional works at: https://uknowledge.uky.edu/toxicology_facpub



Part of the [Genetics and Genomics Commons](#), [Male Urogenital Diseases Commons](#), and the [Oncology Commons](#)

[Right click to open a feedback form in a new tab to let us know how this document benefits you.](#)

Repository Citation

Wang, Ruixin and Liu, Xiaoqi, "Epigenetic Regulation of Prostate Cancer" (2020). *Toxicology and Cancer Biology Faculty Publications*. 97.

https://uknowledge.uky.edu/toxicology_facpub/97

This Review is brought to you for free and open access by the Toxicology and Cancer Biology at UKnowledge. It has been accepted for inclusion in Toxicology and Cancer Biology Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Epigenetic Regulation of Prostate Cancer

Digital Object Identifier (DOI)

<https://doi.org/10.1016/j.gendis.2019.10.018>

Notes/Citation Information

Published in *Genes & Diseases*, v. 7, issue 4.

© 2019 Chongqing Medical University

This is an open access article under the CC BY-NC-ND license (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).



REVIEW ARTICLE

Epigenetic regulation of prostate cancer

Ruixin Wang ^{a,b,*}, Xiaoqi Liu ^{b,c}

^a Department of Biochemistry, Purdue University, West Lafayette, IN, 47907, USA

^b Department of Toxicology and Cancer Biology, University of Kentucky, Lexington, KY, 40536, USA

^c Markey Cancer Center, University of Kentucky, Lexington, KY, 40536, USA

Received 11 March 2019; received in revised form 16 October 2019; accepted 30 October 2019

Available online 9 November 2019

KEYWORDS

DNMT;
Epigenetics;
EZH2;
HDAC;
Phosphorylation;
Prostate cancer

Abstract Prostate cancer (PCa) the second leading cause of cancer death in males in the United State, with 174,650 new cases and 31,620 deaths estimated in 2019. It has been documented that epigenetic deregulation such as histone modification and DNA methylation contributes to PCa initiation and progression. EZH2 (enhancer of zeste homolog 2), the catalytic subunit of the Polycomb Repressive Complex (PRC2) responsible for H3K27me3 and gene repression, has been identified as a promising target in PCa. In addition, overexpression of other epigenetic regulators such as DNA methyltransferases (DNMT) is also observed in PCa. These epigenetic regulators undergo extensive post-translational modifications, in particular, phosphorylation. AKT, CDKs, PLK1, PKA, ATR and DNA-PK are the established kinases responsible for phosphorylation of various epigenetic regulators.

Copyright © 2019, Chongqing Medical University. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

While a majority of studies have demonstrated that accumulation of genetic mutations will result in cancer initiation and progression, epigenetic changes without altering DNA sequences, which have been under intensive research in recent years, also contribute to activation of oncogenes and inactivation of tumor suppressors, and subsequently

leading to the development of cancer.¹ Epigenetics is usually defined as a heritable change in gene expression without alteration in DNA sequence, including three primary epigenetic mechanisms - DNA methylation, covalent modification of histones and non-coding RNAs. It has been noted that heritable epigenetic marks can be dynamically regulated in response to any change in physiological conditions. Therefore, failure of the appropriate maintenance of these marks will highly possibly result in disease states such as cancer.² Epigenetic modifiers refer to adding or removing DNA methylation or histone modifications, which are defined as writers or erasers respectively. Besides writers and erasers, some epigenetic effectors can also be recruited, and affect the final epigenetic programs which we call them “readers”. Increasing data has shown that the

* Corresponding author. Department of Biochemistry, Purdue University, West Lafayette, IN, 47907, USA. Fax: +859 323 1059.

E-mail address: wang2412@purdue.edu (R. Wang).

Peer review under responsibility of Chongqing Medical University.

activity of these epigenetic modifiers are also under regulation of some posttranslational modifications, such as phosphorylation, which might determine the final biological outcome.³ In this review, we will take a comprehensive look of current findings of the regulation of the epigenetic modifiers by phosphorylation during carcinogenesis. We will also further discuss phosphorylation of one critical "writer"- enhancer of zeste homolog 2 (EZH2), which is a histone methyltransferase mainly mediating trimethylation of histone H3 at Lys 27 (H3K27me3), in prostate cancer, and the idea of therapeutic strategies based on EZH2 phosphorylation in prostate cancer treatment.

Epigenetic modification in cancer

DNA methylation aberrations have been firstly and mostly linked to cancer initiation and progression among the epigenetic alterations, which is featured with genome-wide hypomethylation and hypermethylation of clusters of CpGs, known as CpG islands.⁴ The role of global DNA hypomethylation in tumorigenesis has been well established. It occurs in various cancers and becomes an important reason resulting in increased genomic instability and inappropriate activation of oncogenes.⁵ Hypermethylation of CpG islands in the promoters of tumor suppressor genes to silence their expressions has been well noted to contribute to tumorigenesis. Many tumor suppressor genes, which are usually involved in various tumor-associated cellular processes including DNA repair, cell cycle, apoptosis, etc., have been observed to undergo CpG island hypermethylation. These genes, such as cell cycle related gene RB, the DNA repair protein BRCA1, and the tumor suppressor p53, are observed in different types of cancer like esophageal cancer, colorectal and gastric cancers, in which they are commonly mutated.^{6–9}

DNA methylation is maintained by a family of enzymes which are called DNA methyltransferases (DNMTs). There are four members of the DNMT family, including DNMT1, DNMT3A, DNMT3B and DNMT3L. Among them, DNMT1 plays the main role in maintaining the methylation status of DNA; at the same time, DNMT3A and DNMT3B are known to encode the *de novo* methyltransferases which will methylate the unmethylated DNA, while DNMT3L, unlike the other DNMTs, has no enzymatic activity.¹⁰ A number of studies have demonstrated a relationship between alterations of DNMTs and tumorigenesis. Overexpression of DNMTs has been well reported in a variety of human cancers via correlating with aberrant DNA methylation. Consequently, overexpression of DNMTs tends to result in increased metastasis and poor prognosis. Highly expressed DNMT1 has been found in numerous patient specimens such as esophageal squamous cell carcinoma and pancreatic cancer. Similarly, increased DNMT3A or DNMT3B is involved in liver cancer, BRCA1-mutated breast tumor, intestinal neoplasia and prostate tumors.^{11–17} In addition to overexpression of DNMTs, somatic mutations in DNMTs are also reported as an important contributor to malignant transformation. These mutations have been observed in colon cancers or acute myeloid leukemia, thus leading to disruption of normal DNA methylation and subsequently tumor promotion.^{18–20} While deletion of DNMTs in mouse

models has shown a lethal phenotype, several recent studies based on the conditional knockout approach demonstrated that loss of DNMTs also participates in development of peripheral T cell lymphoma (PTCL) or AML.^{21–23}

The N-terminal tails of histones, in which lysine and arginine residues are distributed, are subject to a variety of covalent posttranslational modifications (PTMs), such as acetylation, phosphorylation and methylation.²⁴ PTMs of proteins are highly dynamic in response to the altered contexts to ensure the histone modification in balance which is critical for maintaining genome integrity.²⁵ Many different combinations of PTMs on multiple residues, which defined as "histone code" and regulated by enzymes as "writers" and "erasers", can precisely govern specific cellular responses, such as cell cycle or signal transductions.^{26,27}

Misregulation of histone PTMs, including acetylation, methylation and phosphorylation, have been extensively linked to a variety of cancer types.²⁸ In tumors, such a misregulation results in the abnormal activation of oncogenes or the repression of tumor suppressors. And the PTMs-induced inappropriate activation or inactivation depends on which residues are modified and which type of modifications occur.²⁹ Generally speaking, lysine acetylation can open up chromatin structure and subsequently tend to activate the transcription of its target genes.²⁴ Therefore, histone acetyltransferase (HATs) should promote transcription whereas histone deacetylases (HDACs) should be anti-transcriptional. Alterations and mutations occur on HATs (e.g. MOZ or CBP/EP300) or HDACs have been reported to correlate with a poor clinical outcome in cancer patients.^{30–32} Besides the histone acetylation "writers" HATs or "erasers" HDACs, the histone acetylation "readers" the BET (bromodomain and extraterminal domain) proteins, such as BRD4, show increased expression in a variety of cancers. Consequently, the BET inhibitors, like JQ1, have demonstrated great therapeutic efficacy for the treatment of cancers, such as leukemia.^{33,34} Unlike histone acetylation, histone methylation will not exclusively act as a transcriptional activator or repressor; the activation or inactivation depends upon the open or close of the chromatin structure arising from which residue is modified and the degree of its methylation.³⁵ Similar with histone acetylation, tumor cells are also found common alterations of histone methylation and its histone methyltransferases (HMTs). For example, H3K27 trimethylation contributes to the aberrant silencing of multiple tumor suppressor genes and is associated with poor diagnosis of patients, and correspondingly, its main HMT, EZH2, is overexpressed in these cancers such as prostate cancer and breast cancer.³⁶ Similarly, the dysregulation of other HMTs and the methylation patterns (e.g. G9a and H3K9me3), have been found in cancers as well.³⁷ Histone demethylases have been recently identified to have a linkage to cancer. For example, LSD1, the demethylase for H3K4 and H3K9 residues, has been found overexpressed in many types of cancer.³⁸ Histone phosphorylation is a dynamic process catalyzed by several distinct kinases that are depending on different amino acid residues in histone.³⁹ Histone phosphorylation occurs with the change of many cellular processes, such as cell cycle, DNA damage

repair, and apoptosis, therefore its misregulation often leads to tumorigenesis. Accordingly, the kinases regulating the histone phosphorylation are always found overexpressed in cancers. For example, high PRK1 level, which mediates H3T11 phosphorylation, correlates with high stages of prostate cancer.⁴⁰

Noncoding RNAs (ncRNAs) has also been regarded as a crucial epigenetic modification gaining intensive attention. With technical development, many types of ncRNAs have been identified, including microRNAs (miRNAs), PIWI-interacting RNAs (piRNAs), small nucleolar RNAs (snoRNAs), the complicated long ncRNAs (lncRNAs), and the newly validated circular RNAs (circRNAs), etc.⁴¹ Increasing evidence has shown that ncRNAs are of essential importance for the development of human diseases notably cancers via their crucial regulatory potency in gene expression at the transcriptional, post-transcriptional, or translational processes. Among them, the most deeply studied miRNAs and lncRNAs, aberrantly expressed in a variety of cancers, play oncogenic or tumor suppressive functions in these processes.^{42,43}

As a post-transcriptional regulator of gene expression, miRNAs have been extensively believed abnormally expressed in tumors. The dysregulation of miRNAs would affect several cancer hallmarks for tumor initiation and progression which include sustaining cell cycle and proliferative signaling, avoiding from cell death, and intriguing invasion and metastasis.⁴⁴ Cell-cycle regulation is highly rely on cyclins, cyclin-dependent kinases (CDKs) and their inhibitors. In a variety of human tumors, the CDK inhibitor p27^{Kip1}, p21^{CIP1} and p16^{INK4a} have been found to be directly regulated by miRNAs such as miR-221/222 miR-663, miR-302 and miR-24 which are highly upregulated in tumors.^{45–47} In addition, miRNAs can also regulate expression of CDKs and cyclin not just inhibitors. Cyclin D1 and CDK4 are repressed by miRNA-545 leading to cell-cycle arrest in lung cancer.⁴⁸ MiRNAs have been known to take part in resisting apoptosis. Dysregulation of several miRNAs involved in p53 functions, such as miR-17–92, has been identified and confers tumor cells resistant to chemotherapy-induced cell death.⁴⁹ It has also been shown that miRNAs are fully involved in cancer metastasis. For example, a number of TGF- β -associated miRNAs, such as miR-155, engage in the process of TGF- β -induced epithelial–mesenchymal transition (EMT) which have been widely accepted as a key step in the cancer metastasis.^{50,51}

Similar with miRNAs, increasing research has revealed lncRNAs playing an emerging role in tumor suppression or oncogenesis in recent years. It has been well demonstrated that aberrant lncRNAs affect cancer cell proliferation and metastasis by remodeling chromatin structure and regulating oncogenes or tumor suppressors transcriptionally and post-transcriptionally.⁵² A good example of lncRNAs mediating epigenetic modifications is the HOX antisense intergenic RNA (HOTAIR) which has been applied as a predictive marker for metastasis in a variety of cancer, such as breast cancer.⁵³ HOTAIR working with the chromatin-modifying complexes PRC2 which catalyzes H3K27 trimethylation, affects the transcription of target genes via a PCR2-lncRNA binding.⁵⁴ Besides influencing PRC2 enzymatic function, HOTAIR can also enhance PLK1-mediated proteasomal

degradation of SUZ12 which is a core component of PRC2 complex during hepatitis B virus-induced liver carcinogenesis.⁵⁵

Phosphorylation of epigenetic modifiers in cancer

Epigenetic modifiers, including “writers”, “erasers” and “readers”, play a crucial role in maintaining the dynamic balance of epigenetic modification patterns, usually depending on their activities. It has been well established that these modifiers are also under regulation of post-translational modifications (PTMs), and their activity are affected by these PTMs consequently.³ Among the PTMs, the importance of phosphorylation contributing to epigenetic events in response to environmental changes has been widespread accepted. These phosphorylation is catalyzed and mediated by many kinases including protein kinase B (PKB/Akt), cyclin-dependent kinases (CDKs), polo-like kinase 1(PLK1), protein kinase A (PKA), AMP-activated protein kinase (AMPK), casein kinase 2 (Ck2) and ataxia telangiectasia and Rad3 related kinase (ATR), etc. Phosphorylation-mediated regulation of modifiers may directly activate or suppress their enzymatic activity, or indirectly regulate the interaction between modifiers with other proteins or RNAs, or make chromatin structure tight or loose.⁵⁶ And several epigenetic modifiers have been found either aberrantly hyperphosphorylated or hypophosphorylated in cancer cells, including DNA methyltransferases, histone methyltransferases, histone demethylases, histone acetyltransferases and deacetylases.

DNMTs especially DNMT1 have been known to be phosphorylated to regulate the protein stability and enzymatic activity. AKT and PKC kinases were reported to phosphorylate DNMT1 at Ser127/143 and Ser127, respectively, which will disrupt the interactions of DNMT1 with PCNA and UHRF1 in human cells to promote tumorigenesis.^{57,58} In addition, it has been reported that GSK3 β can interact with DNMT1 and then phosphorylate DNMT1 at Ser410 and Ser414, and finally promote β TrCP-induced proteasomal degradation of DNMT1.⁵⁹ There are not many reports showing the phosphorylation of DNMT3s, however, the kinase CK2 tends to phosphorylate DNMT3A, and decrease the global genomic methylation levels.⁶⁰

The histone methyltransferase EZH2 has been shown to be phosphorylated by several kinases, such as AKT, AMPK, CDKs, or Janus kinase 3 (JAK3), in various types of cancer. AKT-mediated phosphorylation of EZH2 at Ser21 results in loss of methylation of H3K27 and increase of expressions of genes used to be silent by H3K27me3. The AKT-mediated phosphorylation of EZH2 promotes expression of several critical oncogenes, and is involved in the development of prostate cancer, uterine cancer and glioblastoma tumorigenesis.^{61–63} AMPK can also phosphorylate EZH2 at Thr311 to disrupt the polycomb repressive complex 2 (PRC2), in which EZH2 is the core component, and thus suppresses methyltransferase activity in both ovarian and breast cancers.⁶⁴ In addition, CDK1/2 have also been reported to phosphorylate EZH2 at Thr350 and Thr487, which will not only inhibit the enzymatic activity, but also block EZH2

binding to its target region, and will highly increase the risk of tumorigenesis.^{65,66} Phosphorylation of EZH2 by JAK3 induces a noncanonical function of EZH2 to promote transcriptional activation in natural killer/T-cell lymphoma.⁶⁷ Collectively, regulation of EZH2 by phosphorylation is highly correlated with tumorigenesis with regard to its activity. We will further discuss about the role of EZH2 and its phosphorylation in prostate cancer and castration-resistant prostate cancer (CRPC) in the subsequent session.

Although contribution to cancer by the phosphorylation of H3K27 histone methyltransferase has been extensively studied, phosphorylation of histone methyltransferases that add methyl groups to other residues of histone tail, such as H3K4, was also reported. For instance, ATR phosphorylates MLL (H3K4 methyltransferase) on Ser516 in response to environmental stress in the S phase, resulting in its degradation and finally contributing to human MLL leukaemia.⁶⁸ Several studies also demonstrated the unusual phosphorylation events occurring on arginine methyltransferases. For example, in myeloproliferative neoplasms, Janus kinase 2 (JAK2) oncogenic mutant V617F can phosphorylate protein arginine methyltransferase 5 (PRMT5), consequently decreasing its activity and increasing expression of genes that are inhibited by PRMT5.⁶⁹

Histone demethylases also appear to be regulated by phosphorylation, although how phosphorylation affects the activities still remains to be elucidated. Protein kinase A (PKA)-induced phosphorylation of H3K9me2 demethylase PHD finger protein 2 (PHF2) at Ser1056 results in its increased binding to a DNA-binding protein ARID5B, reduction of methylation on ARID5B, and decrease of gene transcription.⁷⁰ CDK1 can catalyze phosphorylation of another demethylase, PHD finger protein 8 (PHF8) at Ser33 and Ser84, leading to disruption of PHF8 with chromatin in acute promyelocytic leukemia.⁷¹

Phosphorylation can also regulate the activities of HATs and HDACs, and therefore affect the gene transcription via histone acetylation patterns. The histone acetyltransferase CBP is phosphorylated by CDK2 in a cell cycle-dependent manner. One study demonstrated that the DNA-dependent protein kinase (DNA-PK) phosphorylates hGCN5, which possesses HAT activity, and the phosphorylation suppresses its HAT activity.⁷² In addition, in response to DNA damage, the HAT activity of activating transcription factor 2 (ATF2) is phosphorylated.⁷³ The PI3K/AKT pathway also stimulates p300 phosphorylation at Ser1834 and its transcriptional activator potential.⁷⁴

Like HMTs, HDACs have been commonly believed that their activities are tightly regulated by phosphorylation. The HDAC family members, including HDAC1, HDAC2, HDAC3, HDAC4, HDAC5, HDAC6 and HDAC8, are phosphorylated by many kinases, such as CK2, PKA, extracellular signal regulated kinase (ERK1/2), etc. within different residues, to influence the structure, stability, acetyltransferase activity, binding with partners, or cellular localization, and ultimately leading to either pro- or anti-tumorigenesis.^{75,76} I will take two recent findings as examples. Activated PI3K/AKT pathway in breast cancer cells can lead to the phosphorylation of the p70 S6 kinase (S6K1) and transcriptionally regulate estrogen receptor α (ER α) expression.⁷⁷ Moreover, c-Jun N-terminal kinase (JNK)-

mediated phosphorylation of HDAC3 in triple-negative breast cancer (TNBC) might affect the sensitivity of HDAC inhibitors in treatment.⁷⁸

EZH2 methyltransferase and its phosphorylation in prostate cancer

In 2019, prostate cancer (PCa) has become the most common cancer with 174,650 newly diagnosed cases and the second leading cause of death associated with cancer or cancer-related factors in males in the United States with estimated 31,620 deaths.⁷⁹ Androgen deprivation therapy (ADT) is the routinely used approach to treat PCa patients. Although patients initially respond to ADT well, castration-resistant prostate cancer (CRPC) eventually occurs in most of these patients after several years and then develops into even worse metastasis.⁸⁰ It has been well established that androgen receptor (AR) signaling is enhanced and plays an important role in CRPC.⁸¹ Subsequently, the AR inhibitors, such as enzalutamide and abiraterone, have been approved by the Food and Drug Administration (FDA) for the treatment of late stage PCa.⁸² However, enzalutamide resistance eventually develops for almost all cases, making the disease almost incurable. Therefore, how to overcome enzalutamide resistance of CRPC has been under intensive research, and new targets and mechanism-based strategies are urgently needed to treat these patients.^{83–85}

As we described above, EZH2, the catalytic subunit of PRC2 complex, plays a critical role in repressing gene expression by mediating H3K27me3. Many studies have demonstrated that there is a tight linkage between EZH2 and oncogenesis, and that EZH2-mediated trimethylation triggers silencing of tumor suppressor genes in cancer.⁸⁶ Besides acting as a transcriptional suppressor, emerging evidence has shown the uncanonical role of EZH2 towards transcription activation of some genes, whose expression seems to be PRC2-independent. EZH2 has been identified as either a direct transcription activator or coactivator binding with other transcription factors to promote expression of several oncogenes. For example, the transcription levels of genes in NOTCH pathway, NF- κ B pathway or Wnt pathway are directly or indirectly regulated by EZH2 in breast cancer and colon cancer, respectively, which are independent of EZH2 methyltransferase activity^{87–90} (see Fig. 1).

In PCa especially in CRPC, EZH2 is overexpressed and promotes cancer cell proliferation and invasion, making EZH2 an attractive anti-cancer drug target. Similar with other types of cancer, aberrant PTMs, such as phosphorylation, of EZH2 are also found in PCa (Fig. 2). For example, AKT-mediated phosphorylation of EZH2 at Ser21 induces a functional switch from a PRC2-dependent transcription repressor to a PRC2-independent transcription coactivator working with AR to promote the development of CRPC.^{61,63} In addition to AKT kinase, CDK1/2 can also phosphorylate EZH2 at Thr350 during S and G2/M phases. Phosphorylation of Thr350 promotes PCa cell proliferation and migration. Consequently, blocking Thr350 phosphorylation is important for abrogation of the oncogenic activity of EZH2.^{65,66}

Polo-like kinase 1 (PLK1), a regulator of various stages of mitosis, has been shown to be overexpressed in various




Figure 1 The dynamic epigenetic modifications on DNA and histone tail. Enzymes coordinately regulate the epigenetic modifications by adding or removing epigenetic hallmarks. Deregulation of the enzymes can lead to oncogenesis.




Figure 2 A model of the EZH2 functional switch by its hyper phosphorylation in CRPC. Deregulation of EZH2 phosphorylation can change EZH2 from a transcriptional repressor depending upon PRC2 to a transcriptional co-activator cooperating with AR which is independent of PRC2, finally contributing to CRPC progression.

types of cancers. Our lab has made a series of discoveries to show that Plk1 plays a critical role in different aspects of PCa, including its initiation, progression and therapy resistance.⁹¹ It was shown that PLK1 directly phosphorylates SUZ12, another component of PRC2 complex, and that PLK1 phosphorylation of SUZ12 abolishes its interaction with EZH2 and the PRC2 function.⁵⁵ Whether PLK1 also directly phosphorylates EZH2 is unknown.

It is worth of mentioning that EZH2 has an influence on DNA methylation by direct association with DNMTs.⁹² DNA methylation abnormal changes can be highly present in advanced stages of prostate cancer, and changes are tied together with alterations of EZH2 activity in silencing tumor suppressors. Thus, if not all, modifications, like PTMs, that affect EZH2 activity will also affect DNA methylation. For example, one study has shown that EZH2 directly interacts with DNMTs and contributes to DNA hypermethylation of selected target genes (particularly GSTP1 and RARB2) by qRT-PCR in 47 primary prostate cancers.⁹³ Additionally, EZH2 and DNMT3B have been shown to mediate

hypermethylation of HOXB13 in prostate cancer cells, and both EZH2 and DNMT3B could be repressed by an anticancer agent, all-trans retinoic acid (ATRA) flowing a reduced methylation of HOXB13.⁹⁴

Conclusions

PCa is the most commonly diagnosed malignant neoplasm of males in the United States, and ADT is an effective treatment for patients with PCa. However, most patients ultimately develop resistance and cancer relapse. Treatment for CRPC is very limited. Therefore, exploring novel cellular mechanisms controlling progression of PCa is very critical for identifying new targets and eventually developing efficient strategies to treat CRPC. Expression of EZH2 is often upregulated in CRPC, thus EZH2 has been proposed as a target for CRPC. Importantly, it has been demonstrated that EZH2 becomes hyperphosphorylated in CRPC cells.⁶³ However, very few studies reported the effect of PLK1-dependent phosphorylation on epigenetic modifications.

Our ongoing study is expected to fill in this knowledge gap by determining whether and how PLK1-dependent phosphorylation of various epigenetic regulators contributes to PCa progression and drug resistance.

Financial information

NIH R01 CA157429 (X. Liu), R01 CA192894 (X. Liu), R01 CA196835 (X. Liu), R01 CA196634 (X. Liu). The work was also partially supported by University of Kentucky Cancer Center (P30 CA177558).

Conflict of Interest

The authors declare no conflict of interests.

References

- Feinberg AP. The epigenetics of cancer etiology. *Semin Cancer Biol.* 2004;14(6):427–432.
- Egger G, Liang G, Aparicia A, Jones P. Epigenetics in human disease and prospects for epigenetic therapy. *Nature.* 2004;429:457.
- Biswas S, Rao CM. Epigenetic tools (The Writers, the Readers and the Erasers) and their implications in cancer therapy. *Eur J Pharmacol.* 2018;837:8–24.
- Jones PA, Baylin SB. The fundamental role of epigenetic events in cancer. *Nat Rev Genet.* 2002;3:415.
- Rodriguez J, Frigila J, Vendrell E, et al. Chromosomal instability correlates with genome-wide DNA demethylation in human primary colorectal cancers. 2006;66(17):8462–9468.
- Clark SJ, Melki J. DNA methylation and gene silencing in cancer: which is the guilty party? *Oncogene.* 2002;21:5380.
- Baylin SB. DNA methylation and gene silencing in cancer. *Nat Clin Pract Oncol.* 2005;2:54.
- Ohtani-Fujita N, Fujita T, Aoike A, et al. CpG methylation inactivates the promoter activity of the human retinoblastoma tumor-suppressor gene. *Oncogene.* 1993;8(4):1063–1067.
- Rideout W, Coetzee G, Olumi A, Jones P. 5-Methylcytosine as an endogenous mutagen in the human LDL receptor and p53 genes. *Science.* 1990;249(4974):1288–1290.
- Jin B, Robertson KD. DNA methyltransferases, DNA damage repair, and cancer. *Adv Exp Med Biol.* 2013;754:3–29.
- Esteller M. *Epigenet Cancer.* 2008;358(11):1148–1159.
- Zhao S, Zhu S, Hao X, Li P, Zhang S. Effects of DNA methyltransferase 1 inhibition on esophageal squamous cell carcinoma. *Dis Esophagus.* 2011;24(8):601–610.
- Peng D-F, Kanai Y, Sawada M, et al. DNA methylation of multiple tumor-related genes in association with overexpression of DNA methyltransferase 1 (DNMT1) during multistage carcinogenesis of the pancreas. *Carcinogenesis.* 2006;27(6):1160–1168.
- Zhao Z, Wu Q, Cheng J, et al. Depletion of DNMT3A suppressed cell proliferation and restored PTEN in hepatocellular carcinoma cell. *J Biomed Biotechnol.* 2010, 737535.
- Butcher DT, Rodenhiser DI. Epigenetic inactivation of BRCA1 is associated with aberrant expression of CTCF and DNA methyltransferase (DNMT3B) in some sporadic breast tumours. *Eur J Cancer.* 2007;43(1):210–219.
- Lin H, Yamada Y, Nguyen S, et al. Suppression of intestinal neoplasia by deletion of Dnmt3b. *Mol Cell Biol.* 2006;26(8):2976–2983.
- Kobayashi Y, Absher D, Gulzar Z, et al. DNA methylation profiling reveals novel biomarkers and important roles for DNA methyltransferases in prostate cancer. *Genome Res.* 2011;21(7):1017–1027.
- Baylin SB, Jones PA. A decade of exploring the cancer epigenome - biological and translational implications. *Nat Rev Cancer.* 2011;11(10):726–734.
- Kanai Y, Ushijima S, Nakanishi Y, et al. Mutation of the DNA methyltransferase (DNMT) 1 gene in human colorectal cancers. *Cancer Lett.* 2003;192(1):75–82.
- Neumann M, Heesch S, Schlee C, et al. Whole-exome sequencing in adult ETP-ALL reveals a high rate of DNMT3A mutations. *Blood.* 2013;121(23):4749–4752.
- Rodenhiser D, Mann M. Epigenetics and human disease: translating basic biology into clinical applications. *CMAJ (Can Med Assoc J) : Canadian Medical Association journal = journal de l'Association medicale canadienne.* 2006;174(3):341–348.
- Haney SL, Upchurch GM, Opavska J, et al. Dnmt3a is a haploinsufficient tumor suppressor in CD8+ peripheral T cell lymphoma. *PLoS Genet.* 2016;12(9). e1006334-e1006334.
- Zheng Y, Zhang H, Wang Y, et al. Loss of Dnmt3b accelerates MLL-AF9 leukemia progression. *Leukemia.* 2016;30:2373.
- Kouzarides T. Chromatin modifications and their function. *Cell.* 2007;128(4):693–705.
- Chi P, Allis CD, Wang GG. Covalent histone modifications-miswritten, misinterpreted and mis-erased in human cancers. *Nat Rev Cancer.* 2010;10(7):457–469.
- Jenuwein T, Allis CD. Translating the Histone Code. *Science.* 2001;293(5532):1074–1080.
- Cohen I, Poręba E, Kamieniarz K, Schneider R. Histone modifiers in cancer: friends or foes? *Genes & Cancer.* 2011;2(6):631–647.
- Baxter E, Windloch K, Gannonet F, Lee J. Epigenetic regulation in cancer progression. *Cell Biosci.* 2014;4, 45-45.
- Garraway Levi A, Lander Eric S. Lessons from the cancer genome. *Cell.* 2013;153(1):17–37.
- Schneider R, Di Cerbo V. Cancers with wrong HATs: the impact of acetylation. *Briefings Funct Genomics.* 2013;12(3):231–243.
- Iyer NG, Özdag H, Caldas C. p300/CBP and cancer. *Oncogene.* 2004;23:4225.
- Dell'Aversana C, Lepore I, Altucci L. HDAC modulation and cell death in the clinic. *Exp Cell Res.* 2012;318(11):1229–1244.
- Busslinger M, Tarakhovskiy A. Epigenetic control of immunity. *Cold Spring Harbor Perspect Biol.* 2014;6(6):a019307.
- Qi J. Bromodomain and extraterminal domain inhibitors (BETi) for cancer therapy: chemical modulation of chromatin structure. *Cold Spring Harbor Perspect Biol.* 2014;6(12):a018663.
- Izzo A, Schneider R. Chatting histone modifications in mammals. *Briefings Funct Genomics.* 2010;9(5–6):429–443.
- Valk-Lingbeek ME, Bruggeman SWM, van Lohuizen M. Stem cells and cancer: the polycomb connection. *Cell.* 2004;118(4):409–418.
- Kondo Y, Shen L, Ahmed S, et al. Downregulation of histone H3 lysine 9 methyltransferase G9a induces centrosome disruption and chromosome instability in cancer cells. *PLoS One.* 2008;3(4). e2037.
- Shi YG, Tsukada Y. The discovery of histone demethylases. *Cold Spring Harbor Perspect Biol.* 2013;5(9):a017947.
- Oki M, Aihara H, Ito T. Role of histone phosphorylation in chromatin dynamics and its implications in diseases. *Subcell Biochem.* 2007;41:319–336.
- Metzger E, Yin N, Wissmann M, et al. Phosphorylation of histone H3 at threonine 11 establishes a novel chromatin mark for transcriptional regulation. *Nat Cell Biol.* 2008;10(1):53–60.
- Lin C-P, He L. Noncoding RNAs in cancer development. *Annu Rev Cell Biol.* 2017;1(1):163–184.
- Iorio MV, Croce CM. Causes and consequences of microRNA dysregulation. *Cancer J.* 2012;18(3):215–222.

43. Gibb EA, Brown CJ, Lam WL. The functional role of long non-coding RNA in human carcinomas. *Mol Cancer*. 2011;10, 38–38.
44. Peng Y, Croce CM. The role of MicroRNAs in human cancer. *Signal Transduct Target Ther*. 2016;1(1):15004.
45. Gillies JK, Lorimer IAJ. Regulation of p27Kip1 by miRNA 221/222 in glioblastoma. *Cell Cycle*. 2007;6(16):2005–2009.
46. Lal A, Kim HH, Abdelmohsen K, et al. p16(INK4a) translation suppressed by miR-24. *PLoS One*. 2008;3(3). e1864.
47. Dolezalova D, Mraz M, Barta T, et al. MicroRNAs regulate p21Waf1/Cip1 protein expression and the DNA damage response in human embryonic stem cells. *Stem Cells*. 2012;30(7):1362–1372.
48. Du B, Wang Z, Zhang X, et al. MicroRNA-545 suppresses cell proliferation by targeting cyclin D1 and CDK4 in lung cancer cells. *PLoS One*. 2014;9(2). e88022.
49. Yan H-l, Xue G, Mei Q, et al. Repression of the miR-17-92 cluster by p53 has an important function in hypoxia-induced apoptosis. *EMBO J*. 2009;28(18):2719–2732.
50. Kalluri R, Weinberg RA. The basics of epithelial-mesenchymal transition. *J Clin Invest*. 2009;119(6):1420–1428.
51. Kong W, Yang H, He L, et al. MicroRNA-155 is regulated by the transforming growth factor beta/Smad pathway and contributes to epithelial cell plasticity by targeting RhoA. *Mol Cell Biol*. 2008;28(22):6773–6784.
52. Sanchez Calle A, Kawamura Y, Takeshita F, Ochiya T. Emerging roles of long non-coding RNA in cancer. *Cancer Sci*. 2018;109(7):2093–2100.
53. Gupta RA, Shah N, Wang K, et al. Long non-coding RNA HOTAIR reprograms chromatin state to promote cancer metastasis. *Nature*. 2010;464(7291):1071–1076.
54. Portoso M, Ragazzini R, Brenčić Ž, et al. PRC2 is dispensable for HOTAIR-mediated transcriptional repression. *EMBO J*. 2017;36(8):981–994.
55. Zhang H, Diab A, Fan H, et al. PLK1 and HOTAIR accelerate proteasomal degradation of SUZ12 and ZNF198 during hepatitis B virus-induced liver carcinogenesis. *Cancer Res*. 2015;75(11):2363–2374.
56. DeCarlo D, Hadden MK. Oncoepigenomics: making histone lysine methylation count. *Eur J Med Chem*. 2012;56:179–194.
57. Hervouet E, Lalier L, Debien E, et al. Disruption of Dnmt1/PCNA/UHRF1 interactions promotes tumorigenesis from human and mice glial cells. *PLoS One*. 2010;5(6). e11333.
58. Estève P-O, Chang Y, Samaranyake M, et al. A methylation and phosphorylation switch between an adjacent lysine and serine determines human DNMT1 stability. *Nat Struct Mol Biol*. 2011;18(1):42–48.
59. Li C, Ebert PJR, Li Q-J. T cell receptor (TCR) and transforming growth factor β (TGF- β) signaling converge on DNA (cytosine-5)-methyltransferase to control forkhead box protein 3 (foxp3) locus methylation and inducible regulatory T cell differentiation. *J Biol Chem*. 2013;288(26):19127–19139.
60. Deplus R, Blanchon L, Rajavelu A, et al. Regulation of DNA methylation patterns by CK2-mediated phosphorylation of Dnmt3a. *Cell Rep*. 2014;8(3):743–753.
61. Cha T-L, Zhou B-P, Xia W, et al. Akt-Mediated phosphorylation of EZH2 suppresses methylation of lysine 27 in histone H3. *Science*. 2005;310(5746):306–310.
62. Bredfeldt TG, Greathouse KL, Safe SH, et al. Xenoestrogen-induced regulation of EZH2 and histone methylation via estrogen receptor signaling to PI3K/AKT. *Mol Endocrinol*. 2010;24(5):993–1006.
63. Xu K, Wu ZJ, Groner AC, et al. EZH2 oncogenic activity in castration-resistant prostate cancer cells is polycomb-independent. *Science*. 2012;338(6113):1465–1469.
64. Wan L, Xu K, Wei Y, et al. Phosphorylation of EZH2 by AMPK suppresses PRC2 methyltransferase activity and oncogenic function. *Mol Cell*. 2018;69(2):279–291.
65. Chen S, Bohrer LR, Rai AN, et al. Cyclin-dependent kinases regulate epigenetic gene silencing through phosphorylation of EZH2. *Nat Cell Biol*. 2010;12(11):1108–1114.
66. Wei Y, Chen YH, Li LY, et al. CDK1-dependent phosphorylation of EZH2 suppresses methylation of H3K27 and promotes osteogenic differentiation of human mesenchymal stem cells. *Nat Cell Biol*. 2011;13(1):87–94.
67. Yan J, Li B, Lin B, et al. EZH2 phosphorylation by JAK3 mediates a switch to noncanonical function in natural killer/T-cell lymphoma. *Blood*. 2016;128(7):948–958.
68. Liu H, Takeda S, Kumar R, et al. Phosphorylation of MLL by ATR is required for execution of mammalian S-phase checkpoint. *Nature*. 2010;467(7313):343–346.
69. Liu F, Zhao X, Perna F, et al. JAK2V617F-Mediated phosphorylation of PRMT5 downregulates its methyltransferase activity and promotes myeloproliferation. *Cancer Cell*. 2011;19(2):283–294.
70. Baba A, Ohtake F, Okuno Y, et al. PKA-dependent regulation of the histone lysine demethylase complex PHF2–ARID5B. *Nat Cell Biol*. 2011;13:668.
71. Arteaga MF, Mikesch JH, Qiu J, et al. The histone demethylase PHF8 governs retinoic acid response in acute promyelocytic leukemia. *Cancer Cell*. 2013;23(3):376–389.
72. Barlev NA, Poltoratsky V, Owen-Hughes T, et al. Repression of GCN5 histone acetyltransferase activity via bromodomain-mediated binding and phosphorylation by the Ku–DNA-dependent protein kinase complex. 1998;18(3):1349–1358.
73. Kawasaki H, Schiltz L, Chiu R, et al. ATF-2 has intrinsic histone acetyltransferase activity which is modulated by phosphorylation. *Nature*. 2000;405:195.
74. Huang W-C, Chen C-C. Akt phosphorylation of p300 at ser-1834 is essential for its histone acetyltransferase and transcriptional activity. *Mol cell Biol*. 2005;25(15):6592–6602.
75. Seto E, Yoshida M. Erasers of histone acetylation: the histone deacetylase enzymes. *Cold Spring Harbor Perspect Biol*. 2014;6(4):a018713.
76. Williams KA. *Phosphorylation of histone deacetylase 6 within its c-terminal region by extracellular signal regulated kinase 1*. Dissertation. University of South Florida; 2013.
77. Citro S, Miccolo C, Meloni L, Chiocca S. PI3K/mTOR mediate mitogen-dependent HDAC1 phosphorylation in breast cancer: a novel regulation of estrogen receptor expression. *J Mol Cell Biol*. 2015;7(2):132–142.
78. Hanigan TW, Aboukhatwa SM, Taha TY, et al. Divergent JNK phosphorylation of HDAC3 in triple-negative breast cancer cells determines HDAC inhibitor binding and selectivity. *Cell Chem Biol*. 2017;24(11):1356–1367.
79. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2019. *CA: Cancer J Clin*. 2019;69(1):7–34.
80. Karantanos T, Corn PG, Thompson TC. Prostate cancer progression after androgen deprivation therapy: mechanisms of castrate resistance and novel therapeutic approaches. *Oncogene*. 2013;32(49):5501–5511.
81. Ferraldeschi R, Welti J, Luo J, et al. Targeting the androgen receptor pathway in castration-resistant prostate cancer: progresses and prospects. *Oncogene*. 2015;34:1745–1757.
82. Rodriguez-Vida A, Galazi M, Rudman S, Chowdhury S, Sternberg CN. Enzalutamide for the treatment of metastatic castration-resistant prostate cancer. *Drug Des Dev Ther*. 2015;9:3325–3339.
83. Liu C, Lou W, Zhu Y, et al. Intracrine androgens and AKR1C3 activation confer resistance to enzalutamide in prostate cancer. *Cancer Res*. 2015;75(7):1413–1422.
84. Li Y, Chan SC, Brand LJ, Hwang TH, Silverstein KA, Dehm SM. Androgen receptor splice variants mediate enzalutamide resistance in castration-resistant prostate cancer cell lines. *Cancer Res*. 2013;73(2):483–489.

85. Kong Y, Cheng L, Mao F, et al. Inhibition of cholesterol biosynthesis overcomes enzalutamide resistance in castration-resistant prostate cancer (CRPC). *J Biol Chem*. 2018;293(37):14328–14341.
86. Tan J-z, Yan Y, Wang XX, Jiang Y, Xu HE. EZH2: biology, disease, and structure-based drug discovery. *Acta Pharmacol Sin*. 2014;35(2):161–174.
87. Shi B, Liang J, Yang X, et al. Integration of estrogen and Wnt signaling circuits by the polycomb group protein EZH2 in breast cancer cells. *Mol Cell Biol*. 2007;27(14):5105–5119.
88. Lee ST, Li Z, Wu Z, et al. Context-specific regulation of NF- κ B target gene expression by EZH2 in breast cancers. *Mol Cell*. 2011;43(5):798–810.
89. Gonzalez ME, Moore HM, Li X, et al. EZH2 expands breast stem cells through activation of NOTCH1 signaling. *Proc Natl Acad Sci*. 2014;111(8):3098–3103.
90. Jung H-Y, Jun S, Lee M, et al. PAF and EZH2 induce Wnt/ β -Catenin signaling hyperactivation. *Mol Cell*. 2013;52(2):193–205.
91. Li J, Karki A, Hodges KB, et al. Cotargeting polo-like kinase 1 and the Wnt/ β -Catenin signaling pathway in castration-resistant prostate cancer. *Mol Cell Biol*. 2015;35(24):4185–4198.
92. Fuks F. DNA methylation and histone modifications: teaming up to silence genes. *Curr Opin Genet Dev*. 2005;15(5):490–495.
93. Hoffmann MJ, Engers R, Florl AR, Otte AP, Muller M, Schulz WA. Expression changes in EZH2, but not in BMI-1, SIRT1, DNMT1 or DNMT3B are associated with DNA methylation changes in prostate cancer. *Cancer Biol Ther*. 2007;6(9):1399–1408.
94. Liu Z, Ren G, Shangguan C, et al. ATRA inhibits the proliferation of DU145 prostate cancer cells through reducing the methylation level of HOXB13 gene. *PLoS One*. 2012;7(7). e40943.