



Development, Isolation and Characterisation of a New Non-Virulent Mimosine-Degrading *Klebsiella pneumoniae* Strain from the Rumen Liquor of German Steers by Using IBT-Goettinger Bioreactor

Aung Aung
University of Veterinary Science, Myanmar

U. ter Meulen
Georg-August-University of Goettingen, Germany

F. Gessler
Miprolab, Germany

H. Böhnel
Georg-August-University of Goettingen, Germany

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/22/1-2/23>

The 22nd International Grassland Congress (Revitalising Grasslands to Sustain Our Communities) took place in Sydney, Australia from September 15 through September 19, 2013.

Proceedings Editors: David L. Michalk, Geoffrey D. Millar, Warwick B. Badgery, and Kim M.

Broadfoot

Publisher: New South Wales Department of Primary Industry, Kite St., Orange New South Wales, Australia

Development, isolation and characterisation of a new non-virulent mimosine-degrading *Klebsiella pneumoniae* strain from the rumen liquor of German steers by using IBT-Goettinger bioreactor

A Aung^{AB}, U ter Meulen^C, F Gessler^D and H Böhnel^{BD}

^A University of Veterinary Science, Yezin, Nay Pyi Taw, Myanmar

^B Institute of Applied Biotechnology in the Tropics, Georg-August-University of Goettingen, Germany

^C Institute of Animal Physiology and Nutrition, Georg-August-University of Goettingen, Kellnerweg 6, D-37077 Germany

^D Miprolab, Marie-Curie-Str. 7, 37079 Goettingen, Germany

Contact email: aung.aaung@gmail.com

Keywords: *Leucaena*, *Klebsiella pneumoniae*, mimosine.

Introduction

The tropical legume *Leucaena leucocephala* (leucaena) has many uses, including: a potential source of firewood and timber; for soil erosion control (Dijkman 1950); to provide shade; to enhance soil fertility; and as a nutritious forage for animal feed (Ruskin 1977). It is widely used as forage for cattle in tropical agriculture (Shelton 1998). In Myanmar, leucaena is used as a protein source in urea-molasses multi-nutrient blocks for ruminants (Ni Ni Maw *et al.* 2004). However, the use of leucaena as ruminant feed is not without problems, because it contains mimosine, a toxic anti-nutritional factor limiting its use as animal feed. Jones (1981) reported the absence of toxicity when leucaena was fed to goats and cattle in Hawaii and Indonesia. According to the low dihydroxypyridine (DHP) in urine of those animals, it was assumed that they could degrade mimosine and DHP. Hawaiian goats, but not Australian goats, could degrade 3,4-DHP ruminally (Jones and Megarrity 1983). Inoculation of susceptible animals with rumen liquor containing mimosine-degrading bacteria protected against DHP toxicity in ruminants (Jones and Lowry 1984). For maintaining mimosine-degrading bacteria, the donor animals should be fed on leucaena continuously and it is expensive to maintain their veterinary care. Hence, we tried to develop mimosine-degrading ruminal bacteria using a fermenter, intending to produce a source of inoculum for the routine control of leucaena toxicosis in ruminants.

Materials and Methods

The donor steer was from the Institute of Animal Physiology and Nutrition, Faculty of Agriculture, Georg-August University of Goettingen, and had never consumed leucaena leaves. Rumen liquor was obtained via a rumen cannula and immediately placed in a conical flask, flushed with carbon dioxide, stoppered, maintained under carbon dioxide, and frozen at 4 °C in a refrigerator for 2 weeks to kill the protozoa. Before subsequent fermentation, rumen liquor was thawed. Fermentation was commenced with 520 mL of rumen liquor. The IBT- Goettinger Bioreactor was a modified form of that used by Böhnel and Zeggu (1993). The medium 98-5 for rumen microbes (Allison *et al.* 1990),

without mimosine, was supplied continuously. Infusion of mimosine started at 25 mg/d, increasing by 25 mg of mimosine every second day for 16 days, when 200 mg/d was infused. Daily samples were taken and mimosine degradability of the sample was assessed by colourimetric assay. Purification of the mimosine-degrading bacterial strain was done from the samples showing mimosine degradability. After that, tests for biochemical properties and molecular identification were conducted. For proof that bacterial strains were non-pathogenic, a mouse bioassay was carried out to determine the lethal dose.

Results and Discussion

Although the rumen liquor of the donor steer was unable to degrade mimosine initially, infusing gradually increased amounts of mimosine for 16 days (maximum of 200 mg/d mimosine) enabled the liquor to degrade mimosine (Fig. 1). It is possible that some ruminal bacteria became adapted to degrading mimosine, as found by Ruskin (1977). According to molecular identification, genotypic characteristics of this isolate were almost identical (94% similarity) to those of reference *Klebsiella pneumoniae* from GenBank entries held by the National Centre for Biotechnology Information (NCBI). Although this isolate is similar to pathogenic *K. pneumoniae*, it has some different




Figure 1. Effects of treatment with mimosine on the ability of rumen liquor to degrade mimosine. Sample 1 was taken on day 15 of treatment with mimosine, when dosage was 200 g/d; sample 2 was taken on day 16 of treatment.

biochemical properties (Biolog test). In this experiment, gradually increased amounts of mimosine infused into rumen liquor induced adaptation in the bacteria first, so they could degrade mimosine. Mice inoculated with recommended dose of *K. pneumoniae* (3.3×10^8 CFU/mL) showed no clinical signs of klebsiellosis, as reported by Thakker *et al.* (2006).

Conclusion

This study has shown that treatment of rumen liquor with mimosine increases the ability of the bacteria to degrade mimosine. *Klebsiella pneumoniae* isolated from this experiment was effective at degrading mimosine. Studies are needed to compare the effectiveness of this bacterium in protecting animals fed leucaena, in comparison with *Synergistes jonesii*. An experiment to measure these parameters will be conducted in Myanmar.

Acknowledgments

The laboratory work and technical support for this study were provided by the Institute of Applied Biotechnology in the Tropics, Georg-August-University of Goettingen, Germany. A. Aung thanks the German Academic Exchange Service (DAAD) for the support of doctoral scholarship in the IPAG programme. We appreciate support from Prof. Dr T. Wolfe for reviewing the manuscript.

References

- Allison MJ, Hammond AC, Jones RJ (1990). Detection of ruminal bacteria that degrade toxic dihydroxypyridine Compounds produced from mimosine. *Applied and Environmental Microbiology* **56**, 590-594.
- Böhnelt H, Zeggu T (1993). Modern adapted biotechnology for the production of rhizobium inoculant for soy beans. *Plant Research and Development* **38**, 81-87.
- Dijkmann DJ (1950). Leucaena – A promising soil-erosion control plant. *Economic Botany* **4**, 337-349.
- Jones RJ (1981). Does ruminal metabolism of mimosine explain the absence of leucaena toxicity in Hawaii? *Australian Veterinary Journal* **57**, 55-56.
- Jones RJ, Megarrity RG (1983). Comparative toxicity responses of goats fed on *Leucaena leucocephala* in Australia and Hawaii. *Australian Journal of Agricultural Research* **34**, 781-790.
- Jones RJ, Lowry JB (1984). Australian goats detoxify the goitrogen 3-hydroxy,4-(1H) pyridone after rumen infusion from an Indonesian goat. *Experientia* **40**, 1435-1436
- Ni Ni Maw, Khin San Mu, Aung Aung, Moe Thida Htun (2004). Proceedings of the annual research conference held in Yangon on June 28-30, 2004 (Livestock and Fishery Science. Myanmar Academy of Agriculture, Forestry, Livestock and Fishery Sciences). pp. 163-175.
- Ruskin FR (1977). Leucaena: Promising forage and tree crop for the tropics. 1st Edn National Academy Press and National Research Council, Washington D.C.
- Shelton HM (1998). The leucaena genus. New opportunities for agriculture (A review of workshop outcomes). In 'Leucaena adaptation, quality and farming systems'. (Eds HM Shelton, RC Gutteridge, BF Mullen, RA Bray) pp. 15-24. (Australian Centre for International Agricultural Research: Canberra)
- Thakker C, Burhanpurwala Z, Rastogi G, Shouche Y, Ranade D (2006). Isolation and characterization of a new osmotolerant, non-virulent *Klebsiella pneumoniae* strain SAP for biosynthesis of succinic acid. *Indian Journal of Experimental Biology* **44**, 142-150.