

University of Kentucky

UKnowledge

Pharmaceutical Sciences Faculty Patents

Pharmaceutical Sciences

5-26-2009

Genes Encoding Several Poly (ADP-Ribose) Glycohydrolase (PARG) Enzymes, the Proteins and Fragments Thereof, and Antibodies Immunoreactive Therewith

Myron Jacobson
University of Kentucky

Elaine L. Jacobson
University of Kentucky

Jean-Christoph Amé

Winston Lin
University of Kentucky, winston.lin@uky.edu

Follow this and additional works at: https://uknowledge.uky.edu/ps_patents



Part of the [Pharmacy and Pharmaceutical Sciences Commons](#)

Right click to open a feedback form in a new tab to let us know how this document benefits you.

Recommended Citation

Jacobson, Myron; Jacobson, Elaine L.; Amé, Jean-Christoph; and Lin, Winston, "Genes Encoding Several Poly (ADP-Ribose) Glycohydrolase (PARG) Enzymes, the Proteins and Fragments Thereof, and Antibodies Immunoreactive Therewith" (2009). *Pharmaceutical Sciences Faculty Patents*. 119.
https://uknowledge.uky.edu/ps_patents/119

This Patent is brought to you for free and open access by the Pharmaceutical Sciences at UKnowledge. It has been accepted for inclusion in Pharmaceutical Sciences Faculty Patents by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.



US007538194B2

(12) **United States Patent**
Jacobson et al.

(10) **Patent No.:** **US 7,538,194 B2**
(45) **Date of Patent:** **May 26, 2009**

(54) **GENES ENCODING SEVERAL POLY (ADP-RIBOSE) GLYCOHYDROLASE (PARG) ENZYMES, THE PROTEINS AND FRAGMENTS THEREOF, AND ANTIBODIES IMMUNOREACTIVE THEREWITH**

(75) Inventors: **Myron K. Jacobson**, Lexington, KY (US); **Elaine L. Jacobson**, Lexington, KY (US); **Jean-Christophe Amé**, Obernai (FR); **Winston Lin**, Lexington, KY (US)

(73) Assignee: **University of Kentucky Research Foundation**, Lexington, KY (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 189 days.

(21) Appl. No.: **11/638,012**

(22) Filed: **Dec. 13, 2006**

(65) **Prior Publication Data**

US 2007/0136833 A1 Jun. 14, 2007

Related U.S. Application Data

(62) Division of application No. 11/058,995, filed on Feb. 15, 2005, now abandoned, which is a division of application No. 09/973,451, filed on Oct. 9, 2001, now Pat. No. 6,906,180, which is a division of application No. 09/302,812, filed on Apr. 30, 1999, now Pat. No. 6,333,148.

(60) Provisional application No. 60/083,768, filed on May 1, 1998.

(51) **Int. Cl.**
C12Q 1/68 (2006.01)

(52) **U.S. Cl.** **530/387.1**; 435/4; 435/6; 435/18; 435/183

(58) **Field of Classification Search** 530/387.1; 435/4, 6, 18, 183
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

6,333,148 B1 * 12/2001 Jacobson et al. 435/4

OTHER PUBLICATIONS

Tavossoli et al. (Biochimica et Biophysica Acta 1985; 827: 228-234).*

* cited by examiner

Primary Examiner—Brandon J Fetterolf

(74) *Attorney, Agent, or Firm*—Fulbright & Jaworski, LLP

(57) **ABSTRACT**

The isolation and characterization of cDNAs encoding poly (ADP-ribose) glycohydrolase (PARG) enzymes and the amino acid sequences of PARGs from several species are described. PARG is involved in the cellular response to DNA damage and its proper function is associated with the body's response to neoplastic disorder inducing agents and oxidative stress. Expression vectors containing the cDNAs and cells transformed with the vectors are described. Probes and primers that hybridize with the cDNAs are described. Expression of the cDNA in *E. coli* results in an enzymatically active protein of about 111 kDa and an active fragment of about 59 kDa. Methods for inhibiting PARG expression or overexpressing PARG in a subject for therapeutic benefit are described. Exemplary of PARG inhibitors are anti-sense oligonucleotides. The invention has implications for treatment of neoplastic disorder, heart attack, stroke, and neurodegenerative diseases. Methods for detecting a mutant PARG allele are also described. Antibodies immunoreactive with PARGs and fragments thereof are described.

8 Claims, 24 Drawing Sheets

FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5

bPARG	(422)	ED...	KRKEQCEMKHQ RTE..	RKIPKYI PPH	SEQ ID NO: 19
hPARG	(421)	ED...	RRKEQWETKHQ RTE..	RKIPKYV PPH	SEQ ID NO: 20
mPARG	(413)	ED...	RRKEQCEVRHQ RTE..	RKIPKYI PPN	SEQ ID NO: 21
CePARG	(29)	HQVPTMKRRRLTEHGNTTESLLKEDPEEPKS			SEQ ID NO: 22
hPARP	(205)	EG...	KRKG D. EVDG. VDEVAKKKSKEKDK		SEQ ID NO: 23
mPARP	(205)	EG...	KRKG D. EVDG. TDEVAKKKSKEKTDK		SEQ ID NO: 24
bPARP	(208)	EG...	KRKG D. EVDG. IDEVTKKKSKEKDK		SEQ ID NO: 25
aPARP	(205)	EG...	KRKG E. EVDG..	NVVAKKKSKEKEK	SEQ ID NO: 26
XIPARP	(204)	EG...	KRKAD. EVDG. HSAATKKIKKEKEK		SEQ ID NO: 27
DmPARP	(202)	EELPDTKRAKM. ELSDTNEEGEKKQR.....			SEQ ID NO: 28
SPARP	(205)	EGVSSAKKAKI. EKIDEEDAASIKELTEKIKK			SEQ ID NO: 29

FIGURE 6




FIGURE 7




FIGURE 8




FIGURE 9




FIGURE 10




FIGURE 11




FIGURE 12




FIGURE 13




FIGURE 14

Strategy to Obtain Homologous PARG Sequences





FIGURE 15

Domain Organization of PARGs



SEQ ID NO:

2 Bparg 1 MSAGPGCEPCTKRPRWDAATA SPPAASDARSF PGRQRRVLD SKDAPVQFRVPPSSSGCALGRAGQERGSATSL
4 Eparg 1 MNAGPGCEPCTKATRGAATTS . PAASDARSF PGRQRRVLD PKDAEVQFRVPPSSPACVPGQAGQERGSATSL
6 Mparg 1 MSAGPGWEPCTKA . RWGAAGT SAPIASDARSF PGRQRRVLD PKDAPVQFRVPPSSPACVSGRAGFERGSATHF
8 Dparg 1 -----
10 Ceparg1 -----

2 Bparg 74 VFKQKTI TSWMDTKG IKTVESKLS EXEHHHTREESMMSSVQKDFYQHHEKLEHVSQ L GFDKSPVEKGTQY
4 Eparg 73 VFKQKTI TSWMDTKG IKTVESKLS EXEHHHTREESMMSSVQKDFYQHHEKLVHVSQ L SLDKSLTEKSTQY
6 Mparg 73 VFKQKTI TSWMDTKG IKTVESK . . . SXEHHHTREESMMSSVQKDFYPHKVEKLEHVPQ L NLDKSPTEKSSQY
8 Dparg 1 -----
10 Ceparg1 -----

2 Bparg 218 LKQHQTAA MCKWQHGHGPHSERLLESEPPAVTLVPEQF SHAHVDS SPKDDHSDTNS EESRDNQFLTHVKL . .
4 Eparg 217 LKQHQTAA MCKWQHGHGKHTEQ LLESEPTVTLVPEQF SHAHIDRSS PQNDHSDTDS EHRDNQFLTTVKL . .
6 Mparg 214 LKQQQTASVCKWQHGHGKHA BQLLASEPPAGTLPKQLSHAHI GQSPETDDHSDTDHEEDRDNQFLTPIKL . .
8 Dparg 1 -----
10 Ceparg1 ----- MSKFIELGDPVTQDEKDYEDYVGVGFAEQVP

2 Bparg 298 . ANAKQTMED EQREARSHQKCGKACHPAEACAGCQEQEFTDVVSEPLSDTGS EDVGTGLKHAHLHRQESSL
4 Eparg 297 . ANAKQTMED EHAAREAKSHQKCSKCHPGEDCASCQODELDVVPKSP LSDVGS EDVGTGSKHDHKLIRQESSCL
6 Mparg 291 . ANTKPTVGDGQ . . ARSHCKSGSRQSKEDCTGCQEQEEVDVLPESPLSDVGAEDI GTGPKHDHKLITGQESSL
8 Dparg 1 -----
10 Ceparg 33 TMKRRKL TEEGNTTESKEDPEEPKSRDVFVSSQSSDESQEDSAENPEIAKEVSENCENLTETLKI SNIESLDN

2 Bparg 396 GHSPPFEKESEPEPMDVDNSKHSCQDSEADEETSPGFDE . QEDSSSAQTANKP SRFPQREADTELKRKRSSAK
4 Eparg 395 GHSPPFEKESEPEPMDVDNSKHSCQDSEADEETSPGFDE . QEDGSSSQTANKP SRFQARDADIEFRKRYSTK
6 Mparg 387 GDSPPFEKESEPEPMDVDNSKHSCQDSEADEETSPVFDE . QEDRSS . QTANKLSSCQAREADGDLRKRKYLTK
8 Dparg 1 MQEFRSHLI.FPIFQ . KVIQS . TANRRR . ASASVL TNRLGK ALCLNCARMSKSPDGGISEIE . . .
10 Ceparg106 VTERSEHTLDN HKSTEPHEE . DVNNKSHIDVAINSEDEDELVL EENHKEWRDGEQVQQLS . . QDLFADD

FIG. 16A

SEQ ID NO:

2 Bparg 468 GGEIRLHFQFEGGESRAGMN.DVNAKRPGSTSSLNVECRNSKQEGKDKSKITDEFMRVPKAEDKRKEQCEMKH
4 Eparg 467 GGEVRLHFQFEGGESRTGMN.DLNAKLPGNISLNVECRNSKQEGKDKSKITDELMLRPLKAEDRKRKEQWETKH
6 Mparg 354 GGEVRLHFQFE.GEMNAGTS.DLNAKPSGNSLNVECRSSKQEGKDKSKITDEFMRISKSEDRRKEQCEVRH
8 Dparg 59 .TEEE.PENLANSL..DDSWRGVSMEA.IHRNRQPFLEENLPPVTAGNLHRVMIQLPIRET..PFR.PIKSPG
10 Ceparg 172 QELIEIPGIMKDTTQTLDITDSEVETAQKMEMIETEADSTFVGEDSKATKIVRTSSSSF...LSFTVSTCEAP

2 Bparg 540 QRTERKIPKYPPHLSPDKKWLGTPIEEM..RRMPCGIRLPLRPSANHTVTRVDLLLRIGEVPK.PFPPTHF
4 Eparg 539 QRTERKIPKYVPPPELSPDKKWLGTPIEEM..RRMPCGIRLPLRPSANHTVTRVDLLLRAGEVPK.PFPPTHY
6 Mparg 425 QRTERKIPKYI PPMLPPEKKWLGTPIEEM..RKMPCGIRLPSLRPSASHTVTRVDLLLRAGEVPK.PFPPTHY
8 Dparg 125 X..WDSEEVRLPCAPESKYPRENPDGS.TTIDFRWEMIERALLQPIKTCHEELQAAIISIHTTIRDQWEFRALE
10 Ceparg 242 AKGRARMIQKELEKEVIAFTEGNLT....LQPDLNKVDPDRNIRI.CTIPNFPASQG...KLRED.NRIG...

2 Bparg 610 KDLWDNKEVXMPCEQNLYPVEDENGE.RAAGSRWELIQTALLHRLTRPQNLKDAIILKINVAISKKWDFDTALI
4 Eparg 609 KDLWDNKEVXMPCEQNLYPVEDENGE.RTAGSRWELIQTALLHKFTRPQNLKDAIILKINVAISKKWDFDTALI
6 Mparg 495 KDLWDNKEVXMPCEQNLYPVEDENGE.RTAGSRWELIQTALLHKFTRPQNLKDAIILKINVAISKKWDFDTALV
8 Dparg 196 QLLDEELDESETRVFFQSILPDMVKIALCLPNICTQPIPLLKQKMNHSITMSQEQIASLLANAFFCTFPRRRA
10 Ceparg 367 PKIVLPQRWRE.FDSRGR...RDSYFYFKRKLGDGILKCIKTTGYFMFVGLLH.....HMWE..FDPDITIK

2 Bparg 682 DFWDXLVEEAEOHLYQSILPDMVKIALCLPNICTQPIPLLKQKMNHSITMSQEQIASLLANAFFCTFPRRRA
4 Eparg 681 DFWDXLVEEAEOHLYQSILPDMVKIALCLPNICTQPIPLLKQKMNHSITMSQEQIASLLANAFFCTFPRRRA
6 Mparg 567 DFWDXLVEEAEOHLYQSILPDMVKIALCLPNICTQPIPLLKQKMNHSITMSQEQIASLLANAFFCTFPRRRA
8 Dparg 269 .KMKSEISSYPDIIHFNRLEFGRSSRKPELKTILFCIFRRV..TERDASNVPVTVVRRSGLPEE.L.IDWS
10 Ceparg 424 ...LPAL.EMYKEMSELVGREEVLEKFAVARIAKTAEDILPERIYRLVGDVE.SATLSHKQCAALVARMF

2 Bparg 755 .KMKSEISSYPDIIHFNRLEFGRSSRKPELKTILFCIFRRV..TEKK...PTGLVTFIRQS.L.ED.F.PEWE
4 Eparg 754 .KMKSEISSYPDIIHFNRLEFGRSSRKPELKTILFCIFRRV..TEKK...PTGLVTFIRQS.L.ED.F.PEWE
6 Mparg 387 .KMKSEISSYPDIIHFNRLEFGRSSRKPELKTILFCIFRRV..TEKK...PTGLVTFIRQS.L.ED.F.PEWE
8 Dparg 337 QSAAPLGDVPLEVDAEGTIEDEGIGLLQVDFAHKYLGGVLGHCVCQVEIRFVICPELLVGLKLFTECLRPFEA
10 Ceparg 491 FA.....RPDSPFS.....FCRILSSDKSICVEKLFLEFTIF.....DKMSMDPPDGAVSF.RLTKMD

FIG. 16B

SEQ ID NO:	2	RCEKLL..TRLEVIYEGTIEGNGQGMLQVDFAHRFVGGVTSAGLVQEEIRFLIHPELIVSRLFIEVLDHNEC
	4	RCEKPL..TRLEVIYEGTIEGNGQMLQVDFAHRFVGGVTSAGLVQEEIRFLIHPELII SRLFIEVLDHNEC
	6	RCEKPL..TRLEVIYEGTIEGNGRGLQVDFAHRFVGGVTSAGLVQEEIRFLIHPELIVSRLFIEVLDHNEC
	8	LVMLGAERYSNYTGIAGSFEWS.....GNFEDSTP.RDSSCRRQTAI VALDALHFA.QSHH...QYREIDL
	10	KDIFNEEW.XDXXLRSLPEVEFFDEMLIEDTAL.CTQVDFAHHEHLGGVLNHGSVQEEIRFLMCPMMVMGMLL
	2	LIITGTEQISEYTGIAETVRWA.....RSHEDRSE.RDDWQRRTEI VAIDALHFR.RYLD...QFVPEKIR
	4	LIITGTEQISEYTGIAETVRWS.....RSHEDRSE.RDDCERRCTEIVAIDALHFR.RYLD...QFVPEKMR
	6	LIITGTEQISEYTGIAETVRWA.....RSHEDRSE.KDDWQRRCTEIVAIDALHFR.RYLD...QFVPEKVR
	8	MERELHKAYIGFVHMMVTPP..PGVATGNWCCGAFGGDSYLKALLQLMVCAQLGRPLAYYTFGNVEFRDDF
	10	CEKMKQLEAISIVGAYVFSYTYGYGHTLKWAELOPNHSRQNTNEFRDRFRGRKLVETIAIDAILFKGSKLDCQT
	2	RELHKAYCGFLRPGVSSEHL SAVATGNWCCGAFGGDARLKALIQYLA AAVAERDVVIFTFGDSELMRDIYSMH
	4	RELHKAYCGFLRPGVSSEHL SAVATGNWCCGAFGGDARLKALIQYLA AAVAERDVVIFTFGDSELMRDIYSMH
	6	RELHKAYCGFLRPGVSEHL SAVATGNWCCGAFGGDARLKALIQYLA AAVAERDVVIFTFGDSELMRDIYSMH
	8	HEMWLLFRNDGTTVQQ.LWS.I LRSYSRLIKEKSSKEPRENKASKKKLYDFI...KEELKVRDVPGE GAS
	10	EQLNKANIIREMKKASIGFMSQGPFTHIP.IVTGWGCGAFNGDKPLKFI IQVIAAGVADRPLHFC SFGEPE
	2	TFLTERKLTVGE.VYKLLRLRYNEECRCSTPGP.....DIKLYPFIYHAVESCTQTTNQPQRTGA-----
	4	IFLTERKLTVGD.VYKLLRLRYNEECRCSTPGP.....DIKLYPFIYHAVESCAETADHSGQRTGT-----
	6	TFLTERKLDVGK.VYKLLRLRYNEECRCSTPGP.....DIKLYPFIYHAVESSAETDMPGQKAGT-----
	8	AEAGSSRVAGLGEKSETS AKSSPELNKQPARPQITITQSTDLLPAQLSQDNSNSSEDDQALLMLSDDEEA
	10	LAAACKKIIERMKQDVTLGKSCFSIFS-----
	2	Bparg 1087 -----
	4	Eparg 1086 -----
	6	Mparg 969 -----
	8	Dparg 684 NAMMEAAASLEAKSSVEISNSTTSKTSSTATKSMGGGRQLSLEMLDTHYEKGSASKRPRKSPNC SKAEG
	10	Ceparg 787 -----

FIG. 16C

SEQ ID NO:		
2	Bparg 1	-----
4	Eparg 1	-----
6	Mparg 1	-----
8	Dparg 410	TDKDEKDDIVD
10	Ceparg 543	-----
2	Bparg 101	-----
4	Eparg 100	-----
6	Mparg 97	-----
8	Dparg 472	-----
10	Ceparg 614	-----
2	Bparg 201	-----
4	Eparg 200	-----
6	Mparg 197	-----
8	Dparg 543	-----
10	Ceparg 687	-----

FIG. 16D

FIGURE 17




FIGURE 18




FIGURE 19




FIGURE 20





FIGURE 21



1

**GENES ENCODING SEVERAL POLY
(ADP-RIBOSE) GLYCOHYDROLASE (PARG)
ENZYMES, THE PROTEINS AND
FRAGMENTS THEREOF, AND ANTIBODIES
IMMUNOREACTIVE THEREWITH**

**CROSS REFERENCE TO RELATED
APPLICATION**

This application is a divisional of application Ser. No. 11/058,995 filed on Feb. 15, 2005, now abandoned which is a divisional of application Ser. No. 09/973,451 filed Oct. 9, 2001, now U.S. Pat. No. 6,906,180, which is a divisional of application Ser. No. 09/302,812, filed on Apr. 30, 1999, now U.S. Pat. No. 6,333,148, which claims the benefits of U.S. Provisional Application No. 60/083,768, filed May 1, 1998. The entire disclosure of U.S. Provisional Application Nos. 60/083,768 and 09/302,812 are incorporated herein by reference.

**STATEMENT REGARDING FEDERALLY
SPONSORED RESEARCH OR DEVELOPMENT**

The present invention was supported in part by the National Institutes of Health (Grant CA43894). The United States Government may have certain rights in the invention.

TECHNICAL FIELD

The present invention relates to poly(ADP-ribose)glycohydrolases (PARGs) and peptides having poly(ADP-ribose) glycohydrolase activity. In addition, the invention also relates to antibodies, including monoclonal antibodies and antibody fragments, that have specific interaction with epitopes present on poly(ADP-ribose) glycohydrolases. Methods of treatment and diagnosis using the poly(ADP-ribose) glycohydrolases, and antibodies specific for poly(ADP-ribose) glycohydrolases are disclosed. The present invention has implications for the treatment of neoplastic disorder, reperfusion following ischemia, neurological disorders, and related conditions.

BACKGROUND OF THE INVENTION

Genomic damage, if left unrepaired, can lead to malignant transformation, or cell death by senescence (aging), necrosis or apoptosis. Among the variables that can affect the ultimate biological consequence of DNA damage to a particular cell are (i) the amount, type, and location of the DNA damage and (ii) the efficiency and bioavailability of the cellular DNA repair mechanism.

The activation of poly(ADP-ribose) polymerase (PARP) by DNA strand breaks is often one of the first cellular responses to DNA damage. PAR catalyzes the conversion of nicotinamide adenine dinucleotide (NAD) to multi-branched polymers containing up to 200 ADP-ribose residues. Increases in polymer levels of more than 100-fold may occur within minutes of DNA damage. Once synthesized, polymers are rapidly turned over, being converted to free ADP-ribose by the action of poly(ADP-ribose) glycohydrolase (PARG) (1). An ADP-ribosyl protein lyase has been proposed to catalyze removal of protein-proximal ADP-ribose monomers (2). FIG. 1 illustrates these processes schematically.

The process of activating PARP upon DNA damage can rapidly lead to energy depletion because each ADP-ribose unit transferred by PARP consumes one molecule of NAD, which in turn, requires six molecules of ATP to regenerate NAD. Additionally, NAD is a key carrier of electrons needed

2

to generate ATP via electron transport and oxidative phosphorylation or by glycolysis. The overactivation of PARP due to substantial DNA damage can significantly deplete the cellular pools of NAD and ATP (3). ADP-ribose polymer metabolism, and thus PARP and PARG have been linked to the enhancement of DNA repair (4), limitation of malignant transformation (5), enhancement of necrotic cell death (6), and involvement in programmed cell death (7). To date, studies of the structure and function of the enzymes of ADP-ribose polymer metabolism have been mainly limited to PARP (8). Little is known about the function and regulation of PARG.

BRIEF SUMMARY OF THE INVENTION

As embodied and broadly described herein, the present invention is directed to nucleic acids molecules, peptides, methods, vectors and antibodies that are related to the poly(ADP-ribose) glycohydrolase (PARG) enzyme.

One embodiment of the invention is directed to an isolated and purified nucleic acid molecule or nucleic acid molecule analog comprising a sequence that encodes a polypeptide having poly(ADP-ribose) glycohydrolase (PARG) activity. The nucleic acid molecule may encode the complete full-length PARG gene or a fragment of the PARG gene. The nucleic acid molecule may be DNA, RNA or peptide nucleic acid (PNA). The nucleic acid molecule can be linear, such as, for example, an isolated fragment or a linear phage DNA. In addition, the isolated nucleic acid molecule may be circular, such as for example in a plasmid. The nucleic acid molecule may also be a single stranded DNA or RNA such as the single stranded DNA or RNA in a single stranded DNA virus or single stranded RNA virus. The nucleic acid molecule may be of yeast, insect or mammalian origin.

The nucleic acid molecule of the invention, may be of mammalian origin, such as, for example of bovine or murine origin. In a preferred embodiment of the invention, the nucleic acid molecule may be of human origin. While the sequence of the nucleic acid molecule is of mammalian origin, the nucleic acid molecule may be replicated in another organism such as an insert in a viral genome, a plasmid in a bacterium or a 2-micron plasmid in a yeast.

Preferably, the nucleic acid molecule has, a high degree of sequence similarity with a sequence shown in SEQ ID NO: 1 (Genbank Accession Number U78975), SEQ ID NO: 3 (Genbank Accession Number AF005043), SEQ ID NO: 5 (Genbank Accession Number AF079557), SEQ ID NO: 7 (Genbank Accession Number AF079556) or SEQ ID NO: 9 (Genbank Accession Number CEF20C5). The high degree of sequence similarity may be, for example, about 70%, preferably about 80%, even more preferably about 90% and most preferably substantially identical such as for example about 100% identity.

The nucleic acid molecule that encodes a polypeptide having poly(ADP-ribose)glycohydrolase (PARG) activity may be single or double stranded nucleic acid molecule of any length such as, for example, about 20 bases in length, about 30 bases in length, about 40 bases in length, about 50 bases in length, about 100 bases in length, about 200 bases in length, about 500 bases in length, about 1000 bases in length, about 1500 bases in length, about 2000 bases in length, about 3000 bases in length. It is understood that "bases" in this patent application means "basepairs" when referring to double stranded nucleic acid molecules and bases when referring to single stranded nucleic acid molecules. In a preferred embodiment of the invention, the nucleic acid molecule may be at least about 1000 base or basepairs long and have at least

about 80% sequence similarity with a sequence shown in SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7 or SEQ ID NO: 9.

In one embodiment of the invention, the nucleic acid molecule may have sequence similarity to one region-of the PARG sequence. The region may be, for example, from about base residue 2113 to about residue 3105 of SEQ ID NO: 3. Alternatively, the region may be, from residue 1240 to about residue 3105 of SEQ ID NO: 3 or from residue 175 to about residue 3105 of SEQ ID NO: 3.

Another embodiment of the invention is directed to the expression and overexpression of PARG in a cell. Expression vectors may mediate the expression of a polypeptide with poly(ADP-ribose) glycohydrolase (PARG) enzyme activity. Expression systems and expression vectors are known in the art. For example, one expression vector may comprise a regulatory sequence which is operatively linked to a nucleotide sequence at least about 1000 base pairs in length, which has at least 70% sequence similarity with a sequence shown in SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7 or SEQ ID NO: 9. In a preferred embodiment, the sequence similarity is at least about 80% identity, more preferably at least about 90% identity and most preferably about 100% identity. The expression vector may be any expression vector that is capable of directing expression of a gene in a host cell including, prokaryotic, eukaryotic, mammalian and viral vector. Examples of such vectors include pCMV-Script cytomegalovirus expression vectors for expression in mammalian cells, pESP and pESC vectors for expression in *S. pombe* and *S. cerevisiae*, pET vectors for expression in bacteria, pSPUTK vectors for high-level transient expression, and pPbac and pMbac vectors for expression in fall army worm (SF9) cells. Such vectors are available commercially from suppliers such as, for example, Invitrogen (Carlsbad, Calif.) or Stratagene (La Jolla, Calif.). In the use of viral vectors, it is understood that defective viral vectors—vectors that are genetically engineered to deliver a gene or gene product to a host but which cannot replicate in a host is preferred. Procedures for the practice of *in vitro* and *in vivo* expression are well known to those of skill in the art and are further available with the specific expression products and cell lines from commercial suppliers.

Another embodiment of the invention is directed to a host cell transformed with a vector containing a nucleic acid molecule with a sequence that encodes a polypeptide having poly(ADP-ribose) glycohydrolase (PARG) activity. The host cell may be any eukaryotic or prokaryotic cell such as, for example a human, murine, rattus, bovine, insect, yeast or bacteria. Specific cell lines are well known to those of skill in the art and are available from suppliers such as the American Tissue Type Collection (ATCC, Manassas, Va.) and Stratagene (La Jolla, Calif.) and the like. A preferred embodiment of the invention is directed to cells transformed with the PARG expression vector which shows an elevated level of PARG relative to non-transformed cells. Especially preferred are cells transformed with an inducible PARG expression vector that have normal or slightly elevated PARG levels before induction and have significantly elevated PARG levels after induction.

An embodiment of the invention is directed to an isolated protein having poly(ADP-ribose) glycohydrolase (PARG) activity. The protein may comprise an amino acid sequence with at least 70% sequence similarity with a sequence shown in SEQ ID NO: 2 (Genbank Accession Number U78975), SEQ ID NO: 4 (Genbank Accession Number AF005043), SEQ ID NO: 6 (Genbank Accession Number AF079557), SEQ ID NO: 8 (Genbank Accession Number AF079556), or

SEQ ID NO: 10 (Genbank Accession Number CEF20C5). The sequence similarity is preferably at least about 80%, more preferably at least about 90% and most preferably substantially identical with a sequence shown in SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, or SEQ ID NO: 10. In a preferred embodiment of the invention, the preferred isolated protein having poly(ADP-ribose) glycohydrolase (PARG) activity and has a molecular weight greater than about 100 kDa.

Another embodiment of the invention is directed to an oligonucleotide which is greater than about 10 bases in length and less than about 1000 bases in length which is complementary to a sequence shown SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, or SEQ ID NO: 9. The oligonucleotide may be, for example, greater than about 20 bases in length, greater than about 30 bases in length, greater than about 40 bases in length, greater than about 50 bases in length, greater than about 100 bases in length, greater than about 200 bases in length or greater than about 300 bases in length. The oligonucleotide, which may be optionally labeled with a detectable marker, may be DNA, RNA or PNA. A detectable marker may be, for example, a radioactive isotope such as ³²P or ¹²⁵I, an epitope such as FLAG.

One preferred oligonucleotide is an antisense oligonucleotide directed to the mRNA of PARG. Antisense oligonucleotide as a method of suppression is well known to those in the art. For example, the phosphorothioate oligonucleotide, ISIS 2922, has been shown to be effective against cytomegalovirus retinitis in AIDS patients (9). It is thus well known that oligonucleotides, when administered to animals and humans, can have a useful therapeutic effect. In a preferred embodiment, the oligonucleotide is at least about 10 nucleotides in length, such as, greater than about 20 bases in length, greater than about 30 bases in length, greater than about 40 bases in length, greater than about 50 bases in length, greater than about 100 bases in length, greater than about 200 bases in length or greater than about 300 bases in length. In another preferred embodiment, the oligonucleotide has a ribozyme activity.

Another embodiment of the invention is directed to an isolated polypeptide of at least 6 amino acid residues in length and having a molecular weight less than about 65 kDa, which has at least about 80% sequence similarity with a sequence shown in any one of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8 or SEQ ID NO: 10. The polypeptide may be, for example, at least about 10 amino acids in length, at least about 20 amino acids in length, at least about 30 amino acids in length, at least about 40 amino acids in length, at least about 50 amino acids in length, at least about 75 amino acids in length, at least about 100 amino acids in length, at least about 150 amino acids in length, at least about 250 amino acids in length or at least about 500 amino acids in length or more.

In a preferred embodiment, the polypeptide has a molecular weight less than about 40 kDa and has at least about 90% sequence similarity with a sequence shown in any one of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8 or SEQ ID NO: 10. The polypeptide preferably has poly(ADP-ribose) glycohydrolase (PARG) activity or is immunogenic and elicits antibodies immunoreactive with a poly(ADP-ribose) glycohydrolase (PARG) enzyme. In a more preferred embodiment, the polypeptide comprises an amino acid sequence substantially identical with SEQ ID NO: 4 from about residue 647 to about residue 977.

Another embodiment of the invention is directed to an isolated polypeptide of at least 10 amino acid residues in length and which has at least about 80% sequence similarity

with a sequence shown in any one of SEQ ID-NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8 or SEQ ID NO: 10. Preferably, the polypeptide is at least about 20 amino acids in length, such as, for example at least about 30 amino acids, about 40 amino acids, about 50 amino acids, about 100 amino acids, about 200 amino acids and about 500 amino acids in length.

Another embodiment of the invention is directed to an antibody immunoreactive with an isolated polypeptide of at least about 6 amino acid residues in length and having a molecular weight less than about 65 kDa, which has at least about 80% sequence similarity with a sequence shown in any one of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8 or SEQ ID NO: 10. In a preferred embodiment, antibody is immunoreactive with a polypeptide with a molecular weight less than about 40 kDa and has at least about 90% sequence similarity with a sequence shown in any one of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8 or SEQ ID NO: 10. In another preferred embodiment, the antibody is immunoreactive with a polypeptide comprising an amino acid sequence substantially identical with SEQ ID NO: 4 from about residue 647 to about residue 977.

Another embodiment of the invention is directed to a method of detecting a polypeptide having PARG activity comprising the steps of contacting the polypeptide with an antibody immunoreactive with an isolated polypeptide of at least about 6 amino acid residues in length and having a molecular weight less than about 65 kDa, which has at least about 80% sequence similarity with a sequence shown in any one of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8 or SEQ ID NO: 10, and determining whether the antibody immunoreacts with the polypeptide.

Another embodiment of the invention is directed to a method of preventing, treating, or ameliorating a disease condition or disorder in an individual comprising the step of administering a therapeutically effective amount of a poly (ADP-ribose) glycohydrolase (PARG) inhibitor or activator to the individual. The disease condition or disorder may be any condition associated with responses to DNA damage, examples of which include a neoplastic disorder, a myocardial infarction, a vascular stroke or a neurodegenerative disorder. The PARG inhibitor or activator may be a small molecule inhibitor or activator of PARG or may be an antisense oligonucleotide that can hybridize in vivo to messenger RNA encoded by a PARG gene. PARG based treatment may be directed to new methods for preventing, treating or ameliorating disorders associated with DNA damage. These disorders include neoplastic disorders, inborn genetic errors, myocardial infarctions, vascular strokes, aging, and neurodegenerative disorders such as Alzheimer's disease, Huntington's disease, Parkinson's disease, and neurotoxicity generally.

Another embodiment of the invention is directed to the identification of novel PARG modulators which can activate or inhibit DNA repair and/or apoptosis. A PARG modulator is a compound that can activate or inhibit PARG. These modulators are preferably more efficacious and do not have the known side effects of present modulators. One method of identifying an agent that inhibits or activates poly(ADP-ribose) glycohydrolase (PARG) activity comprise the steps of providing a liquid medium that contains a polypeptide having PARG activity contacting the polypeptide with a candidate agent, in the presence of a reference compound having affinity for the polypeptide, under predetermined assay conditions, and determining the affinity of the candidate agent for the polypeptide relative to the reference compound. Thus, the modulation activity of the candidate agent relative to the

reference compound is determined. In this method, the polypeptide may be immobilized on a solid support. Further, the polypeptide may be generated in vitro by culturing a cell transformed with a nucleic acid molecule encoding PARG under conditions effective to express the polypeptide.

Another embodiment of the invention is directed to a method of identifying a mutant PARG allele in an individual comprising the step of obtaining genomic material from the individual; digesting the genomic material with a restriction enzyme having a recognition site inclusive of the mutant allele; fractionating the restriction fragments obtained from the digestion; and comparing the fractionation pattern with that obtained for a normal allele, thereby determining the presence or absence of the mutant allele. The fractionating step may be performed with electrophoresis.

Another embodiment of the invention is directed to a method of identifying a mutant PARG allele in an individual comprising the steps of hybridizing an oligonucleotide with genomic material from the individual, which oligonucleotide hybridizes under predetermined hybridization conditions to a region immediately 5' of a predetermined mutation site in the PARG alleles with the 3' terminus of the oligonucleotide complementary to an unmutated PARG allele; extending the oligonucleotide using PCR amplification; and determining the degree to which extension occurs, thereby determining the presence or absence of the mutant allele. The PCR extension reaction may be performed at a temperature above about 50° C. The determination may be performed by conducting electrophoresis (using for example, acrylamide at about 4% to about 10% or agarose and low melting temperature agarose from about 0.8% to about 4%) on the products of PCR amplification.

Another embodiment of the invention is directed to a method of screening molecules for PARG modulating activity (inhibition or activation) comprising the steps of providing a purified PARG enzyme; assaying the enzyme in the presence of a molecule to be screened; and comparing the activity of the PARG enzyme in the presence of the molecule to the activity of the PARG enzyme in the absence of the molecule.

Another embodiment of the invention is directed to a method of gene therapy comprising the step of delivering an oligonucleotide having a sequence complementary to at least a portion of a polynucleotide encoding a PARG enzyme to a cell to be treated. In the method, the oligonucleotide may have a sequence complementary to a sequence encoding a C-terminal portion of a PARG enzyme. Further, in the gene therapy method, the oligonucleotide may further comprise a ribozyme.

Another embodiment of the invention is directed to a method of delivering to a cell surface, an oligonucleotide having a sequence complementary to at least a portion of a polynucleotide encoding a PARG enzyme to a cell to be treated. In the method, the oligonucleotide may have a sequence complementary to a sequence encoding a C-terminal portion of a PARG enzyme. Further, in the method, the oligonucleotide may further comprise a ribozyme. The portion of a polynucleotide encoding a PARG enzyme may be, for example, the polynucleotide encoding the N terminus third of PARG, the middle third of PARG, or the C terminus third of PARG. The portion of a polynucleotide may encode a smaller part of PARG such as the N terminus 10% of PARG, the C terminus 10% of PARG, or any 10% portion in between such as from 10% to 20%, from 20% to 30%, from 30% to 40%, from 40% to 50%, from 50% to 60%, from 60% to 70%, from 70% to 80%, from 80% to 90%. The percent value used means a percent of the linear amino acid sequence. Thus, for a 1000 amino acid protein, the N terminus 10 percent is from

amino acid 1 to 100; 10% to 20% percent would be from amino acid 100 to 200 and so on. For a 970 amino acid protein, the N terminal 10% would be from amino acid 1 to 97; 10% to 20% would be from amino acids 98 to 194 amino acids.

Another embodiment of the invention is directed to a method of sensitizing a cell to a chemotherapeutic agent comprising the step of contacting the cell with a molecule that modulates the activity of a PARG enzyme. The molecule may be an oligonucleotide having a sequence complementary to at least a portion of a polynucleotide encoding a PARG enzyme. For example, the oligonucleotide may have a sequence complementary to a sequence encoding a C-terminal portion of a PARG enzyme. The portion of a polynucleotide encoding a PARG enzyme may be, for example, the polynucleotide encoding the N terminus third of PARG, the middle third of PARG, or the C terminus third of PARG. The portion of a polynucleotide may encode a smaller part of PARG such as the N terminus 10% of PARG, the C terminus 10% of PARG, or any 10% portion in between such as from 10% to 20%, from 20% to 30%, from 30% to 40%, from 40% to 50%, from 50% to 60%, from 60% to 70%, from 70% to 80%, from 80% to 90%. The oligonucleotide may further comprise a ribozyme. The method may be used, for example, as a method of treating a diseased cell characterized by the presence of DNA strand breaks. In the treatment, the cell is contacted with a molecule that modulates an enzymatic activity of a PARG enzyme.

Another embodiment of the invention is directed to a pharmaceutical composition comprising an oligonucleotide having a sequence complementary to at least a portion of a polynucleotide encoding a PARG enzyme. The produced molecule may be an oligonucleotide having a sequence complementary to at least a portion of a polynucleotide encoding a PARG enzyme. For example, the oligonucleotide may have a sequence complementary to a sequence encoding a C-terminal portion of a PARG enzyme. The oligonucleotide may comprise a ribozyme activity.

Another embodiment of the invention is directed to a virus that causes the production of an oligonucleotide having a sequence complementary to a polynucleotide encoding a PARG enzyme. This may be, for example, a viral vector which after the infection of a host cell, causes the production of an antisense RNA of PARG. The molecule may be an oligonucleotide having a sequence complementary to at least a portion of a polynucleotide encoding a PARG enzyme. For example, the oligonucleotide may have a sequence complementary to a sequence encoding a C-terminal portion of a PARG enzyme. The oligonucleotide may further comprise a ribozyme activity.

Other embodiments and advantages of the invention are set forth, in part, in the description that follows and, in part, will be obvious from this description and may be learned from the practice of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 depicts the cellular biochemical process that occurs after DNA damage.

FIG. 2 depicts the SDS-PAGE analysis of purified bovine thymus PARG.

FIG. 3 depicts the alignment of the DNA sequences of two PCR products and eight λ gt11 cDNA clones used to identify the cDNA coding for bovine PARG.

FIG. 4 depicts a northern blot analysis of bovine kidney cells mRNA transcripts.

FIG. 5 depicts an alignment of the putative bipartite NLS of bovine, human, and murine PARG and comparison with the bipartite NLS of PARP from different organisms.

FIG. 6 depicts expression of bPARG enzyme activity in *E. coli* (10).

FIG. 7 depicts a Southern blot analysis of bovine DNA probed with PARG cDNA.

FIG. 8 depicts activity gel autoradiogram of *E. coli* expressed bovine PARG.

FIG. 9 depicts the analysis by anion exchange HPLC of material released from ADP-ribose polymers by PARG action.

FIG. 10 depicts the SDS-PAGE analysis of the purification of *E. coli* expressed GST-PARG.

FIG. 11 depicts a schematic representation of the portions of the bovine PARG cDNA expressed as GST fusion constructs.

FIG. 12 depicts the cloning of the 1.8 kb PCR EcoRI fragment encoding for the 65 kDa catalytic domain of PARG.

FIG. 13 depicts an autoradiogram of an activity gel of GST-PARG fusion constructs expressed in *E. coli* and PARG expressed in baculovirus.

FIG. 14 depicts a schematic representation of the strategy used to isolate cDNA molecules encoding PARG from various organisms.

FIG. 15 depicts the domain organization of PARGs from different organisms.

FIG. 16 depicts an amino acid sequence alignment of bovine, murine, human, *drosophila* and *C. elegans* PARG enzymes.

FIG. 17 depicts a western blot of recombinant PARGs.

FIG. 18 depicts western blots of natural and recombinant expressed PARG.

FIG. 19 depicts the characterization of PARG by Western Blot in mouse cells of different PARP genotypes.

FIG. 20 depicts a partial restriction map of the mouse PARG locus.

FIG. 21 depicts a schematic representation of the strategy used to create PARG knockout mice.

DETAILED DESCRIPTION OF THE INVENTION

DEFINITIONS

List of Abbreviations

ADP	adenosine diphosphate
ADPR	ADP-ribose
AMP	adenosine monophosphate
ASPCR	allele-specific PCR
bp	base pair(s)
bPARG	bovine PARG
CePARG	<i>C. elegans</i> PARG
dPARG	<i>Drosophila melanogaster</i> PARG
DTT	dithiothreitol
GSH-Sepharose	Glutathione-Sepharose 4B
GST	glutathione-S transferase
hPARG	human PARG
HPLC	high pressure liquid chromatography
ICE	interleukin-1 b converting enzyme
IPTG	isopropyl- β -D-thiogalactoside
kb	kilobase pair(s)
MDBK	Madin-Darby bovine kidney cells
mPARG	murine PARG
NAD	nicotinamide adenine dinucleotide
NLS	nuclear location signal

PADPR DHB-Sepharose poly(ADP-ribose)-dihydroxyboronyl-Sepharose
 PAGE polyacrylamide-gel electrophoresis
 PARG poly(ADP-ribose)glycohydrolase
 PARP poly(ADP-ribose)polymerase [EC 2.4.2.30]
 PCR polymerase chain reaction
 PEG-6,000 polyethylene glycol 6,000
 PEG polyethylene glycol
 PMSF phenylmethylsulfonyl fluoride
 PR-AMP phosphoribosyl-adenosine monophosphate
 RFLP restriction fragment length polymorphism
 SDS sodium dodecyl sulfate
 SSCP single-strand conformation polymorphism
 TPCK Trypsin: L-1-tosylamido-2-phenylethyl chloromethyl ketone.

An “agonist” as defined herein refers to a molecule which, when bound to PARG, increases or prolongs the effect of PARG. Agonist may include proteins, nucleic acid molecules, carbohydrates, or any other molecules that bind to and modulate the effect of PARG.

An “allele” or “allelic sequence”, as defined herein refers to an alternative form of PARG. Alleles may result from at least one mutation in the nucleic acid molecule sequence and may result in altered mRNAs or polypeptides whose structure or function may or may not be altered. Any given natural or recombinant gene may have none, one, or many allelic forms. Common mutational changes which give rise to alleles, are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

An “ortholog” as defined herein refers to a nucleotide or amino acid sequence that is related to a reference nucleotide or amino acid sequence through speciation, and is therefore identical or structurally similar to the reference sequence.

A given nucleotide or amino acid sequence is said to be “substantially identical” with another sequence when the compared sequences have the same residues in the same order, excepting for any degeneracy (nucleotides) and conservative substitutions (amino acids).

A “regulatory sequence” of an expression vector is a DNA sequence necessary for inducing transcription of a gene, and includes a functional promoter and/or enhancer sequence. The term “operatively linked” as used herein means that a first nucleotide sequence, such as a regulatory element, is fused in frame with a second nucleotide sequence so as to afford a faithful transcription of the entire nucleotide sequence, which upon translation yields the desired protein.

The term “immunoreactivity” and related terms refers to the ability of antibodies and fragments thereof to bind to particular regions (antigens) presented by polypeptides and proteins, presented to the antibodies either as immunogens or targets. Typically, the binding affinity of the antibodies for their antigen is in the range 10^5 to 10^{11} , with higher affinities being preferred.

The term “specific immunoreactivity” refers to the ability of antibodies and fragments thereof to bind to particular regions (antigens) presented by polypeptides and proteins, presented to the antibodies either as immunogens or targets and not to unrelated antigens. For example, an antibody with specific immunoreactivity to actin will bind actin but would not bind another protein, such as a polymerase, which do not share epitopes with actin.

The term “nucleic acid molecule” refers to DNA, RNA and nucleic acid molecule analogs such as PNA and the like. PNA or “Peptide Nucleic Acid” is a nucleic acid molecule analog that has a neutral “peptide-like” backbone with nucleobases

that allow the molecule to hybridize to complementary RNA or DNA with higher affinity and specificity than corresponding oligonucleotides. PNA can be made to be more resistant to normal nucleases and are especially desirable, for example, in gene therapy. PNA is known to one of skill in the art and can be purchased or custom synthesized in numerous commercial laboratories including PerSeptive Biosystems, Inc. (Framingham, Mass.).

The term “modulate” means to activate or inhibit. For example, a PARG modulator may activate or inhibit PARG activity. “Modulation activity” means the amount of activation or inhibition. For example, a compound that increase PARG (or any other enzyme) activity by 10% will have a modulation activity of 10%. Conversely, a compound that decreases PARG activity by 10% will have a modulation activity of -10%.

As used herein, a given nucleotide or amino acid sequence is said to have a defined percentage of sequence similarity with another sequence when the two sequences differ by no more than the specified sequence similarity, including conservative substitutions, insertions, and deletions. Degenerate codons do not result in a change in amino acid upon translation, therefore, it is appreciated that identical amino acids can be encoded by several equivalent codons. The term “homology” and “sequence similarity” should have the same meaning for the purpose of this patent. Similarity parameters may be any generally acceptable parameter. For the purposes of this patent, percent similarity between two polymers such as nucleic acid molecules and polypeptides is preferably defined by Karlin and Altschul (11). The similarity algorithms of Karlin and Altschul are well known to those of skill in the art as exemplified by their adoption by the National Center for Biological Information. For nucleic acid molecule sequence searching, one desirable set of parameters would M (score for a pair of matching residues) at 5; N (score for mismatching residues) at -4; W (word length) at 11. For proteins, it is well known that some amino acids are similar and that substitution would be conservative. That is, for example, the replacement of an acidic amino acid with another acidic acid would be considered a conservative mismatch while the replacement of an acidic amino acid with a basic amino acid would be considered a more divergent mismatch. Preferably, the parameters for a desirable protein similarity determination are expressed in the sequence similarity matrix BLOSUM62 as described in Henikoff & Henikoff (12). Other similarity matrices that are also preferred in the invention are PAM40, PAM120 and PAM250 as described in Altschul (13).

The rapid synthesis of ADP-ribose polymers that occurs in response to DNA strand breaks is accompanied by very rapid polymer turnover, indicating that PARP and PARG activities are closely coordinated as cells respond to DNA damage. While PARP has been widely studied, information concerning structure and function relationships of PARG is much more limited. The present invention discloses the isolation of a cDNA encoding the bovine, human, murine and *drosophila* PARG and their deduced amino acid sequences.

The availability of PARP cDNA has allowed a number of molecular genetic approaches to study the function(s) of ADP-ribose polymer metabolism and the availability of PARG cDNA should allow the design of additional molecular genetic approaches for studying this metabolism. For example, disruption of the gene encoding PARG in mice containing a normal PARP gene will allow the determination of whether other cellular enzymes can replace PARG in the turnover of ADP-ribose polymers and/or whether development of animals will occur in the absence of PARG. Alternatively, disruption of the PARG gene in mice containing a

disrupted PARP gene may provide insights for the coordinated function of PARP and PARG.

One embodiment of the invention is directed to a deoxyribonucleic acid (DNA) molecule that encodes a polypeptide having poly(ADP-ribose) glycohydrolase (PARG) activity. Preferably, the molecule is of mammalian origin, such as, for example, of human origin.

In a preferred embodiment, a DNA molecule of the invention comprises a nucleotide sequence with at least about 70% sequence similarity with a sequence shown in SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, and SEQ ID NO: 9. Higher degrees of sequence similarity, such as about 80%, about 90%, and about 100% are preferred. Most preferred is a DNA molecule comprising a nucleotide sequence substantially identical with any one of sequence shown in SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, and SEQ ID NO: 9. It is preferred that a DNA molecule of the present invention comprises at least about 1000 nucleotides and has a nucleotide sequence with at least 80% sequence similarity with a sequence of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, and SEQ ID NO: 9. Most preferably, the DNA molecule consist of a nucleotide sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, and SEQ ID NO: 9.

For a DNA molecule of the present invention based on a human PARG gene it is preferred that the molecule comprises a nucleotide sequence that shows similarity to the sequence shown in SEQ ID NO: 3 from about residue 2113 to about residue 3105. More preferably, the sequence similarity is from about residue 1240 to about residue 3105. Still more preferably, the DNA molecule comprises a nucleotide sequence similarity to the coding sequence for the full-length hPARG as shown in SEQ ID NO: 3 from about residue 175 to about residue 3105.

A DNA molecule of the present invention affords probes and primer molecules that can be used in hybridization assays and PCR amplification. An exemplary oligonucleotide is less than about 1000 residues in length and comprises a nucleotide sequence at least about 10 residues long to ensure hybridization. Preferably, the at least about 10 residue region of the oligonucleotide is complementary to a sequence shown in SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, and SEQ ID NO: 9. Typically, the oligonucleotide will be a DNA molecule, which can be labeled by any method as desired, for example, with a radiolabel, a fluorescence label, or chemi-luminescent label.

Another embodiment of the invention is directed to a nucleic acid molecule that hybridizes to in a nucleic acid blot (Southern blot, Northern blot) to a sequence of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, or SEQ ID NO: 9 under stringent hybridization conditions. A nucleic acid blot may be made using techniques defined in Molecular Cloning, Second Edition, Sambrook et al., Cold Spring Harbor Press, Cold Spring Harbor, N.Y. DNA to be analyzed may be separated in agarose or acrylamide gels. The DNA may be transferred to nylon or nitrocellulose membrane using techniques known to those in the art. Stringent hybridization condition may be for example, prehybridizations 42° C. in 50% formamide, 0.25 M sodium phosphate buffer, pH 7.2, 0.25 M NaCl, 7% SDS, 1 mM EDTA for 10 hours, 100 ug denatured salmon sperm DNA, hybridization at 42° C. in 50% formamide, 0.25 M sodium phosphate buffer, 100 ug denatured salmon sperm DNA, pH 7.2, 0.25 M NaCl, 7% SDS, 1 mM EDTA, 1 ng/ml probe with a specific activity of 10⁹ cpm/ug DNA, for 16 hours. The probe may comprise any contiguous sequence from SEQ ID NO: 1, SEQ ID NO: 3,

SEQ ID NO: 5, SEQ ID NO: 7, or SEQ ID NO: 9. Preferably, said contiguous sequence is at least about 50 bases long, more preferably, the contiguous sequence is at least about 75 bases long, such as at least about 100 bases, at least about 200 bases long or at least about 300 bases long. The complete sequence of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, or SEQ ID NO: 9. Methods of labeling probes to with radioactive labels are known to those of skill in the art.

Method of washing after stringent hybridization are known. A stringent washing may comprise, for example, two washes at in 2×SSC, 0.1% SDS. for 15 minutes each at room temperature; two washes in 0.2×SSC, 0.1% SDS for 15 minutes each at room temperature; and a final three washes in 0.2×SSC, 0.1% SDS for 15 minutes each at 60° C. The final wash may be increased in temperature for reduced background. For example, the final wash may be a final three washes in 0.2×SSC, 0.1% SDS for 15 minutes each at 65° C. or a final three washes in 0.2×SSC, 0.1% SDS for 15 minutes each at 68° C.

If a radioactive probe is used, hybridization may be monitored using known techniques such as autoradiogram or a two dimensional measurement of radioactivity.

An anti-sense oligonucleotide is also afforded by the present invention. The anti-sense molecule is typically less than about 1000 residues in length to ensure ease of synthesis, and hybridizes to an RNA molecule, e.g., messenger RNA, which has at least 70% sequence similarity with a sequence of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9. Preferably, the anti-sense molecule is at least about 10 nucleotides in length to ensure hybridization with mRNA. Even more preferably, the anti-sense molecule may be at least about 15 nucleotides in length such as, for example, at least about 20 nucleotides in length; at least about 30 nucleotides in length; at least about 50 nucleotides in length; at least about 75 nucleotides in length; at least about 100 nucleotides in length; at least about 150 nucleotides in length; at least about 200 nucleotides in length; at least about 500 nucleotides in length; at least about 1000 nucleotides in length; or at least about 1500 nucleotides in length. It is also preferred that the molecule has a ribozyme activity so that it can degrade the mRNA that it binds to.

An antisense oligonucleotide may be used therapeutically to inhibit translation of mRNA encoding PARG. Synthetic antisense oligonucleotides may be produced, for example, in a commercially available oligonucleotide synthesizer. This invention provides a means to therapeutically alter levels of expression of a human or other mammalian PARG by the use of a synthetic antisense oligonucleotide drug that inhibits translation of mRNA encoding PARG. Synthetic antisense oligonucleotides, or other antisense chemical structures designed to recognize and selectively bind to mRNA, are constructed to be complementary to portions of the nucleotide sequence shown in SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7 and SEQ ID NO: 9. An antisense oligonucleotide may be designed to be stable in the blood stream for administration to patients by injection, or in laboratory cell culture conditions, for administration to cells removed from the patient. The antisense may be designed to be capable of passing through cell membranes in order to enter the cytoplasm and nucleus of the cell by virtue of physical and chemical properties of the antisense oligonucleotide which render it capable of passing through cell membranes (e.g., by designing small, hydrophobic antisense oligonucleotide chemical structures) or by virtue of specific transport systems in the cell which recognize and transport the antisense oligonucleotide into the cell. In addition, the antisense oligonucleotide can be designed for administration only to

certain selected cell populations by targeting the antisense oligonucleotide to be recognized by specific cellular uptake mechanisms which bind and take up the antisense oligonucleotide only within certain selected cell populations. For example, the antisense oligonucleotide may be designed to bind to transporter found only in a certain cell type, as discussed-above. The antisense oligonucleotide may be designed to inactivate the PARG mRNA by (1) binding to the PARG mRNA and thus inducing degradation of the mRNA by intrinsic cellular mechanisms such as RNase I digestion, (2) by inhibiting translation of the mRNA target by interfering with the binding of translation-regulating factors or of ribosomes, or (3) by inclusion of other chemical structures, such as ribozyme sequences or reactive chemical groups, which either degrade or chemically modify the target mRNA. Synthetic antisense oligonucleotide drugs have been shown to be capable of the properties described above when directed against mRNA targets (14). In addition, coupling of ribozymes to antisense oligonucleotides is a promising strategy for inactivating target mRNA (15). In this manner, an antisense oligonucleotide directed to PARG may serve as a therapy to reduce PARG expression in particular target cells of a patient and in any clinical condition that may benefit from reduced expression of PARG.

It is known by those in the art that as a result of the degeneracy of the genetic code, a multitude of nucleotide sequences encoding PARG, some bearing minimal homology to the nucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates a nucleic acid molecule that encodes a polypeptide consisting of an amino acid sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8 and SEQ ID NO: 10. The invention contemplates each and every possible variation of nucleotide sequence that could be made by selecting combinations based on possible codon choices that would encode the oligopeptides disclosed herein. These combinations are made in accordance with the standard triplet genetic code as applied to the nucleotide sequence of naturally occurring PARG and all such variants are to be considered as being specifically disclosed.

Although nucleic acid molecules which encode PARG and its variants preferably hybridizes under high stringency conditions to the nucleotide sequence of the naturally occurring PARG gene under appropriate conditions of stringency, it may be advantageous to produce nucleotide sequences encoding PARG or its derivatives possessing a substantially different codon usage. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding PARG and its derivatives and variants without altering the produced amino acid sequence include the production of RNA transcripts having more desirable properties, such as greater half-life, than transcripts produced from the naturally occurring sequence.

In order to express a biologically active or immunologically active PARG, the nucleic acid molecule encoding PARG or functional equivalents, may be inserted into appropriate expression vector, such as, for example a vector which contains the necessary elements for the transcription and translation of the inserted coding sequence. Thus, another aspect of the present invention is an expression vector comprising a regulatory sequence operatively linked to nucleic acid molecule comprising a nucleotide sequence disclosed herein. For example, an expression vector can contain a nucleotide

sequence at least about 1000 base pairs in length, which has at least about 70%, about 80%, or higher, sequence similarity with a sequence shown in SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, and SEQ ID NO: 9.

Methods that are known to those skilled in the art may be used to construct expression vectors containing sequences encoding PARG and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination.

A variety of expression vector/host systems may be utilized to contain and express sequences encoding PARG. These include, for example, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with virus expression vectors (e.g., baculovirus), insects infected with virus expression vectors (e.g., fall army worm infected with baculovirus); plant cell systems transformed with virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus; TMV) or with bacterial expression vectors (e.g., Ti or bacterial plasmids); or animal cell systems. The invention is not limited by the host cell employed.

Prokaryotic expression systems are commercially available from a number of suppliers worldwide. Prokaryotic expression vectors provide a convenient system to synthesize proteins. If it is desired to express a protein with characteristics such as immunogenic properties, 3D conformation, and other features exhibited by authentic PARG, the protein may be expressed in an eukaryotic protein expression system. The eukaryotic expression systems are numerous and include mammalian, amphibian, plant, insect, and yeast expression systems.

Yeast hosts that can be used for expression include *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Pichia pastois*, *Hansela polymorpha*, *Kluyveromyces lactis*, and *Yarrowia lipolytica*. Yeast hosts offer the advantages of rapid growth on inexpensive minimal media and ease in large-scale production using bioreactors. Another advantage of yeast is the ability to direct expression to cytoplasmic localization or for extracellular export.

Most yeast vectors for protein expression are derivatives of the *S. cerevisiae* 2 μ (two micron) plasmid. Yeast vectors include pYES and pEST from Stratagene (La Jolla, Calif.). Constitutive gene expression by the yeast plasmid cassette can be mediated by well known promoters such as the glyceraldehyde-3-phosphate dehydrogenase promoter (TDH3); the triose phosphate isomerase promoter (TPI1); the phosphoglycerate isomerase promoter (PGK1); the alcohol dehydrogenase isozyme II (ADH2) gene promoter; GAL1 and GAL10 promoters; the metallothionein promoter from the CUP1 gene (induced by copper sulfate); and the PHO5 promoter (induced by phosphate limitation). Proper termination of yeast transcripts is known to those in the art. Termination signals may include the MF-alpha-1, TPI1, CYC1, and PGK1 genes. These termination signals may be spliced onto the 3' end of the insert to provide proper termination.

Insect expression systems include baculovirus based vectors designed to express foreign proteins in a number of insect hosts and insect cell line hosts. Insect and insect cell lines may be of *Drosophila melanogaster*, *Aedes albopictus*, *Spodoptera frugiperda*, and *Bombyx mori* origin. Numerous expression systems comprising cells, vectors, hosts and the like can be purchased from a variety of commercial sources.

The control elements or regulatory sequences necessary for the proper expression of the insert, in this case PARG, may comprises promoters, enhancers (including both proximal

and distal control elements) which interact with the host proteins to carry out transcription and translation. Such elements may vary in their strength and specificity and are known to those in the art. Depending on the vectors system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, may be used. For example, the LacZ promoter may be used in a bacterial cell; the baculovirus polyhedrin promoter may be used in an insect cell; plant promoters such as heat shock promoters, and storage protein promoters, plant virus promoters and the like may be used in a plant cell. In a mammalian cell expression system, an SV40 promoter or EBV promoter may be used.

Methods and protocols for both prokaryotic and eukaryotic expression systems are generally known to those in the art. Further, the cells, vectors, growth medium may be purchased from commercial suppliers. The catalogs and product literature of commercial suppliers provide detailed protocols to enable the expression of proteins in prokaryotic and eukaryotic systems including bacterial, yeast, insect, insect cell, and mammalian cell systems. The product literature and catalogs of Clontech (Palo Alto, Calif.), Invitrogen (Carlsbad, Calif.), Life Technologies (Rockville, Md.), Novagen (Madison, Wis.), Pharmigen (San Diego, Calif.), Quantum Biotechnologies (Montreal, Quebec, Canada), and Stratagene (La Jolla, Calif.) are incorporated herein by reference.

A further aspect of the invention is isolated proteins and protein fragments having poly(ADP-ribose) glycohydrolase (PARG) activity. Such a protein can comprise an amino acid sequence with sequence similarity of at least about 70%, about 80% or higher to a sequence shown SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, and SEQ ID NO: 10. For example, the full-length bovine PARG has a molecular weight greater than about 100 kDa, thereby distinguishing it from previously known PARGs. The protein may be purified, for example, from cell lysates using the antibodies of the invention. The purification may be through an antibody column.

PARG polypeptides are another aspect of the invention. Polypeptides of PARG may be used, for example, to generate antibodies in an immunogenic procedure. To be effective it is preferred that the polypeptides are at least about 6 amino acid residues in length, such as for example, at least about 10 amino acids in length, at least about 20 amino acids in length, at least about 30 amino acids in length, at least about 50 amino acids in length, at least about 75 amino acids in length, at least about 100 amino acids in length, at least about 150 amino acids in length, at least about 200 amino acids in length, or at least about 400 amino acids in length. In one embodiment, the polypeptide has a molecular weight less than about 65 kDa and with at least about 80% sequence similarity with a sequence shown in SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, and SEQ ID NO: 10. The polypeptide may consist of the sequence set forth in SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, or SEQ ID NO: 10.

The polypeptide of the invention may be conjugated to a larger molecule, such as, for example, keyhole limpet hemocyanin (KLH), to increase the immunogenicity of the polypeptide. The increased immunogenicity of the polypeptide will, in turn, increase the yield of antibody. Preferably, the polypeptide has a molecular weight less than about 40 kDa and with at least about 90% sequence similarity with a sequence shown in SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, and SEQ ID NO: 10. The polypeptide can also be used in a wide variety of assays, e.g., as a competitor of antigen in a liquid sample in an antibody-based assay. Therefore, it is preferred that the polypeptide has poly

(ADP-ribose)glycohydrolase (PARG) activity. A particularly preferred polypeptide is of human origin and comprises an amino acid sequence substantially identical with SEQ ID NO: 4 from about residue 647 to about residue 977—the C terminus catalytic region of the enzyme. Longer sequences more inclusive of the natural molecule are of course also contemplated.

The invention also encompasses PARG variants and alleles. A preferred PARG variant is one having at least 80% and more preferably at least 90% amino acid similarity to the amino acid of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8 and SEQ ID NO: 10 and which retains at least one biological, immunological or other functional characteristic or activity of PARG. A most preferred PARG variant is one having at least 95% amino sequence similarity or identity to human PARG (SEQ ID NO: 3).

Antibodies to PARG may be generated using numerous established methods that are well known in the art. One example of such a method is described in the Examples. Generated antibodies may include, for example, polyclonal, monoclonal, chimeric, single chain, Fab fragments, Fab' fragments, Fab'(2) fragments, and fragments produced by a FAB expression library. Humanized antibodies and single chain antibodies may also be produced after the amino acid sequence of effective antibodies are determined.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others, may be immunized by injection with PARG or any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, for example, Freund's mineral gels such as aluminum hydroxide, and surface-active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, and dinitrophenol. Among adjuvant used in humans, BCG (*bacilli Calmette-Guerin*) and *Corynebacterium parvum* are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to PARG have an amino acid sequence consisting of at least five amino acids and more preferably at least about 10 amino acids, such as for example about 20 amino acids or about 40 amino acids. It is also preferable that they are identical to a portion of the amino acid sequence of the natural PARG. Short stretches of PARG amino acids may be fused with those of another protein such as keyhole limpet hemocyanin and antibodies may be produced against the chimeric molecule.

Antibodies may be produced by inducing in vivo production in the lymphocyte population of a living animal or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in published procedures (16).

Antibody fragments that contain specific binding sites for PARG may be generated. For example, such fragments include the F(ab')₂ fragment, Fab fragment, Fab' fragment which can be produced by enzymatic digestion of the antibody molecule. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity (Huse, W. D. (1989) Science 254, 1275-1281).

Therapeutic Methods

A method of preventing, treating, or ameliorating a disease condition in a patient, which disease state is affected by the level of PARG expression is also contemplated. This method entails administering a therapeutically effective amount of a poly(ADP-ribose)glycohydrolase (PARG) inhibitor or acti-

vator to the individual. Particularly, implicated disease states are neoplastic disorder, myocardial infarction, vascular stroke and neurodegenerative disorders.

In one embodiment, antisense oligonucleotides for PARG may be used alone or in combination with other chemotherapeutic agents to treat neoplastic disorder. The anti-sense oligo is designed to hybridize in vivo to messenger RNA expressed by the organism. The use of anti-sense molecules in a therapeutic setting is described, for example, by S. Agrawal, *Anti-sense Therapeutics*, Humana Press. Currently favored protocols call for the oligo to have ribozyme activity in an effort to degrade the mRNA. These methods are described, for example, in *Therapeutic Application of Ribozymes*, K. Scanlon, ed., Humana Press. Therefore, in one embodiment, an antagonist of PARG may be administered to a subject to prevent or treat neoplastic disorder.

PARG levels may be enhanced to suppress DNA repair and increase a cell's susceptibility to chemotherapy drugs. Therefore, in another embodiment, a PARG enhancer is administered to a subject along with a chemotherapeutic drug as a treatment for neoplastic disorder.

Neoplastic disorders that can be treated by PARG elevation and chemotherapy include benign and malignant neoplasm such as, for example, adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, hyperplasia and hypertrophy. Neoplastic disorders may include, in particular, neoplastic disorders of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus. For the purposes of this invention, a neoplastic disorder is any new and abnormal growth; specifically a new growth of tissue in which the growth is uncontrolled and progressive. Malignant cancer is a subset of neoplastic disorders which show a greater degree of anaplasia and have the properties of invasion and metastasis.

The synthesis of effective anti-sense inhibitors is known. Numerous approaches have been previously described and generally involve altering the backbone of the polynucleotide to increase its stability in-vivo. Exemplary oligonucleotides and methods of synthesis are described in U.S. Pat. Nos. 5,661,134; 5,635,488; and 5,599,797 (phosphorothioate linkages), U.S. Pat. Nos. 5,587,469 and 5,459,255 (N-2 substituted purines), U.S. Pat. No. 5,539,083 (peptide nucleic acids) and U.S. Pat. Nos. 5,629,152; 5,623,070; and 5,610,289 (miscellaneous approaches). The disclosures of each of these references are incorporated herein by reference.

Significantly, the present invention discloses a method of identifying an agent that inhibits or activates poly(ADP-ribose) glycohydrolase (PARG) activity. Such method comprises (i) providing a liquid medium that contains a polypeptide of the present invention; (ii) contacting the polypeptide with a candidate agent, in the presence of a reference compound having affinity for the polypeptide, under predetermined assay conditions; and (iii) determining the affinity of the candidate agent for the polypeptide relative to the reference compound, thereby determining the inhibition or activation activity of the candidate agent relative to the reference compound. These determinations can be facilitated by immobilizing the polypeptide on a solid support. Alternatively, the polypeptide can be generated in vitro by culturing a cell transformed with a PARG gene under conditions effective to express the polypeptide.

Combination therapies are also afforded by the present invention in which a PARG inhibitor or activator is administered in combination with a chemotherapeutic or a "clot-

busting" drug. The clot-busting drug may be, for example, tissue plasminogen activator (t-PA) or streptokinase.

In some cases it may be desired to overexpress PARG in the cells of an organism in order to achieve the correct PARP/PARG balance. In this context of gene therapy, it is desired to stably transfect target cells with a vector, such as, for example, a viral or a DNA (nucleic acid) vector, so that the desired gene is overexpressed. Gene therapy vector systems and protocols are well known and are described, for example, in the *Internet Book of Gene Therapy* (17) Anti-sense and ribozyme approaches to cancer gene therapy are described in chapters 7-9 of the *Internet Book of Gene Therapy*, and are incorporated herein by reference. Another reference is *Gene Therapy Protocols*, P. Robbins, ed., Humana Press. Furthermore, gene therapy methods have advanced greatly and are well documented in numerous issued US patents. Gene therapy may be practiced, for example, by substituting a nucleic acid molecule of the invention with the nucleic acid molecule described in the methods referred to in any issued US patents directed to gene therapy (18).

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

Diagnostic Methods

Methods of genotyping an individual for a mutant PARG allele are also afforded by the present invention. A number of protocols are available for identifying a mutant allele as described herein once the nucleotide sequence encoding PARG is known. Some exemplary methods are restriction fragment length polymorphism (RFLP), allele-specific PCR (ASPCR) and single-strand conformation polymorphism (SSCP). Armed with this information, the genetic susceptibility of an individual to an above-mentioned disease condition can be assessed.

An allele-specific method for identifying point mutations by differential PCR amplification is described by (19). A non-electrophoretic method of genotyping with allele-specific PCR employs a dye specific for double-stranded DNA (20). A method of detecting mutations referred to as single-stranded conformation polymorphism (SSCP) is presently widely employed (21). A hybrid of SSCP and Sanger dideoxy sequencing, called dideoxy fingerprinting (ddF) has recently been described (22).

Other methods of identifying allelic mutations are known to the skilled artisan. Probably the most commonly used method of genotyping is restriction fragment length polymorphism (RFLP) (23), which employs one or more restriction enzymes to identify mutant alleles occurring within a restriction site. This method has been used extensively in forensic applications and is employed commercially by such companies as Helix Biotech, Inc. Reliagene Technologies, Inc. and GenTest Laboratories, Inc. Accordingly, an instant mutant PARG allele can be detected by RFLP methods, optionally by one of these commercial entities. The above methods are most effective in the detection of homozygotes for the defective allele.

An RFLP method of identifying a mutant PARG allele in an individual entails: (i) obtaining genomic material from the individual; (ii) digesting the genomic material with a restriction enzyme having a recognition site inclusive of the mutant allele; (iii) fractionating the restriction fragments obtained from the digestion, e.g., by electrophoresis; and (iv) comparing the fractionation pattern with that obtained for a normal allele, thereby determining the presence or absence of the mutant allele.

An ASPCR method of identifying a mutant PARG allele in an individual entails: (i) hybridizing an oligonucleotide with genomic material from the individual; (ii) attempting to extend the oligonucleotide using PCR amplification; and (iii) determining the degree to which extension occurs, thereby determining the presence or absence of the mutant allele. In this method, it is preferred that the oligonucleotide hybridizes under predetermined hybridization conditions to a region immediately 5' of a predetermined mutation site in the PARG allele with the 3' terminus of the oligonucleotide complementary to an unmutated PARG allele. In these protocols, the PCR extension reaction is generally attempted at a temperature above about 50° C., more preferably above about 60° C.

A variety of protocols including ELISA, RIA and FACS for measuring PARG levels are known in the art and provide a basis for diagnosing altered or abnormal levels of PARG expression. Normal or standard values for PARG expression may be established by combining body fluids and tissue biopsies from normal mammalian subjects, rupturing the cells or permeating the cells, combining the cells with antibody under conditions suitable for complex formation. The amount of standard complex formation may be quantified by various methods but preferably by photometric means. Quantities of PARG expressed in subject, control, and disease sample are compared to standard values to determine between normal, reduced or enhanced levels of PARG.

A still further aspect of the invention pertains to an antibody immunoreactive with a polypeptide of the present invention. Preferably the antibodies are specifically immunoreactive with the polypeptides of this invention such as, SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, and SEQ ID NO: 10. Frequently it is desired to label the antibody, e.g., with a radiolabel, fluorescent or epitope label, to permit visualizing the antibody. Thus, antibodies immunoreactive with the PARG of this invention are afforded, which can be used to study features of PARG heterogeneity and possible modes of regulation. The high degree of sequence similarity between bovine PARG, human PARG and murine PARG permits eliciting antibodies to PARG of one species, which are found to be cross-reactive with PARGs from other organisms. These antibodies are valuable in characterizing PARG in-vivo under defined physiological conditions in many different organisms.

Accordingly, a method of detecting a polypeptide having PARG activity, for example, a diagnostic assay, entails: (i) contacting the polypeptide with an aforementioned antibody of the invention; and (ii) determining whether the antibody immunoreacts with the polypeptide. Binding can be ascertained in an sandwich assay, as is well known, due to the ability of the antibodies to immunoreact with an epitope of PARG. Preferably, monoclonal antibodies, such as those prepared by the method of Kohler and Milstein (24) and labeled antigens effective in competing with the polypeptide, are employed. Exemplary assays are disclosed in U.S. Pat. No. 4,375,110, the disclosure of which is incorporated herein by reference.

The present invention includes immunoreactive fragments of a PARG enzyme. Immunoreactive fragments can be fragments that can elicit an immune response that recognizes a PARG enzyme. Alternatively, immunoreactive fragments can be fragments that are specifically bound by an antibody that specifically binds a PARG enzyme. Any of variety of methods may be employed in order to identify contiguous peptide fragments of a PARG enzyme that comprise immunoreactive sequences. PARG enzymes may be fractionated by proteases, cyanogen bromide, etc. and the resultant fragments assessed for their capacity to specifically bind anti-PARG antibodies.

In an alternative embodiment, one or more synthetic peptides may be prepared in order to locate contiguous amino acid sequences that are immunoreactive. The peptides may have a sequence that includes a series of contiguous amino acids that are identical to a series of contiguous amino acids of a PARG enzyme. The peptides may be of about six amino acids to about 500 amino acids in length. The peptide may also include sequences that are not identical to sequences of a PARG so long as it includes at least about six contiguous amino acids that are identical to about six contiguous amino acids of a PARG enzyme. In a preferred embodiment, the peptide will be about 50 amino acids in length. In other preferred embodiments the length of the peptide may be from about six amino acids to about 30 amino acids.

The peptides of the present invention may comprise amino acid sequences that elicit antibodies that specifically bind to the peptide or to a PARG enzyme. Alternatively, the peptides may contain sequences that are specifically bound by anti-PARG antibodies. Peptides that are bound by anti-PARG antibodies may identified through the use of Epitope Scanning™ strategy (Cambridge Research Biochemicals, Inc.). Thus, the linear sequence of amino acids of a particular PARG enzyme is used to construct a set of peptides of defined length which overlap other members of the set by one or more residues. The peptides may be any length; however, lengths of from about 6 to 25 amino acids are preferred. In selecting the length, a general consideration is that antibodies that recognize linear native epitopes constitute approximately 60-70% of the anti-protein antibody population (25).

The number of overlapping amino acids will generally be more than half of the length of the peptides. That is, if the peptides are about 20 amino acids long, the overlap may be 11 or more amino acids long. In preferred embodiments, each peptide will be selected such that the number of overlapping amino acid residues in adjacent peptides is from about (n-1) to (n-3), where "n" is the number of amino acids in the peptide. An overlap of (n-1) is particularly preferred. Thus, in a particularly preferred embodiment, a first peptide may have the amino acid sequence of residues 1-10 of a PARG enzyme, a second peptide may have the amino acid sequence of residues 2-11 of the same PARG enzyme, a third peptide may have the sequence of residues 3-12 of the same PARG enzyme and so on until the entire sequence of the PARG enzyme has been synthesized in fragments.

The peptides may be synthesized using any means known to those of skill in the art. In a preferred embodiment, the peptide will be synthesized using an automated synthesizer such as a multipin peptide synthesis system. Such systems or peptides synthesis services are commercially available from suitable providers known to those skilled in the art.

To identify suitable peptides, each peptide is introduced into a well of a microtiter plate, and assayed for its ability to bind to antibodies elicited by a PARG enzyme. Such assays may be conducted in various ways known to those skilled in the art. One suitable assay is conducted by immobilizing a peptide on the surface of a well and then contacting the peptide with a solution containing an anti-PARG antibody. After washing, the well is contacted with a labeled antibody that specifically binds to the anti-PARG antibody. Thus, the presence of label in the well indicates that the anti-PARG antibody bound to the immobilized peptide. Another preferred method of determining the ability of the peptide to be specifically recognized by anti-PARG antibodies is a competitive ELISA.

Once a particular peptide has been found to bind to anti-PARG antibodies, the peptide can be used to elicit monospecific antibodies. By immunizing an experimental animal with

a single peptide containing a single antigenic determinant, the antibodies elicited will all specifically bind to the same antigenic determinant even though the antibodies are not monoclonal.

Where desired, the peptides can be modified to increase their immunogenicity. Thus, they may be modified to contain an amino-terminal and/or a carboxyl-terminal cysteine or lysine residue with or without spacer arms. The peptides may be conjugated to carriers such as bovine serum albumin, ovalbumin, human serum albumin, KLH (keyhole limpet hemocyanin) or tetanus toxoid. The use of human serum albumin is preferred over ovalbumin or bovine serum.

The peptides, alone or conjugated to a carrier, may be themselves capable of eliciting an antibody response when administered to an experimental animal. Alternatively, the peptides, alone or conjugated to a carrier, may be administered in conjunction with an adjuvant. Those skilled in the art will understand that a variety of materials may function as adjuvants. Examples of possible adjuvants include, but are not limited to, Freund's complete adjuvant, Freund's incomplete adjuvant, lipopolysaccharide (LPS) and the like. Any material that increases the immune response to a fragment of a PARG enzyme may be used as an adjuvant.

The ability to produce large amounts of active PARG enzyme permits, for the first time, the large scale screening of chemical libraries for molecules capable of inhibiting or activating PARG enzymatic activity. The screening may be conducted using any assay for PARG known to those skilled in the art. In a preferred embodiment, the screen may be conducted using the TLC based assay described by Ménard, et al. (26). A known amount of PARG will be incubated under standardized conditions with [³²P]-poly(ADPR) in the presence of inhibitor or activator. After an appropriate period of time, the reaction will be stopped and the reaction mixture separated on PEI-F cellulose TLC plates. The TLC plates may be developed in an appropriate solvent system such as methanol followed by 0.3N LiCl. The amount of ADPR released in the reaction will be quantified and the effect of the inhibitor or activator on enzymatic activity will be determined. Typical reaction conditions are 50 mM potassium phosphate (pH 7.5) at 37° C. in the presence of 25 μM [³²P]-poly(ADPR). The concentration of the inhibitor or activator can be varied as necessary to determine the K_i value of the inhibitor or activator according to standard procedures.

Another embodiment of the invention is directed to a method of altering the response of the cell to a genotoxic stress by modulating the concentration of ADPR polymers. As discussed above, the metabolism of ADPR polymers is critical in determining the fate of cells subjected to genotoxic stress. The modulation can be either an increase or a decrease in the concentration of the polymers. In one embodiment of the present invention, the concentration of ADPR polymers can be decreased by the use of a gene therapy vector expressing a high level of PARG. In another embodiment of the present invention, the concentration of polymers can be increased by inhibiting the enzymatic activity of the PARG enzyme by the addition of inhibitors or activators identified as described above. Alternatively, the concentration of ADPR polymers can be increased by interfering with the endogenous expression of PARG enzymes using antisense oligonucleotide technology.

Knowledge of the nucleotide sequence of the PARG gene permits the preparation of antisense therapeutics containing sequences complementary to the mRNA of PARG gene. The preparation and delivery of antisense therapeutics is well known to those skilled in the art. For example, antisense therapeutics have been used to treat neoplastic disorder as

exemplified by Smith, U.S. Pat. No. 5,248,671, specifically incorporated herein by reference. Additional examples of antisense therapeutics are provided by Miller, U.S. Pat. Nos. 4,511,713 and 4,757,055, specifically incorporated herein by reference.

In the present invention, an oligonucleotide having a sequence complimentary to the mRNA of the PARG gene will be prepared. Such an oligonucleotide is said to be an antisense oligonucleotide with respect to the PARG gene. The oligonucleotide may be RNA or DNA or a may contain both RNA and DNA portions. The oligonucleotide may contain modified bonds so as to enhance the stability of the oligonucleotide and render it more resistant to the action of cellular nucleases. For example, the oligonucleotide may be constructed with phosphorothioate nucleotides, phosphonate nucleotides and other types of modified nucleotides known to those skilled in the art. The structure of the oligonucleotide may be altered so as to include other types of bonds that do not naturally occur in oligonucleotides. For example, adjacent nucleosides might be joined using linear alkyl chains, peptide bonds or other types of structures. The only limitation is that the resulting oligonucleotide remains capable of hybridizing to the target PARG mRNA.

The antisense oligonucleotides may be delivered by any means customarily used in the art. For example, the oligonucleotide may be delivered in neutral liposomes, cationic liposomes or by ballistic high speed injection. Alternatively the DNA sequence encoding the antisense oligonucleotide may be inserted into a gene vector and the vector may be introduced into target cells. The vector may be any type of gene therapy vector known to those skilled in the art. Preferred embodiments include, plasmid vectors and viral vectors. Viral vectors are seen to include those vectors customarily used for gene therapy applications including, but not limited to, retroviral vectors, vaccinia virus vectors, herpes virus vectors, adenovirus vectors and adeno-associated virus vectors. Upon introduction of the vector into target cells, the vector will direct expression of a nucleic acid molecule comprising the appropriate sequence to hybridize with the mRNA encoding a PARG enzyme. In a preferred embodiment, introduction of the vector into the target cell will result in the production of an RNA molecule that hybridizes with the mRNA of a PARG enzyme and also includes one or more additional RNA sequences capable of functioning as a ribozyme. The ribozyme portion of the molecule will cause the cleavage of the mRNA encoding the PARG enzyme thereby preventing the production of PARG.

Therapeutics of this type may be used to treat a wide variety of conditions. In one embodiment, an antisense therapeutic will be used to treat neoplastic disorder. In a preferred embodiment, an antisense therapeutic of the present invention will be delivered in combination with a currently known chemotherapeutic agent. In general, chemotherapeutic agents function by disrupting the integrity of DNA in target cells. Since the recovery of a cell from such DNA disruption is highly dependent upon the normal ADPR polymer metabolism, the presence of the antisense therapeutic will have the effect of chemosensitizing the neoplastic cells by disturbing the ratio PARG and PARP.

In another preferred embodiment, the antisense oligonucleotides of the present invention may be used to treat a variety of conditions caused by genotoxic oxidative stress. Examples include cardiac disorders, neuronal disorders, reperfusion injury, neurotoxicity, Alzheimer's disease, Huntington's disease and Parkinson's disease. It has been shown that inhibition of ADPR polymer synthesis provides protection against cellular damage caused by nitric oxide injury. Zhang,

et al., U.S. Pat. No. 5,587,384, specifically incorporated herein by reference, teach that decreasing the amount of ADPR polymers formed can result in protection against nitric oxide induced neurotoxicity. As discussed above, decreasing the amount of ADPR polymers in the cell can be accomplished by the introduction of gene therapy vector expressing PARG, thus, the present invention can be used to treat neurodegenerative conditions resulting from oxidative stress.

CONCLUSION

The synthesis and rapid turnover of ADP-ribose polymers is an immediate cellular response to DNA damage. Reported here is the isolation and characterization of cDNAs encoding various poly(ADP-ribose) glycohydrolase (PARG) enzymes responsible for ADP-ribose polymer turnover. PARG was isolated from bovine thymus, yielding a protein of approximately 59 kDa. Based on the sequence of oligopeptides derived from the enzyme, polymerase chain reaction products and partial cDNA clones were isolated and used to construct a putative full-length cDNA. The cDNA of approximately 4.1 kb pairs predicts expression of a protein of approximately 111 kDa, nearly twice the size of the isolated protein. A single transcript of approximately 4.3 kb pairs is detected in bovine kidney poly(A)⁺ RNA, consistent with expression of a protein of 111 kDa. Expression of the cDNA in *Escherichia coli* results in an enzymatically active protein of 111 kDa and an active fragment of 59 kDa. Analysis of restriction endonuclease fragments from bovine DNA by Southern hybridization indicate that PARG is encoded by a single copy gene. Taken together, the results indicate that previous reports of multiple PARGs can be explained by proteolysis of an 111-kDa enzyme. The deduced amino acid sequence of the bovine PARG shares little or no sequence similarity with differing types of known proteins; however, it contains a putative bipartite nuclear location signal as would be predicted for a nuclear protein. The availability of cDNA clones for PARG should facilitate structure-function studies of the enzyme and its involvement in cellular responses to genomic damage.

Other embodiments and advantages of the invention are set forth, in part, in the description that follows and, in part, will be obvious from this description and may be learned from practice of the invention.

EXAMPLES

Example 1

Purification of Bovine PARG

PARG was purified from bovine thymus tissue (Pel-Freez, Rogers, A K) by modifications of previously published procedures (27). The enzyme was isolated up to the polyethylene glycol (PEG)-6,000 fractionation step as described previously (28). However, DNA-agarose and heparin-Sepharose chromatographic steps used previously were omitted, and the PEG-6,000 fraction was applied directly to an affinity matrix of poly(ADP-ribose)-dihydroxyboronyl-Sepharose (PADPR DHB-Sepharose). The active fractions eluted from PADPR DHB-Sepharose (25 ml) were pooled, placed in dialysis tubing, concentrated against dry PEG-20,000 to approximately 12 ml, and dialyzed against 2 liters of 20 mM potassium phosphate buffer, pH 8.0, 0.1% Triton X-100, 5 mM β -mercaptoethanol, 0.1 mM thioglycolic acid, 0.4 M KCl (buffer A). The sample was loaded onto a 1.0 \times 11-cm Toyopearl AF-Red (Supelco) column, and PARG was eluted with an 80-ml linear gradient of 0.4-2 M KCl in buffer A. The active

fractions, eluting at approximately 1.25 M KCl, were pooled, placed in dialysis tubing, concentrated against solid sucrose to approximately 9 ml, and dialyzed against 20 mM potassium phosphate buffer, pH 7.2, 0.75 M KCl, 0.1% Triton X-100, 10% glycerol, 5 mM β -mercaptoethanol, 0.1 mM thioglycolic acid. PARG activity was determined as described by Menard and Poirier (29), and protein content was determined by the method of Bradford (30). The final preparation was quantified by SDS-PAGE (31) and Coomassie Blue staining to compare the intensity of the protein band with a known amount of bovine serum albumin (32).

The purification procedure for the bovine thymus PARG summarized in Table 1 is typical for results obtained from six separate preparations of the enzyme. Purification from 500 g of bovine thymus achieved approximately 50,000-fold purification and yielded approximately 20 μ g of purified protein. An aliquot of the purified enzyme was precipitated with trichloroacetic acid, washed with acetone, resuspended in SDS-PAGE sample buffer, separated on a 10% SDS-PAGE gel, and stained with Coomassie Blue. Analysis of the final preparation of SDS-PAGE revealed that more than 95% of the protein migrated at an apparent molecular mass of approximately 59 kDa (FIG. 2). In FIG. 2, an aliquot of the purified enzyme was precipitated by TCA, washed with acetone, resuspended in SDS-PAGE sample buffer, separated on a 10% SDS-PAGE gel and stained with Coomassie blue. The positions of molecular weight marker proteins are shown.

TABLE 1

Purification of PARG from bovine thymus					
Step	Protein mg	Total activity units	Specific activity units/mg protein	Yield %	Purification -fold
Crude extract	27,800	57,400	2.06	100	1.0
Protamine sulfate	12,500	58,000	4.64	101	2.3
Ammonium sulfate	4,480	30,000	6.70	52	3.3
CM-Sepharose	171	19,100	112	33	55
PEG 6000	23.0	7,530	327	13	160
PADPR-DHB-Sepharose	1.30	6,730	5,180	12	2,500
Toyopearl AF-Red	0.023	2,260	98,300	4	48,000

Example 2

Peptide Sequencing

Prior to proteolytic fragmentation, the purified bPARG (40 μ g in 100 μ l of 0.4 M ammonium bicarbonate buffer, pH 8.0, 8 M urea) was incubated in a final concentration of 2.2 mM dithiothreitol at 56 $^{\circ}$ C. for 15 min. Iodoacetamide was added to a final concentration of 2.0 mM, and the sample was incubated at 25 $^{\circ}$ C. for 15 min. After dilution with an equal volume of water, 1.5 units of immobilized L-1-tosylamido-2-phenylethyl chloromethyl ketone-treated trypsin (Pierce Chemical, Rockford, Ill.) was added, and the sample was incubated at 37 $^{\circ}$ C. for 18 h with gentle rotary shaking. Finally, the mixture was subjected to centrifugation at 16,000 \times g for 5 min to separate the tryptic fragments from the immobilized trypsin. The tryptic fragments were adjusted to 0.05% in trifluoroacetic acid and separated on a 4.6 mm \times 25 cm, Microsorb MV, C₄ reversed-phase HPLC column (Rainin) eluted with an 80-min linear gradient from 4 to 44%

acetonitrile in 0.05% trifluoroacetic acid. Four oligopeptide fractions, with approximate elution times of 61, 63, 68, and 75 min, were selected for peptide sequence analysis by the Edman degradation method. Amino acid sequence data of four oligopeptides, designated by their approximate HPLC elution times from the reversed-phase column, are shown in Table II.

TABLE II

Amino acid sequence of oligopeptides derived from bPARG			
Oligopeptide	Amino Acid Sequence		SEQ ID NO:
	10	20	30
68	LFTEVLHDHNE	CLIIITGTEQY	SEYTGVAETY R
			SEQ ID NO: 11
63	AYCGFLRPGV	SSENLSAVAT	GNXGCGAFG
			SEQ ID NO: 12
61	FLINPELIVS	R	
			SEQ ID NO: 13
75	IALXLPNIXT	QPIPLL	
			SEQ ID NO: 14

Example 3

cDNA Cloning

To obtain cDNA clones encoding bovine PARG, PCR amplification experiments were followed by the screening of two different bovine cDNA libraries. FIG. 3 depicts the alignment of the DNA sequences of two PCR products and eight λ gt11 cDNA clones used to identify the cDNA coding for bovine PARG. The two PCR products and clones 1 and 2 were obtained from the bovine thymus cDNA library. Clones 3-8 were obtained from the bovine kidney cDNA library. The positions of restriction sites used in this study are shown, and the top diagram shows the consensus clone, denoting the relative location of the coding regions for oligopeptides, 75, 61, 68, and 63 as well as the open reading frame and noncoding regions. For each of the cDNA inserts characterized, the sequence of both strands was determined by the dideoxynucleotide chain termination method using SequenaseTM (U.S. Biochemical Corp., Cleveland, Ohio).

The first step leading to the isolation of cDNA clones was to synthesize two multi-degenerate 17-mer primers, GAYCAYAAAYGARTGYYT (SEQ ID NO: 15) and CKRIANGTYTCNGCRTA (SEQ ID NO: 16) (where Y represents T/C, R is A/G, K is T/G, and N is A/T/C/G), based on two regions of the SEQ ID NO: 11; "DHNECL" (amino acids 7 to 12 of SEQ ID NO: 11) and "YAETYR" (amino acid 26 to amino acid 31 of SEQ ID NO: 11) (Table II). Using the multidegenerate primers and an oligo(dT)-primed bovine thymus cDNA λ gt11 library BL1019b from Clontech (Palo Alto, Calif.), PCR amplification generated a 74-bp DNA fragment with a deduced amino acid sequence identical to the corresponding region of oligopeptide 68. Next, two specific 24-mer oligonucleotide primers, ATCATCACAGGTACTGAGCAGTAC (SEQ ID NO: 17) and GCCTGTGTATTCACTGTACTGCTC (SEQ ID NO: 18), based on the sequence of this 74-bp DNA were used in combination with λ gt11 forward and reverse primers to amplify PCR products 1 and 2 from the bovine thymus library. PCR product 1 contained 231 bp of sequence including the region encoding the N-terminal region of oligopeptide 68 (SEQ ID NO: 11) and the entire

sequence of oligopeptide 61 (SEQ ID NO: 13). PCR product 2 contained 757 bp, which included a sequence encoding the C-terminal region of oligopeptide 68 (SEQ ID NO: 11) and the entire sequence of oligopeptide 63 (SEQ ID NO: 12).

The sequence information obtained from PCR products 1 and 2 was used to isolate cDNA clones obtained by the screening of bovine thymus and bovine kidney cDNA libraries. A 518-bp EcoRI-HindIII fragment from PCR product 2 was used as a probe to screen approximately 1×10^6 independent clones from the bovine thymus library. Two positive cDNA clones (clones 1 and 2) were isolated, which overlapped PCR products 1 and 2. However, attempts to obtain clones from the bovine thymus library that contained sequence 5' to clone 2 were unsuccessful. Thus, a 231-bp EcoRI-KpnI fragment from clone 2 was used as a probe to screen approximately 5×10^5 independent clones of the bovine kidney 5' stretch plus cDNA λ gt11 library BL3001b (Clontech, Palo Alto, Calif.). Three positive cDNA clones (clones 3-5) were obtained, all of which contained sequence 5' to clone 2. Each of these clones also contained a sequence encoding oligopeptide 75. Clones 1-5 provided multiple overlapping sequences in the 3'-terminal portion of a consensus cDNA, but additional clones were sought to obtain overlapping sequences for the 5'-terminal region. Thus, a 436-bp EcoRI-KpnI fragment located at the 5' end of clone 3 was used as a probe to screen approximately 6×10^5 independent clones of the bovine kidney library. Clones 6-8 provided overlapping sequences for the 5'-terminal region. The full-length cDNA was constructed by ligating a 3.9-kb XbaI-NsiI fragment from pWL11 (clone 1 cDNA insert in pTZ18R (33)) and a 3.0-kb NsiI-XbaI fragment from pWL13 (clone 4 cDNA insert in pTZ 18R). The resulting plasmid, termed pWL30, contained the 4,070-bp full-length cDNA.

FIG. 3 shows an alignment of the DNA sequences of two PCR products and eight λ gt11 cDNA clones used to identify the cDNA coding for bovine PARG. The two PCR products and Clones 1 and 2 were obtained from the bovine thymus cDNA library. Clones 3 through 8 were obtained from the bovine kidney cDNA library. The position of restriction sites used in this study is shown and the top diagram shows the consensus clone, denoting the relative location of the coding regions for oligopeptides 75, 61, 68, and 63 as well as the open reading frame and non coding regions.

The nucleotide sequence of cDNA coding for bovine PARG is shown in the sequence listing as SEQ ID NO: 1. The deduced amino acid sequence of the enzyme is shown in the sequence listing as SEQ ID NO: 2. The four oligopeptides sequenced from purified enzyme is within SEQ ID NO: 2. They are IALCLPNICTQPIPLLK (amino acid 601 to 617, SEQ ID NO: 2); LINPELIVSR (amino acid 761 to 770, SEQ ID NO: 2); LFTEVLHDHNECLIIITGTEQYSEYTGVAETYR (amino acid 771 to 801, SEQ ID NO: 2) and AYCGFLRPGV PSENLSAVAT GNWCGCGAFGGDAR (amino acid 849 to 880, SEQ ID NO: 2). The combined nucleotide sequence of Clones 1 through 8 predicted a full-length cDNA clone of 4,070 bp containing 257 bp of 5'-non-coding sequence, a single open reading frame of 2,931 bp (beginning at the ATG at position 258 of SEQ ID NO: 1) and a 3'-non-coding region of 882 bp, and the deduced amino acid sequence which predicts a protein of 977 amino acids and a molecular weight of 110.8 kDa.

Analysis of the Sequence of Bovine PARG

The cDNA clone (SEQ ID NO: 1) has features typical of cDNAs that code for mammalian proteins. These include (i) an oligo A (putative poly(A)+) sequence at the 3'-end, (ii) a polyadenylation signal (AATAAA) 12 bp upstream from the oligo A sequence, (iii) a sequence of ATTTA in the 3'-untranslated region thought to play a role in selective mRNA degradation in mammalian cells (34), (iv) a single open reading frame, and (v) a nucleotide sequence around the first start codon commonly found at known sites of initiation of translation (35). The evidence that the cDNA clone constructed represents a full-length or nearly full-length clone for PARG is shown by the observation that hybridization of poly(A)+ RNA from bovine kidney cells with the cDNA showed a single band of hybridization of approximately the same size as the cDNA under stringent hybridization conditions (set forth above) (FIG. 4).

The nucleotide sequence encoding bovine PARG indicates that PARG shares little or no sequence similarity with other known sequences. A search of sequence data banks has failed to reveal significant sequence similarity with any sequences coding for known proteins. A strong sequence similarity has been observed with human and rat cDNA clones that likely represent partial clones for PARG from these species. Examination of protein sequence databases such as Genbank and SwissPro also has shown that the deduced amino acid sequence of PARG lacks any sequence similarity with known proteins. However, the amino acid sequence shares a significant similarity with a protein sequence from *Caenorhabditis elegans* that may represent the PARG protein from this organism (36).

The deduced amino acid sequence of PARG has been examined for a number of structural motifs that can be predicted from the primary amino acid sequence. The expressed PARG protein was observed to be able to form dimers stable to SDS-PAGE conditions. In that regard, residues 871-907 show significant homologies to known leucine zipper dimerization sequences (37).

Another motif identified is a putative bipartite nuclear localization signal (NLS) (38). It is interesting that PARP also contains a bipartite NLS (39). FIG. 5 compares deduced amino acid sequences in the NLS region of the bovine PARG, and regions of putative PARG sequences from human, mouse and *C. elegans*, with the NLS region of PARP from seven different organisms. Conserved residues are noted in bold and the amino acid distances are from the amino terminal methionine residue. Abbreviations and references for the sequences shown are as follows: bPARG, bovine PARG (SEQ ID NO: 19); hPARG, human PARG (SEQ ID NO: 20); mPARG, murine PARG (SEQ ID NO: 21); CePARG, *Caenorhabditis elegans* PARG (SEQ ID NO: 22); hPARP, human PARP (SEQ ID NO: 23; 40); mPARP, murine PARP (SEQ ID NO: 24; 41); bPARP, bovine PARP (SEQ ID NO: 25, 42); aPARP, chicken PARP (SEQ ID NO: 26; 43); XIPARP, *Xenopus laevis* PARP (SEQ ID NO: 27; 44); DmPARP, *Drosophila melanogaster* PARP (SEQ ID NO: 28; 45); SpPARP, *Sarcophaga peregrina* PARP (SEQ ID NO: 29; 46). In FIG. 5, conserved residues are noted in boldface type, and the amino acid distances are from the amino-terminal methionine residue. Sequence alignment of putative bipartite nuclear localization signal of bovine, human and murine PARG compared to the nuclear localization signal of PARP from different organisms. The putative NLS of PARG fulfills the criteria for bipartite NLS in that it contains conserved acidic and basic amino acid residues at

two different locations each within the region of sequence similarity to the NLS of PARP (47).

A surprising finding was that the bovine PARG cDNA clone codes for a protein of approximately 111 kDa, which is nearly twice the size of the PARG protein isolated from bovine thymus (FIG. 2). It indicates that PARG contains a protease sensitive site that, following proteolysis, yields a protein fragment of approximately 59 kDa that still retains enzymatic activity. Several pieces of evidence favor this possibility. (i) Expression of the carboxyl terminal portion of the cDNA resulted in enzymatic activity (FIG. 6, bar 5). (ii) All of the oligopeptides sequenced were located in the carboxyl terminal half of the protein (FIG. 3, FIG. 6 and Table 2). (iii) The only protein, other than 59 kDa protein detected in the thymus preparation was approximately 111 kD (FIG. 2). (iv) The PARG activity expressed in bacteria was sensitive to proteolysis, yielding a protein of approximately 56 kD (FIG. 6). (v) The cleavage site in PARG is in the region of the putative NLS and the PARP NLS is located in a protease sensitive site (48). Taken together with the data suggesting that bovine PARG appears to be coded for by a single copy gene (FIG. 7), proteolysis seems likely to explain the presence of PARG activity of molecular weight of approximately 74 kDa and 59 kDa in bovine thymus preparations (49). Likewise, a similar mechanism could explain previous reports of a PARG of 74 kDa isolated from nuclear fractions of guinea pig liver and human placenta (50) and a PARG of 59 kDa isolated from postnuclear fractions of guinea pig liver (51).

While proteolysis of a larger protein to yield smaller proteins retaining PARG activity seems likely to explain the size heterogeneity of PARG previously reported, it remains to be determined if proteolysis normally occurs in vivo or whether it occurs during purification of the enzyme. While the results presented here show that a full-length protein can be expressed containing PARG activity (FIG. 8), the molecular size of PARG in vivo also remains to be determined. If PARG occurs as a larger protein, an interesting possibility is that the amino terminal region may be involved in the regulation of enzymatic activity.

Example 5

Expression of bPARG in *Escherichia coli*

To determine whether the isolated cDNA encoded PARG, bPARG was expressed using two different bacterial expression systems, the pTrcHis Xpress System™ (Invitrogen, Carlsbad, Calif.), in which the expressed protein contains a leader poly-histidine sequence, and the glutathione S-transferase (GST) gene fusion system (Pharmacia Biotech Inc., Piscataway, N.J.). For expression in the pTrcHis Xpress system, three different DNA fragments were amplified and inserted into the pTrcHis expression plasmid. Constructs A and B contained the entire opening reading frame of 110.8 kDa, which together with the fusion partner predicted a protein of about 115 kDa. Construct B also contained the 3'-untranslated region of the clone. Construct A, containing the cDNA sequence-3 to 2,946, was prepared by subcloning a 2.9kb XhoI-EcoRI DNA fragment amplified from pWL30 with primers WIN34 (GCTGCGGGTCTCGACGATGAGT-GCGGGC) (SEQ ID NO: 30) and WIN15 (CGGTCTA-GAATTCAGTGGCTCCTCAGGC) (SEQ ID NO: 31). Construct B, containing the cDNA sequence-3 to 3,813, was prepared by subcloning a 3.8-kb XhoI-EcoRI DNA fragment amplified from pWL30 with primers WIN34 (SEQ ID NO: 30) and WIN33 (CCGGAATTCGGGTTTTTTGTTAAT-

GAAAATTATTAAC) (SEQ ID NO: 32). Construct C, containing cDNA sequence 964-2,946, was prepared by subcloning a 2.0-kb DNA fragment amplified from pWL13 with primers WIN14 (TCAGAGCAGATGAACTCGAGCAGTC-CAGG) (SEQ ID NO: 33) and WIN15 (SEQ ID NO: 31). Since the isolated PARG of approximately 59 kDa contained enzymatic activity, construct C contained only the 75-kDa carboxyl-terminal region of the PARG, which predicted a fusion protein of approximately 79 kDa.

For expression experiments of bPARG as a GST fusion protein, an insert containing the cDNA sequence from position 1138 to 2946 was prepared by subcloning a 1.8-kb EcoRI-EcoRI fragment amplified from pWL30 with the oligonucleotide CCAATTTGAAGGAGGAATCCCCGCCGC-CACCATGAATGATGTGAATGCCAAACGACCTGGA (SEQ ID NO: 34) and WIN15 (SEQ ID NO: 31) as primers. The resulting DNA fragment was inserted into the EcoRI site of the pGEX-2T expression vector, and the plasmid was used to transform *E. coli* NM522 cells.

For expression experiments, bacterial cultures were grown at 37° C. in 1% Bacto-tryptone, 0.5% yeast extract, and 0.5% NaCl to a density of approximately 0.6 A₆₀₀/ml and were induced with 1 mM isopropyl-β-D-thiogalactoside (IPTG). Cells were collected by centrifugation, and crude extracts were prepared by sonication (10 A₆₀₀/ml) in 10 mM sodium phosphate buffer, pH 7.2, 150 mM NaCl, 0.5 mg/ml lysozyme, 0.1 mg/ml phenylmethylsulfonyl fluoride, 1 mM EDTA, 0.7 μg/ml pepstatin A, 0.5 μg/ml leupeptin, and 1 μg/ml aprotinin. Cell extracts were subjected to centrifugation, and the supernatant fraction was used for assay. PARG assay conditions were as described previously (52). Following incubations, portions of reaction mixture were analyzed by thin layer chromatography or subjected to anion exchange HPLC.

Using a thin layer chromatography assay that measures release of [³²P]ADP-ribose from [³²P]ADP-ribose polymers (53), PARG activity was detected in extracts from cells transformed by each of the constructs. FIG. 6 shows results obtained with constructs B and C. Reaction mixtures contained approximately 15,000 cpm of [³²P]ADP-ribose polymers, and the cpm shown represent ADP-ribose released from the ADP-ribose polymers. Bar 1, a strain transformed by pTrcHis without an insert but induced with 1 mM IPTG for 5 h at 37° C. A strain containing construct B is shown without the addition of IPTG (bar 2) or after the addition of 1 mM IPTG for 1.5 h (bar 3) or 5 h (bar 4). A strain containing construct C 5 h after induction by IPTG is shown in the absence (bar 5) and presence (bar 6) of 167 μM ADP-hydroxymethylpyrrolidine diol (54). No activity was detected in cells transformed with the empty vector, but activity was detectable without induction by IPTG, indicating a leaky lac promoter. The addition of IPTG resulted in a time-dependent increase of up to approximately 4.5-fold in enzymatic activity. FIG. 6 also shows that the enzymatic activity was strongly inhibited by the presence of ADP-hydroxymethylpyrrolidine diol, a specific inhibitor of PARG (55).

In FIG. 9, material released from ADP-ribose polymers by anion exchange HPLC was analyzed. Extracts from a strain containing construct B were incubated with [³²P]ADP-ribose polymers (56), and a portion was analyzed by anion exchange HPLC as described. The elution times for AMP, ADPR, and PR-AMP are indicated by arrows. The material analyzed was PARG expressed in *E. coli*. The results indicated that the material released from ADP-ribose polymers is exclusively ADP-ribose by strong anion exchange HPLC (FIG. 9), demonstrating that the cell extracts did not contain any other

ADP-ribose polymer-degrading enzymes such as phosphodiesterase, which catalyzes the formation of AMP and phosphoribosyl-AMP (57).

Anion exchange HPLC utilized a Whatman Partisil SAX column equilibrated with 7 mM potassium phosphate buffer pH 4.0, at a flow rate of 1 ml/min. The sample was diluted in the same buffer, applied to the column, and eluted with a 30-min linear gradient from 7 mM potassium phosphate buffer, pH 4.0 to 250 mM potassium phosphate buffer, 0.5 M KCl, pH 4.0.

To determine the size of the expressed enzymatic activity, an activity gel assay (58) was used. Activity gel assays for bPARG were done by casting polyacrylamide gels with auto-modified PARP containing [³²P]ADP-ribose polymers as described previously (59). Following electrophoresis, PARG was renatured by incubating the gels at 25° C. in 5 volumes of 50 mM sodium phosphate buffer, pH 7.5, 50 mM NaCl, 10% glycerol, 1% Triton X-100, 10 mM β-mercaptoethanol, changing the buffer every 3 h for a total of five changes. After an additional incubation at 37° C. for 3 h, gels were dried, and PARG activity was detected following autoradiography as a clear band on a black background. Cell extracts containing PARG fused to GST were examined for binding to glutathione-Sepharose 4B (GSH-Sepharose) (Pharmacia Biotech Inc.) according to the specifications of the manufacturer. No bands were produced from extracts from the IPTG-induced pTrcHisB vector that did not contain an insert. Extracts from cells transformed with a construct containing a PARG insert showed bands at approximately 15 and 59 kDa (FIG. 8). During storage at 4° C., cell extracts lost activity migrating at the higher molecular weight, while the activity at approximately 59 kDa increased.

Expression of bPARG in the pTrcHisB expression vector did not result in detectable amounts of protein by staining the Coomassie Blue. Thus, another construction was designed to overexpress a 69-kDa carboxyl-terminal region of the PARG as a fusion with GST, which allows convenient protein purification by affinity chromatography on a GSH-Sepharose column. Two hours after induction with IPTG, strong expression of a protein migrating at approximately 90 kDa was observed. This protein bound to GSH-Sepharose and was eluted by GSH. The construct contained a thrombin cleavage site between the GST and the 69-kDa region of PARG, and the treatment of the material bound to GSH-Sepharose with thrombin resulted in the release of a protein that migrated at approximately 59 kDa. This result suggests that the protein purified from the bovine thymus may be larger than suggested by its migration on SDS-PAGE. The result of this experiment is presented in FIG. 10. Lane 1 shows extract from uninduced cells; lane 2 shows extract from cells induced with 1 mM IPTG for 2 hours; lane 3 shows proteins in extracts from cells shown in lane 2 that bound to GSH-Sepharose; lane 4 shows material released from GSH-Sepharose by treatment with thrombin.

In addition to the GST fusion construct described above, several other GST fusion proteins have been made. FIG. 11 shows the portions of the bovine PARG gene that have been expressed. The top line represents the structure of bovine PARG mRNA containing the open reading frame encoding the 111 kDa PARG protein. The different parts of PARG that have been cloned in expression vectors are represented with the size of the resulting expressed recombinant proteins. The expression of the 65 kDa catalytic domain of PARG (starting at the amino acid MNDV) in pGEX-2T as a fusion protein with glutathione-S-transferase (29 kDa) is detailed. Among the constructs, only the clone designed to express a protein of 69 kDa starting at amino acid +380 from the sequence of

31

bovine PARG (bPARC_{MNDV}) allowed high level expression as a fusion protein with glutathione-S transferase (GST). A 1.8 kb PCR EcoRI fragment encoding for the 65 kDa catalytic domain of PARG was cloned into the EcoRI site of pGEX-2T giving pGEX-2T-bPARC_{MNDV}. This construction results in the expression of a fused polypeptide consisting of the sequence of GST. Amino acids derived from the polylinker and thrombin site and the 65 kDa domain (FIG. 12).

In addition to various constructs designed to express PARG in *E. coli*, a recombinant baculovirus expressing a functional PARG has been constructed using the methodology of Summers and Smith as set out in U.S. Pat. No. 4,879,236 which is specifically incorporated herein by reference.

bPARC_{MNDV} was cloned in baculovirus transfer vector pVL1393 using the EcoRI site. The recombinant vector was constructed as follows. An insert containing the cDNA sequence from position 1138-2946 of bovine PARG was prepared by subcloning a 1.8 kb EcoRI fragment amplified from pWL30 using oligonucleotides CCAATTTGAAGGAG-GAATTCGCCGCCACCATGAATGATGT-GAATGCCAAACG ACCTGGA (SEQ ID NO: 34) and GCGTCTAGAATTCACCTGGCTCCTCAGGC (SEQ ID NO: 31, WIN15). The resulting fragment was inserted into the EcoRI site of the pVL1393 baculovirus transfer vector. The amplification introduced a Kozak consensus sequence (gaattccgccgccaccatgaa SEQ ID NO: 35) at the start site of translation to enhance expression of the recombinant protein. The resulting recombinant plasmid was cotransfected with linearize Baculogold™ baculovirus DNA (Pharming, San Diego, Calif.) into SF9 cells according to the manufacturers instructions. Recombinant viruses isolated using standard techniques. Overexpression of the recombinant protein was confirmed by Western blot and the results displayed in FIG. 13 demonstrate that the 65 kDa domain expressed in *E. coli* contained enzymatic activity (lane 2) migrating with the same apparent molecular weight as the enzyme purified from bovine thymus (lane 1). Likewise, a construct expressing bPARC_{MNDV} domain in SF9 insect cells infected with recombinant baculovirus showed activity (lane 4) migrating with the same apparent molecular weight.

Example 6

Northern Blot Analysis

An surprising feature of the consensus full-length cDNA clone was that it predicted expression of a protein of approximately 111 kDa (FIG. 3, SEQ ID NO: 1, and SEQ ID NO: 2), while the enzymatically active PARG from thymus had a molecular weight of approximately 59 kDa (FIG. 2). To determine the size of the RNA transcript for PARG, total RNA and poly(A)+ RNA were isolated from bovine kidney (MDBK) cells and annealed using Clone 4 as the hybridization probe.

Total cytoplasmic RNA and poly(A)+ RNA were isolated from bovine kidney MDBK cells (ATCC #CCL22) using TRIzol reagent (Gibco/BRL) following the manufacturer's recommendations. After the RNA was fractionated, it was then transferred to nylon membranes and hybridized with Clone 4 (FIG. 3) radiolabeled by a random hexamer priming method (21). The results are presented in FIG. 4. Total RNA (5 µg, lanes 1A and 1B) and poly(A)+ RNA (4 µg, lanes 2A and 2B) were separated on a denaturing agarose gel (60). Panel A shows the ethidium bromide stained gel and panel B shows the autoradiogram of a Northern blot analysis using a random primed, ³²P-labeled DNA probe constructed from Clone 4 (FIG. 3). A single transcript of approximately 4.3 kb

32

was detected in the poly(A)+ RNA (FIG. 4, lane 2). Thus, the transcript size was consistent with the expression of a 111 kDa PARG protein.

Example 7

Southern Blot Analysis of PARG Genomic Complexity

Previous studies have reported that PARG isolated from nuclear fractions had a molecular weight of approximately 75 kDa (61), while PARG isolated from whole cell homogenates or postnuclear supernatant fractions had a molecular weight of approximately 59 kDa (62). These results suggest that either two or more genes may code for PARG or that proteolysis generates lower molecular weight forms from higher molecular weight forms. The cDNA isolated encoded a protein considerably larger than any PARG proteins previously described, consistent with the possibility that the different forms of PARG are derived from a single form by proteolytic cleavage. To test the hypothesis that PARG is encoded by a single copy gene, the genomic complexity of the PARG gene was analyzed by a Southern hybridization experiment.

Total genomic DNA was prepared from bovine thymus tissue as described previously (63) and DNA (10 µg) was digested with EcoRI, BglII, XbaI or PstI, fractionated on a 1% agarose gel, transferred to a nylon membrane (Hybond N+, Amersham), and hybridized using an 828 bp HindIII fragment of Clone 1 radiolabeled as described for clone 4 above (64). Pre-hybridizations and hybridizations were carried out at 42° C. in 50% formamide, 0.25 M sodium phosphate buffer, pH 7.2, 0.25 M NaCl, 7% SDS, 1 nM EDTA. The blot was annealed with a ³²P-labeled DNA probe corresponding to the carboxyl terminal region of the PARG protein.

The results of the Southern blot analysis are presented in FIG. 7. Genomic DNA was digested with four different restriction enzymes, EcoRI (lane 1), BglII (lane 2), XbaI (lane 3) and PstI (lane 4), none of which cleave within the carboxyl terminal region of the PARG cDNA. Following electrophoresis, the restriction digests were subjected to hybridization with a probe that corresponded to the carboxyl terminal region of the PARG cDNA. The analysis displayed in FIG. 7 shows that, in each restriction digest, the probe hybridized primarily with a single restriction fragment. The fainter signals likely reflect the presence of introns in the PARG gene. This result indicates that PARG is encoded by a single copy gene in the bovine genome.

Example 8

Isolation and Characterization of PARGs from Other Species

The isolation and characterization of bovine cDNA encoding poly(ADP-ribose)glycohydrolase (PARG) has been described above. Using the information provided by the sequencing of bovine PARG, various tools were used, including public sequence databases searches and screening of cDNA libraries using PARG specific probes, to clone and sequence the cDNA and determine the primary structure of PARG from human, mouse, *Drosophila* and *Caenorhabditis elegans*. Mammalian sequences newly obtained using this combined strategy show high sequence similarity to bovine PARG (bPARG), whereas the sequences of *Drosophila* and *C. elegans* only display significant homologies in the region responsible of the catalytic activity of the protein.

The strategy followed to obtain cDNAs coding for proteins with sequence similarity to bovine PARG is summarized in FIG. 14. dBEST, GenBank, SwissProt and PIR databases were searched for PARG like-sequences at the nucleotide or amino acid level using the programs BLASTn, TBLASTn (Altschul et al., 1990) respectively, available at the NIH site on the Worldwide Web, and also included in the sequence analysis package from the Genetic Computer Group, Inc. (GCG) (Madison, Wis.), version 9.1. Both programs perform pair-wise sequence comparisons on multiple nucleotide or amino acid sequences. PARG multiple sequence comparisons obtained with these programs are very similar. Box-shading of the amino acids in the multi-sequence alignment was obtained using the program BOXSHADE (K. Hofmann and M. D. Baron). The first step involved extensive searching for sequences with bPARG similarity in various databases. As a result of this search several partial nucleotide sequences sharing extensive homologies with bPARG cDNA were obtained from the dBEST database (65). These sequences were the result of random cloning and sequencing of partial cDNAs clones obtained from mRNAs expressed in various tissues and organisms. Among them, partial cDNAs coding for PARG from human and mouse were available. One of these human clones was particularly interesting as its sequence (2500 bp long) overlapped the coding sequence of bovine PARG from aa470 to aa977 (Carboxy terminus end) and contained all the 3' untranslated region of the human PARG cDNA. This clone (No. 50859; GenBank accession number: H17209) was requested and freely obtained from the IMAGE Consortium (in collaboration with Washington University School of Medicine in St. Louis, Mo. and Merck & Co., info@image.llnl.gov). The sequence of the clone was then completed. This partial cDNA permitted design of a radiolabeled probe (fragment HindIII—KpnI of 677 bp) specific to human PARG (SEQ ID NO: 36).

Example 9

Cloning and Sequencing

The cloning procedures used in this work generally known and are also described in details in the book, *Molecular Cloning: A Laboratory Manual* (Maniatis et al., 1982). DNA sequencing was performed using the dideoxynucleotide method of Sanger (Sanger et al., 1977). Chemical reagents were purchased from Sigma (St. Louis, Mo.). Restriction enzymes, T4 DNA ligase were from New England Biolabs, Inc. (Beverly, Mass.), T7 DNA polymerase Sequenase from US Biochemical (Cleveland, Ohio), Calf Intestine Phosphatase from Boehringer, Mannheim (Indianapolis, Ind.). The phagemid pTZ18/19R is from Pharmacia (Piscataway, N.J.). The labeled nucleotides α -[³⁵S]-dATP and α -[³²P]-dCTP were purchased from ICN (Costa Mesa, Calif.). Human thymus and murine liver 5'-stretch cDNA libraries cloned in the vector λ gt 10 were from Clontech (Palo Alto, Calif.).

A single, isolated colony of C600Hfl *E. coli* strain was picked and grown in 5 ml of Luria-Bertani medium (LB)+10 mM MgSO₄+0.2% maltose overnight at 37° C. in a shaker. The bovine library lysate was diluted 1:250,000 and incubated with the C600Hfl bacterial overnight culture and 1 \times lambda dilution buffer. Next, LB soft top agar+10 mM MgSO₄ was added, and the entire mixture was quickly poured onto 90 mm LB agar+10 mM MgSO₄ plates. The plates were cooled briefly at room temperature to allow the inoculum to soak into the agar before they were incubated at 37° C. for 6-7 hr. The number of clear plaques was counted to determine the titer.

Plaques containing the entire library that had been plated were transferred to nitrocellulose or nylon membranes. The filters were then washed in a 1.5 M NaCl/0.5 M NaOH solution to lyse the cells. This was followed by a 5 min wash in neutralizing solution (1.5 M NaCl/1 M Tris buffer pH 8). Finally, the filters were rinsed in 0.2 \times SSPE (30 mM NaCl/2 mM sodium phosphate buffer pH 7.2/0.2 mM EDTA) (Sambrook et al., 1992). The filters were then dried and baked in a 80° C. oven for 2 hr to fix the lysed plaques onto the filters.

Radioactive probes were prepared using a random hexamer priming method. Pre-hybridizations and hybridizations were carried out at 42° C. in 50% formamide, 0.25 M sodium phosphate buffer, pH 7.2, 0.25 M NaCl, 7% SDS, 1 mM EDTA.

Example 10

Specific Methods Used for Library Screening

All the cloning procedures used in obtaining the additional PARG cDNAs and determining their sequences were performed essentially as described for the bovine PARG cDNA and sequence. Human thymus and murine liver 5'-stretch cDNA libraries cloned in the vector λ gt 10 were from Clontech (Palo Alto, Calif.).

Library plating and titering: A single, isolated colony of C600Hfl *E. coli* strain was picked and grown in 5 ml of Luria-Bertani medium (LB)+10 mM MgSO₄+0.2% maltose overnight at 37° C. in a shaker. The library lysate was diluted 1:250000 and incubated with the C600Hfl bacterial overnight culture and 1 \times lambda dilution buffer. Next, LB soft top agar+10 mM MgSO₄ was added, and the entire mixture was quickly poured onto 90 mm LB agar+10 mM MgSO₄ plates. The plates were cooled briefly at room temperature to allow the inoculum to soak into the agar before they were incubated at 37° C. for 6-7 hr. The number of clear plaques was counted to determine the titer.

Plaque lifts: Plaques containing the entire library that have been plated are transferred to nitrocellulose or nylon membranes. The filters are then washed in a 1.5 M NaCl/0.5 M NaOH solution to lyse the cells. This is followed by a 5 min wash in neutralizing solution (1.5 M NaCl/1 M Tris buffer pH 8). Finally, the filters are rinsed in 0.2 \times SSPE (30 mM NaCl/2 mM sodium phosphate buffer pH 7.2/0.2 mM EDTA) (66). The filters are then dried and baked in a 80° C. oven for 2 hr to fix the lysed plaques onto the filters.

Making a radioactive probe and Hybridizations: Radioactive probes were prepared using a random hexamer priming method. Pre-hybridizations and hybridizations were carried out at 42° C. in 50% formamide, 0.25 M sodium phosphate buffer, pH 7.2, 0.25 M NaCl, 7% SDS, 1 mM EDTA. This partial cDNA allowed to design a radiolabeled probe (fragment HindIII—KpnI of 750 bp long) specific to human PARG.

Example 11

Screening of a Human Thymus 5'-stretch cDNA Libraries

Multiple screenings of a human thymus 5'-stretch cDNA library were performed to complete the cloning of human PARG cDNA. For each screening a new probe was designed and used to screen approximately one million recombinants of the library. During each round of screening, overlapping clones were isolated at high stringency conditions and sub-cloned into the EcoRI site of pTZ18/19R phagemid using

35

standard techniques. The different positive clones (J5, C, E1, E2, M, M', M'', P', P'', Of, O2) were characterized by restriction analysis, subcloned into the appropriate restriction sites of pTZ18/19R as necessary and sequenced in both strand using the dideoxynucleotide method. The probe used to complete the cloning of the human cDNA library is shown is SEQ ID NO: 37. Finally, a full-length cDNA sequence was assembled which encodes the human PARG. The sequence of the cDNA encoding human PARG is presented in the sequence listing as SEQ ID NO: 3 and the amino acid sequence of human PARG is presented in the sequence listing as SEQ ID NO: 4.

The human PARG sequence shares extensive amino acid sequence homologies with bovine PARG with more than 89% identity. The sequence similarity is also high at the nucleotide level particularly in the region coding for the protein (174ATG-TGA3104). Surprisingly the 5'-untranslated region of the human sequence displays a completely different sequence with an extensive sequence similarity with highly repeated polymorphic DNA sequences found in the human genome such as Alu repetitive elements or variable number of tandem repeats (VNTR).

Example 12

Screening of Mouse Liver 5'-stretch cDNA Libraries

To isolate a PARG cDNA from the mouse liver cDNA library, a probe was designed from the human cDNA clone coding for PARG. Analysis of the bovine and human sequences revealed that PARG was highly conserved between these two species, suggesting that it might also be conserved in the mouse. Based on the restriction map of the human cDNA clone, a region in the human clone was selected, located where the active site of the protein is encoded, that exhibited near identity to its counterpart in the bovine clone. This region, consisting of approximately 800 bases, was excised from the entire human clone by digestion with the restriction endonuclease, HindIII, then purified by agarose gel separation and radiolabeled by random priming.

This probe was used to screen a mouse liver 5'-stretch cDNA library. One clone consistently hybridized with the probe. After two rounds of screening to ensure the purity of the clone the 2.5 kb insert was subcloned into the plasmid pTZ19R and sequenced. Comparison with the sequence of bovine and human PARG showed that this clone had the partial sequence that has extensive similarities to the two other mammalian sequences covering almost entirely the coding region from nucleotide -10 to a few nucleotides from the end of the coding region. A second screen was performed to obtain the missing part of the cDNA using a radioactive probe specifically designed to hybridize with the region the most 3' of the previous clone to increase the chance to get the missing part of the cDNA.

With this new probe, the same mouse liver cDNA library was screened to obtain a second clone, containing an insert that was about 3 kb. This clone was purified, subcloned and sequenced. The sequence showed that this second clone starts at amino acids 634, extends toward the stop codon to approximately 900 nucleotides into the 3' non-coding region.

A search of the dBEST database turned up one significant match to a 400 bp fragment cloned from mouse muscularis. This fragment had an exact match to the very tail end of the second clone and exceeded it by 34 bases. This extra extension contained the oligo A sequence as well as the polyadenylation signal. Because there was an exact match, the cDNA sequence was completed using this information coming from

36

the database. The complete cDNA sequence of murine PARG is presented in the Sequence Listing as SEQ ID NO: 5 and SEQ ID NO: 6.

Example 13

Obtaining the *Drosophila* PARG cDNA

Among the clones obtained from DNA databases searches were several clones from the *Drosophila* genome sequencing project (European *Drosophila* Genome Sequencing Consortium) as well as the *Drosophila* expression sequence TAG sequencing project (67). The EST clone was requested from the University of California Berkeley and obtained. Because the sequence published in the dBEST database was only partial, its sequence was completed in our laboratory and compared to a genomic sequence, part of the distal X chromosome of *Drosophila melanogaster* submitted by Murphy et al., August 1997 which presumably contains the gene of *Drosophila* PARG. The 768 aa shares less homologies with only 40% identity (48% similarity) mainly located in the catalytic domain of the protein. The domain organization of the protein is also very different with an unknown domain of 20 kDa located Carboxy terminus of the highly conserved active domain. (See FIG. 15). The sequence of the cDNA encoding the *Drosophila* PARG is presented in the Sequence Listing as SEQ ID NO: 7 and the amino acid sequence of the *Drosophila* PARG is presented in the Sequence Listing as SEQ ID NO: 8.

Example 14

Obtaining the *C. elegans* PARG Sequence

This sequence has been obtained by searching the GenBank database with the mammalian PARG protein sequence. A sequence with PARG similarity was found in the cosmid F20C5 (Accession number: Z68161, SEQ ID NO: 38) derived from the *C. elegans* genomic DNA (68). The overall sequence conservation (726aa, MW 83129 Da) with the other PARG sequences is as follows: 32% similarity and 22% identity with the mammalian PARG and 39% similarity and 30% identity with the *Drosophila* PARG. The sequence is presented in the Sequence Listing as SEQ ID NO: 38 (Genbank accession number CEF20C5). SEQ ID NO: 38 contains 12 exons as follows: exon 1 from 3591 to 3635; exon 2 from 3681 to 4121; exon 3 from 5065 to 5235; exon 4 from 5930 to 6152; exon 5 from 6200 to 6267; exon 6 from 7246 to 7338; exon 7 from 7386 to 7553; exon 8 from 7738 to 7853; exon 9 from 8153 to 8435; exon 10 from 8487 to: 8610; exon 11 from 8662 to 8952; and exon 12 from 9383 to: 9540. The coding sequence of the CePARG protein, which is publicly available from Accession number: Z68161, is referred to in the Sequence Listing as SEQ ID NO: 9. Its corresponding amino acid sequence is referred to in the Sequence Listing as SEQ ID NO: 10. The amino acid sequence of the *C. elegans* PARG is presented on the alignment (FIG. 16)

Example 15

Cloning and Overproduction of the Carboxyl-terminus 69 kDa Domain of Bovine PARG (bPARG) in *E. coli*

As described, above, bovine PARG is encoded by a messenger of 4 kb predicting a protein of 110 kDa, almost twice the size of the purified enzyme (65 kDa). It is also demon-

strated that bPARG can be expressed in *E. coli* as an active enzyme either as a 110 kDa or a 65 kDa protein. This result combined with other evidence implies that the active site of PARG is located in the carboxyl-terminal part of the protein. FIG. 11 is a schematic representation of the different clones we have expressed in bacteria. Among them, only the clone designed to express a protein of 69 kDa starting at the amino acid +380 from the sequence of bovine PARG (bPARG_{MNDV}) allowed high level expression as a fusion protein with glutathione-S transferase (GST).

The heterologous expression of bPARG_{MNDV} was conducted as represented in FIG. 12. The 1.8 kb cDNA encoding the 69 kDa carboxyl-terminal part of bovine PARG was amplified by PCR and cloned in the EcoRI site-of pGEX-2T vector (Pharmacia) in fusion with GST giving the pGEX-2T-bPARG_{MNDV} plasmid. *E. coli* NM522 cells transformed with the pGEX-2T-bPARG_{MNDV} were induced by addition of IPTG, resulting in expression of a 90 kDa fusion protein. The fusion protein can be conveniently purified using Glutathione-Sepharose and the bPARG_{MNDV} can be released by treatment with thrombin while the GST protein remains bound to the beads of GSH-Sepharose. In this manner milligram amounts of protein can be routinely obtained.

Example 16

Characterization of the Purified 65 kDa Domain and the Generation of Antibodies

The purified bPARG_{MNDV} was characterized by activity gel assays (69) by casting polyacrylamide gels with automodified PARP containing [³²P]ADP-ribose polymers. The results demonstrate that the 65 kDa domain expressed in *E. coli* contained enzymatic activity migrating with the same apparent molecular weight as the enzyme purified from bovine thymus. Likewise, a construction expressing bPARG_{MNDV} domain in SF9 insect cells infected with recombinant baculovirus showed activity migrating with the same apparent molecular weight.

The availability of PARG cDNA allows the development of new molecular tools to study this enzyme in its cellular context. Until this work, it was not possible to obtain PARG in sufficient quantities to produce antibodies against the protein. The antibody raised against bovine PARG is able to recognize PARG from other organisms and, thus, will be valuable in characterizing PARG in vivo under defined physiological conditions in many different organisms.

Antibodies against bPARG_{MNDV} overexpressed in *E. coli* were raised in rabbits using the procedure described by Vaitukaitis (70). Specific high affinity antibodies are generated by administration of small doses of immunogens intradermally over a wide anatomic area of the animal. Rabbits were immunized by three injections of 10-50 µg of the Mr 65,000 protein band excised from a preparative SDS polyacrylamide gel. Titer and affinity of sera harvested weekly were followed by conventional methods. Peak affinity was attained in 8 to 10 weeks after primary immunization. For each animal, a pre-immune serum was retained as a control.

FIG. 17 shows a Western blot experiment demonstrating the specificity of the resulting. PARG anti-serum against the purified bPARG from thymus (lane 1), SF9 protein extract expressing 65 kDa-bPARG_{MNDV} in recombinant baculovirus (lane 2), recombinant 65 kDa-PARG_{MNDV} purified by treatment with thrombin from GSH-Sepharose (lane 3), and an *E. coli* crude extract expressing the fusion protein GST-65 kDa-

PARG_{MNDV} (lane 4). The pre-immune serum did not show reactivity against any of these fractions even at a low dilution (1/250).

Antibodies directed against the 45 kDa -terminal have also been generated using the same strategy used to generate antibodies against the catalytic domain. This involved the over-expression of the 45 kDa protein domain in *E. coli* in a construct designed for easy purification, followed by injection of the purified protein into rabbits. The heterologous expression of PARG45 was conducted by cloning a part (1.1 kb) of the coding region of the cDNA, generated by PCR amplification of the region located between the ATG(267) codon and nucleotide 1400 in the bovine sequence, into the Eco RI site of the bacterial expression vector pGEX-2T (Pharmacia) in fusion with glutathione-S-transferase. *E. coli* NM522 cells transformed with this construct were induced by addition of IPTG, resulting in expression of a 72 kDa fusion protein. The fusion protein was purified using glutathione Sepharose and the PARG45 was released by treatment with thrombin, while the GST protein remained bound to the GSH Sepharose beads. In this manner milligram amounts of protein were obtained. Antibodies against PARG45 overexpressed in *E. coli* were raised in rabbits using the procedure described (71). Specific high affinity antibodies were generated by administration of small doses of immunogens subcutaneously over a wide area of the animal. Rabbits were immunized by three injections of 10-50 µg of the 45 kDa protein band excised from a preparative SDS-polyacrylamide gel. Titer and affinity of sera harvested weekly were followed by conventional methods. Peak affinity was attained in 8 to 10 weeks after primary immunization. For each animal, a pre-immune serum was retained as a control.

Example 17

Conservation of PARG in Tissues and Organisms

Tissue and cell extracts from different origins were homogenized in a cold hypotonic lysis buffer containing a cocktail of protease inhibitors and sonicated. SDS and β-mercaptoethanol were added to insure inactivation of any remaining active proteases. Thirty µg of protein from each extract was analyzed by Western-blot using the anti-PARG antibody (FIG. 18). In all of the fractions from bovine tissues, PARG was observed as a major band at 65 kDa. However, less intense, discrete proteins of higher molecular weight were also detected. These proteins may correspond to different forms of PARG; the band of highest molecular weight (about 115 kDa) found in thymus extract likely corresponds to the full-length of PARG (111 kDa) as deduced from the cDNA. Multiple species were detected in cell extracts from mouse fibroblasts, rat PC 12 cells, and SF9 insect cells. This result shows that the sequence of PARG is well conserved phylogenetically. Moreover, the conservation includes multiple molecular forms of the protein.

Example 18

Regulation of the Expression of PARG

In the metabolism of ADP-ribose polymers, the activities of PARP and PARG are closely related. Soon after polymer has been synthesized by PARP following DNA damage, it is extensively degraded by PARG. The net result is that the polymer has a very short half life. The close relationship between the two proteins suggests a possible mode of regulation in which PARG expression depends on the presence of

PARP. In order to test if the presence or the absence of PARP influences the expression of PARG, a Western Blot experiment was performed with cell extracts from mouse fibroblasts of different PARP genotypes (72).

Cell extract (30 µg) from mouse cells with PARP+/, PARP+/- and PARP-/- genotypes were separated by SDS-PAGE, transferred to a membrane and probed with the antibodies indicated. The results are shown in FIG. 19. In FIG. 19, purified PARG (50 ng) from bovine thymus (lane 1), 30 µg of protein of a total extract from PARG recombinant baculovirus infected SF9 cells (lane 2), 150 ng of purified recombinant PARG produced in the bacteria (lane 3) and 30 µg of protein of a crude extract from *E. coli* NM522 transformed with pGEX-2T-bPARG_{MNDV} 2 h after induction by IPTG (lane 4) were separated on a 0.1% SDS-12% polyacrylamide gel, then transferred on nitrocellulose, and incubated with a 1/5000 dilution of the rabbit polyclonal antiserum raised against the 65 kDa domain of bPARG. Proteins were revealed by immunofluorescence with the ECL detection kit (Amersham) and autoradiography. Panel A is a western blot of PARP in cells of varying PARP genotype showing the results of the analysis using anti-PARP antibodies. The amount of PARP expressed varies as expected dependant upon the genotype of the cell line with the PARP-/- cell line producing no detectable amount of PARP. Panel B is western blot of PARG from various tissues using an anti-PARG antibody. It shows that the level of PARP is variable. The amount of PARG present in the cell extracts was not dependent upon the PARG genotype of the cell. Further support for this view is provided by the results of the PARG activity assay presented in panel C. The specific activity of PARG detected in the extracts showed no significant difference among the three genotypes.

Example 19

Preparation PARG Gene Ablation (Knockout) Animals

One embodiment of the present invention is experimental animals with targeted mutations in the PARG gene. These animals may be constructed using standard techniques and the cDNA sequence of the PARG. In the following example, a mouse containing a targeted mutation in the PARG gene is constructed. Those skilled in the art will readily appreciate that other experimental animals, including but not limited to rats, guinea pigs, hamsters and the like, may be constructed using similar techniques. The construction of animals with disrupted genes may be accomplished using standard techniques such as those described by Moreadith (73). Further, cells lines, construction kits, and protocols for knockout mice are available from commercial suppliers such as Stratagene (Stratagene 1999 catalog, La Jolla, Calif.). Commercial services such as Lexicon Genetics (The Woodlands, Tex.) and Chrysalis DNX Transgenic Sciences (Princeton, N.J.) also offer complete ES cell knockout mice production services.

A genomic clone of the murine PARG enzyme may be isolated from a genomic library by screening with a probe derived from the cDNA sequence of PARG. A mouse 129/SV genomic library (Stratagene) containing mouse genomic sequences in λ phage was screened using a 2.49 kb fragment of the mouse PARG cDNA as a probe. A partial restriction map of one positive clone thus isolated, R1, is provided in FIG. 20. The R1 clone contains the genomic sequence corresponding to the 5'-most end of the murine cDNA. The clone was subcloned into pBluescript as three fragments. The plasmid containing the 5'-end contained a 2.8 kb NotI-EcoRI fragment and was designated p2.8R. The fragment containing the central portion contained a 3.5 kb EcoRI fragment and was designated p3.5R. The plasmid containing the 3'-end of the gene contained a 7.0 kb EcoRI-NotI fragment and was

designated 7.0R. Sequencing the resulting plasmids revealed that p2.8R contained no sequences corresponding to the cDNA, p3.5R contains a 1.5 kb promoter region and/or untranslated region and exon 1 coding for 72 amino acids including the initiation ATG codon and p7.0R contains at least 4 additional exons. Gene targeting vectors may be constructed using both p3.5R and p7.0R.

A gene targeting vector may contain one or more selection genes flanked by genomic sequences. The targeting vector is introduced into the genome by homologous recombination resulting in the incorporation of the selection gene into the genome of the cell. The mouse PARG gene was targeted using a "conditional" inactivation procedure outlined in FIG. 21. This approach allows the production of viable animals even if the disrupted gene results in a lethal phenotype since the gene is not disrupted until a second "conditional" recombination event is induced.

A lox-P sequence may be inserted into the first intron. A cassette expressing the neomycin resistance gene (neo) and the thymidine kinase gene (TK) flanked by two additional lox-P sites may be placed in intron 2. In the presence of Cre recombinase, recombination will occur between two lox-P sites thereby deleting the genomic sequences present between the sites. A MC1-DTA cassette is ligated at the 3'-end of the vector to reduce random integration of the vector into the genome.

The targeting vector may be introduced into embryonic stem cells by any method known to those skilled in the art such as transfection, lipofection or electroporation. In a preferred embodiment, the targeting vector will be introduced into embryonic stem cells by electroporation. After homologous recombination, cells containing the neo gene will be selected for using G418. Selected cells will then be analyzed by PCR and Southern blot.

To generate mutant alleles of PARG, the positive embryonic stem cell clones identified will be transfected with a plasmid expressing Cre recombinase. The action of Cre recombinase will result in three different mutant alleles. Mutant allele I contains a deletion in exon 2 but still maintains the selection genes neo and TK. Mutant allele II contains the genomic sequence for exon 2 flanked by two lox-P sites (exon 2 is said to be "floxed") and does not contain the selection genes. Mutant allele III has a deletion of the genomic sequences and does not contain the selection genes.

Mice containing each of the three mutant alleles may be constructed by microinjecting embryonic stem cells containing the mutant allele into blastocytes resulting in the production of chimeric and mutant mice. Mice homozygous in mutant allele I or III will be null mutants in that they will be unable to express a functional PARG enzyme due to the loss of required genomic sequences. In the absence of Cre recombinase, mice containing mutant allele II will express a wild type protein. In the presence of the Cre recombinase, the PARG will lose exon 2, thus producing an inactive protein. To inactivate the gene, these mice will be bred to mice expressing Cre recombinase under the control of a tissue specific promoter. This will result in mice expressing PARG in some tissues and not expressing PARG in others. Mice homozygous in mutant allele II, will be valuable for evaluating the role of PARG in specific tissues.

Although the present invention has been described with reference to certain examples for purposes of clarification and illustration. It should be appreciated that certain obvious improvements and modifications can be practiced within the scope of the appended claims and their equivalents. Other embodiments and uses of the invention will be apparent to those skilled in the art from consideration of the specification

and practice of the invention disclosed herein. All U.S. patents GenBank sequence listings, and other references noted herein for whatever reason are specifically incorporated by reference. The specification and examples should be considered exemplary only with the true scope and spirit of the invention indicated by the following claims.

1. Miwa, M. et al. (1971) *J. Biol. Chem.* 246, 6362-6364;
- Ueda, K. et al. (1972) *Biochem. Biophys. Res. Commun.* 46, 516-523
2. Oka, J. et al. (1984) *J. Biol. Chem.* 259, 986-995.
3. Gaal, J. et al. (1987) *Trends in Biol. Sci.*, 12, 129.
4. Nudka, N. et al. (1980) *Eur. J. Biochem.* 105, 525-530; Jacobson, E. et al. (1985) *Carcinogenesis* 6, 715-718; Küipper, J. et al. (1990) *J. Biol. Chem.* 265, 18721-18724; Ding, R. et al. (1992) *J. Biol. Chem.* 267, 12804-12812.
5. Jacobson, E. et al. (1985) in *ADP-ribosylation of Proteins* (Althaus, F. R., Hilz, H., and Shall, S., eds) pp. 277-283, Springer-Verlag, Berlin; Lubet, R. et al. (1986) *Carcinogenesis* 7, 71-75; Kasid, U. et al. (1986) *Carcinogenesis* 7, 327-330.
6. Berger, N. (1985) *Radiat. Res.* 101, 4-15.
7. Kaufmann, S. et al. (1993) *Cancer Res.* 53, 3976-3985; Lazebnik, Y. et al. (1994) *Nature* 371, 346-347.
8. de Murcia, G. et al. (1994) *Trends Biochem. Sci.* 19, 172-176.
9. BioWorld Today, Apr. 29, 1994, p. 3.
10. Thomassin, H. et al. (1990) *Nucleic Acids Res.* 18, 4691-4694.
11. Karlin, Samuel and Stephen F. Altschul (1993). *Proc. Natl. Acad. Sci. USA* 90:5873-7
12. Henikoff and Henikoff *Proc. Natl. Acad. Sci. USA* 89:10915-19, 1992
13. Altschul, Stephen F. (1991). *J. Mol. Biol.* 219:555-65.
14. Cohen, J. S., 1989; Weintraub, H. M., 1990
15. N. Sarver et al., 1990
16. Orlandi, R. et al., (1989) *Proc. Natl. Acad. Sci.* 86, 3833-3837; Winter, G et al., (1991) *Nature* 349, 293-299
17. R. Sobol and K. Scanlon eds., available at www.appleton-lange.com.
18. e.g., at least, but not limited to U.S. Pat. No. 5,797,870, U.S. Pat. No. 5,804,383, U.S. Pat. No. 5,670,161, U.S. Pat. No. 5,645,829, U.S. Pat. No. 5,741,486, U.S. Pat. No. 5,836,905, U.S. Pat. No. 5,843,069, U.S. Pat. No. 5,827,216, U.S. Pat. No. 5,871,464, U.S. Pat. No. 5,702,384, U.S. Pat. No. 5,810,888, U.S. Pat. No. 5,787,900, U.S. Pat. No. 5,752,515, U.S. Pat. No. 5,674,192, U.S. Pat. No. 5,658,955, U.S. Pat. No. 5,656,465, U.S. Pat. No. 5,547,932, U.S. Pat. No. 5,873,904, U.S. Pat. No. 5,792,651, U.S. Pat. No. 5,772,888, U.S. Pat. No. 5,641,750, U.S. Pat. No. 5,641,749, and U.S. Pat. No. 5,626,561
19. Kaltenbock, B. et al. (1998) *Biotechniques*, 24, 202-206.
20. Moran, P. et al. (1998) *Biotechniques*, 24, 206-212.
21. Orita, M. et al. (1989) *PNAS USA*, 86, 2766-2770.
22. Liu, Q. et al. (1998) *Biotechniques*, 24, 140-147.
23. *Proceeding from the Sixth International Symposium on Human Identification* 1995 (ISBN 1-882274-55-5).
24. Kohler and Milstein, *Nature*, 256:495-497 (1975).
25. Geysen, et al. (1984) *Proc. Natl. Acad. Sci. USA* 81:3998-4002.
26. Ménard, et al. (1987) *Biochem. Cell Biol.* 65, 668-673.
27. Hatakeyama, K. et al. (1986) *J. Biol. Chem.* 261, 14902-14911.
28. Thomassin, H. et al. (1990) *Nucleic Acids Res.* 18, 4691-4694.
29. Ménard, L. et al. (1987) *Biochem. Cell Biol.* 65, 668-673.
30. Bradford, M. (1976) *Anal. Biochem.* 72, 248-254.
31. Laemmli, U. (1970) *Nature* 227, 680-685.
32. Althaus, F. et al. (1987) *Molecular Biology, Biochemistry and Biophysics*, Vol. 37, Springer-Verlag, Berlin.
33. Mead, D. et al. (1986) *Protein Eng.* 1, 67-74.
34. Shaw, et al. (1986) *Cell* 46, 659-667.
35. Kozak (1987) *Nucleic Acids Res.* 15, 8125-8148.
36. Wilson, et al. (1994) *Nature (London)* 368, 32-38.
37. Brendel, et al. (1992) *Proc. Natl. Acad. Sci. USA* 89, 2002-2006.
38. Robbins, et al. (1991) *Cell* 64, 615-623.
39. Schreiber, et al. (1992) *EMBO J.* 11, 3263-3269.
40. Uchida, K. et al. (1987) *Biochem. Biophys. Res. Commun.* 148, 617-622, Cherney, B. et al. (1987) *Proc. Natl. Acad. Sci. U.S.A.* 84, 8370-8374; Kurosaki, T. et al. (1987) *J. Biol. Chem.* 262, 15990-15997.
41. Huppi, K. et al. (1989) *Nucleic Acids Res.* 17, 3387-3401.
42. Saito, I. et al. (1990) *Gene (Amst.)* 90, 249-254.
43. Ittel, M.-E. et al. (1991) *Gene (Amst.)* 102, 157-164.
44. Saulier-Le Drean, B. (1992) *Poly(ADP-ribose) Polymerase in Xenopus lae-vis*, Ph.D. thesis, Université De Rennes, France.
45. Uchida, K. et al. (1993) *Proc. Natl. Acad. Sci. U.S.A.* 90, 3481-3485.
46. Masutani, M. et al. (1994) *Eur. J. Biochem.* 220, 607-614.
47. Schreiber, et al. (1992) *EMBO J.* 11, 3263-3269.
48. Lazebnik, et al. (1994) *Nature (London)* 371, 346-347.
49. Brochu, et al. (1994) *Biochem. Biophys. Acta* 1219, 342-350.
50. Tanuma, et al. (1986) *J. Biol. Chem.* 261,965-969; Uchida, et al. (1993) *J. Biol. Chem.* 268, 3194-3200.
51. Maruta, et al. (1991) *Biochemistry* 30, 5907-5912.
52. Ménard, L. et al. (1987) *Biochem. Cell Biol.* 65, 668-673.
53. Ménard, L. et al. (1987) *Biochem. Cell Biol.* 65, 668-673.
54. Slama, J. et al. (1995) *J. Med. Chem.* 38,389-393; Slama, J. et al. (1995) *J. Med. Chem.* 38, 4332-4336.
55. Slama, J. et al. (1995) *J. Med. Chem.* 38, 389-393; Slama, J. et al. (1995) *J. Med. Chem.* 38, 4332-4336.
56. Ménard, L. et al. (1987) *Biochem. Cell Biol.* 65, 668-673.
57. Althaus, F. et al. (1987) *Molecular Biology, Biochemistry and Biophysics*, Vol. 37, Springer-Verlag, Berlin.
58. Brochu, G. et al. (1994) *Anal. Biochem.* 218, 265-272.
59. Brochu, G. et al. (1994) *Anal. Biochem.* 218, 265-272.
60. Moreadith, et al. (1997) *J. Mol. Med.* 75, 208-216.
61. Tanuma, et al. (1986) *J. Biol. Chem.* 261, 965-969 and Uchida, et al. (1993) *J. Biol. Chem.* 268, 3194-3200.
62. Hatakeyama, et al. (1986) *J. Biol. Chem.* 261, 14902-14911; Thomassin, et al. (1990) *Nucleic Acids Res.* 18, 4691-4694; and Maruta, et al. (1991) *Biochemistry* 30, 5907-5912.
63. Sambrook, et al. (1992) *MOLECULAR CLONING: A LABORATORY MANUAL*, Cold Spring Harbor Laboratory Cold Spring harbor, N.Y.
64. Feinberg, et al. (1983) *Anal. Biochem.* 132, 6-13.
65. Boguski, 1995
66. Sambrook, et al. (1992) *MOLECULAR CLONING: A LABORATORY MANUAL*, Cold Spring Harbor Laboratory Cold Spring Harbor, N.Y.
67. BDGP/HHMI *Drosophila* EST Project, University of California Berkeley, EST@fruitfly.berkeley.edu
68. Nematode Sequencing Project, Sanger Centre, Hinxton, Cambridge CB10 1RQ, England and Department of Genetics, Washington University, St. Louis, Mo. 63110, USA. E-mail: jes@sanger.ac.uk
69. Brochu G. et al. (1994) *Anal. Biochem.* 218, 265-272.
70. Vaitukaitis (1981) *Methods in Enzymology* 73, 46-52.
71. Vaitukaitis (1981) *Methods in Enzymology* 73, 46-52.
72. Wang, et al. (1995) *Genes & Dev.* 9, 509-520.
73. Moreadith, et al. (1997) *J. Mol. Med.* 75, 208-216.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 38

<210> SEQ ID NO 1

<211> LENGTH: 4070

<212> TYPE: DNA

<213> ORGANISM: Bos taurus

<220> FEATURE:

<400> SEQUENCE: 1

```

accggaagt gaacgaagcc cgaatcagaa cggctcatcc tgaggctggt aggggtgccg 60
tggaagaggg aaggcaggcg tctggatagg gcctggttcg ggaggctgtc agagcaggag 120
ctgcagaagc agtcagcggc agagggggca tggtgccggg aggcaccgag gagggggcgc 180
agtccgtccc tcccagggtt agtgaatgag gctctacgcc cgggctggcc cggagactca 240
gtgctgctgg tcccagcatg agtgccggcc cggctgtga gccctgcacc aagcgacccc 300
gctgggacgc cgctgcaact tctccgccgg cgcctcggga cgcccgagc tccccggca 360
ggcagaggcg cgtcctcgat tccaaggacg ctccgggtca gttcagggtc cgccttcct 420
cgtcaggctg cgccctgggc cgggcgggac agcaccgagg cagcgccacc tctctgttt 480
tcaaacagaa gactataacc agttggatgg aactaaagg aatcaagaca gttgaatcag 540
aaagtttgca tagtaaaaga aacaacaata caagagaaga atccatgatg agttctgtac 600
aaaaagataa cttttatcaa cataacatgg aaaaattaga aatgtttct cagctaggtt 660
ttgataagtc accagttgaa aaaggtacac agtatgtgaa gcagcatcag actgccccta 720
tgtgtaagtg gcagaatgaa gggccacact cagaacggct tttggaaagt gaacctccag 780
cggtaactct ggtaccagag cagttcagta atgctaagt cgatcagtcg tccccaaagg 840
atgatcacag tgacacaaat agtgaggaga gtagagataa tcagcagttt ttgacacatg 900
taaagcttgc gaatgcaaag cagacgatgg aagatgaaca gggcagagaa gccagaagcc 960
accagaagtg tggcaaggct tgccatcctg cagaagcctg tgcagggtgt cagcaggagg 1020
agacagacgt ggtgtccgag agccccttgt cggacactgg ctctgaggat gttggtactg 1080
gactgaaaaa tgccaacaga ttgaatagac aagaaagtag tctaggaaat tctcctccat 1140
ttgaaaaga aagtgaacct gagtcaccaa tggatgtaga taattccaaa aatagttgtc 1200
aggattcaga agcagatgaa gagacaagtc caggttttga tgaacaggaa gatagcagtt 1260
ctgctcaaac agcaataaaa ccttcaaggt toccaaccaag agaagctgac actgagttga 1320
ggaagcggtc ctctgctaag ggaggtgaga ttcgattaca tttccaattt gaaggaggag 1380
agagtcgagc tggaatgaat gatgtgaatg ccaaacgacc tggaagtact tctagcctga 1440
atgtagagtg cagaaattct aagcaacatg ggagaaagga ttctaaaaac acagatcatt 1500
tcatgagagt gcccacaaagc gaggacaaaa gaaaagaaca atgtgaaatg aaacatcaaa 1560
gaacagaagc gaagatccct aaatacatc cacctcacct ttctccagat aagaaatggc 1620
ttggaactcc tattgaggag atgaggagaa tgccaagggtg tgggatccgg ctgcctccct 1680
tgagaccatc tgccaatcac acagtgacta ttcgggtaga tcttttgcca ataggagaag 1740
ttctaaacc tttccaaca cattttaag atttgggga caacaagcat gttaagatgc 1800
cttggtcaga acaaaacttg tacctctgtg aagatgagaa tggtagcgga gctgcaggca 1860
gccggtggga actcattcag actgcacttc tcaacagget cactcggcc cagaacctga 1920
aggatgctat tctgaagtac aatgtggcat attctaagaa atgggacttt acagctttga 1980

```


-continued

ttgatttctg ggataaggta ctagaagaag cagaagctca acacttgat cagtccatct	2040
tgctgatgat ggtgaaaatt gcaactctgtc tgccaaatat ttgtaccag ccaataccac	2100
tcctgaaaca gaagatgaat cattccatca caatgtcaca ggaacagatt gccagtcttt	2160
tagctaatgc tttcttctgc acgtttccac gacgcaatgc caagatgaaa tcagagtatt	2220
ccagttatcc agatattaac ttcaatcggg tgtttgaagg acgttcatca aggaaaccag	2280
agaagcttaa aacgctcttc tgctacttta gaagagtcac agagaaaaa cccactgggt	2340
tggtgacatt cacaagacag agtcttgaag attttccaga gtgggaaaga tgtgaaaaac	2400
tcctgactcg actgcatgtc acttaacgaag gtacataga aggaaacggc cagggcatgc	2460
tacaggtgga ttttgcaaac cgtttcgttg gaggtggtgt aaccagtgcg ggacttgtgc	2520
aagaagaaat ccgcttttta atcaaccctg agttgattgt ttcacggctc ttcactgagg	2580
tgctggatca caatgaatgt cttatcatca caggtactga gcagtacagt gaatacacag	2640
gctatgccga aacataaccgc tggggcccgga gccatgaaga caggagcga agggacgact	2700
ggcagaggcg cagcactgag atcgtcgcca tcgacgcct ccaactcaga cgctacctcg	2760
accagttgt gcccgagaag atcagacggg agcttaacaa ggcttactgt ggatttcttc	2820
gtectggagt ttcttcagag aacctgtctg cagtggctac aggaaactgg ggctgtggtg	2880
cctttggggg tgatgctaga ctaaaagcct taatacagat cctggcagct gctgtagctg	2940
agcgagacgt ggtttatttc acctttgggg actcagaact gatgagagac atttacagca	3000
tgcatacatt cctcactgag aggaaactga ctggtggaga agtatataag ctgctgctac	3060
gatattacaa tgaagaatgc agaaactgct ccacccccgg accagacatc aagctttatc	3120
cattcatata ccatgcagtt gagtcctgta cacagaccac caaccagccg ggacaaagga	3180
cgggggectg aggagccaag tgactagacg ctccccactt gtgtaacaag aagggtgac	3240
gtgtgaactg acatgatatc catgtgtata taatccgctt ttgtaggcaa ggatgcagtc	3300
ccttccgccc atgcagctgt cagtacatct ggcctcctc catcccgact tacatagact	3360
gagacatact ttgtttcttt ttttttctat ttcagcctg attcttttat ttttctttct	3420
tttgccatc agacttcttg tgaatttca tcagagtttg tgctcagcct ggcaggtgctc	3480
ttttttgatg cctaaatata caaatcacct ctgcagctag cagatgccac ggaaggtggt	3540
ggaaccctag gagctgtaac tgagtctgtc gcagatctcc ctctgagcct ctcaccctca	3600
ccctattatc attgtggtgg tggaggtttt ttgatttttg aaataagagt tgggtttggt	3660
aaataatata gatctcctag gtttaagatt ttatatata gaatactttt caaaaagtta	3720
ttttgagata tcacctttat ttgtaatggt aatttgctg tccttttcc cctgatcaat	3780
ttgtattgac tgtttttgga aattgaccac aatgaaagga aatagagaa taagagtttc	3840
ccaaatggtg tttaaaaaca aacaggttca agacacgca aggacctcgt ttctgggat	3900
ttttttcttt tttctttttt tgaattagga ttattgtttg ttcttgggtg cttgagacat	3960
attcatataa ccaaagttaa ggaactggga acttcgtggt gattgtgaca tattgaagtt	4020
tctctggtac tcaaaggtta tgtagttaat aaattttcat taacaaaaa	4070

<210> SEQ ID NO 2

<211> LENGTH: 977

<212> TYPE: PRN

<213> ORGANISM: Bos taurus

<220> FEATURE:

<400> SEQUENCE: 2

-continued

Met	Ser	Ala	Gly	Pro	Gly	Cys	Glu	Pro	Cys	Thr	Lys	Arg	Pro	Arg	Trp	1	5	10	15
Asp	Ala	Ala	Ala	Thr	Ser	Pro	Pro	Ala	Ala	Ser	Asp	Ala	Arg	Ser	Phe	20	25	30	
Pro	Gly	Arg	Gln	Arg	Arg	Val	Leu	Asp	Ser	Lys	Asp	Ala	Pro	Val	Gln	35	40	45	
Phe	Arg	Val	Pro	Pro	Ser	Ser	Ser	Gly	Cys	Ala	Leu	Gly	Arg	Ala	Gly	50	55	60	
Gln	His	Arg	Gly	Ser	Ala	Thr	Ser	Leu	Val	Phe	Lys	Gln	Lys	Thr	Ile	65	70	75	80
Thr	Ser	Trp	Met	Asp	Thr	Lys	Gly	Ile	Lys	Thr	Val	Glu	Ser	Glu	Ser	85	90	95	
Leu	His	Ser	Lys	Glu	Asn	Asn	Asn	Thr	Arg	Glu	Glu	Ser	Met	Met	Ser	100	105	110	
Ser	Val	Gln	Lys	Asp	Asn	Phe	Tyr	Gln	His	Asn	Met	Glu	Lys	Leu	Glu	115	120	125	
Asn	Val	Ser	Gln	Leu	Gly	Phe	Asp	Lys	Ser	Pro	Val	Glu	Lys	Gly	Thr	130	135	140	
Gln	Tyr	Leu	Lys	Gln	His	Gln	Thr	Ala	Ala	Met	Cys	Lys	Trp	Gln	Asn	145	150	155	160
Glu	Gly	Pro	His	Ser	Glu	Arg	Leu	Leu	Glu	Ser	Glu	Pro	Pro	Ala	Val	165	170	175	
Thr	Leu	Val	Pro	Glu	Gln	Phe	Ser	Asn	Ala	Asn	Val	Asp	Gln	Ser	Ser	180	185	190	
Pro	Lys	Asp	Asp	His	Ser	Asp	Thr	Asn	Ser	Glu	Glu	Ser	Arg	Asp	Asn	195	200	205	
Gln	Gln	Phe	Leu	Thr	His	Val	Lys	Leu	Ala	Asn	Ala	Lys	Gln	Thr	Met	210	215	220	
Glu	Asp	Glu	Gln	Gly	Arg	Glu	Ala	Arg	Ser	His	Gln	Lys	Cys	Gly	Lys	225	230	235	240
Ala	Cys	His	Pro	Ala	Glu	Ala	Cys	Ala	Gly	Cys	Gln	Gln	Glu	Glu	Thr	245	250	255	
Asp	Val	Val	Ser	Glu	Ser	Pro	Leu	Ser	Asp	Thr	Gly	Ser	Glu	Asp	Val	260	265	270	
Gly	Thr	Gly	Leu	Lys	Asn	Ala	Asn	Arg	Leu	Asn	Arg	Gln	Glu	Ser	Ser	275	280	285	
Leu	Gly	Asn	Ser	Pro	Pro	Phe	Glu	Lys	Glu	Ser	Glu	Pro	Glu	Ser	Pro	290	295	300	
Met	Asp	Val	Asp	Asn	Ser	Lys	Asn	Ser	Cys	Gln	Asp	Ser	Glu	Ala	Asp	305	310	315	320
Glu	Glu	Thr	Ser	Pro	Gly	Phe	Asp	Glu	Gln	Glu	Asp	Ser	Ser	Ser	Ala	325	330	335	
Gln	Thr	Ala	Asn	Lys	Pro	Ser	Arg	Phe	Gln	Pro	Arg	Glu	Ala	Asp	Thr	340	345	350	
Glu	Leu	Arg	Lys	Arg	Ser	Ser	Ala	Lys	Gly	Gly	Glu	Ile	Arg	Leu	His	355	360	365	
Phe	Gln	Phe	Glu	Gly	Gly	Glu	Ser	Arg	Ala	Gly	Met	Asn	Asp	Val	Asn	370	375	380	
Ala	Lys	Arg	Pro	Gly	Ser	Thr	Ser	Ser	Leu	Asn	Val	Glu	Cys	Arg	Asn	385	390	395	400
Ser	Lys	Gln	His	Gly	Arg	Lys	Asp	Ser	Lys	Ile	Thr	Asp	His	Phe	Met	405	410	415	
Arg	Val	Pro	Lys	Ala	Glu	Asp	Lys	Arg	Lys	Glu	Gln	Cys	Glu	Met	Lys				

-continued

420					425					430					
His	Gln	Arg	Thr	Glu	Arg	Lys	Ile	Pro	Lys	Tyr	Ile	Pro	Pro	His	Leu
	435						440					445			
Ser	Pro	Asp	Lys	Lys	Trp	Leu	Gly	Thr	Pro	Ile	Glu	Glu	Met	Arg	Arg
	450					455					460				
Met	Pro	Arg	Cys	Gly	Ile	Arg	Leu	Pro	Pro	Leu	Arg	Pro	Ser	Ala	Asn
465					470					475					480
His	Thr	Val	Thr	Ile	Arg	Val	Asp	Leu	Leu	Arg	Ile	Gly	Glu	Val	Pro
				485					490					495	
Lys	Pro	Phe	Pro	Thr	His	Phe	Lys	Asp	Leu	Trp	Asp	Asn	Lys	His	Val
			500					505					510		
Lys	Met	Pro	Cys	Ser	Glu	Gln	Asn	Leu	Tyr	Pro	Val	Glu	Asp	Glu	Asn
		515					520					525			
Gly	Glu	Arg	Ala	Ala	Gly	Ser	Arg	Trp	Glu	Leu	Ile	Gln	Thr	Ala	Leu
	530					535					540				
Leu	Asn	Arg	Leu	Thr	Arg	Pro	Gln	Asn	Leu	Lys	Asp	Ala	Ile	Leu	Lys
545					550					555					560
Tyr	Asn	Val	Ala	Tyr	Ser	Lys	Lys	Trp	Asp	Phe	Thr	Ala	Leu	Ile	Asp
				565					570						575
Phe	Trp	Asp	Lys	Val	Leu	Glu	Glu	Ala	Glu	Ala	Gln	His	Leu	Tyr	Gln
			580					585					590		
Ser	Ile	Leu	Pro	Asp	Met	Val	Lys	Ile	Ala	Leu	Cys	Leu	Pro	Asn	Ile
		595					600					605			
Cys	Thr	Gln	Pro	Ile	Pro	Leu	Leu	Lys	Gln	Lys	Met	Asn	His	Ser	Ile
	610					615					620				
Thr	Met	Ser	Gln	Glu	Gln	Ile	Ala	Ser	Leu	Leu	Ala	Asn	Ala	Phe	Phe
625					630					635					640
Cys	Thr	Phe	Pro	Arg	Arg	Asn	Ala	Lys	Met	Lys	Ser	Glu	Tyr	Ser	Ser
				645					650					655	
Tyr	Pro	Asp	Ile	Asn	Phe	Asn	Arg	Leu	Phe	Glu	Gly	Arg	Ser	Ser	Arg
			660					665					670		
Lys	Pro	Glu	Lys	Leu	Lys	Thr	Leu	Phe	Cys	Tyr	Phe	Arg	Arg	Val	Thr
		675					680					685			
Glu	Lys	Lys	Pro	Thr	Gly	Leu	Val	Thr	Phe	Thr	Arg	Gln	Ser	Leu	Glu
	690					695					700				
Asp	Phe	Pro	Glu	Trp	Glu	Arg	Cys	Glu	Lys	Leu	Leu	Thr	Arg	Leu	His
705					710					715					720
Val	Thr	Tyr	Glu	Gly	Thr	Ile	Glu	Gly	Asn	Gly	Gln	Gly	Met	Leu	Gln
				725					730					735	
Val	Asp	Phe	Ala	Asn	Arg	Phe	Val	Gly	Gly	Gly	Val	Thr	Ser	Ala	Gly
			740					745					750		
Leu	Val	Gln	Glu	Glu	Ile	Arg	Phe	Leu	Ile	Asn	Pro	Glu	Leu	Ile	Val
		755					760					765			
Ser	Arg	Leu	Phe	Thr	Glu	Val	Leu	Asp	His	Asn	Glu	Cys	Leu	Ile	Ile
		770					775				780				
Thr	Gly	Thr	Glu	Gln	Tyr	Ser	Glu	Tyr	Thr	Gly	Tyr	Ala	Glu	Thr	Tyr
785					790					795					800
Arg	Trp	Ala	Arg	Ser	His	Glu	Asp	Arg	Ser	Glu	Arg	Asp	Asp	Trp	Gln
				805					810					815	
Arg	Arg	Thr	Thr	Glu	Ile	Val	Ala	Ile	Asp	Ala	Leu	His	Phe	Arg	Arg
			820					825					830		
Tyr	Leu	Asp	Gln	Phe	Val	Pro	Glu	Lys	Ile	Arg	Arg	Glu	Leu	Asn	Lys
		835					840					845			

-continued

Ala Tyr Cys Gly Phe Leu Arg Pro Gly Val Ser Ser Glu Asn Leu Ser
850 855 860

Ala Val Ala Thr Gly Asn Trp Gly Cys Gly Ala Phe Gly Gly Asp Ala
865 870 875 880

Arg Leu Lys Ala Leu Ile Gln Ile Leu Ala Ala Val Ala Glu Arg
885 890 895

Asp Val Val Tyr Phe Thr Phe Gly Asp Ser Glu Leu Met Arg Asp Ile
900 905 910

Tyr Ser Met His Thr Phe Leu Thr Glu Arg Lys Leu Thr Val Gly Glu
915 920 925

Val Tyr Lys Leu Leu Leu Arg Tyr Tyr Asn Glu Glu Cys Arg Asn Cys
930 935 940

Ser Thr Pro Gly Pro Asp Ile Lys Leu Tyr Pro Phe Ile Tyr His Ala
945 950 955 960

Val Glu Ser Cys Thr Gln Thr Thr Asn Gln Pro Gly Gln Arg Thr Gly
965 970 975

Ala

<210> SEQ ID NO 3
<211> LENGTH: 4069
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:

<400> SEQUENCE: 3

ggcgtctggg aagtgaggag cgtctctgcc tggcagaggc tgcaatctct gcactttggg 60
gggccaaggc aggcgctgag aaggacgcgc agtccatctc tctcaggtta gtgaaatgag 120
gctctccgcg gggccggccc ggggacagtg cgctgctggt cccagcatga atgcgggccc 180
cggctgtgaa ccctgcacca aagcgaccgc ctggggcgcc gctacaactt cgccggctgc 240
ttcggacgcc cggagcttcc cgagcaggca gaggcgcgtc ctcgacccca aggacgetca 300
cgtgcagtcc agggctccac cgtcctcgcc agcctgcgtc ccagggcagg cgggacagca 360
cagaggcagc gccacctcgc ttgttttcaa acaaaagact attaccagtt ggatggacac 420
taaaggaatc aagacagcgg aatcagaaaag tttggatagt aaagaaaaca acaatacaag 480
aatagaatcc atgatgagtt ctgtacaaaa agataacttt taccaacata atgtagaaaa 540
attagtaaat gtttctcagc taagtcttga taagtcactc actgaaaaaa gtacacagta 600
tttgaaccag catcagactg cagcaatgtg taagtggcaa aatgaaggga aacacacgga 660
gcagcttttg gaaagtgaac ctcaaacagt aacctggtg ccagagcagt ttagtaatgc 720
taacattgat cggtcacctc aaaatgatga tcacagtgc acagatagtg aagagaatag 780
agacaatcaa cagtttctca caactgtaaa gcttgcaaat gcaaagcaga ctacggaaga 840
tgaacacgcc agagaagcca aaagccacca gaagtgcagc aagtcttgcc atcctgggga 900
agactgtgca agttgtcagc aagatgagat agacgtggtg ccaaagagtc cattgtcaga 960
tgttggtctc gaggatggtg gtactgggtc aaaaaatgac aacaaattga ttagacaaga 1020
aagttgccta ggaaattctc ctccatttga gaaggaaagt gaaccggaat caccgatgga 1080
tgttgataat tctaaaaata gttgtcaaga ctcaagaagca gatgaggaga caagtccagg 1140
ttttgatgaa caagaagatg gtagtctcct ccaaacagca aataaacctt caaggttcca 1200
agcaagagac gctgacattg aatttaggaa acggtactct actaagggcg gtgaagttag 1260
attacatttc caatttgaag gaggagagag tcgcactgga atgaatgatt taaatgctaa 1320

-continued

actacctgga aatatttcta gcctgaatgt agaatgcaga aattctaagc aacatggaaa	1380
aaaggattct aaaatcacag atcatttgat gagactgccc aaagcagagg acagaagaaa	1440
agaacagtgg gaaaccaaac atcaaagaac agaaaggaag atccctaaat acgttccacc	1500
tcacctttct ccagataaga agtggccttg aactcccatt gaggagatga gaagaatgcc	1560
tcggtgtggg atccggctgc ctctcttgag accatctgcc aatcacacag taactattcg	1620
ggtagatcct ttgcgagcag gagaagtcc taaacctttt ccaacacatt ataaagattt	1680
gtgggataac aagcatgtta aaatgccttg ttcagaacaa aatttgtacc cagtgaaga	1740
tgagaatggt gagcgaactg cggggagccg gtgggagctc attcagactg cacttctcaa	1800
caaatttaca cgaccccaaa acttgaagga tgctattctg aaatacaatg tgcatattc	1860
taagaaatgg gactttcacg ctttgatcga tttctgggat aaggtacttg aagaagcaga	1920
agctcaacat ttatatcagt ccactctgcc tgatatggtg aaaattgcac tctgtctgcc	1980
aaatatttgc acccagccaa taccactcct gaaacagaag atgaatcatt ccatcacaat	2040
gtcgcaggaa cagattgcc a gtcttttagc taatgcttcc ttctgcacat ttccacgacg	2100
aaatgctaag atgaaatcgg agtattctag ttaccagac attaacttca atcgattgtt	2160
tgagggacgt tcatcaagga aaccggagaa acttaaaacg ctcttctgct actttagaag	2220
agtcacagag aaaaaacctg ctgggttggg gacatttaca agacagagtc ttgaagattt	2280
tcagaatgg gaaagatgtg aaaaaccctt gacacgattg catgtcactt acgaaggtac	2340
catagaagaa aatggccaag gcatgctaca ggtggatttt gcaaatcgtt ttgttgagg	2400
tggtgtaacc agtgcaggac ttgtgcaaga agaaatccgc tttttaatca atcctgagtt	2460
gattatttca cggtcttca ctgaggtgct ggatcacaat gaatgtctaa ttatcacagg	2520
tactgagcag tacagtgaat acacaggcta tgctgagaca tatcgttggg cccggagcca	2580
cgaagatggg agtgaaggg acgactgcga gcggcgctgc actgagatcg ttgccatcga	2640
tgctcttca ttcagacgct acctcgatca gtttgtgcct gagaaaatga gacgcgagct	2700
gaacaaggct tactgtggat ttctcctcc tggagtttct tcagagaatc tttctgcagt	2760
ggccacagga aactggggct gtggtgcctt tgggggtgat gccaggtaa aagccttaat	2820
acagatattg gcagctgctg cagctgagcg agatgtggtt tatttcacct ttggggactc	2880
agaattgatg agagacattt acagcatgca cattttcctt actgaaagga aactcactgt	2940
tgagatgtg tataagctgt tgctaogata ctacaatgaa gaatgcagaa actgttccac	3000
cctggacca gacatcaagc tttatccatt catataccat gctgtcgagt cctgtgcaga	3060
gaccgctgac cattcagggc aaaggacag gacctgagga gccgagcga tagcatctcc	3120
tcccacctcc caccagagac gtccctgctt agctgtcagg tgtaatatat gaattgactt	3180
aagttaatat aaatgtgtac ataatccaca tttgtagtca aggacgcaat ctcttccaca	3240
catgtgcagt tgtcagttgg tacatctaaa ctccctccat cctgactcac gtggacttag	3300
atatgttttg tttctatttt ctctatttc agtttttcat tctttgatgt ttatttcttt	3360
tgccatcag atctcttggt aaatcccatg gaaggttggt ctcagctgct gggctctctt	3420
cttctgccc atatattata ccagttgctt ctgcagcccg cagatgccc a gcgatgccc	3480
gaaacaagt tgaatccca ggaatctctt taactgattt tgctaaaaat ctccctgtga	3540
gccttccact caactcttaa tatgcttga ttgtttaagt ttttaattc tgaaaattaa	3600
taattaggtt tttttcata tgtgttgc ataatgcaacc tcctaggtta aaatagtttc	3660

-continued

```

ttttatctgta 3720
atggttgtc 3780
ctgagatgaa 3840
gattccaaga 3900
gactaatgtt 3960
gaacttcagt 4020
atgaattttc 4069

```

```

<210> SEQ ID NO 4
<211> LENGTH: 976
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:

```

```

<400> SEQUENCE: 4

```

```

Met Asn Ala Gly Pro Gly Cys Glu Pro Cys Thr Lys Ala Thr Arg Trp
1          5          10          15
Gly Ala Ala Thr Thr Ser Pro Ala Ala Ser Asp Ala Arg Ser Phe Pro
20          25          30
Ser Arg Gln Arg Arg Val Leu Asp Pro Lys Asp Ala His Val Gln Phe
35          40          45
Arg Val Pro Pro Ser Ser Pro Ala Cys Val Pro Gly Gln Ala Gly Gln
50          55          60
His Arg Gly Ser Ala Thr Ser Leu Val Phe Lys Gln Lys Thr Ile Thr
65          70          75          80
Ser Trp Met Asp Thr Lys Gly Ile Lys Thr Ala Glu Ser Glu Ser Leu
85          90          95
Asp Ser Lys Glu Asn Asn Asn Thr Arg Ile Glu Ser Met Met Ser Ser
100         105         110
Val Gln Lys Asp Asn Phe Tyr Gln His Asn Val Glu Lys Leu Val Asn
115         120         125
Val Ser Gln Leu Ser Leu Asp Lys Ser Leu Thr Glu Lys Ser Thr Gln
130         135         140
Tyr Leu Asn Gln His Gln Thr Ala Ala Met Cys Lys Trp Gln Asn Glu
145         150         155         160
Gly Lys His Thr Glu Gln Leu Leu Glu Ser Glu Pro Gln Thr Val Thr
165         170         175
Leu Val Pro Glu Gln Phe Ser Asn Ala Asn Ile Asp Arg Ser Pro Gln
180         185         190
Asn Asp Asp His Ser Asp Thr Asp Ser Glu Glu Asn Arg Asp Asn Gln
195         200         205
Gln Phe Leu Thr Thr Val Lys Leu Ala Asn Ala Lys Gln Thr Thr Glu
210         215         220
Asp Glu His Ala Arg Glu Ala Lys Ser His Gln Lys Cys Ser Lys Ser
225         230         235         240
Cys His Pro Gly Glu Asp Cys Ala Ser Cys Gln Gln Asp Glu Ile Asp
245         250         255
Val Val Pro Lys Ser Pro Leu Ser Asp Val Gly Ser Glu Asp Val Gly
260         265         270
Thr Gly Ser Lys Asn Asp Asn Lys Leu Ile Arg Gln Glu Ser Cys Leu
275         280         285
Gly Asn Ser Pro Pro Phe Glu Lys Glu Ser Glu Pro Glu Ser Pro Met
290         295         300

```

-continued

Asp Val Asp Asn Ser Lys Asn Ser Cys Gln Asp Ser Glu Ala Asp Glu
 305 310 315 320
 Glu Thr Ser Pro Gly Phe Asp Glu Gln Glu Asp Gly Ser Ser Ser Gln
 325 330 335
 Thr Ala Asn Lys Pro Ser Arg Phe Gln Ala Arg Asp Ala Asp Ile Glu
 340 345 350
 Phe Arg Lys Arg Tyr Ser Thr Lys Gly Gly Glu Val Arg Leu His Phe
 355 360 365
 Gln Phe Glu Gly Gly Glu Ser Arg Thr Gly Met Asn Asp Leu Asn Ala
 370 375 380
 Lys Leu Pro Gly Asn Ile Ser Ser Leu Asn Val Glu Cys Arg Asn Ser
 385 390 395 400
 Lys Gln His Gly Lys Lys Asp Ser Lys Ile Thr Asp His Leu Met Arg
 405 410 415
 Leu Pro Lys Ala Glu Asp Arg Arg Lys Glu Gln Trp Glu Thr Lys His
 420 425 430
 Gln Arg Thr Glu Arg Lys Ile Pro Lys Tyr Val Pro Pro His Leu Ser
 435 440 445
 Pro Asp Lys Lys Trp Leu Gly Thr Pro Ile Glu Glu Met Arg Arg Met
 450 455 460
 Pro Arg Cys Gly Ile Arg Leu Pro Leu Leu Arg Pro Ser Ala Asn His
 465 470 475 480
 Thr Val Thr Ile Arg Val Asp Leu Leu Arg Ala Gly Glu Val Pro Lys
 485 490 495
 Pro Phe Pro Thr His Tyr Lys Asp Leu Trp Asp Asn Lys His Val Lys
 500 505 510
 Met Pro Cys Ser Glu Gln Asn Leu Tyr Pro Val Glu Asp Glu Asn Gly
 515 520 525
 Glu Arg Thr Ala Gly Ser Arg Trp Glu Leu Ile Gln Thr Ala Leu Leu
 530 535 540
 Asn Lys Phe Thr Arg Pro Gln Asn Leu Lys Asp Ala Ile Leu Lys Tyr
 545 550 555 560
 Asn Val Ala Tyr Ser Lys Lys Trp Asp Phe Thr Ala Leu Ile Asp Phe
 565 570 575
 Trp Asp Lys Val Leu Glu Glu Ala Glu Ala Gln His Leu Tyr Gln Ser
 580 585 590
 Ile Leu Pro Asp Met Val Lys Ile Ala Leu Cys Leu Pro Asn Ile Cys
 595 600 605
 Thr Gln Pro Ile Pro Leu Leu Lys Gln Lys Met Asn His Ser Ile Thr
 610 615 620
 Met Ser Gln Glu Gln Ile Ala Ser Leu Leu Ala Asn Ala Phe Phe Cys
 625 630 635 640
 Thr Phe Pro Arg Arg Asn Ala Lys Met Lys Ser Glu Tyr Ser Ser Tyr
 645 650 655
 Pro Asp Ile Asn Phe Asn Arg Leu Phe Glu Gly Arg Ser Ser Arg Lys
 660 665 670
 Pro Glu Lys Leu Lys Thr Leu Phe Cys Tyr Phe Arg Arg Val Thr Glu
 675 680 685
 Lys Lys Pro Thr Gly Leu Val Thr Phe Thr Arg Gln Ser Leu Glu Asp
 690 695 700
 Phe Pro Glu Trp Glu Arg Cys Glu Lys Pro Leu Thr Arg Leu His Val
 705 710 715 720

-continued

Thr	Tyr	Glu	Gly	Thr	Ile	Glu	Glu	Asn	Gly	Gln	Gly	Met	Leu	Gln	Val
				725					730					735	
Asp	Phe	Ala	Asn	Arg	Phe	Val	Gly	Gly	Gly	Val	Thr	Ser	Ala	Gly	Leu
			740					745					750		
Val	Gln	Glu	Glu	Ile	Arg	Phe	Leu	Ile	Asn	Pro	Glu	Leu	Ile	Ile	Ser
		755					760					765			
Arg	Leu	Phe	Thr	Glu	Val	Leu	Asp	His	Asn	Glu	Cys	Leu	Ile	Ile	Thr
	770					775					780				
Gly	Thr	Glu	Gln	Tyr	Ser	Glu	Tyr	Thr	Gly	Tyr	Ala	Glu	Thr	Tyr	Arg
785					790					795					800
Trp	Ser	Arg	Ser	His	Glu	Asp	Gly	Ser	Glu	Arg	Asp	Asp	Cys	Glu	Arg
				805					810					815	
Arg	Cys	Thr	Glu	Ile	Val	Ala	Ile	Asp	Ala	Leu	His	Phe	Arg	Arg	Tyr
			820					825					830		
Leu	Asp	Gln	Phe	Val	Pro	Glu	Lys	Met	Arg	Arg	Glu	Leu	Asn	Lys	Ala
		835					840					845			
Tyr	Cys	Gly	Phe	Leu	Arg	Pro	Gly	Val	Ser	Ser	Glu	Asn	Leu	Ser	Ala
	850					855					860				
Val	Ala	Thr	Gly	Asn	Trp	Gly	Cys	Gly	Ala	Phe	Gly	Gly	Asp	Ala	Arg
865					870					875					880
Leu	Lys	Ala	Leu	Ile	Gln	Ile	Leu	Ala	Ala	Ala	Ala	Ala	Glu	Arg	Asp
				885				890						895	
Val	Val	Tyr	Phe	Thr	Phe	Gly	Asp	Ser	Glu	Leu	Met	Arg	Asp	Ile	Tyr
			900					905					910		
Ser	Met	His	Ile	Phe	Leu	Thr	Glu	Arg	Lys	Leu	Thr	Val	Gly	Asp	Val
		915					920						925		
Tyr	Lys	Leu	Leu	Leu	Arg	Tyr	Tyr	Asn	Glu	Glu	Cys	Arg	Asn	Cys	Ser
	930					935					940				
Thr	Pro	Gly	Pro	Asp	Ile	Lys	Leu	Tyr	Pro	Phe	Ile	Tyr	His	Ala	Val
945					950					955					960
Glu	Ser	Cys	Ala	Glu	Thr	Ala	Asp	His	Ser	Gly	Gln	Arg	Thr	Gly	Thr
				965					970					975	

<210> SEQ ID NO 5

<211> LENGTH: 3814

<212> TYPE: DNA

<213> ORGANISM: Mus musculus

<220> FEATURE:

<400> SEQUENCE: 5

```

gggggactgt gtgctgctgg tcccagcatg agtgccggcc cgggctggga gccctgcacg      60
aaagcgcgct ggggcgccgc tggaacttct ggcgccactg cctcggactc cgggagcttc      120
cctggcaggc agaggcgtgt tctcgacccc aaggacgctc cgtccagtt cagggtccct      180
ccgtcctcgc cagcctcgct ctggggcgcg gggggaccgc acagaggcaa cgccacctcg      240
ttgttttca aacaaaagac tattactact tggatggata ctaaaggacc caagacagct      300
gaatcagaaa gtaaaagaaa caacaataca agaattgact ccatgatgag ttctgtgcag      360
aaagataact ttaccacaca taagtgga aaattggaaa atgttcctca gctaaatctt      420
gataaatcac ccacagaaaa gagttcacag tatttgaacc aacagcagac tgcgagtgtg      480
tgcaagtggc agaatgaagg gaagcatgca gaacagcttt tggcaagtga gcctcccggc      540
gggactccgc taccaagca gcttagtaat gctaacattg gtcagtcacc ccacactgat      600
gaccacagtg acacagatca tgaagaagac agagacaatc agcagtttct tacacctata      660

```


-continued

aaacttgcaa atacaaagcc aacagtagga gatgggcagg ccagaagcaa ctgtaagtgc	720
agtggatctc gccagctctgt gaaagactgt acaggctgtc aacaggagga ggtggatgtg	780
ctaccagaga gtcctttgtc agatgttggg gccgaggaca ttggaactgg accaaaaaat	840
gacaacaaat tgactggaca agaaagcagc ctagggtgatt cgcctccatt tgagaaagaa	900
agtgagcctg agtcaccaat ggatgtagac aactcgagaa acagttgtca agattcagaa	960
gcagatgaag aaacaagtcc agtctttgat gagcaagatg atcgttcttc ccaaacagca	1020
aataaaacttt caagttgcc aagcaagagaa gctgatggcg atcttaggaa acggtatttg	1080
actaagggaa gtgaagttag attgcatttc caatttgaag gagaaaataa tgctgggacc	1140
agtgacttaa atgccaaagcc atctggaaac tcttctagcc ttaatgtaga gtgtagaagt	1200
tccaagcagc atggaaaaag ggattctaaa attacagatc atttcatgag aatttccaag	1260
tcagaggaca gaagaaaaga acaatgtgaa gtcagacatc aaagaacaga aaggaagatt	1320
ccaaaataca tcccacctaa cctccctcca gagaagaagt ggctgggaac tcctattgag	1380
gaaatgagaa aaatgcctcg gtgtgggac catttgcctt ccttaagacc atctgcaagt	1440
cacacagtga ctgttcgggt agaccttctg agagcaggag aggttccgaa acctttcca	1500
acacattaca aagattttgt ggataacaaa catgtgaaaa tgccttgttc ggaacaaaac	1560
ttgtaccctg tggaagatga gaatggtag cgaactgcag ggagtaggtg ggagctcatt	1620
cagactgcac ttctcaacaa attcacacga cccagaact tgaaggatgc gattctgaaa	1680
tacaatgtgg catattctaa gaaatgggac tttacagctt tggttgattt ctgggataag	1740
gtacttgaag aagcagaggc ccaacattta taccagtcca ttttacctga catggtgaaa	1800
attgcactct gtctgcaaaa tatttgcacc cagccaatac cactcctgaa acagaagatg	1860
aatcattctg tcacgatgtc acaggaacag atcgccagtc ttttagctaa tgctttcttc	1920
tgacatttc cccgacggaa tgccaagatg aaatcggagt attctagtta cccagacatt	1980
aacttcaatc ggttgtttga aggacgttca tcaaggaaac cagaaaact gaaaacactc	2040
ttctgctact ttcgaaagat cacagagaaa aaacctacag gattggtgac atttacaaga	2100
cagagctctg aagattttcc agaatgggaa aggtgtgaaa agcctctgac acgcttacac	2160
gtcacttacg aggtaccat agaaggcaac ggccgaggca tgctacaggt ggattttgca	2220
aatcgttttg ttggaggtgg tgtgactggt gcgggacttg tacaagaaga aatcagattt	2280
ttaatcaatc ctgaatgat tgtttcacgg ctgttctact aggtgctgga tcacaatgag	2340
tgtcttatta tcacaggtac tgaacagtac agtgaataca caggctatgc tgaacttat	2400
cgttgggccc gaagccatga agatgggagt gaaaaggacg attggcagcg gcgctgcacg	2460
gagatcgttg ccattgacgc acttcacttc agacgctacc tcgatcagtt tgtgctgag	2520
aaagtgagac gtgagcttaa caaggcttac tgcggattcc tccgtcctgg agttccttct	2580
gaaaaatctt ctgcagtggc cacgggaaac tggggctgtg gtgcctttgg gggtagcgt	2640
agattaaag ccttaataca gatcctggca gctgctgcgg ctgaactgga cgtggtttat	2700
ttcacctttg gggactcaga gttgatgaga gacatttaca gcatgcacac tttccttacc	2760
gagaggaagc tggatgttgg aaaagtgtac aagttattgc ttagatacta caatgaagaa	2820
tgacaaaact gttccacccc tggaccagac atcaagcttt atccattcat ataccatgct	2880
gttgagtcaa gtgcagagac cactgacatg ccaggacaga aggcaggcac ctgaggaaca	2940
agtgactagg acctcctctc aaagagacat cctatttgaa atgtgggggtg tgatgtctga	3000
attgactgaa tctgatctaa gtgtgtatat aatccacatt tgtaatcaag gatgcagtct	3060

-continued

```

cttctgcata tgcagtgtt tcttgttcat cctggaggac atgccttag acatggcttc 3120
ttcaattttt cttctcttc agtctttatt ctttgatttt tttttccaa cttgatttct 3180
tgggaaaact caagaaaggt tgcactcagc ttctagatct ttctcttct gtctgtgtgt 3240
tgtccagact gctttgggg ctagcagata ccatcacact tggaggaagt tacaaatcca 3300
gaaatctgag tttgtgcag atttacctgt gagcttctca ctcccaacc ttgttaggct 3360
tgtgtgtct acattttcaa ttttgaagt tgaagtttt cttatgttac ttaatgctag 3420
tatcttttag gctaaaacta tttctattt aaggcagact aattccagt ttctctttg 3480
aaacatcct cctataagta acggttttt tctctcttt tccccagcg ctattttaga 3540
agctggccaa gaggaagaa aatgtagaat aaaaggattt tcctcggatg ctataagaa 3600
gccagttca agagegttg ggtttttgt ttttcaaga cttgttttc ctttgcagct 3660
agggtgagt cttgttctgt ggtgctgagg gcatagctct gtaaccaag gtctttgtg 3720
gagactgat gctgatttgt acatatgaa gtttctctg caggaaat tagagtaat 3780
aaatttcatt aataaatcat ttgtcagaaa aaaa 3814

```

```

<210> SEQ ID NO 6
<211> LENGTH: 968
<212> TYPE: PRT
<213> ORGANISM: Mus musculus
<220> FEATURE:

```

```

<400> SEQUENCE: 6

```

```

Met Ser Ala Gly Pro Gly Trp Glu Pro Cys Thr Lys Ala Arg Trp Gly
 1           5           10          15
Ala Ala Gly Thr Ser Ala Pro Thr Ala Ser Asp Ser Arg Ser Phe Pro
          20           25           30
Gly Arg Gln Arg Arg Val Leu Asp Pro Lys Asp Ala Pro Val Gln Phe
          35           40           45
Arg Val Pro Pro Ser Ser Pro Ala Cys Val Ser Gly Arg Ala Gly Pro
          50           55           60
His Arg Gly Asn Ala Thr Ser Phe Val Phe Lys Gln Lys Thr Ile Thr
 65           70           75           80
Thr Trp Met Asp Thr Lys Gly Pro Lys Thr Ala Glu Ser Glu Ser Lys
          85           90           95
Glu Asn Asn Asn Thr Arg Ile Asp Ser Met Met Ser Ser Val Gln Lys
          100          105          110
Asp Asn Phe Tyr Pro His Lys Val Glu Lys Leu Glu Asn Val Pro Gln
          115          120          125
Leu Asn Leu Asp Lys Ser Pro Thr Glu Lys Ser Ser Gln Tyr Leu Asn
          130          135          140
Gln Gln Gln Thr Ala Ser Val Cys Lys Trp Gln Asn Glu Gly Lys His
 145          150          155          160
Ala Glu Gln Leu Leu Ala Ser Glu Pro Pro Ala Gly Thr Pro Leu Pro
          165          170          175
Lys Gln Leu Ser Asn Ala Asn Ile Gly Gln Ser Pro His Thr Asp Asp
          180          185          190
His Ser Asp Thr Asp His Glu Glu Asp Arg Asp Asn Gln Gln Phe Leu
          195          200          205
Thr Pro Ile Lys Leu Ala Asn Thr Lys Pro Thr Val Gly Asp Gly Gln
 210          215          220
Ala Arg Ser Asn Cys Lys Cys Ser Gly Ser Arg Gln Ser Val Lys Asp

```

-continued

225					230															240
Cys	Thr	Gly	Cys	Gln	Gln	Glu	Glu	Val	Asp	Val	Leu	Pro	Glu	Ser	Pro					
				245					250					255						
Leu	Ser	Asp	Val	Gly	Ala	Glu	Asp	Ile	Gly	Thr	Gly	Pro	Lys	Asn	Asp					
			260					265					270							
Asn	Lys	Leu	Thr	Gly	Gln	Glu	Ser	Ser	Leu	Gly	Asp	Ser	Pro	Pro	Phe					
		275					280					285								
Glu	Lys	Glu	Ser	Glu	Pro	Glu	Ser	Pro	Met	Asp	Val	Asp	Asn	Ser	Arg					
	290					295					300									
Asn	Ser	Cys	Gln	Asp	Ser	Glu	Ala	Asp	Glu	Glu	Thr	Ser	Pro	Val	Phe					
305					310				315						320					
Asp	Glu	Gln	Asp	Asp	Arg	Ser	Ser	Gln	Thr	Ala	Asn	Lys	Leu	Ser	Ser					
				325					330					335						
Cys	Gln	Ala	Arg	Glu	Ala	Asp	Gly	Asp	Leu	Arg	Lys	Arg	Tyr	Leu	Thr					
			340					345					350							
Lys	Gly	Ser	Glu	Val	Arg	Leu	His	Phe	Gln	Phe	Glu	Gly	Glu	Asn	Asn					
		355					360					365								
Ala	Gly	Thr	Ser	Asp	Leu	Asn	Ala	Lys	Pro	Ser	Gly	Asn	Ser	Ser	Ser					
	370					375					380									
Leu	Asn	Val	Glu	Cys	Arg	Ser	Ser	Lys	Gln	His	Gly	Lys	Arg	Asp	Ser					
385					390					395				400						
Lys	Ile	Thr	Asp	His	Phe	Met	Arg	Ile	Ser	Lys	Ser	Glu	Asp	Arg	Arg					
				405				410						415						
Lys	Glu	Gln	Cys	Glu	Val	Arg	His	Gln	Arg	Thr	Glu	Arg	Lys	Ile	Pro					
			420					425					430							
Lys	Tyr	Ile	Pro	Pro	Asn	Leu	Pro	Pro	Glu	Lys	Lys	Trp	Leu	Gly	Thr					
		435					440					445								
Pro	Ile	Glu	Glu	Met	Arg	Lys	Met	Pro	Arg	Cys	Gly	Ile	His	Leu	Pro					
	450					455					460									
Ser	Leu	Arg	Pro	Ser	Ala	Ser	His	Thr	Val	Thr	Val	Arg	Val	Asp	Leu					
465					470					475				480						
Leu	Arg	Ala	Gly	Glu	Val	Pro	Lys	Pro	Phe	Pro	Thr	His	Tyr	Lys	Asp					
				485					490					495						
Leu	Trp	Asp	Asn	Lys	His	Val	Lys	Met	Pro	Cys	Ser	Glu	Gln	Asn	Leu					
			500					505					510							
Tyr	Pro	Val	Glu	Asp	Glu	Asn	Gly	Glu	Arg	Thr	Ala	Gly	Ser	Arg	Trp					
		515					520					525								
Glu	Leu	Ile	Gln	Thr	Ala	Leu	Leu	Asn	Lys	Phe	Thr	Arg	Pro	Gln	Asn					
	530					535					540									
Leu	Lys	Asp	Ala	Ile	Leu	Lys	Tyr	Asn	Val	Ala	Tyr	Ser	Lys	Lys	Trp					
545					550					555					560					
Asp	Phe	Thr	Ala	Leu	Val	Asp	Phe	Trp	Asp	Lys	Val	Leu	Glu	Glu	Ala					
				565					570					575						
Glu	Ala	Gln	His	Leu	Tyr	Gln	Ser	Ile	Leu	Pro	Asp	Met	Val	Lys	Ile					
			580					585					590							
Ala	Leu	Cys	Leu	Pro	Asn	Ile	Cys	Thr	Gln	Pro	Ile	Pro	Leu	Leu	Lys					
		595					600					605								
Gln	Lys	Met	Asn	His	Ser	Val	Thr	Met	Ser	Gln	Glu	Gln	Ile	Ala	Ser					
	610						615					620								
Leu	Leu	Ala	Asn	Ala	Phe	Phe	Cys	Thr	Phe	Pro	Arg	Arg	Asn	Ala	Lys					
625					630					635					640					
Met	Lys	Ser	Glu	Tyr	Ser	Ser	Tyr	Pro	Asp	Ile	Asn	Phe	Asn	Arg	Leu					
				645					650					655						

-continued

Phe Glu Gly Arg Ser Ser Arg Lys Pro Glu Lys Leu Lys Thr Leu Phe
 660 665 670
 Cys Tyr Phe Arg Arg Val Thr Glu Lys Lys Pro Thr Gly Leu Val Thr
 675 680 685
 Phe Thr Arg Gln Ser Leu Glu Asp Phe Pro Glu Trp Glu Arg Cys Glu
 690 695 700
 Lys Pro Leu Thr Arg Leu His Val Thr Tyr Glu Gly Thr Ile Glu Gly
 705 710 715 720
 Asn Gly Arg Gly Met Leu Gln Val Asp Phe Ala Asn Arg Phe Val Gly
 725 730 735
 Gly Gly Val Thr Gly Ala Gly Leu Val Gln Glu Glu Ile Arg Phe Leu
 740 745 750
 Ile Asn Pro Glu Leu Ile Val Ser Arg Leu Phe Thr Glu Val Leu Asp
 755 760 765
 His Asn Glu Cys Leu Ile Ile Thr Gly Thr Glu Gln Tyr Ser Glu Tyr
 770 775 780
 Thr Gly Tyr Ala Glu Thr Tyr Arg Trp Ala Arg Ser His Glu Asp Gly
 785 790 795 800
 Ser Glu Lys Asp Asp Trp Gln Arg Arg Cys Thr Glu Ile Val Ala Ile
 805 810 815
 Asp Ala Leu His Phe Arg Arg Tyr Leu Asp Gln Phe Val Pro Glu Lys
 820 825 830
 Val Arg Arg Glu Leu Asn Lys Ala Tyr Cys Gly Phe Leu Arg Pro Gly
 835 840 845
 Val Pro Ser Glu Asn Leu Ser Ala Val Ala Thr Gly Asn Trp Gly Cys
 850 855 860
 Gly Ala Phe Gly Gly Asp Ala Arg Leu Lys Ala Leu Ile Gln Ile Leu
 865 870 875 880
 Ala Ala Ala Ala Ala Glu Arg Asp Val Val Tyr Phe Thr Phe Gly Asp
 885 890 895
 Ser Glu Leu Met Arg Asp Ile Tyr Ser Met His Thr Phe Leu Thr Glu
 900 905 910
 Arg Lys Leu Asp Val Gly Lys Val Tyr Lys Leu Leu Leu Arg Tyr Tyr
 915 920 925
 Asn Glu Glu Cys Arg Asn Cys Ser Thr Pro Gly Pro Asp Ile Lys Leu
 930 935 940
 Tyr Pro Phe Ile Tyr His Ala Val Glu Ser Ser Ala Glu Thr Thr Asp
 945 950 955 960
 Met Pro Gly Gln Lys Ala Gly Thr
 965

<210> SEQ ID NO 7
 <211> LENGTH: 2781
 <212> TYPE: DNA
 <213> ORGANISM: Drosophila melanogaster
 <220> FEATURE:

<400> SEQUENCE: 7

tcgaagtgtg tggatattat aaagtgcgat attcatcaca gctatcgctc atccccaaaa 60
 caccggtatg caagaattca ggtcacactt gatttttccg atattccaaa aggtttacca 120
 atctacggca aatcgccgca gagcaagtgc atccgtgctg accaatcgac tcggcaaggc 180
 tttgtgetta aactgcgcc aagatgtcgaa gtcgcccgat ggcgggattt ccgaaataga 240
 aacggaggag gagccggaat atctggcgaa ctcctagat gattcgtggc gtggagtttc 300

-continued

catggaggct atacatcgta atcggcagcc tttcgaattg gagaatttgc caccagtgac	360
tgccggcaat ctccaccggg ttatgtacca gctgcccaatt cgtgaaacac cgccacgccc	420
ctacaaatca cgggaaagt gggactccga gcatgtgctg ctgccctgtg cgcccagtc	480
gaaatatccg agggagaatc cggatggcag caccaccatc gattttcgtc gggaaatgat	540
cgaaacgagcc cttctgcagc ccataaagac gtgtgaggaa ctgcaggcgg cgataatc	600
atataatacc acctataggg atcagtggca ctttcgtgcc cttcatcaac ttctcgacga	660
ggactggac gagagcgaac cacgggtttt cttcgaggat ctattgccgc gcattatccg	720
attggcattg cggctaccgg acttgattca atcgccagtt ccgctgctca agcaccacaa	780
gaacgcctca ttgagcctga gccaacagca gatctcctgc ctgttgcca atgccttctt	840
gtgcacgttt ccccgaagaa acacctcaa gaggaagtcc gactacagca cttttccaga	900
catcaacttt aacagccttt accaatcgac gggaccggca gttctggaga agcttaaatg	960
cattatgcac tattttcgtc gcgtgtgtcc cacagagcgg gatgccagca atgtgccac	1020
cgggtggtga acctttgttc gtcggagcgg attgccgaa catctgatcg actggagcca	1080
aagtgcggcg ccggtgggtg atgtgccatt gcacgtggat gccaggggaa caatcgagga	1140
tgagggcatt ggactgctgc aagtagactt tgccaacaaa tatttgggtg gcggtgtctt	1200
gggacatggc tgcgttcagg aggagatacg ctttgttacc tgtccggagc tatttggggg	1260
taaactcttt acggagtgtc tgcgaccatt cgaggccctg gtgatgttg gcgccgaaag	1320
gtatagtaac tatacgggat atgccggaag cttcgagtgg tccggcaact ttgaggattc	1380
aacgccaaga gatagctcag gtcgtcgaca aacggccatt gtggcaatcg atgccctaca	1440
ttttgccag tcacatcacc aatatcgcca ggatctcatg gaaagggagc tgaacaaggc	1500
gtacattgga tttgttcaact ggatggtgac gccgccaccg ggtgtggcaa ctggtaactg	1560
gggttgcggc gcattcggcg gtgactccta tctgaaagcc ctgctgcaac ttatggtctg	1620
cgcccagttg ggcagacctt tggcctacta tacctttgga aatgtggagt ttagggatga	1680
ttttcatgaa atgtgctgtg tgtttcgaaa tgacgggact acggtgcagc agctttggag	1740
tattttaagg tcgtacagta ggcttattaa ggagaagagc tccaaggagc cgcgtgagaa	1800
taaggcatcc aaaaagaagc tatatgattt tattaagag gaacttaaga aggtcagaga	1860
tgtgcccgga gagggagcat ccgccgaagc tggaagctct agagtagctg gattaggcga	1920
aggaaaatca gaaacatcag cgaaatcctc gccagaactc aacaagcaac ccgccgacc	1980
gcaaatcacc ataacgcaac aaagtaccga tctattgccc gcgcaattat cgcaagataa	2040
ctctaattct tcggaagatc aggcccttct tatgctgtcg gacgatgagg aggccaatgc	2100
catgatggag gccgctagtc tggaggctaa aagcagcgtg gaaataagca acagcagcac	2160
aacgtccaaa acgagcagta cagccacgaa atcaatgggt tcaggtggcc gccagttgag	2220
tctgctcgag atgctggaca cccattatga aaagggttcg gcctcgaaga ggccacgaaa	2280
atcaccacaac tgcagcaagg ctgagggttc agcaaagagt cgtaaggaga tcgatgtgac	2340
cgacaaggac gaaaaggagc atattgttga ctaggtgata ttgcaactaca ggattgttac	2400
tgcccccaaa aattgaagag gtataaaatg tattgtagat aactttaagg acatatttag	2460
ggcattttaa agtaggatca ttgtaagtcg aataaagtga aatTTTTTTT ttttttaaat	2520
tatactattc taactctgaa agacaatttt actgttaaat ttgtataaca ttcgaattaa	2580
ttaataaat ttgttatatc atgcaaatct agcttttatt atgcgaaatt tgtagttaa	2640

-continued

```

gccagtaaag tttcttttta ttttaaccgaa accttttgtt tattttatatt gaccacaaca 2700
agaacatcaa caacaacaac cacgaaaaaa aagcgaatat atatttggtt gttcgtatat 2760
atatatatat ctaagcagat c 2781

```

```

<210> SEQ ID NO 8
<211> LENGTH: 768
<212> TYPE: PRT
<213> ORGANISM: Drosophila melanogaster
<220> FEATURE:

```

```

<400> SEQUENCE: 8

```

```

Met Gln Glu Phe Arg Ser His Leu Ile Phe Pro Ile Phe Gln Lys Val
 1          5          10          15
Tyr Gln Ser Thr Ala Asn Arg Arg Arg Ala Ser Ala Ser Val Leu Thr
          20          25          30
Asn Arg Leu Gly Lys Ala Leu Cys Leu Asn Cys Ala Arg Met Ser Lys
          35          40          45
Ser Pro Asp Gly Gly Ile Ser Glu Ile Glu Thr Glu Glu Glu Pro Glu
          50          55          60
Asn Leu Ala Asn Ser Leu Asp Asp Ser Trp Arg Gly Val Ser Met Glu
 65          70          75          80
Ala Ile His Arg Asn Arg Gln Pro Phe Glu Leu Glu Asn Leu Pro Pro
          85          90          95
Val Thr Ala Gly Asn Leu His Arg Val Met Tyr Gln Leu Pro Ile Arg
          100          105          110
Glu Thr Pro Pro Arg Pro Tyr Lys Ser Pro Gly Lys Trp Asp Ser Glu
          115          120          125
His Val Arg Leu Pro Cys Ala Pro Glu Ser Lys Tyr Pro Arg Glu Asn
          130          135          140
Pro Asp Gly Ser Thr Thr Ile Asp Phe Arg Trp Glu Met Ile Glu Arg
          145          150          155          160
Ala Leu Leu Gln Pro Ile Lys Thr Cys Glu Glu Leu Gln Ala Ala Ile
          165          170          175
Ile Ser Tyr Asn Thr Thr Tyr Arg Asp Gln Trp His Phe Arg Ala Leu
          180          185          190
His Gln Leu Leu Asp Glu Glu Leu Asp Glu Ser Glu Thr Arg Val Phe
          195          200          205
Phe Glu Asp Leu Leu Pro Arg Ile Ile Arg Leu Ala Leu Arg Leu Pro
          210          215          220
Asp Leu Ile Gln Ser Pro Val Pro Leu Leu Lys His His Lys Asn Ala
          225          230          235          240
Ser Leu Ser Leu Ser Gln Gln Gln Ile Ser Cys Leu Leu Ala Asn Ala
          245          250          255
Phe Leu Cys Thr Phe Pro Arg Arg Asn Thr Leu Lys Arg Lys Ser Glu
          260          265          270
Tyr Ser Thr Phe Pro Asp Ile Asn Phe Asn Arg Leu Tyr Gln Ser Thr
          275          280          285
Gly Pro Ala Val Leu Glu Lys Leu Lys Cys Ile Met His Tyr Phe Arg
          290          295          300
Arg Val Cys Pro Thr Glu Arg Asp Ala Ser Asn Val Pro Thr Gly Val
          305          310          315          320
Val Thr Phe Val Arg Arg Ser Gly Leu Pro Glu His Leu Ile Asp Trp
          325          330          335
Ser Gln Ser Ala Ala Pro Leu Gly Asp Val Pro Leu His Val Asp Ala

```

-continued

340					345					350					
Glu	Gly	Thr	Ile	Glu	Asp	Glu	Gly	Ile	Gly	Leu	Leu	Gln	Val	Asp	Phe
				355											365
Ala	Asn	Lys	Tyr	Leu	Gly	Gly	Gly	Val	Leu	Gly	His	Gly	Cys	Val	Gln
				370											380
Glu	Glu	Ile	Arg	Phe	Val	Ile	Cys	Pro	Glu	Leu	Leu	Val	Gly	Lys	Leu
															400
Phe	Thr	Glu	Cys	Leu	Arg	Pro	Phe	Glu	Ala	Leu	Val	Met	Leu	Gly	Ala
				405											415
Glu	Arg	Tyr	Ser	Asn	Tyr	Thr	Gly	Tyr	Ala	Gly	Ser	Phe	Glu	Trp	Ser
				420											430
Gly	Asn	Phe	Glu	Asp	Ser	Thr	Pro	Arg	Asp	Ser	Ser	Gly	Arg	Arg	Gln
				435											445
Thr	Ala	Ile	Val	Ala	Ile	Asp	Ala	Leu	His	Phe	Ala	Gln	Ser	His	His
															460
Gln	Tyr	Arg	Glu	Asp	Leu	Met	Glu	Arg	Glu	Leu	Asn	Lys	Ala	Tyr	Ile
															480
Gly	Phe	Val	His	Trp	Met	Val	Thr	Pro	Pro	Pro	Gly	Val	Ala	Thr	Gly
				485											495
Asn	Trp	Gly	Cys	Gly	Ala	Phe	Gly	Gly	Asp	Ser	Tyr	Leu	Lys	Ala	Leu
				500											510
Leu	Gln	Leu	Met	Val	Cys	Ala	Gln	Leu	Gly	Arg	Pro	Leu	Ala	Tyr	Tyr
				515											525
Thr	Phe	Gly	Asn	Val	Glu	Phe	Arg	Asp	Asp	Phe	His	Glu	Met	Trp	Leu
															540
Leu	Phe	Arg	Asn	Asp	Gly	Thr	Thr	Val	Gln	Gln	Leu	Trp	Ser	Ile	Leu
															560
Arg	Ser	Tyr	Ser	Arg	Leu	Ile	Lys	Glu	Lys	Ser	Ser	Lys	Glu	Pro	Arg
				565											575
Glu	Asn	Lys	Ala	Ser	Lys	Lys	Lys	Leu	Tyr	Asp	Phe	Ile	Lys	Glu	Glu
				580											590
Leu	Lys	Lys	Val	Arg	Asp	Val	Pro	Gly	Glu	Gly	Ala	Ser	Ala	Glu	Ala
				595											605
Gly	Ser	Ser	Arg	Val	Ala	Gly	Leu	Gly	Glu	Gly	Lys	Ser	Glu	Thr	Ser
															620
Ala	Lys	Ser	Ser	Pro	Glu	Leu	Asn	Lys	Gln	Pro	Ala	Arg	Pro	Gln	Ile
															640
Thr	Ile	Thr	Gln	Gln	Ser	Thr	Asp	Leu	Leu	Pro	Ala	Gln	Leu	Ser	Gln
				645											655
Asp	Asn	Ser	Asn	Ser	Ser	Glu	Asp	Gln	Ala	Leu	Leu	Met	Leu	Ser	Asp
				660											670
Asp	Glu	Glu	Ala	Asn	Ala	Met	Met	Glu	Ala	Ala	Ser	Leu	Glu	Ala	Lys
				675											685
Ser	Ser	Val	Glu	Ile	Ser	Asn	Ser	Ser	Thr	Thr	Ser	Lys	Thr	Ser	Ser
															700
Thr	Ala	Thr	Lys	Ser	Met	Gly	Ser	Gly	Gly	Arg	Gln	Leu	Ser	Leu	Leu
															720
Glu	Met	Leu	Asp	Thr	His	Tyr	Glu	Lys	Gly	Ser	Ala	Ser	Lys	Arg	Pro
				725											735
Arg	Lys	Ser	Pro	Asn	Cys	Ser	Lys	Ala	Glu	Gly	Ser	Ala	Lys	Ser	Arg
				740											750
Lys	Glu	Ile	Asp	Val	Thr	Asp	Lys	Asp	Glu	Lys	Asp	Asp	Ile	Val	Asp
				755											765

-continued

```

<210> SEQ ID NO 9
<211> LENGTH: 2181
<212> TYPE: DNA
<213> ORGANISM: Caenorhabditis elegans
<220> FEATURE:

<400> SEQUENCE: 9
atgagcaaga agtttatcga actgggtgat cctgtcactc aagacgagaa agactacgaa      60
gactatgtcg gagttgggtt cgccgatcaa gtcccgacaa tgaaggcg gaagttgaca      120
gaacatggaa atactacaga atcaaaagaa gatcctgaag agccaaaaag ccgtgacgta      180
tttgtctcct cgcagtcaag tgatgagagt caagaagatt cggctgaaaa tccggagatc      240
gctaaaagaag tgtcagaaaa ttgtgaaaa ctgacagaaa ctctcaaaat ttctaataat      300
gagagtttgg acaatgttac tgaagatct gaacacactc ttgataatca caaaagtact      360
gaaccaatgg aagaagatgt aaacaacaag tccaatattg acgttgcgat taattctgac      420
gaggatgatg aactgttctt ggaagagaat aataaagaaa tgagggatgg agaacaagta      480
caacagttgt cacaggattt attcgtgat gatcaagagc taattgaata tccaggaatt      540
atgaaagaca ctacaactca actggatata acagattctg aagtggagac tgctcaaaaa      600
atggaaatga ttgaagaaac tgaagcagat tcgacatttg taggcgagga ttcaaaagct      660
acgaaaactg tgaggacatc cagttcaagt ttcctgtcaa ctgtttcaac atgcgaagcc      720
cctgcaaaag gacgagcaag aatgtatcaa aaagagttgg aaaagcatgt gattgcattt      780
actgagggaa atctcact acaaccagat ttgaacaaag ttgatcccg cagaaactat      840
cgatatgtga caattccgaa ctttccagct tccaagaa aacttcgaga agataatcga      900
tatggcccaa aatcgtttt gcctcaaaaga tggcgagaat ttgattcgag gggccgtaga      960
agagactcat atttctatct caaacgtaag ctcgatggat atttgaaatg ctacaaaaca     1020
actggatatt ttatgtttgt tggacttttg cacacatgt gggaaattga cccagacatc     1080
acataaaac tgccagcact ggaaatgat tacaagaga tgcggaact tgttggtaga     1140
gaagagggtt tggaaaaatt tgcacgagtt gcccgcatcg caaaaactgc tgaagatatt     1200
ctgccagagc gaatttatcg tcttgttgg gacgtcgaat cagctacctt gagccacaag     1260
caatgtgctg cacttgttgc gagaatggtt tttgccgac cggacagtcc tttcagtttc     1320
tgccgaatc tctcgtctga taaatctatt tgtgtggaga aacttaaatt cctgttctc     1380
tatttcgaca aaatgtcaat ggatccaccg gatggtgccg tcagttttag acttcaaaaa     1440
atggataaag atacgttcaa cgaagagtgg aaagataaaa aattacgttc tcttcctgaa     1500
gttgaattct ttgatgaaat gcttatgaa gacacagctc tctgtacaca agttgatttt     1560
gccaacgaac atcttggtyg cggagtttta aatcatgggt ctgttcagga ggagatccgt     1620
ttcttgatgt gtccagaaat gatggttggg atgttgttgt gcgagaaaat gaaacaactg     1680
gaagcgattt caattgttgg agcttacgtt ttcagttctt atactggtta tggtcatact     1740
ctaaaatggg cagaacttca accaaatcat tctcgtcaga atacaaacga atttcgagat     1800
cgttttggac gtcttcgggt agaaaactatt gcaatcgatg caattctgtt caaaggatca     1860
aaattagatt gtcagacgga gcagttaaac aaagcaata tcattagga aatgaagaaa     1920
gcatctatcg gattcatgag ccagggaccg aaattcaca atattccaat tgttactgga     1980
tggtggggat gtggagcatt taatggggac aagccactga agttcataat ccaagtaatt     2040
gctgccggag tcgctgatcg tccacttcat ttctgttcat ttggagaacc cgagcttgcc     2100

```


-continued

gcaaagtgca agaaaattat agaacgaatg aaacagaagg acgtaacact tggtaagtca 2160
 tgtttttcaa tcttcagttg a 2181

<210> SEQ ID NO 10
 <211> LENGTH: 726
 <212> TYPE: PRT
 <213> ORGANISM: Caenorhabditis elegans
 <220> FEATURE:

<400> SEQUENCE: 10

Met Ser Lys Lys Phe Ile Glu Leu Gly Asp Pro Val Thr Gln Asp Glu
 1 5 10 15
 Lys Asp Tyr Glu Asp Tyr Val Gly Val Gly Phe Ala His Gln Val Pro
 20 25 30
 Thr Met Lys Arg Arg Lys Leu Thr Glu His Gly Asn Thr Thr Glu Ser
 35 40 45
 Lys Glu Asp Pro Glu Glu Pro Lys Ser Arg Asp Val Phe Val Ser Ser
 50 55 60
 Gln Ser Ser Asp Glu Ser Gln Glu Asp Ser Ala Glu Asn Pro Glu Ile
 65 70 75 80
 Ala Lys Glu Val Ser Glu Asn Cys Glu Asn Leu Thr Glu Thr Leu Lys
 85 90 95
 Ile Ser Asn Ile Glu Ser Leu Asp Asn Val Thr Glu Arg Ser Glu His
 100 105 110
 Thr Leu Asp Asn His Lys Ser Thr Glu Pro Met Glu Glu Asp Val Asn
 115 120 125
 Asn Lys Ser Asn Ile Asp Val Ala Ile Asn Ser Asp Glu Asp Asp Glu
 130 135 140
 Leu Val Leu Glu Glu Asn Asn Lys Glu Met Arg Asp Gly Glu Gln Val
 145 150 155 160
 Gln Gln Leu Ser Gln Asp Leu Phe Ala Asp Asp Gln Glu Leu Ile Glu
 165 170 175
 Tyr Pro Gly Ile Met Lys Asp Thr Thr Thr Gln Leu Asp Ile Thr Asp
 180 185 190
 Ser Glu Val Glu Thr Ala Gln Lys Met Glu Met Ile Glu Glu Thr Glu
 195 200 205
 Ala Asp Ser Thr Phe Val Gly Glu Asp Ser Lys Ala Thr Lys Thr Val
 210 215 220
 Arg Thr Ser Ser Ser Ser Phe Leu Ser Thr Val Ser Thr Cys Glu Ala
 225 230 235 240
 Pro Ala Lys Gly Arg Ala Arg Met Tyr Gln Lys Glu Leu Glu Lys His
 245 250 255
 Val Ile Ala Phe Thr Glu Gly Asn Leu Thr Leu Gln Pro Asp Leu Asn
 260 265 270
 Lys Val Asp Pro Asp Arg Asn Tyr Arg Tyr Cys Thr Ile Pro Asn Phe
 275 280 285
 Pro Ala Ser Gln Gly Lys Leu Arg Glu Asp Asn Arg Tyr Gly Pro Lys
 290 295 300
 Ile Val Leu Pro Gln Arg Trp Arg Glu Phe Asp Ser Arg Gly Arg Arg
 305 310 315 320
 Arg Asp Ser Tyr Phe Tyr Phe Lys Arg Lys Leu Asp Gly Tyr Leu Lys
 325 330 335
 Cys Tyr Lys Thr Thr Gly Tyr Phe Met Phe Val Gly Leu Leu His Asn
 340 345 350

-continued

```

Met Trp Glu Phe Asp Pro Asp Ile Thr Tyr Lys Leu Pro Ala Leu Glu
 355                               360                   365

Met Tyr Tyr Lys Glu Met Ser Glu Leu Val Gly Arg Glu Glu Val Leu
 370                               375                   380

Glu Lys Phe Ala Arg Val Ala Arg Ile Ala Lys Thr Ala Glu Asp Ile
 385                               390                   395                   400

Leu Pro Glu Arg Ile Tyr Arg Leu Val Gly Asp Val Glu Ser Ala Thr
 405                               410                   415

Leu Ser His Lys Gln Cys Ala Ala Leu Val Ala Arg Met Phe Phe Ala
 420                               425                   430

Arg Pro Asp Ser Pro Phe Ser Phe Cys Arg Ile Leu Ser Ser Asp Lys
 435                               440                   445

Ser Ile Cys Val Glu Lys Leu Lys Phe Leu Phe Thr Tyr Phe Asp Lys
 450                               455                   460

Met Ser Met Asp Pro Pro Asp Gly Ala Val Ser Phe Arg Leu Thr Lys
 465                               470                   475                   480

Met Asp Lys Asp Thr Phe Asn Glu Glu Trp Lys Asp Lys Lys Leu Arg
 485                               490                   495

Ser Leu Pro Glu Val Glu Phe Phe Asp Glu Met Leu Ile Glu Asp Thr
 500                               505                   510

Ala Leu Cys Thr Gln Val Asp Phe Ala Asn Glu His Leu Gly Gly Gly
 515                               520                   525

Val Leu Asn His Gly Ser Val Gln Glu Glu Ile Arg Phe Leu Met Cys
 530                               535                   540

Pro Glu Met Met Val Gly Met Leu Leu Cys Glu Lys Met Lys Gln Leu
 545                               550                   555                   560

Glu Ala Ile Ser Ile Val Gly Ala Tyr Val Phe Ser Ser Tyr Thr Gly
 565                               570                   575

Tyr Gly His Thr Leu Lys Trp Ala Glu Leu Gln Pro Asn His Ser Arg
 580                               585                   590

Gln Asn Thr Asn Glu Phe Arg Asp Arg Phe Gly Arg Leu Arg Val Glu
 595                               600                   605

Thr Ile Ala Ile Asp Ala Ile Leu Phe Lys Gly Ser Lys Leu Asp Cys
 610                               615                   620

Gln Thr Glu Gln Leu Asn Lys Ala Asn Ile Ile Arg Glu Met Lys Lys
 625                               630                   635                   640

Ala Ser Ile Gly Phe Met Ser Gln Gly Pro Lys Phe Thr Asn Ile Pro
 645                               650                   655

Ile Val Thr Gly Trp Trp Gly Cys Gly Ala Phe Asn Gly Asp Lys Pro
 660                               665                   670

Leu Lys Phe Ile Ile Gln Val Ile Ala Ala Gly Val Ala Asp Arg Pro
 675                               680                   685

Leu His Phe Cys Ser Phe Gly Glu Pro Glu Leu Ala Ala Lys Cys Lys
 690                               695                   700

Lys Ile Ile Glu Arg Met Lys Gln Lys Asp Val Thr Leu Gly Lys Ser
 705                               710                   715                   720

Cys Phe Ser Ile Phe Ser
 725

```

```

<210> SEQ ID NO 11
<211> LENGTH: 31
<212> TYPE: PRT
<213> ORGANISM: Bos taurus
<220> FEATURE:

```

-continued

<400> SEQUENCE: 11

Leu Phe Thr Glu Val Leu Asp His Asn Glu Cys Leu Ile Ile Thr Gly
 1 5 10 15
 Thr Glu Gln Tyr Ser Glu Tyr Thr Gly Tyr Ala Glu Thr Tyr Arg
 20 25 30

<210> SEQ ID NO 12

<211> LENGTH: 29

<212> TYPE: PRT

<213> ORGANISM: Bos taurus

<220> FEATURE:

<400> SEQUENCE: 12

Ala Tyr Cys Gly Phe Leu Arg Pro Gly Val Ser Ser Glu Asn Leu Ser
 1 5 10 15
 Ala Val Ala Thr Gly Asn Xaa Gly Cys Gly Ala Phe Gly
 20 25

<210> SEQ ID NO 13

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Bos taurus

<220> FEATURE:

<400> SEQUENCE: 13

Phe Leu Ile Asn Pro Glu Leu Ile Val Ser Arg
 1 5 10

<210> SEQ ID NO 14

<211> LENGTH: 16

<212> TYPE: PRT

<213> ORGANISM: Bos taurus

<220> FEATURE:

<400> SEQUENCE: 14

Ile Ala Leu Xaa Leu Pro Asn Ile Xaa Thr Gln Pro Ile Pro Leu Leu
 1 5 10 15

<210> SEQ ID NO 15

<211> LENGTH: 17

<212> TYPE: DNA

<213> ORGANISM: Bos taurus

<220> FEATURE:

<400> SEQUENCE: 15

gaycayaayg artggyt

17

<210> SEQ ID NO 16

<211> LENGTH: 17

<212> TYPE: DNA

<213> ORGANISM: Bos taurus

<220> FEATURE:

<400> SEQUENCE: 16

ckrtangtyt cngcrta

17

<210> SEQ ID NO 17

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Bos taurus

<220> FEATURE:

<400> SEQUENCE: 17

-continued

 atcatcacag gtactgagca gtac 24

<210> SEQ ID NO 18
 <211> LENGTH: 24
 <212> TYPE: DNA
 <213> ORGANISM: Bos taurus
 <220> FEATURE:

<400> SEQUENCE: 18

gcctgtgtat tcaactgtact gctc 24

<210> SEQ ID NO 19
 <211> LENGTH: 26
 <212> TYPE: PRT
 <213> ORGANISM: Bos taurus
 <220> FEATURE:

<400> SEQUENCE: 19

Glu Asp Lys Arg Lys Glu Gln Cys Glu Met Lys His Gln Arg Thr Glu
 1 5 10 15

Arg Lys Ile Pro Lys Tyr Ile Pro Pro His
 20 25

<210> SEQ ID NO 20
 <211> LENGTH: 26
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:

<400> SEQUENCE: 20

Glu Asp Arg Arg Lys Glu Gln Trp Glu Thr Lys His Gln Arg Thr Glu
 1 5 10 15

Arg Lys Ile Pro Lys Tyr Val Pro Pro His
 20 25

<210> SEQ ID NO 21
 <211> LENGTH: 26
 <212> TYPE: PRT
 <213> ORGANISM: Mus Musculus
 <220> FEATURE:

<400> SEQUENCE: 21

Glu Asp Arg Arg Lys Glu Gln Cys Glu Val Arg His Gln Arg Thr Glu
 1 5 10 15

Arg Lys Ile Pro Lys Tyr Ile Pro Pro Asn
 20 25

<210> SEQ ID NO 22
 <211> LENGTH: 32
 <212> TYPE: PRT
 <213> ORGANISM: Caenorhabditis elegans
 <220> FEATURE:

<400> SEQUENCE: 22

His Gln Val Pro Thr Met Lys Arg Arg Lys Leu Thr Glu His Gly Asn
 1 5 10 15

Thr Thr Glu Ser Leu Leu Leu Lys Glu Asp Pro Pro Glu Pro Lys Ser
 20 25 30

<210> SEQ ID NO 23
 <211> LENGTH: 26
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:

-continued

<400> SEQUENCE: 23

Glu Gly Lys Arg Lys Gly Asp Glu Val Asp Gly Val Asp Glu Val Ala
 1 5 10 15

Lys Lys Lys Ser Lys Lys Glu Lys Asp Lys
 20 25

<210> SEQ ID NO 24

<211> LENGTH: 26

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<220> FEATURE:

<400> SEQUENCE: 24

Glu Gly Lys Arg Lys Gly Asp Glu Val Asp Gly Thr Asp Glu Val Ala
 1 5 10 15

Lys Lys Lys Ser Arg Lys Glu Thr Asp Lys
 20 25

<210> SEQ ID NO 25

<211> LENGTH: 26

<212> TYPE: PRT

<213> ORGANISM: Bos taurus

<220> FEATURE:

<400> SEQUENCE: 25

Glu Gly Lys Arg Lys Gly Asp Glu Val Asp Gly Ile Asp Glu Val Thr
 1 5 10 15

Lys Lys Lys Ser Lys Lys Glu Lys Asp Lys
 20 25

<210> SEQ ID NO 26

<211> LENGTH: 25

<212> TYPE: PRT

<213> ORGANISM: Galus galus

<220> FEATURE:

<400> SEQUENCE: 26

Glu Gly Lys Arg Lys Gly Glu Glu Val Asp Gly Asn Val Val Ala Lys
 1 5 10 15

Lys Lys Ser Arg Lys Glu Lys Glu Lys
 20 25

<210> SEQ ID NO 27

<211> LENGTH: 26

<212> TYPE: PRT

<213> ORGANISM: Xenopus laevis

<220> FEATURE:

<400> SEQUENCE: 27

Glu Gly Lys Arg Lys Ala Asp Glu Val Asp Gly His Ser Ala Ala Thr
 1 5 10 15

Lys Lys Lys Ile Lys Lys Glu Lys Glu Lys
 20 25

<210> SEQ ID NO 28

<211> LENGTH: 25

<212> TYPE: PRT

<213> ORGANISM: Drosophila melanogaster

<220> FEATURE:

<400> SEQUENCE: 28

Glu Glu Leu Pro Asp Thr Lys Arg Ala Lys Met Glu Leu Ser Asp Thr

-continued

1	5	10	15	
---	---	----	----	--

Asn Glu Glu Gly Glu Lys Lys Gln Arg
 20 25

<210> SEQ ID NO 29
 <211> LENGTH: 31
 <212> TYPE: PRT
 <213> ORGANISM: Sarcophaga peregrina
 <220> FEATURE:

<400> SEQUENCE: 29

Glu Gly Val Ser Ser Ala Lys Lys Ala Lys Ile Glu Lys Ile Asp Glu
 1 5 10 15

Glu Asp Ala Ala Ser Ile Lys Glu Leu Thr Glu Lys Ile Lys Lys
 20 25 30

<210> SEQ ID NO 30
 <211> LENGTH: 28
 <212> TYPE: DNA
 <213> ORGANISM: Bos taurus
 <220> FEATURE:

<400> SEQUENCE: 30

gctgcgggtc tcgacgatga gtgcgggc 28

<210> SEQ ID NO 31
 <211> LENGTH: 29
 <212> TYPE: DNA
 <213> ORGANISM: Bos taurus
 <220> FEATURE:

<400> SEQUENCE: 31

gcgtctagaa ttcaactggc tectcaggc 29

<210> SEQ ID NO 32
 <211> LENGTH: 38
 <212> TYPE: DNA
 <213> ORGANISM: Bos taurus
 <220> FEATURE:

<400> SEQUENCE: 32

ccggaattcg ggtttttgt taatgaaat ttattaac 38

<210> SEQ ID NO 33
 <211> LENGTH: 29
 <212> TYPE: DNA
 <213> ORGANISM: Bos taurus
 <220> FEATURE:

<400> SEQUENCE: 33

tcagagcaga tgaactcgag cagtcacag 29

<210> SEQ ID NO 34
 <211> LENGTH: 61
 <212> TYPE: DNA
 <213> ORGANISM: Bos taurus
 <220> FEATURE:

<400> SEQUENCE: 34

ccaatttgaa ggaggaattc ccgccgccac catgaatgat gtgaatgcca aacgacctgg 60
 a 61

<210> SEQ ID NO 35

-continued

```

<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE: Kozak consensus sequence
<223> OTHER INFORMATION: Synthesized by oligonucleotide synthesizer

<400> SEQUENCE: 35
gaattcccg cgcaccatg aa 22

<210> SEQ ID NO 36
<211> LENGTH: 674
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:

<400> SEQUENCE: 36
agaagaaaat ggccaaggca tgctacaggt ggattttgca aatcgtttg ttggagggtg 60
tgtaaccagt gcaggacttg tgcaagaaga aatccgcttt ttaatcaatc ctgagttgat 120
tatttcacgg ctcttactg aggtgctgga tcacaatgaa tgtctaatta tcacaggtac 180
tgagcagtac agtgaataca caggctatgc tgagacatat cgttgggtccc ggagccacga 240
agatgggagt gaaagggagc actgcgagcg gcgctgcact gagatcgttg ccatcgatgc 300
tcttcaactc agacgctacc tcgatcagtt tgtgcctgag aaaatgagac gcgagctgaa 360
caaggcttac tgtggatttc tccgtcctgg agtttcttca gagaatcttt ctgcagtggc 420
cacaggaaac tggggctgtg gtgcctttgg gggatgatgcc aggttaaaag ccttaataca 480
gatattggca gctgctgcag ctgagcgaga tgtggtttat ttcaccttg gggactcaga 540
attgatgaga gacatttaca gcatgcacat tttccttact gaaaggaac tcactgttgg 600
agatgtgtat aagctgttgc tacgatacta caatgaagaa tgcagaaact gttccacccc 660
tggaccagac atca 674

<210> SEQ ID NO 37
<211> LENGTH: 200
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:

<400> SEQUENCE: 37
aaaaatagtt gtcaagactc agaagcagat gaggagacaa gtccaggttt tgatgaacaa 60
gaagatggta gttcctccca aacagcaaat aaaccttcaa ggtccaagc aagagacgct 120
gacattgaat ttaggaaacg gtactctact aagggcggtg aagttagatt acatttccaa 180
tttgaaggag gagagagtcg 200

<210> SEQ ID NO 38
<211> LENGTH: 29793
<212> TYPE: DNA
<213> ORGANISM: Caenorhabditis elegans
<220> FEATURE:

<400> SEQUENCE: 38
gatctcgaag taaaaactca cgcagaaaga gtcctcctc ctttagcatg agaatccaac 60
tttgaatga taactctggc aacatcaaca gtttgagaga aagcacgtgc ttgggcttca 120
caagcttgtc caatagaagc atccatcaca aaaacaacat tatctggtgt aactgcgttg 180
gaaacttggg gcatttcttc gaaaagtgaa gcttcttgc tgtgacgacc tgatgtatca 240
acaatgatga tttcgaaccc ttcttgctgc aaaacaaata ttattaaacc attttctgt 300

```

-continued

gataaattac cgtgaatttt tctactcctt cggcggcaat ttttacgggg tcaatttcag	360
agtatgatcc atagaaggga atacgagctt ttgtggcatt ttgctttaat tgatcaaaag	420
ctccagcacg gaatgtatcg gcacagatca gacatgtttt ccatcctttt ctttggtagt	480
aatcgcctat ctgaacttga aaagtgttga aaagtgttg gaagtttact aattaaaaaa	540
tataatgttt gatggtgtgt gagctttcta ttgtaattca tggaacgaac cttggtacaa	600
gtcgtagtgt taccggaacc ttgaagacca acaaacatga aaacgttgcg acgtcctttt	660
gttgggtgta aaggagtac accaggatcc acaagcttca gcagttcatt gaatactgtc	720
ttctgaatgt accgacgttt gtttgcctct ccgacgatct cttcgaaatt aatcgctttt	780
ctgaaaaat ttattaaatt taaatcctaa atagcgtaaa aatttacttc acgttgcctt	840
taagttgctt tacaagacga atatgaacat cagattcaat aagagctgta cagacttctt	900
tcagcatcaa atccagctcc ccctcattga taacgggtct ctgaccgagc tttccgatcg	960
catttcggat tttccgcccc aaatcggcca aaaccatttt gaactgaaat ttgaaatgct	1020
ttaatttggt taagcataga attaaacgcy ttttaaatcg agagcaccat aaaaacagtt	1080
tggagaaaaa tcgataattc ttgtaggaga ttcagtcctt gtggttttct tcggcttctt	1140
aatcattttt tgacgacata gtggtatttc acaataggtt ttttcaagac acaacagatt	1200
tttcacaaag agtagagaag aaatggaaaa ctgtagattt cttctcgaag agccgagaaa	1260
ggcaaggat tggaagtta aaaaggtaat gtttctttat tcttttttca aaacaataat	1320
aaatggaaaa tatatattta tagataacaa tttcagacag ttaaaatcac gtgaaaaatt	1380
caaatttcaa cacaaaaatt gacgagtgga accccgttgt tgcgccttga agagtaacgc	1440
ttgcgcgttt gacgatttta ttgacgcgtt tctggtgcat gcgggaaatt ttttattttc	1500
aacttttttc ctgtttggtt atcctttttt aattgaattc tcatgatttg aaagcttga	1560
aaaaatattt tttgctcaaa aacatgcgtt ttgtaaaaca ttgattagat tcaaggcaat	1620
taatggattt ttgcacgttc caaaaaaaag gaaattcatt ttttgaaaat tttgataatt	1680
taataatgaa aaatgttcca tagatttatt caatgccatc cttctctata atctcgaact	1740
tccgcatcct tcaactgtgg tagaggattt tgcaatacca tatagtcgta ataataaact	1800
ttagtgaaca aatccaagac atcagctcct gagtaaatga atgatttata aaaactgctg	1860
attttctcgt aggaagaaaag agaatcagct aataatcctg cgttgcctat tctgtcaggc	1920
cgcttaaatg ttaaaaaata aaaacgtttt aagctaattt tgtatgtcta gaaactctaa	1980
ctcacaagca tttctgcata cgccggatta gttggttttg caaaaagcga gtaacttaca	2040
aaagtgaatt tttgattcat ctcttcocatt tcacaaaacc aattttgtgg tacgtatttc	2100
atatgatcct catccacttt tttagttttt gaatgtattt gtgtgagttg tgtccagatt	2160
tgaataagat aacatctcag atccaacttg caattgaagc aagaacgatc ttctctgaaa	2220
ttttatatga ccttaaaactt tatacttttg tagtttcgtc gatatctgat cgttcagttg	2280
tataggtatg tacatctcta ggtttatgtg ctacacgaaa atataatttg ttttacctaa	2340
cacacgcac cataaaatga tctacaaatc gttcaattgg atcctgtcct ggaataaata	2400
atttccaatt cgtaaagttt gcattcaact cattttctcg tttcaaatcg tcgatatccg	2460
caaaatattg tagtgaatca ctatcacaca ctctgaaaag cacaatattc atatttcgta	2520
gttaataatg aacctcacga ttcattcatta aatttctctt ggagcccgcga taatacttgc	2580
tgcccaatta aagtatcagt ttcacagatt gcagttctat catttccgat agcctcaaat	2640
aagatttaat cttaagcgag tgttctgatc aatttaaata tttgatactc accgcaagtt	2700

-continued

tcttcgaaac	ttgttcgaaa	gctggaat	tagaatatcc	ttcaaaactt	tttctctcgc	2760
cctcatcaag	ccataataag	ttttgatcag	caatatattc	gaataaatta	gtctctgata	2820
aatctcgtat	cacaatcttt	ttttctactc	taaagaatac	aattttgata	agaatgataa	2880
taattataat	tataatagtt	cgctcgtgag	ttgatgaaga	ccacataatt	agtttaatgg	2940
caagctatgc	aacttgttga	atactaatag	gacttagcaa	atcttatctt	gaaccttttt	3000
cattcgaag	aaaaatgaga	tcgaatctcg	ttcaaaactgt	ggagtagtca	gttaagaaac	3060
ttgtttctag	tttgtgagga	gacactggag	aacgtgaaag	tattacccat	acgcaatatt	3120
tttgccgga	aaaatcgggt	acccggcttc	gacacgacag	tttttaaaac	ttgtaaatag	3180
gtatgtaaaa	gaaaacttta	attttaaacg	tggtgtttcg	gaattttcat	cgttttgtca	3240
tagttattct	acaaataatt	atttatgaaa	aaaaaactaa	aatataacta	taataacacc	3300
tgaatattaa	caaatcgtac	gaaaaaaaaac	tatgaaaaaa	atggatgaaa	attccgcagc	3360
aacgagagtt	tgaattttca	gtattcttta	aaggcttacc	gatttcaata	aatagtgaca	3420
ctgaaaattg	tagtttttaa	actagttggt	tagtatcatc	aaatattcaa	tccttcaaaa	3480
attcctcaat	attaacgtat	tttctcta	tgtcttcatt	atctaaaaaa	aagttgcaat	3540
atatttttcc	aggcagaaat	agactttcac	aaaacacatc	gacacttcga	atgagcaaga	3600
agtttatcga	actgggtgat	cctgtcactc	aagacgtag	ttatagtttt	tattacttga	3660
acattatcat	ctttttacag	gagaaagact	acgaagacta	tgtcggaggt	ggtttcgcgc	3720
atcaagtccc	gacaatgaaa	aggcgggaagt	tgacagaaca	tggaaatact	acagaatcaa	3780
aagaagatcc	tgaagagcca	aaaagccgtg	acgtatttgt	ctcctcgag	tcaagtgatg	3840
agagtcaaga	agattcggct	gaaaatccgg	agatcgctaa	agaagtgtca	gaaaattgtg	3900
aaaatctgac	agaaactctc	aaaatttcta	atattgagag	tttgacaat	gttactgaaa	3960
gatctgaaca	cactcttgat	aatcacaaaa	gtactgaacc	aatggaagaa	gatgtaaaaa	4020
acaagtccaa	tattgacggt	gcgattaatt	ctgacgagga	tgatgaactt	gttctggaag	4080
agaataataa	agaaatgagg	gatggagaac	aagtacaaca	ggtcaggaaa	ttttacaagt	4140
gaatgaata	agttaatcac	caaatgaaat	aaggacattt	cccatcagaa	aggtcttctg	4200
aattttaggt	gtaatgttaa	ttttttgctg	tagtttttcc	cattgtttga	aatttttgcc	4260
aaaattagtt	attgcatacc	cttcatgttt	ttgaagattg	tttaggaatg	agaaaacatt	4320
ttggacgctt	ttattattag	gacaccaaac	actttttggt	gaaaaaacag	ctcgtttaa	4380
aaaagctttt	ttcaaaaaat	ctgacgcaag	gcttgtgaat	tttcgttttc	ccctgatttt	4440
taaaatttct	cctaaagttt	tttgctaata	tttttcgcta	tcgcgtaatt	tactagttaa	4500
tcaacaaaaa	attttttttt	tttcatagat	tttttataag	tttttgaaaa	catagattta	4560
aaacttaaac	ttaaattttg	acaaggcgag	aggaaaaaat	taaaaattgc	tgaacattca	4620
gatgccgggt	accttatttt	tggttcaaaa	atcccaatat	tacgcgtctg	ggttatagtc	4680
atttgccttt	attaaattaa	tgggttctct	tggaaaagta	agttctgttt	tgttttcagc	4740
ttatcacttc	atcaaacgga	aggaaagggt	gattaaggaa	agtaaacata	ttttatgttg	4800
ttcttctcac	ttcctccatt	tcgcaataat	ataactcgag	aaatatagaa	ttttgttcga	4860
agttttcttt	ttccttcaac	attttaaaata	ttgttagtat	taccagaaaa	aatagaaaaa	4920
atcgaagaaa	tttgcaaaaa	agcagacgta	gaggctacgt	acttcttaag	cacgcccttt	4980
ttcttttaaa	ttgttcgggt	cgtaccgaga	tccggtacct	tattttacaa	cgttttctgt	5040

-continued

tccaaaaata	ataatgtact	gcagttgtca	caggatttat	tcgctgatga	tcaagagcta	5100
attgaatata	caggaattat	gaaagacact	acaactcaac	tgatataaac	agattctgaa	5160
gtggagactg	ctcaaaaaat	ggaaatgatt	gaagaaactg	aagcagattc	gacatttgta	5220
ggcgaggatt	caaaagttag	acaaaatcat	tctgacaagg	attcctgcga	gcaactcagtc	5280
aagagcgagt	cacggcaact	cggtccaaaa	ccattttctaa	ttagtaaaact	ctcaaaaacc	5340
acaactaaat	agcttaaaac	ctttgtaaat	tagcttattt	ttgctaatta	gcaatgattt	5400
taagctaatt	agttgtggtt	tttgagagtt	tactaattag	aaatggtttt	ggaccgagtt	5460
gccgtgactc	gctcttgact	gagcacaagc	aaacttttgt	ggatggtgag	aatcagcggc	5520
aaagtggcac	tactagttag	gaagttgagc	cagattctca	gattaatttg	gtaagacaaa	5580
gaaaataata	atthttattac	ccagatgcat	atthttcatga	ttctgatgca	aaaaatcgg	5640
taccgatct	ggatactaca	atthttgtaa	aatgcgaaaa	ggtttgcacc	tttaaaaaga	5700
actgcaattt	caaacacttg	ttgctgtgga	ttgtttatcg	gtttttaata	ttttttgggtg	5760
agagtaaagt	agaaaagcga	gttcccgcat	tatctgtgtg	cgatttggaa	tacagtactt	5820
ttcaaagacg	cacaccattt	tgcataatac	aaacatttgt	cgtgtcgaaa	cgggtaccg	5880
tgatthttgca	ttaaaagttg	caaaatttca	catagthttt	ataatthagg	ctacgaaaaac	5940
tgtgaggaca	tccagttcaa	gtttcctgtc	aactgtttca	acatgcgaag	cccctgcaaa	6000
aggacgagca	agaatgtatc	aaaaagagtt	ggaaaagcat	gtgattgcat	ttactgaggg	6060
aaatctcaca	ctacaaccag	atthtgaacaa	agttgatccc	gacagaaact	atcgatattg	6120
tacaattccg	aactttccag	cttcccgaag	aagtacgttg	ttcaataaaa	catactaggt	6180
atataattaa	ttatthtcaga	acttccgagaa	gataatcgat	atggcccaaa	aatcgthttg	6240
cctcaaaagat	ggcgagaatt	tgattccgta	catttctatt	gaattaatta	tatactactt	6300
actagaaaca	ccatggagaa	agaatgcaaa	aaattgaatt	ttaaaaacta	atthtttaat	6360
tttggtcaaa	ttthtcagtt	gaatthtaac	caaaatgaaa	actgcgacca	atcaatgact	6420
tttcaaaatc	actthttcaac	caatcaaacg	gagtggtctg	ggctcgaaga	cgctgattgg	6480
ttcggaaatg	ggcgtggttt	ctcatthttg	agggaaattca	aaaaaaggca	tttggtcaca	6540
gttgaaaatc	atgthtttcaa	aagatgcatt	thttatthct	tctcgattht	thttgattht	6600
ctthttgtgt	atthttcgaat	ttaaaagttg	tgtagtcgaa	thttthttatt	gctthattag	6660
actcaaaatt	thctgaaaaac	gccaaatthc	ataatgaaac	thcttgaaaa	ctcttcagca	6720
aaaagttatg	acggctcaaa	aaatggccta	aaattagtha	agattggaga	thtgaccgac	6780
thgtcaatgt	cgcagcggct	ggaaaacaatt	thttthtga	tcaccgtcaa	atthttaagta	6840
tacaacttga	ttatthttcgc	thttaaactt	tatthtagtha	thttaaaagtc	gatggaccggc	6900
gagthtttggc	tcaaaaaaat	taaaaatctc	gccgtccatc	gattthttaa	taccttaatc	6960
aagaataaaa	caaaaggttag	gcaacttgta	tattcaaaat	thgacggtga	thgcaactth	7020
taactaattt	caggccattt	thtgaccgct	cataacttht	thctaaaaag	thttcaagaa	7080
gtthcattat	gaaatthcgg	gtthtcagac	aatthtgagt	ctaataagga	aataaaaaaa	7140
atthcagtag	accaccttht	agaaaattht	ggatthccgc	tacgctaata	cacctthaat	7200
caaaaatatt	tgaagttatt	caaagthaaa	gaattatatt	thcagagggg	ccgtagaaga	7260
gactcatatt	tctatthtcaa	acgtaagctc	gatggatatt	tgaaatgcta	caaaacaact	7320
ggatathttt	tgtthttgtt	aagthtttga	aatacaatthc	gtthtgaagat	thactctatt	7380
thcagggact	thtgcaaac	atgtgggaat	thgaccaga	catcacatata	aaactgccag	7440

-continued

cactggaaat	gtattacaaa	gagatgtcgg	aacttggtgg	tagagaagag	gttttgaaa	7500	
aatttgcacg	agttgcccgc	atcgcaaaaa	ctgctgaaga	tattctgccca	gaggtatgat	7560	
ttatgagata	tacagcattt	cctctaatag	tattgcatat	aaacatttca	ctttgaggtt	7620	
atatcttggg	ttattttaaa	aatatcaata	aatacaaaac	aatagaaaaa	tgataaaaaa	7680	
acattttgtc	agttgataat	ttgggtatag	tattcattca	taatttgatt	tttttagcga	7740	
atztatcgtc	ttgttgggtga	cgtcgaatca	gctaccttga	gccacaagca	atgtgctgca	7800	
cttgttgcca	gaatgttttt	tgcccgcacg	gacagtccct	tcagtttctg	ccgggtgagta	7860	
atacaagaat	gctcatat	ttagaatcaa	tatttgcaag	gaactttaat	cttacgtacg	7920	
tcttaagatg	agcatttttcg	cacatatcct	acgcgcacga	gtctcgacac	gcgaacatcg	7980	
agcttctgta	actcgtatca	atctacaagc	cgttattaca	tcagttttta	atgaatttta	8040	
agaaaatcgt	gcaaaagtag	tgctgagagc	cattcgcgta	agatatgggtg	agatttatca	8100	
tttttagacg	tctagtggat	atctaacaaa	actttataca	ttttatttc	agaattctct	8160	
cgctcgataa	atctat	gtggagaaac	ttaaattcct	gttcaactat	tccgacaaaa	8220	
tgcaaatgga	tccaccggat	ggtgcccgtca	gttttagact	tacaaaaatg	gataaagata	8280	
cgttcaacga	agagtggaaa	gataaaaaat	tacgttctct	tcctgaagtt	gaattccttg	8340	
atgaaatgct	tattgaagac	acagctctct	gtacacaagt	tgattttgcg	aacgaacatc	8400	
ttggtggcgg	agttttaaat	catgggtctg	ttcaggtagt	tatttaaagg	aatataagaa	8460	
tttgaagttt	tatttttttt	atgcaggagg	agatccgttt	cttgatgtgt	ccagaaatga	8520	
tggttggaat	gttgttgtgc	gagaaaaatga	aacaactgga	agcagatttca	attggtggag	8580	
cttacgtttt	cagttcttat	actggttatg	gtaagtctag	actttcaaaa	aaaactgttc	8640	
caatatgtca	atatatttca	ggtcatactc	taaaatgggc	agaacttcaa	ccaaatcatt	8700	
ctcgtcagaa	tacaaacgaa	tttcgagatc	gttttgagc	tcttcgggta	gaaactattg	8760	
caatcgatgc	aattctgttc	aaaggatcaa	aattagattg	tcagacggag	cagttaaaca	8820	
aagcaaatat	cattagggaa	atgaagaag	catctatcgg	attcatgagc	cagggaccga	8880	
aattcacaaa	tattccaatt	gttactggat	ggtggggatg	tggagcattt	aatggggaca	8940	
agccactgaa	gtgtatgtta	tttcattcgt	taaatattga	agatggagga	gagtgaatgg	9000	
ggat	tttgct	tcttttgcaa	aatggcctcc	ctatgtacct	gaaaaaaaaa	tgaaaaaatc	9060
gagaaatatt	gaaaacaaa	caacgaattt	ttcacaattt	tgctaaatt	tttgaatttt	9120	
cgcaaaaatc	ggaatcagcg	attcgtcca	cccatttttc	cgccaatcat	ttataatgtg	9180	
cgagactcaa	aaactgat	tggctagaaa	gtggcgttag	cttcttattt	cggaggaat	9240	
tcaaataggg	aagttaatct	aaatataaac	aatctcgtta	aaaaatgttt	cttttttcaa	9300	
tcttccctat	ttgtttaaat	ttttcttttt	aaagatcgtc	taaaagctac	cagtatctga	9360	
ttcaattatc	ggtttttttc	agtcataatc	caagtaattg	ctgccggagt	cgctgatcgt	9420	
ccacttcatt	tctgttcatt	tggagaaccc	gagcttgcg	caaagtgcaa	gaaaaattata	9480	
gaacgaatga	aacagaagga	cgtaaacatt	ggtaagtc	gtttttcaat	cttcagttga	9540	
tttgaaaaag	ttgtatcgag	ttggaacag	cttttaatct	aaattctgct	aacttacagg	9600	
catgctat	agtatgataa	acaacaccgg	cttgccacat	aagcactttg	aattctacgt	9660	
cttcgataga	atcttactt	atctcagtag	ttcgggaagat	gttgagctt	cgaaatcatc	9720	
accttcagta	tcccagcat	aattcgaatc	gcccacacgg	ccataaagac	cggttccttt	9780	

-continued

cgattaaatt	ctgttaaata	tgcatgctcc	gtctttaaaa	aatcagtcce	cgtattttaa	9840
acgttttgat	tttaattgtc	atattattat	ccgaaattag	tatactcgcc	gtcatgaaag	9900
cccagagatat	ctagttcgca	agtcagaaat	ttttcggagc	atcgtcgtga	tatatgaata	9960
aatacattcc	tgtttttcac	aagtgtagt	tgaaaccaat	ccatgcagac	gtttatttct	10020
gaattaattt	tgaacagat	ttcagagaca	gtgaggttga	cattagatat	gggcaagtaa	10080
caataacagc	agggcagtta	ttatgattat	ggatgctgat	ataggaaagt	cagaacagta	10140
taatcgacga	gaataaaaag	agatgagaag	ataggcgaga	ataaagaacg	ttaacgaaaa	10200
tactgaaga	gctacatttc	caacagaata	agaaatgtag	ttggaaatcc	ctaatcaaac	10260
agaaaagcga	gaaatcatga	ctttcgagat	aaagagattt	atctgcaaac	aattcttgaa	10320
cataaaatta	aagcaccaca	gactgtccaa	attataaaat	cagtttctcg	ctacagtctg	10380
gggtactct	agttccattc	aaaaacttct	tgcaaacaaa	gagaaataaa	cagacttgta	10440
cgggacacat	ataaaatcta	agcatgcttt	gaaaagcggg	gaacatacga	tctattcggg	10500
gatatacata	tatatatata	tatttcatct	catctagagg	atcaccatcg	ttactcatca	10560
aattggttgg	tgtggtgaa	gttatgaaaa	gagcaatttt	aaccgaaaat	caccaaaca	10620
gaaaccaaat	taatgataa	tcgacgagaa	tcatgatgag	atgatgattt	gcttctagca	10680
gaagtttaga	agcacatgct	atcattcatg	ctcacgatga	cgataggttc	gttatgcatt	10740
cttgaagcca	atgacacttc	cattgctcct	tctcttgcgc	tcacacaatt	tccattctcg	10800
tcgtaaatcg	ttcgactttc	gaatatccac	atcttaccgg	gcggcacttt	tggcctgggt	10860
tggcagatct	gaataaaat	ctttcataa	tttaaagtct	gatatcccg	gaaacaatag	10920
ctgaattgaa	acagaagaat	aaatctcacc	tgaatatcca	cgtttgcaag	tgaagtttct	10980
ttcaaaaacg	gctgtgttcc	ctcatagaga	tcactgtaac	cgtaggagaa	tggacggagc	11040
aagtcttca	gcttctcat	gtgctgttca	tttgccggtc	tcacgtgctt	ctttacggga	11100
gcagtggcgg	cagcatacgc	tgatttagat	gctcgcatag	tgctattcct	acgactttga	11160
ggcttcttt	cattgttcta	atgctggttc	atgtggaacg	gtgctgtttg	aaatcctcgc	11220
atgacttgag	acgattgatg	atgctgatgt	ggaggaatgt	ctgcaagatt	atagttatgt	11280
attaaaaatc	aaaaatttgt	gtggttccca	ttttaaaata	aaaaaaaaat	atttttacgc	11340
actttgctga	ggcaaccgat	aactatttcc	tcgctggcga	ctacttctct	tattgtgagc	11400
attatagctc	atgttctcat	gattgagttg	acctgaacga	tcaaggttaa	aactaggcct	11460
aaaactagtc	aaaattactg	agtttctcct	tccacgtcgt	ctgtcgagca	ggctccgagt	11520
acatttttac	tggaaaaacta	taataaatta	caaaaatcac	gccgaaaatg	gggaaaagaa	11580
ttgaaaaatt	gaaggaacac	agaacatttt	ttcaatgcgt	ctctcacggt	cgagactact	11640
gtattcgtgg	tgagacccaa	ctccctcata	aaagcatgcy	cctttagttt	tttaatttaa	11700
ttcatgttgc	caatattggc	caattaattt	caagagactc	tgattgaaag	tgttataaatt	11760
aaactacata	tatttaagct	ttcagcattt	ttttcaatgc	acttgagacg	caaattgaat	11820
aatcaggcac	gtaatgtgtt	ttcagggagc	actataaatt	gtacctttgc	tatccagtgg	11880
gttctttaat	tttccattc	caatcgattt	tttctccac	tctggcagtt	tctttgtcat	11940
cactggacga	gggcattgga	atgggagatg	attcatgtga	caatccacac	atcctgcaat	12000
aatgacatta	tttttataaa	atgttaagat	gatatgctta	ccaggagtaa	atatcatatc	12060
cttttcttta	ttagttggct	tagccttgcy	gccacgttta	ccatttgaca	ttatagttac	12120
ctgaaaatcc	aaaaaattag	atattcaaaa	aggtataaat	ataatataaa	tgcgatttgg	12180

-continued

taaatacggg tgtaatgggc aaccattct atacaggaaa accaaaaaat tcccgcaaaa 12240
 ttatTTTTT ccgaataaaa tgatctactt tgttttatgg tgcgctcta tgtggtatga 12300
 cccttcgatt agtagataga aaagaaaaag gaatgtacga gaatcgtt tattatttat 12360
 ttttgaaaa atcccagaga cataaaaaat cacacagaaa agggaaacag ttttctgac 12420
 aatgttcaaa agtttggtt caatcagcac taataatgtg aaaggtaacc gtatcaatag 12480
 tgatatttc ttattaaaa actgttcgag actacaagaa ggctgaaaa agcccgcaac 12540
 gagactaaa ttcgaaattt cgaattaggt ttaaagatc agaagatcgg cagaaaagta 12600
 tctgataaaa atataagaaa tcggaatagg aatgcatga ggaggtagaa atatggtgaa 12660
 gagatacaga agaagtaggt aagatcggat gaacttgaag cacttttga gatttttgat 12720
 ggtgaagtty gtggatgtag acgttctatg gaacatctga aaattaagat ttttctaaaa 12780
 cacattttc atagaatata atagaatgcc aaatagagaa actagactta cttgaatttc 12840
 tttcgatttc tgtcttcaa ccttctaact gaaatcaact ttcgacgtgt tctcggtgtt 12900
 tcaacaacac catcaacaga aactcagca ccaattcag catcggaatc atcagaagaa 12960
 gactcatcgg aatccaata gaaatggat ttagtattca tcaattcaaa agaatccaat 13020
 gatactgtcg attcagcaag ttggactgaa cttgatggtt gactacgaac ccattgaggg 13080
 cgtcgaggca gaagtcgaga gtatgaggat gcaacgtgga ttgatgatga cgtcaacaat 13140
 ctttgggtgt gagatgaaga agtggtgtat gcagatgtt acagacggaa tggagatgag 13200
 tgaagagcaa gaagacatct gaaaatttga aacgttgtt atgtggacag tactgtaaag 13260
 atcttacctt ggatcataac tacttgccct ctgttttctc ttctctgac ttctacttaa 13320
 aagcatttcc gtctcgatc tccggttact tgaaaatcca actccagaat tttcagcaca 13380
 aagctgctct cccgaaccgt agactgttgc accacgttga ggggttgaca aggatctgaa 13440
 atcagatggt taaagcatgg caagtagagc aacaatgtta accaaaattt ctgaaacttt 13500
 ttcgaaata gtcaaaaatt gacaataact cagtttcacc tatcatagtt ttggaagtca 13560
 accaaaaatt tttgaaattt cataaaaatt ccaactttc taaaaatttg gaagattgat 13620
 atgattgata tgaagattt tatatattt ttaacctggc agacgatact tcaccattaa 13680
 agacacacat gtggagaaga attattttac ttttagtaat ccaacgtttg cacttacctt 13740
 ggagcatgca agcttttagt cattaagct ggaattctag atggagtct tcttgggtgc 13800
 gacattggtt aaataaacat tctgtgttca ttgattgatg atgacgtcat agaaccacgc 13860
 ccagatgaca atggattacg gtatgcatca gaatcagtag attcattcaa tttctagtc 13920
 atttctctg ttttctgaa aattaattt taattaaaga tctaacaaaa atctgacct 13980
 tacattaata agataatcaa catattctaa ctcatctac gtttcattat tttctaattc 14040
 tggctcttc tcatcgaacc gttcggggc attgtgtcgt tgcgggcttg accgtttttt 14100
 gaatttctga aatgttttcc atgcaattt tgttcttatt tgtgtgcat atacagtga 14160
 aatcaaaaac tagtacaaac taattcgtt tagtaataa aaaatcgtg taaaatctca 14220
 gcaaagccaa gatcttggcg ggtccttata tccaagttt gttgccattt tatttcagat 14280
 attcttttcg aaagtcagaa aatttgaatt tagaatcga tggaccatt tcttgtttt 14340
 ttttgttga ttttttaact gtacttttt cgtcagcata tttttcact attaaaacag 14400
 aatattcatg acaataatc cacaaaaaa cgtactttaa tatcatagtc gattggttca 14460
 gaattggaac gagaacctc gacgcgtcga ttgtcagatt ctcgattgat ggacgacgtg 14520

-continued

ctgactgaaa atttctggat tgaaaaaata ttcaaatgaa aaaataaatg agaaactcaa 14580
agtctaaaaa atgaatgttg ttaataacga atatttctga tgagaagagg atagagaaaa 14640
aaaaagagtc taataaaatg catgtgatat cctgcataaa aatcccttct ttttccacta 14700
atccttcgct caattcattc aaatagaact ttgatttcta ttagagtga ggttggttga 14760
acaattttaa taaattaaca ataagccata aaacctcgaa acgtaccatc atcattgagt 14820
ttgaaaaagt ggacggatcc gagtcagtc cctctggaac aaatcgttcc agagcactga 14880
aaacgacaac gttctcccca cagaatcgga ttgtctctc gggaaattgc gcctcgacaa 14940
acgatcctga acctgaaaat tttcgatttt tgtaagctca atggatttta aactgaaaat 15000
gtagtcaaga agtcaagaaa aactgatgga gttctaaatt cgggtgttagg ttttggaaaag 15060
atcgtcaaac aaacaaatgc ataaaactag gtagggaaca aatagtgaaa tagaaaaatg 15120
aaaggcgaca actgccggga gcaagagtac acacaaagaa aaaaagtgc ggaagagcac 15180
agagagcgtc agtccatcag aactgcatag ataaatagat aaagagaaac atgaaacata 15240
aggccacccc ggagagacga caggccagtt ttccggtgaa gatgagagtg cgagaattag 15300
ataagaaaac ggaattgtg atgaaacttt ttcaatcaca acttctagaa ttataagaga 15360
cacctaaagt aattagataa gtgttttaag tgatatttta gattcactgt atcatgttta 15420
aaaaagatat ttcaaaaata tatacctgat ataggaggcc tcctctgagc accgaattgt 15480
tctcgagctg tttccacgag catccgctca cacattgaca taggccgtcg acagccagga 15540
gttgccacct gaataaataa ttattcaatt taaacctaat ttagtaatgg taactttgta 15600
aatgatgggt gatagctcat ataaaattg aattggttct aaagtatac aaattttaat 15660
ttcggtaaaa cttatgaact gtacttttga gttatactat tacaataata ttacccaaat 15720
tattgtatc agatttttgt aatcagtact aacagatttt aggcaacgtc ctgccagaaa 15780
catgggaata tatttgagca gtttttagta agttgccaca gcttgataa gggaaattgta 15840
tcaaatgta cttaatactt tctaagcact gacatagtga actacaaaag tcggtattat 15900
acaatgccac tacaataaaa aatattcaga attcgactga aaaatgagaa aaggacaac 15960
tgaattggac acacgatgac gtgattttca agaacacaaa aaaaagaaaa agaaatcgaa 16020
aatgttggtt gcctcttttt ctttttctat atgagctaga atctcgaatg catgcctaatt 16080
ggagccactc gctctcgttt ctctaagtct cttctacca gtcttttgc caaaaattgc 16140
gatgtcgcag gcgtcccgtt tccgocgtc acggagacac cactatcggc accagatgat 16200
cgtgtaaaga caccgtcttc gttgacttca attgctgaaa taagaggaat tagttttgaa 16260
ttggaaatct gattaataaa agtccoccta ttcaatctaa ttaattttta aacacaaaac 16320
ttactattgt ttaatgggtg tgacgaatta gaagaagttg attgaaatgc gttaacattc 16380
cattcaaagc tatcttcatt ccattctgct cgttctttta tcctttcact cactctcga 16440
gggatgaagt tttcaacaat aagaagcctg aaaacttata ttattctatt aaaaaaatg 16500
aacatcaaat ctaacgaaa gaaaattctt ctggggggaa aaggagagaa ttgtgagaat 16560
aaagaacctg cgctgtcggg atcaaattac actatttgaa ttcaattag aatcagaaag 16620
aaagtgaag aatgaaaat gagtgagaat ctattaaatt gtaattgaga tatcactgaa 16680
cttacttcaa cttcaattcc ttgctcaatt cogcaattgt ctgatccaga tcttgctgtt 16740
cgtcggaaata tgctccagac acatcacgaa tctcattgcy agcctgaaac attcacaac 16800
cttatcttga cacctggtac atctgaagtc aaacctgtct caactttatc aacatcttct 16860
ttaaactttt tgtctttgcc tccacotctg tccttaaatc cgaatgtt tgtttcaaat 16920

-continued

```

ccacagtatc ctcttctcgc cgttccaatg cttcaacat tctctttcc cttcgttttt 16980
gttcggcaag ttctcttctc ttcttttcca gttgtgcatg ttgttctttt gtccttgatt 17040
ctaacccttc atcttcttca gatcctacga taagtcgact ttgaatgttt gctattcttt 17100
cagcaaccceg tgcttgttct atccgttctc tttctaacag atcatgcttc tcttgaattt 17160
ctcttattaa tcgatccttt tcatgtttta tcagtgaatc atctttttga attgcttcaa 17220
tatcatcttc aagttttgct cgttcagcat cataaaaaact ttgagttgct ccatccctcg 17280
atcttgctct tctctgcttc agttgctcac ggagcatttc aatttcttct tgggaattctc 17340
gcagtaaage atccttagga tcttcattaa ttttcggttg attcttgatg tttttagctc 17400
gatttgcata tcgtaatgta ccaagtgtct cctcaaaatt gtaacttgca ggtccaatac 17460
aagcaacatc aactgtcttt gaatttccac cgagagaatc ttgaagaagt cgagtcagtt 17520
ttgaatctcg ataaggaata tgggcagatt tcgcatccac caatgcactg attacatttc 17580
caagagccga taatgaaaga ttgattttcg tagcttcttt aaacttttcg ccagttgctc 17640
ctgttttcga ttgccgttct gaaccagcta aatctacaag atttagtcga ccaactgtaa 17700
tatgactttc tccgtcttca ccaattcggg aacattcaac agtaatgata aagatagcgt 17760
gggaacgaga cgaatgctca ttcattgttg ttcgcctgaa aattttagta aaatcaaatc 17820
caacggcgac cacagaaata acttacccta cagaccgatg cccatttctc cgaatcatca 17880
cttcgtgtat ttcacctact gtccttgta attttgactg aaactttgaa atttatagtc 17940
gtcttctatt tcagaaaaact atcaettacc gttaaactct tcacataaac tcctccatct 18000
ggacgttctt taatttctaa tttcttattc gattcggctt ctaataaatc tcgaagttec 18060
tcctaaagat ttcatttttg taaatcacac atcctaacgc cttacctgat aaatttccaa 18120
atagctagct ctaactaaat actcttgatt atgtgatgct gccatgtgct caaaaatag 18180
gtcaatacac ttatagatga cacctcgttg ttctggatcc gatgattttc cttccattgt 18240
gtgagtcttt ccagttccag tttgaccata tgcaaaaaatc gtggcattat atccgttttag 18300
aacggaatca actagatctc gaaaggtttc ttcatataga tccgattgtg tggaaactata 18360
aaatatatat ttttaaaaaa gagaactcat aaaatcataa acataaaatt gtggagaaat 18420
aattttgaaa aataactaata tttctatagc aggtgaaaaa aaagtgatgt actcctagaa 18480
ataaataatc ttacttttca tcataaatg catcgaatgt aaaatccttc gatggctcat 18540
cttgctcttt tggatttttc agctcaattt gcccacgttg tggtcgcata tgtactatc 18600
tgaaaatgat cgaaatttca aaatataaat atttcaaact ttacttactt tgaataatta 18660
tttgcatttt cttgtgaaga taacggtcga catctcaca ttacctgtaa acataaataa 18720
atacatttat ttgaattttg aaatgtataa aactggatta tgaaattttt aagctggtgg 18780
ttttgtatg agaagtaacg aaaaaagta caatttactt agagtcttgt gatttttctt 18840
tcaaatgcaa aactcaactg aatcataaat agtgatgctt cgaaaagtgt ttagaggaaa 18900
attgtatttt tagtaaaaac taatatacgt tttggactta aaaaaaatt atgttaaaac 18960
ttgaaaaatt acgtttatta gtgcttatat taaaatacgg tttcaaatata atttaaaatt 19020
aaaaaactc accttttggc caaaatcaga cattttagaa actagcatgt actttattac 19080
gttgaatata acttatgttg gaaaatggaa aatttgaaga caggtgaatt ttagtttttt 19140
ttctttttcg tacttctaaa aaatacttca tttattttac attttgagaa ctaatttttg 19200
aacatgtttc gaacaaaaaa aaagattttg aaaaccccaa aaaaacttac tttgacagtc 19260

```

-continued

tctgttttg	aagattttt	cattatttcc	accattttt	gtcactaaat	atttgctat	19320
caatgtaggt	gtcaaggaaa	atthtgggtc	attcctgatt	tagtgagagt	ggtctggaac	19380
ttaagaagat	tagtttaatg	tggaaaaata	atcatattgt	atcgagaaac	ggaattttga	19440
agcaataacc	gctagagaaa	gtgactaaaa	accagaaatt	gtagtctgtg	aatttcaata	19500
tttttggtt	tatgtcacat	ctggacaatc	ggaaaaatat	gcatacattt	gaaattttta	19560
gaaatatttt	gaattaactt	taaaggaaaa	aatgcatta	aaaagattga	aaacatcatt	19620
gacgttgaaa	aatggagaaa	atthtctaatt	tctcatcaaa	atattaaaat	attaaagttc	19680
ttcaataata	tgaaaatgtg	aataaaatgt	ctaaataagc	aaaaaaaaaca	gatcctattc	19740
attataaaat	gttcacacaa	gtgttacatt	tcgtacaaag	aagtactaaa	acggatggac	19800
taaagtaata	ttgtcactcc	cgaaaagacg	aggaagaagt	aatcggaaga	agatgtcggg	19860
agatgagtga	tagtaaaaat	acgaagagac	gcagatagag	agtttgagag	aaggagactt	19920
ctggagggaat	aaaaggtggt	ttcaagatgg	gggacagaga	gggagagggt	taaaagagca	19980
caaatgtgc	ataatatcga	tcctgcgcag	ttgagagacg	cagacaatgt	gaagaatgga	20040
gcatatgttt	ctagtgaaca	ctcagaagta	gttgttcatt	tgtccgaaac	tttgaaaca	20100
tatacatttt	aaacttgacg	tttttgaatt	ataaagggat	ggagggtcct	caaaaagtaa	20160
tcatagacat	gtgtagattt	taaattaaaa	cacaactaga	cataggatga	atcagaagct	20220
taccataaca	ttgttgattt	atthaaaaat	gagaaaaagt	aaaattccgg	atagtcttct	20280
ttgaaaaaat	tcacagagaa	gttataatgt	ttgatgatat	tcactgattt	gtaatacatt	20340
attagtagca	tggcttctat	gtatatagac	tatttttata	tcacatacat	gaaaaagggt	20400
taaggcatgc	gccagggcct	gaaaaacgat	ctacctacca	ggggagctct	agctcttagt	20460
tattaattca	agagactttt	gaaacttgat	tttttgagat	tttattcaat	gatttggtta	20520
aaaaaaaaaga	ttatttgcaa	aaattacaaa	ttttaatgtc	tatactctga	catcggttta	20580
gaacaatttt	agacaggctg	caatgaaagc	aatgaaataa	aatttccttg	aaattataat	20640
agagaatcag	taaaatgttg	cagattattt	gaaaaatgat	gcaagaattc	gcagaaaaat	20700
cagtgaagca	gaaaagtgcg	acaggagacc	gaagtctaaa	aaagtgaatt	atgaataaaa	20760
acaaatcatg	tgactggata	taattgaagg	tcttgattcg	gaaaagataa	ttggagctct	20820
ttgcttaggc	caggctctag	atattttatt	gaagcttttc	agaaatgttc	aaaattatca	20880
ggaaacagttc	tctttgcaat	ttctctatgg	ctcaactacc	agggttttc	ctttttcttc	20940
aaaaagtaga	atthtaaat	ataatthtaa	aatthaaata	ccaagcaaaa	aatcatatac	21000
tcatcatatc	atgtgatcat	atcatataat	catatagggc	tcgttctttt	ttttttcaaa	21060
aaatataaaa	tttactagaa	ccaagcatat	gacaataaaa	tattttgaat	tcactttaat	21120
gggaaaaaaa	caagaaaatt	tcattaacat	tattgaaaac	atcgttgcca	ataggaatgt	21180
agaaaatcaa	atcaaaaatca	agtgagatta	ggaaagaatc	gaaattaggt	agaattggaa	21240
aatctcgatt	ttttaagttg	gattcttaca	cgattttttc	gggatatttt	tcatttttat	21300
ttttagtagt	ttcagcctag	acggctgaga	attcttttca	aaccttccaa	tttcaaagag	21360
attcttccat	aatthaaat	aatthtcatt	cgatattagc	atccattata	tacgtatgat	21420
tcccttttta	aaatcgatc	tccttttcaa	ctgactcatc	acttaagaat	tgttgagtca	21480
tcaactgata	gtgagcagac	accaacaacc	atctctttag	ttccgttcc	gtttatttta	21540
ttttggaatc	taacatattc	aagaaaatta	acttgaaatt	agaataaatg	tttcttgcta	21600
gatttttttg	tcataagtat	ttcttatttg	gattataatt	ttcatctcga	aatcgtagag	21660

-continued

```

agtttttcac tatttttttt tgagttctaa acacttcctt cctcatcgat gatgaagttt 21720
ttgacaaate aactagtttt ttactcatat ctacatcaa tctatgattt tattcaaaaa 21780
cagttaaatt tttttaacga aattaaatg gtcacggac cgagcaaaag ctttcagaat 21840
caactgcttc tttaaattct ttaaaattca atcaactttt cgtgtccaaa gtcacaaact 21900
acacctttca aaaaatattt ctacattatt tgcccacatc ttggcacagt tttcttgcca 21960
ttcttcaata ttttcttctc tgcgtttccc aactcttat tttctgactg ttgacttttc 22020
cattgtatag actcaatttt actttcgttt tttcaatttt tttttctgcg aagttcgggtg 22080
ttaaactccc attttgcaat attaaaaatt tcaatattgc cgtttttggc ttgaatctat 22140
taaaatgatg ctgttttttt ttcagaaaag accaaaacat gccagatgat attccaaaat 22200
tgccacgaca cagagaaaag aaaaatcagc cgaaaggttt gaaaaattta gaaaaatctg 22260
aagttacttt ttaattctt tagacacacc ttggaacaa caaaaactgc ctgctttacg 22320
gcctcattat aacataactt cagcaattcc agttactctg ataacaggag tagccacggt 22380
ggcaatggga attgctcttt atttcggaca taatggatgt gagtttttag agtttattat 22440
ccccaaaaa aaaaatcaaa ttactcttcc ctggtaataa gtaagaaaaa gctaaagaaa 22500
acaaatttct tgtcaaaatt ttacattgta aaccgatagc aacaaaaaac aagtgtcata 22560
aaaaactgta gaaaatcgat aattttgcta caatttcaca aagctaaaaa atatttttta 22620
ttttaccgtg ttagtaccgg aatgttctgc acttgagcct tactattagt tacacaaaat 22680
ggatcaattt tgagcaattt gttgtgaatc tgacaataat tagtcctatt gatatagctt 22740
taggccactc attcgtgttc gtaattttct ttttccttga acttgtaaag gtacagtttt 22800
tgaaaacagg gatgtagtcc aagtagtcaa atattgattc ttgtagcatt agaacaagag 22860
attgtgtaca cggattgtgc tctttcaaat ggaacacaag cttcacgaat tatgagaact 22920
gaaatgggaa atcaaacatt taaatgtgca tatacaatta ctttgaatga cgattatact 22980
gtaagttgag ttttaatttt taaatcatca aagaaaacat atgtatattt ttgcgaagga 23040
aattttggat ctggctcttag gatgaaacga cattgtaaca ttttgattaa agagccctta 23100
gttggaaagt agtgtatctg gtaaaaaac aattcgaaaa tatttaacca aatattgata 23160
aagcctaggt tgaacctgct ctgcagtccc taatttttca cattattttt cttcaaaata 23220
ttactatgat atttcaaagc cgggggtacc atcttaaaat catcatttgc aagtatcaca 23280
attaatgttc aacattacag ggccaagtga agttttatta cggctcttcc aagttctatc 23340
aaaaaatcgc attatacttc aactcacgaa acgatcaaca gctacgtgga aaagttactg 23400
aaactgacgg atgtgatcca ttagaatagc tggatgttaa tggaactaaa gttccattg 23460
cgcgtgtgg gaaagtggct gattcaatgt ttaacggtgc atttcaattg attgcttaat 23520
ttcagtattg caacattttt cattttttat aatacatcta acttcaaaat ttgttttttt 23580
ttcagatacc ttcgaattat tttatatcaa tgataaagcc tcaaacgagg taacacgggt 23640
tccatggaca actcgtggag tactcgggtc aactgaaatg aaaagaaaat tcagaaatcc 23700
gattcagcgc gaaaaccaga cattatgtga tgtgtttgcg gttgaaatga aataagaaaa 23760
aaaaataaatt aaactccatc ttttagggaa caatgcctcc gccatcatgg agatatccga 23820
tctgtcaatt gggactaaac agtattgatc cagatgttgg cattggtttc gagaacattg 23880
attttatggt ttggatgaag gttgcagctc ttccaaaatt cagaaaactg tatagaatac 23940
tgaatcgaca agttgatatg ttcagtaatg gattacctaa aggacaatat cagttgacca 24000

```

-continued

ttaattacag	tatgtttatg	ttaatgttga	atztatgtat	ttatgcaaaa	aatttactgc	24060
aaaagttcac	aataattcca	cccaaacctg	cttaaatatg	gagatgcaag	ttttttgttt	24120
cagataaaca	gtggctccaa	aaaaccaatc	tttgttataa	aacctcacia	aaatttctcg	24180
atatttcttt	attatgggtc	aaacttttga	gaaaaaaggg	aaatttagaa	aattctttca	24240
agcgaaattg	tcaaaathtt	tcaaaaccaa	atttgatttt	ccagatttat	tttttgcga	24300
cttgacaata	gtaaagaaaa	aacaagttga	atttttctat	atgaattctt	atagctgaac	24360
atttttgatc	aatttgaaaa	taatcaatag	acaatttttc	tccatactac	tgattttcag	24420
actatccagt	ggatatgtat	tggggcgaca	agtacttcgt	tatagccaat	gaaaactggg	24480
ttggaccag	gaatctgttt	ctaccagtaa	tctatttggg	tgttggaaca	ttcttacttc	24540
tctgtactat	tctcttcata	ttgatttggg	taaaacagag	actgtcgagg	gttcatccaa	24600
catgaattgg	aaaaactaat	tgaaaataga	cggatgaact	tcaaatgtgt	ttacaagagt	24660
tgaagtctca	aaataagctg	gtagcatgta	ttgtacggga	acagatttgt	atactttgct	24720
ttgtaataaa	aataaaatgt	tattatatta	gtctgtaatt	ttatgtatag	ttcaatttaa	24780
ttgaaataca	taatacccc	ttcagtttat	caattaaagc	tccaactatc	atctcgctgg	24840
tgagattaat	tgctcgagtga	gggcatctga	aatgtaaatt	taaaattaca	aaataaataa	24900
ttgtaagtgc	tatcagatat	aacaaatgat	catttaatta	aggaggaaaa	acaaaacatt	24960
aatttaaaaa	atztatcaaa	aaacaaaaaa	aaacgggtcaa	atattttttc	aaacaaacia	25020
agtaaagcta	atztatatta	aagttgatct	aaactactgg	tggttaggca	tactatagtt	25080
gattttcaacg	ggaagaagcg	caaatcagca	agtgtacatt	gtgttctgaa	aaattgaaat	25140
tcaacagttg	aatataagta	gaaactctac	ctattgctaa	catttattgc	aattcttctg	25200
tgttttgaac	aatatcgaga	tgcctccatc	catcgataaa	ttccgtaga	tttgatgaca	25260
tctcatctac	agcttccaat	tctccaatta	tctgatcctt	cagtttcagc	ttaatatcaa	25320
acgatttttg	aatttctcgg	atatttgctt	cataaactct	cgagatttca	gatttactct	25380
gttgaaatctc	tttaaatgtc	tctgaaatct	tcttctcaag	aactcttttc	tgataatata	25440
gctcagtgat	ttctcgagtt	ctttctttca	taatcatttc	agtattcatt	tgttcttctt	25500
taacattttc	ttcaagtttt	tcaatctttc	gtacataatc	acagaaatga	tcaaccacct	25560
gcaaccattg	cgggtcattt	cgcatagttt	tgagtcctcc	gggttggtca	agaattgcca	25620
caagactttc	tgctctgttc	agtttcagtt	tttctaagt	ttgctcaagt	gggaattctc	25680
tgacctcctg	gtttgcctct	tgggaaggat	cagaaathtt	ttcagaatgg	aacgtcaaaa	25740
tttcttggtc	caaatgggga	tatgtttctac	tagtcccttg	actagaagtc	tcaactagaga	25800
ttttaagtgt	cagttctcga	acattacgct	ccaatttcgg	agaatttcca	gattcagtg	25860
tcacatgact	ctttaatttg	acaatctcat	cattcttatac	aaagatttga	ttttccagtg	25920
cagagacctt	cacttgacia	tcttttgtct	tccatgacia	ttcatctgcc	aacatcttga	25980
tcatgttctc	attcgaatca	atcgtttttt	tcatatcttt	tatcatgtct	tcatcagttt	26040
tgattgtaac	attttgcttt	gaaatttcac	gtttcacaga	attcaagca	attttcaaag	26100
aattgttgaa	attttgcgag	atcaaacctt	caacaccttc	ggatttatca	ttcttgccaa	26160
cattccgatac	attatttagt	tcagttgata	tagaattggg	atcatcaact	gagaatactg	26220
ttaaattctg	tggttgaagt	tccaaaaatt	tctccatgac	tttttcaata	ctatccttct	26280
gataaacatt	gaaggattct	tgaagcattt	taatctcttc	attctttttc	ttgagttcgg	26340
aaaaacagaat	atcttttctt	ttttggaatt	ccactgtcac	taagagataa	tcatctttgt	26400

-continued

```

tttccaaaga tgtaaccagc tgcgatggat tcaaactttc caagtttttc gtaagatagt 26460
caacttcgct ttcaagtttt tgaatgatct gtttgttgtc ttccatcttt tttgttataa 26520
gtataggatc cttgaaataa agagaaaacg tgactatgat cttgtcaata gtttccagca 26580
agtggaaaat gtcataataa tctccatcat ttaatacttt taatttgtcc aaaagtgtat 26640
ccatcaactt ctcttgtttc ggtgtttcat caccaattaa tattccaccg taaccgttaa 26700
ccggatatgg cgacaaatca taaaattggt tttgaagatc ctcatattta gttttgagaa 26760
ttgagaaatc ttcatctctc attatcaaca acttcttcaa tcttctagct tctgtctctg 26820
cttttatttg gaaaatttcg aattgcattt cagctagtat tatctcatct tcttgttcat 26880
cgattaattt ctcaacttct tcttttaatt tttgatattt tgatctttt tctgccatag 26940
cattttgaaa atcttctagt gttgcaaact gatattcagt ggatgtactc gtggattcct 27000
gagtcgaaat ttcagtgctc ttgtttatct gttttacttc ggagcttctg actattttct 27060
ctggagtcca aaacttgtct acttccaaaa aatgtgtttt tttgcttctg aaaaacatat 27120
attaagtaac atctttaaga tattcaggtg cacttacatt tttgaaatat ttggtgacaa 27180
actttgaatt atcaatctga attctctctc ggttccagtg aagcaagcat aattctgaaa 27240
ataaaaatta cagcttttga aaccaatgaa acgaaacaac tattgtattt aaaaaatgct 27300
cacttcaact ccattctctt ccaccgctgc ttcttttttc acacttttcc agtttatcaa 27360
ttaaaaaatc aagtttctgt tgttcaggtg aaggctgaga tgctgtgaac gacatagttc 27420
tgaaaatagt taatttaaat gtagcagaaa aatcttttct agaaagtaaa aaaaatcagt 27480
aaaaacaagt actaagagaa attgaataaa ccaatcacia taatgacttc ttaacaagct 27540
gaaaaataat gcaatagcaa agaaaaacga gtagtctcgg taactccata gtacattatt 27600
tcgttattgg gatcatcata tcatttatgt atgaggatat tatgagtaa ttctaataac 27660
ccgagagtaa aggcaaaaaa tagcatggag tgaaaaaacg gatcaagcaa agaaatcgtg 27720
ttaactttta taacatctag ttgacactgt cagaccaaaa acttaataaa attttcaact 27780
gtacataaca gctagctgaa actgtaattt aattttatat tccctcggtc aattctagct 27840
aaatagcga ttctgagcta agccttcatt tcaaaattaa caaaaaaat gcaatgaaat 27900
ttcacttgtt acataacagc tagctgaaac tgtaatttaa ttttatattc tctcggtaa 27960
ttctagctaa attagcgatt ctgagctaag acttcatttc aaaaaataca aaaaaaatga 28020
attgaaattt tcaactgtac ataacagcta gctgaaactg caatttaatt ttatattcct 28080
tcagtcaatt tcagctaaat tagcaatttt gagctaagtg ttgtgtttc ttaaaacaat 28140
gcaaatttgg atggtttttc gtgttcagtg aacaacaaa caaacaaaa aattctggta 28200
aataaccaca agctgaaact gtgagataat tttttagtga ccattgagtg actgctcata 28260
gacagtggtc tggaattaag actagaatga ttatctctca tgataacata ttatacagag 28320
aagttgggaa gaatgtaggt cattgtaaag cgacagacag gtcgcattga tcaaagagaa 28380
tataagtcga actctttcgt ttggttaact gagggccaat gttatttctt attagggaaa 28440
attaacattt aaggagcaaa ggattgcaaa caaatgcga taagatatat gattatagta 28500
ttttatcttt tgtaagtgtt gccataattt cagtaacgaa aaaaaataca aggcaatttt 28560
agatgttagg aaaatcgaat ttgtctgact agccaacgaa tgttctcaat tgaagttatt 28620
gttctttttt aagatgtttt catacaaatt agtcagtttt cgaagcttca gccacactta 28680
tccgaattga gcaatttcaa aactattttt tgtaaaataa aatacatctc cgaaaaattta 28740

```

-continued

```

catcgagttc ccaacaatac tgtatggata gaaaatacct accaatactg cacatgaaac 28800
gctctgaaaa taatcgmeta ggaaatgaga accttttaaa tataaatga gcacaataag 28860
taatactaac tttattgaga aagaacataa ttgttatgag aatagttttt aaatgagggtg 28920
agaacagaa tatcctgag aataagtga gatacttgaa aatttgtgaa atagtaataa 28980
gtaaatggtt ttcacattag tataaacaat gacagagtca cgcaaaagta cgggaaacat 29040
atgaagttta taatacagtg cagtacagaa aaggtacaaa gtttacaaga atacaattgt 29100
tttttaaaaa taattttttg ttgaaggctt aaggtaatac gattaaagag ctactttctt 29160
ccaatacгаа gttgaattta aaatttaaaa ggaaaaaagg aaaaaatta aaaagcatat 29220
gaaaaatcgg ggcgcathtt tagtgcaaaa aattagatgg catttatttt atcccatcca 29280
tctgaatctt cactgtgtgt ggattttatg tegtcatctt gatcgatcat tgtatcatca 29340
gcttctcctt cttgattgat aagaagacct tgcagttttt ccgaaagtcc gaaatcttc 29400
aaatccttct ctctcaatgc atcatgcatc ttctgaattt cagcggatcg ttcgctattt 29460
tgaataagtt ccatcagaca ctcaattttg ctatcttttt ccatgatttc tcttttatga 29520
tttgcaatct gttcttcttt tgattcaatc tctctctttg aattggctga aataaaagaa 29580
aatgcttaca gatgtgtgta aaacccttag aaaactttca caagcttacc tgtcaatact 29640
tcaaattgcc ccaataagtt gtgcttcac tcttcagttc gaagttaag atcttcaact 29700
gatgtattaa gcgtggcttt ttctgctgta gtgtttgcaa gttgcatctc taacgccatg 29760
acggtcgagt tatgttgatc caaaatatga ctg 29793

```

We Claim:

1. A method of screening candidate molecules for ability to modulate activity of an enzyme which catalyzes release of ADP-ribose from an ADP-ribose polymer, comprising the steps of:

providing a purified enzyme encoded by a nucleotide sequence that is at least 80% identical to the nucleotide sequence as set forth in SEQ ID NO: 1, wherein the enzyme has a molecular weight greater than 100 kilodaltons as determined by SDS-PAGE;

assaying the enzyme in the presence of a candidate molecule to be screened; and

comparing the activity of the enzyme in the presence of the molecule to the activity of the enzyme in the absence of the molecule, wherein a difference therebetween indicates that said molecule modulates activity of said enzyme.

2. The method of claim 1, wherein said enzyme has the amino acid sequence set forth in SEQ ID NO: 2, 4, or 6.

3. The method of claim 1, wherein said enzyme is encoded by the nucleotide sequence of SEQ ID NO: 1, 3, or 5.

4. A method of screening candidate molecules for ability to modulate activity of an enzyme which catalyzes release of ADP-ribose from an ADP-ribose polymer, comprising:

providing a purified enzyme which comprises the amino acid sequences set forth in SEQ ID NOS: 11, 12, 13, and 14 and has a molecular weight greater than 100 kilodaltons as determined by SDS-PAGE, assaying the enzyme in the presence of a candidate molecule to be screened, and

comparing the activity of the enzyme in the presence of the molecule to the activity of the enzyme in the absence of the molecule, wherein a difference therebetween indicates that said molecule modulates activity of said enzyme.

5. The method of claim 4, wherein said enzyme is encoded by a nucleic acid molecule, the complementary sequence of which hybridizes, under stringent conditions, to the nucleotide sequence of SEQ ID NO: 1.

6. The method of claim 4, wherein said enzyme is encoded by a nucleic acid molecule having the nucleotide sequence set forth in SEQ ID NO: 1, 3, or 5.

7. The method of claim 4, wherein said enzyme has the amino acid sequence set forth in SEQ ID NO: 2, 4, or 6.

8. The method of claim 4, wherein said enzyme comprises amino acids 647-977 of SEQ ID NO: 4.

* * * * *