

University of Kentucky

UKnowledge

Physiology Faculty Publications

Physiology

10-31-2018

Hyperhomocysteinemia as a Risk Factor for Vascular Contributions to Cognitive Impairment and Dementia

Brittani R. Price

University of Kentucky, brittani.price15@uky.edu

Donna M. Wilcock

University of Kentucky, donna.wilcock@uky.edu

Erica M. Weekman

University of Kentucky, emweek2@uky.edu

Follow this and additional works at: https://uknowledge.uky.edu/physiology_facpub



Part of the [Cardiovascular Diseases Commons](#), [Cardiovascular System Commons](#), [Cognitive Neuroscience Commons](#), [Geriatrics Commons](#), [Hemic and Lymphatic Diseases Commons](#), [Medical Neurobiology Commons](#), [Neurology Commons](#), [Neurosciences Commons](#), and the [Physiology Commons](#)
Right click to open a feedback form in a new tab to let us know how this document benefits you.

Repository Citation

Price, Brittani R.; Wilcock, Donna M.; and Weekman, Erica M., "Hyperhomocysteinemia as a Risk Factor for Vascular Contributions to Cognitive Impairment and Dementia" (2018). *Physiology Faculty Publications*. 134.

https://uknowledge.uky.edu/physiology_facpub/134

This Review is brought to you for free and open access by the Physiology at UKnowledge. It has been accepted for inclusion in Physiology Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Hyperhomocysteinemia as a Risk Factor for Vascular Contributions to Cognitive Impairment and Dementia

Digital Object Identifier (DOI)

<https://doi.org/10.3389/fnagi.2018.00350>

Notes/Citation Information

Published in *Frontiers in Aging Neuroscience*, v. 10, 350, p. 1-12.

© 2018 Price, Wilcock and Weekman.

This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](#). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.



Hyperhomocysteinemia as a Risk Factor for Vascular Contributions to Cognitive Impairment and Dementia

*Brittani R. Price, Donna M. Wilcock and Erica M. Weekman**

Department of Physiology, Sanders-Brown Center on Aging, University of Kentucky, Lexington, KY, United States

Behind only Alzheimer's disease, vascular contributions to cognitive impairment and dementia (VCID) is the second most common cause of dementia, affecting roughly 10–40% of dementia patients. While there is no cure for VCID, several risk factors for VCID, such as diabetes, hypertension, and stroke, have been identified. Elevated plasma levels of homocysteine, termed hyperhomocysteinemia (HHcy), are a major, yet underrecognized, risk factor for VCID. B vitamin deficiency, which is the most common cause of HHcy, is common in the elderly. With B vitamin supplementation being a relatively safe and inexpensive therapeutic, the treatment of HHcy-induced VCID would seem straightforward; however, preclinical and clinical data shows it is not. Clinical trials using B vitamin supplementation have shown conflicting results about the benefits of lowering homocysteine and issues have arisen over proper study design within the trials. Studies using cell culture and animal models have proposed several mechanisms for homocysteine-induced cognitive decline, providing other targets for therapeutics. For this review, we will focus on HHcy as a risk factor for VCID, specifically, the different mechanisms proposed for homocysteine-induced cognitive decline and the clinical trials aimed at lowering plasma homocysteine.

OPEN ACCESS

Edited by:

Albert Gjedde,
University of Southern Denmark,
Denmark

Reviewed by:

Raluca Elena Sandu,
University of Medicine and Pharmacy
of Craiova, Romania
Roxana Octavia Carare,
University of Southampton,
United Kingdom

*Correspondence:

Erica M. Weekman
emweek2@uky.edu

Received: 06 June 2018

Accepted: 16 October 2018

Published: 31 October 2018

Citation:

Price BR, Wilcock DM and
Weekman EM (2018)
Hyperhomocysteinemia as a Risk
Factor for Vascular Contributions
to Cognitive Impairment
and Dementia.
Front. Aging Neurosci. 10:350.
doi: 10.3389/fnagi.2018.00350

Keywords: hyperhomocysteinemia, vascular cognitive impairment and dementia, B vitamins, homocysteine, dementia

INTRODUCTION

Vascular contributions to cognitive impairment and dementia (VCID) are defined as the conditions arising from vascular brain injuries that induce significant changes to memory, thinking, and behavior. It is the leading cause of dementia behind only Alzheimer's disease (AD); however, there is increasing awareness of the co-morbidity of VCID and AD (Bowler et al., 1998; Zekry et al., 2002; Langa et al., 2004; Jellinger and Attems, 2010). Roughly 60% of AD patients have VCID, and it is thought that vascular injuries act as an extra "hit" to the brain that lowers the threshold for cognitive impairment in persons with AD pathology (Schneider and Bennett, 2010; Vemuri and Knopman, 2016). Also, it is suggested that patients with both AD pathology and VCID have a shorter time to dementia and their rate of cognitive decline is faster (Schneider and Bennett, 2010; Vemuri and Knopman, 2016). Recent studies have also shown that vascular injury precedes AD pathologies, highlighting a role for the vasculature in AD progression (Canobbio et al., 2015; Janota et al., 2016).

While there is no cure for VCID, several studies have identified risk factors that can be modified to reduce risk of developing VCID. A major, yet underrecognized, modifiable risk factor for VCID is hyperhomocysteinemia (HHcy). Defined as elevated plasma levels of homocysteine, a non-protein forming amino acid, HHcy has been identified as a risk factor for cardiovascular disease,

stroke, VCID, and AD (Graham et al., 1997; Bostom et al., 1999; Eikelboom et al., 1999; Beydoun et al., 2014). Studies have shown that serum homocysteine levels are inversely related to cognitive function in patients with dementia and elevated levels are more common among VCID patients than among AD patients (Miller et al., 2002; Clarke et al., 2003). Elevated plasma homocysteine is also associated with hippocampal atrophy, white matter lesions, and lacunar infarcts (Vermeer et al., 2002; Firkbank et al., 2010). In the clinic, it is clear that HHcy plays a role in VCID; however, the mechanisms of homocysteine-induced cognitive impairment and the clinical implications of reducing homocysteine remain unclear. This review paper will focus on proposed mechanisms of homocysteine in the brain, and the clinical trials aimed at lowering homocysteine levels.

HOMOCYSTEINE METABOLISM

Homocysteine is produced in all cells and involved in the metabolism of cysteine and methionine (Selhub, 1999). Normal levels of homocysteine range between 5 and 15 $\mu\text{mol/L}$. Levels between 15 and 30 $\mu\text{mol/L}$ are considered mild, levels at 30–100 $\mu\text{mol/L}$ are moderate and levels above 100 $\mu\text{mol/L}$ are considered severe HHcy. During normal metabolism, ATP activates methionine to form *S*-adenosylmethionine (SAM). SAM is a methyl donor to several different receptors and forms *S*-adenosylhomocysteine (SAH) as a by-product of this methyl reaction. SAH can then be hydrolyzed to form homocysteine. Homocysteine can also go through two different re-methylation processes to form methionine again. In one pathway, folate is reduced to tetrahydrofolate which is then converted to 5, 10-methylenetetrahydrofolate. Methylenetetrahydrofolate reductase (MTHFR) reduces 5, 10-methylenetetrahydrofolate to 5-methyltetrahydrofolate. Finally, 5-methyltetrahydrofolate and the essential cofactor vitamin B12 add a methyl group to homocysteine to form methionine again. In an alternative pathway, betaine-homocysteine *S*-methyltransferase (BHMT) uses betaine synthesized from choline as a methyl group to convert homocysteine back to methionine.

Homocysteine can also go through a transsulfuration pathway to form cysteine. Serine can be enzymatically added to homocysteine by cystathionine beta synthase (CBS) and vitamin B6 to form cystathionine (Locasale, 2013). Cystathionine can then be cleaved by cystathionine gamma lyase (CGL) to form cysteine. While cysteine can be converted back to cystathionine, cystathionine cannot be converted to homocysteine again. The homocysteine metabolic pathway is shown in **Figure 1**.

MECHANISMS OF HOMOCYSTEINE-INDUCED COGNITIVE IMPAIRMENT


Posttranslational Modification of Proteins

As mentioned above, homocysteine is produced in all cells; however, its conversion to cysteine or back to methionine does

not. The brain lacks both CGL and BHMT, making it dependent on the folate cycle for re-methylation of homocysteine to methionine (Sunden et al., 1997). While this makes the brain especially vulnerable to raised levels of homocysteine, the mechanisms of homocysteine toxicity in the brain remain unclear, with several different mechanisms proposed. Some studies suggest the post-translational modification of proteins by homocysteine, termed homocysteinylation, contributes to its toxicity, especially since the degree of homocysteinylation is proportional to increased level of plasma homocysteine (Jakubowski, 1999; Jakubowski et al., 2000; Perla-Kajan et al., 2007). In the presence of adenosine triphosphate, methionyl-tRNA synthase catalyzes the conversion of homocysteine to homocysteine-thiolactone, which has been shown to homocysteinylate proteins and alter their functions. Specifically, homocysteine thiolactone acts as a Na/K ATPase inhibitor in the hippocampus and cortex of rat brain cells, thus changing the membrane potential of neurons (Rasic-Markovic et al., 2009).

Oxidative Stress

Other studies suggest homocysteine induces cellular damage via oxidative stress. As mentioned above, during normal homocysteine metabolism, cysteine is produced. Cysteine is a precursor for glutathione, which is a tripeptide that ultimately reduces reactive oxygen species. Without homocysteine conversion to cysteine, either due to CBS mutations or a diet lacking in vitamin B6, glutathione levels decrease, leading to increased reactive oxygen species and ultimately oxidative stress. Homocysteine metabolism is also regulated by the redox potential in a cell since several enzymes involved in its metabolism are regulated by the oxidative status (Zou and Banerjee, 2005). In one instance, the activity of methionine synthase is lowered when reactive oxygen species are high (Zou and Banerjee, 2005). Studies have also shown an increase in neurodegeneration due to homocysteine-related oxidative stress. In cultured embryonic cortical neurons and differentiated SH-SY-5Y human neuroblastoma cells grown in folate free media, there was an increase in cytosolic calcium, reactive oxygen species, and apoptosis (Ho et al., 2003). A significant increase in homocysteine was also found and inhibiting formation of homocysteine prevented the increase in reactive oxygen species. The increase in reactive oxygen species due to HHcy also alters smooth muscle function and promotes proliferation of smooth muscles cells (Welch and Loscalzo, 1998). Homocysteine has also been shown to inhibit endothelial nitric oxide synthase (eNOS) activity in cultured aortic endothelial cells from adult mice (Jiang et al., 2005) and humans (Jiang et al., 2005). In a genetic mouse model of HHcy where the CBS gene is absent, homozygote knockout mice show reduced eNOS activity compared to wildtype mice (Jiang et al., 2005). While the decreased activity of eNOS can affect oxidative stress, it also inhibits endothelial-dependent vasodilation. Taken together with the changes in vascular smooth muscle cells, these data provide further insight into how homocysteine is a risk factor for VCID.



AMPA and NMDA Receptors

Another proposed mechanism for homocysteine neurodegeneration involves homocysteine's role as an agonist for AMPA (both metabotropic and ionotropic) and NMDA receptors. Homocysteic acid, an oxidative product of homocysteine that is released in response to excitatory stimulation, acts as an excitatory neurotransmitter by activating the NMDA receptor (Cuenod et al., 1990). Activation of both AMPA and NMDA receptors leads to increased intracellular calcium, which in turn leads to activation of several kinases (Robert et al., 2005). Overstimulation of these receptors due to HHcy can then lead to increased free radicals and caspases, which leads to apoptosis (Mattson and Shea, 2003) and neurodegeneration. Using an NMDA antagonist can block the neurotoxic effects of homocysteic acid in the brain (Olney et al., 1987).

Cerebrovascular

The study of animal models has also lent insight into the mechanisms of homocysteine toxicity and its role in VCID. Several animal models have shown that high plasma levels of homocysteine are sufficient to cause cognitive deficits and vascular adverse events in the brain. Induction of HHcy in an animal model can be achieved via genetic manipulation or diet. Genetic manipulation of either CBS or MTHFR can produce mouse models of HHcy. In humans, deficiencies in CBS result in

elevated plasma levels of homocysteine and thrombosis and are the most common cause of hereditary HHcy. CBS[±] heterozygote mice have a 50% lower CBS activity compared to wildtype mice and develop mild HHcy (Watanabe et al., 1995). These mice show endothelial damage, thickened cerebral arteriolar walls, mild hypertension, and blood-brain barrier dysfunction (Baumbach et al., 2002; Weiss et al., 2003; Kamath et al., 2006). In humans, there are several polymorphisms in MTHFR that produce HHcy and neurological conditions such as a progressive demyelinating neuropathy and cognitive impairment (Clayton et al., 1986; Hyland et al., 1988; Surtees et al., 1991). Chen et al. (2001) deleted the MTHFR gene to create a mouse model of HHcy that exhibits motor and gait abnormalities within 5 weeks after birth. MTHFR^{-/-} homozygotes also present with some loss of function in cerebral vessels and abnormal lipid deposition in the aorta and disruption of the laminar structure of the cerebellum with no obvious changes in the cortex or cerebrum (Neves et al., 2004).

Unlike MTHFR and CBS knockout mice, dietary induction of HHcy allows for age related HHcy to be studied. Dietary induction of HHcy in mice and rats can be achieved through a reduction in the essential cofactors needed for homocysteine conversion (folate, vitamins B6, and B12) or enrichment in methionine, which increases the conversion of methionine to homocysteine. A combination of these diets or even a diet of increased homocysteine can also be used to induce HHcy. Troen

et al. (2008) showed that feeding mice a B vitamin deficient diet resulted in cognitive impairment on the Morris water maze and rarefaction of brain capillaries. In another animal model, 6-month-old Sprague–Dawley rats were placed on a diet deficient in folate for 8 weeks. By the end of the 8 weeks, the rats on the folate deficient diet had increased homocysteine levels, ultrastructural changes to cerebral capillaries, endothelial damage, swelling of pericytes, basement membrane thickening, and fibrosis (Kim et al., 2002). Cognitive impairments, decreased acetylcholine in the brain and microhemorrhages were seen in rats that were fed a diet high in homocysteine for 5 or 15 months (Pirchl et al., 2010).

Our lab has also recently developed a model of VCID by inducing HHcy in order to investigate the mechanisms of homocysteine-induced cognitive impairment. We placed 3-month-old C57BL6 mice on a combination diet that is deficient in folate and vitamins B6 and B12 and enriched in methionine (Sudduth et al., 2013) for 3 months. At the end of the 3 months, plasma homocysteine levels reached moderate levels in the mice on the homocysteine diet ($82.93 \pm 3.561 \mu\text{mol/L}$ compared to $5.89 \pm 0.385 \mu\text{mol/L}$ in the control mice). When tested on the radial arm water maze for behavioral deficits, these mice exhibited significant cognitive impairments in spatial memory. Prussian blue staining and magnetic resonance imaging showed microhemorrhages were the main cerebrovascular pathology induced by the HHcy diet. The mice on the HHcy diet also had an increase in several pro-inflammatory cytokines along with an increase in matrix metalloproteinase 9 (MMP9) activity. MMP9 has been shown to degrade tight junctions, leading to microhemorrhages and dystroglycans, and the pro-inflammatory cytokines, tumor necrosis factor alpha (TNF α), and interleukin 1 beta (IL-1 β), stimulate its transcription (Galis et al., 1994; Vecil et al., 2000; Michaluk et al., 2007; Candelario-Jalil et al., 2011; Klein and Bischoff, 2011). Previous studies have also shown homocysteine can induce MMP9 release from mouse cerebral microvessel endothelial cells (Shastry and Tyagi, 2004). Based on this data, another possible mechanism for homocysteine-induced cognitive impairment could be the pro-inflammatory mediated increase in MMP9 leading to tight junction degradation, microhemorrhages, and, finally, cognitive impairment.

Astrocytes

In addition to the pathologies listed above, we have also shown that astrocytic end-feet are disrupted in the mice on the HHcy diet (Sudduth et al., 2017). In the brain, astrocytes make up 50% of the cells and their processes, termed astrocytic end-feet, sheath arterioles, and capillaries. The main function of astrocytic end-feet is to maintain ionic and osmotic homeostasis in the brain (Simard and Nedergaard, 2004). To do this, astrocytes have aquaporin four water channels and several potassium channels located at their end-feet. In our mice on the HHcy diet, we found a significant decrease in these channels, as well as other structural markers located at the end-foot. These decreases in the end-foot channels occur after 10 weeks on the HHcy diet. Cognitive deficits and microhemorrhages are also seen starting at 10 weeks on diet. Interestingly, increases

in the pro-inflammatory cytokines, TNF α , and IL-1 β , occur after only 6 weeks on diet. We had also previously shown that MMP9 was significantly increased in mice on the HHcy diet (Sudduth et al., 2013). Taken together, we hypothesize that another mechanism of homocysteine-induced cognitive impairment involves the inflammatory-MMP9 pathway. In our hypothesis, homocysteine increases TNF α and IL-1 β expression, which in turn activates MMP9, which degrades dystroglycans, a key structural component that anchors the astrocytic end-foot to the basal lamina of the vessels. This disruption of the astrocytic end-foot leads to impaired ionic and osmotic buffering and eventual cognitive impairment.

While several mechanisms of homocysteine-induced cognitive impairment and neurodegeneration have been proposed and discussed here, it is unlikely that homocysteine acts through only one of these mechanisms. Homocysteine may act through several, if not all of these mechanisms. It is also unclear whether the high levels of homocysteine or the lack of B vitamins is the main cause behind the cognitive impairment seen in hyperhomocysteinemic patients. Discussed next are the clinical implications of HHcy and the potential therapeutics tested in clinical trials to lower homocysteine levels and improve cognition.

HYPERHOMOCYSTEINEMIA IN THE CLINICAL SETTING

Extensive clinical data support the role of HHcy as a risk factor for VCID. Given that normal and abnormal values are set by individual clinical laboratories, mild-moderate HHcy is loosely defined by clinical standards (Moll and Varga, 2015). However, plasma homocysteine concentrations ranging from 15 and 100 $\mu\text{mol/L}$ are uniformly considered to be indicative of clinically relevant HHcy. Gibson et al. (1964) reported vascular anomalies in patients with homocystinuria (elevated concentration of homocysteine in both plasma and urine), and McCully (1969) introduced his homocysteine hypothesis which connected HHcy with an increased risk of atherosclerosis (Abraham and Cho, 2010). To date, HHcy continues to serve as a widely recognized risk factor for coronary artery disease (CAD), peripheral vascular disease, myocardial infarction (MI), and cerebrovascular disease (CVD; Maron and Loscalzo, 2009). Of particular importance here is the association between HHcy and CVD. CVD can manifest as a stroke, white matter disease, cerebral large vessel disease (atherosclerosis), and cerebral small vessel disease (arteriosclerosis), all of which can independently induce cognitive impairment ranging from subtle deficits to frank dementia (Troen et al., 2008; Maron and Loscalzo, 2009; Hainsworth et al., 2016). Furthermore, HHcy has been associated with hippocampal and white matter atrophy in older subjects with mild hypertension, as well as an increased rate of hippocampal atrophy and cognitive decline in elderly patients (Clarke et al., 1998; Firkbank et al., 2010). As suggested by the variety of cellular actions of homocysteine described above, there is no shortage of candidate mechanisms by which HHcy induces cognitive impairment despite known etiologies.

Hyperhomocysteinemia vs. Homocystinuria

Both genetic mutations and dietary vitamin deficiencies can affect homocysteine levels resulting in HHcy. Several polymorphisms (notably C677T and A1298C) have been identified in the MTHFR gene in humans, which can induce severe HHcy (>100 $\mu\text{mol/L}$, termed homocystinuria) by limiting conversion of homocysteine back to methionine (Moll and Varga, 2015; Hainsworth et al., 2016). While rare, these polymorphisms induce progressive demyelinating neuropathy and cognitive impairment (Clayton et al., 1986; Hyland et al., 1988; Surtees et al., 1991). That being said, deficiencies in CBS, the rate-limiting enzyme of the aforementioned transsulfuration pathway, are the most common cause of homocystinuria and may result in thrombosis and low levels of cysteine (Sacharow et al., 1993). In contrast to HHcy, homocystinuria is a rare autosomal recessive metabolic disorder characterized by severely elevated plasma homocysteine and subsequently elevated urine homocysteine concentrations. Patients suffering from homocystinuria present with developmental delay, osteoporosis, ocular abnormalities, thromboembolic disease, and severe premature atherosclerosis (Poloni et al., 2018). Given that less marked elevations in plasma homocysteine (i.e., HHcy) are much more common, homocystinuria will not be further discussed in this review. Less marked elevations in plasma homocysteine, referred to as HHcy, may be attributed to factors such as smoking, aging, renal failure, and low dietary levels of folate and vitamins B6 and B12 (Hainsworth et al., 2016).

Prevalence of B Vitamin Deficiency

As suggested, clinical mild–moderate HHcy is common, especially in elderly patients, with the majority of cases resulting from insufficient B vitamin status (Joosten et al., 1993; Troen et al., 2008). The association of B vitamin status and normal central nervous system function dates back to 1849 when Addison reported on the “wandering mind” of patients with pernicious anemia (Smith and Refsum, 2016). Reports of insufficient B vitamin status with concomitant induction of HHcy trace back to a landmark report by the Framingham Heart Study in 1993. A cohort of 1041 elderly participants (418 men, 623 women) between the ages of 67 and 96 showed that plasma homocysteine becomes elevated due to dietary deficiencies in B6 and folic acid and decreased absorption of B12 (Selhub, 2006; McCully, 2007).

According to the Framingham report, daily intake of 3 mg vitamin B6 and 400 μg of folic acid are required to prevent elevations in plasma homocysteine concentration (Selhub, 2006; McCully, 2007). In support of these amounts of dietary B vitamins, the Nurses’ Health Study revealed that similar levels of dietary B6 and folic acid prevent mortality and morbidity from heart disease (Rimm et al., 1998; McCully, 2007). In the United States, mandatory fortification of grains with folic acid was authorized in 1996 and fully implemented in 1998 (Crider et al., 2011). Prior to fortification of grain products, intakes of B6 and folic acid were well below the recommended quantities (McCully, 2007). By contrast, with the exception of those

partaking in a vegan diet, vitamin B12 intake is typically adequate. However, in those >65 years of age lack of gastric acidity, decreased intrinsic factor synthesis by gastric mucosal cells, and history or presence of *H. pylori* infection may contribute to inadequate B12 absorption (McCully, 2007). Not to mention, the aging process itself is associated with decreased ability to absorb B vitamins, which can lead to a gradually rising plasma homocysteine concentration (estimated at 1 $\mu\text{mol/L/decade}$) (McCully, 2007). Literature now suggests between 5 and 30% of the general population, and 25% of those with vascular diseases, to be affected by HHcy (Selhub, 2006; Peng et al., 2015; Yeh et al., 2016). Granted, because blood homocysteine panels are generally ordered only when patients experience a MI or stroke without traditional risk factors, the aforementioned prevalence of HHcy in the general population is likely skewed and possibly underestimated.

HHcy, B Vitamin Status, and Cognition

Regardless, public significance of HHcy in the elderly population should not be ignored given that it is easily treatable with B vitamin fortification and serves as a modifiable risk factor for development of cognitive decline, dementia, and AD. A number of early cross-sectional studies relating HHcy or insufficient B vitamin status to cognitive impairment led to generation of the hypotheses suggesting a causal link. In effort to address whether the hypothesis that HHcy induces cognitive impairment is correct, a number of clinical trials have assessed B vitamin refortification with cognitive endpoints. These vitamin refortification trials are outlined in **Table 1**. Additionally, several meta-analyses of these intervention trials have been conducted (Wald et al., 2010; Ford and Almeida, 2012; Clarke et al., 2014). Upon review, the general consensus suggests that homocysteine-lowering by B vitamin refortification has no significant effect on individual or global cognitive domains despite three trials (FACIT, WAFACS, VITACOG) supporting a beneficial effect. However, when interpreting the results of these trials one needs to consider the fact that many were compromised by the challenges of performing a cognitive clinical trial (cohort age, B vitamin status of said cohort, trial duration, statistical power, etc.).

Limitations of Clinical Trials

As suggested, a number of factors related to trial design and implementation must be considered. First and foremost, the hypothesis being tested should be considered. Assuming the hypothesis is that homocysteine-lowering supplementation with B vitamins slows and/or prevents cognitive decline, those randomized to the placebo arm of the trial must exhibit cognitive decline. As reviewed in **Table 1**, as well as the aforementioned meta-analyses, the majority of trials conducted fail to report significant cognitive decline in those randomized to the placebo arm. Meaning these trials are limited to showing only that B vitamin treatment does not worsen cognition. Additionally, the age range of trial participants must be considered. Referring to the hypothesis above, if cognition is a study measure the age of the participants should reflect the timeframe in which cognitive decline and dementia occur. Duration of the intervention must also be considered given that elderly individuals exhibiting

TABLE 1 | B vitamin clinical trials.

| Reference | Sample size | Mean participant age (years) | Selection criteria | Intervention | Duration of intervention (years) | Cognitive domains assessed | Additional Study Endpoints | Study Conclusion | Study Limitations |
|--|---|------------------------------|--|--|----------------------------------|---|---|---|---|
| Lewerin et al., 2005 | n = 209; 126 intervention, 69 placebo | 76 | Community-dwelling men and women | Placebo vs. 500 µg B12, 800 µg folic acid, 3 mg B6 | 0.3 | Short-term memory, long-term memory, perceptual speed, verbal ability, spatial ability, inductive reasoning | Plasma [Hcy], postural-locomotor-manual (PLM) test | Despite normalization of plasma [Hcy], B vitamin supplementation did not improve cognitive performance. | No reported, significant decline in cognition observed in participants randomized to the placebo arm. |
| Eussen et al., 2006 | n = 195; 65 intervention, 65 placebo | 82–83 | Patients with suspected, mild B12 deficiency | Placebo vs. 1000 µg B12 vs. 1000 µg B12, 400 µg folic acid | 0.5 | Attention, construction, sensorimotor speed, short-term memory, executive function | Plasma [Hcy] | Supplementation with B12 alone or in combination with folic acid does not improve cognitive performance. | No reported, significant decline in cognition observed in participants randomized to the placebo arm. |
| Stott et al., 2005 | n = 185; 23–24/group | 72–76 | Patients with ischemic vascular disease | Placebo vs. 400 µg B12, 2.5 mg folic acid, 25 mg B6, 25 mg B2 (alone and in various combinations) | 1 | TICS-M, information processing | Plasma [Hcy] | Despite normalization of plasma [Hcy], B vitamin supplementation did not improve cognitive performance. | No reported, significant decline in cognition observed in participants randomized to the placebo arm. |
| VISP Trial Toole et al., 2004 | n = 3680; 1853 low-dose intervention, 1827 high-dose intervention | 66 | Patients with history of non-disabling cerebral infarction | Low-dose: 6 µg B12, 20 µg folic acid, 0.2 mg B6 vs. high-dose: 400 µg B12, 2.5 mg folic acid, 25 mg B6 | 1.8 | MMSE | Plasma [Hcy], recurrent cerebral infarction, CHD, death | Despite normalization of plasma [Hcy] in both the low-dose and high-dose intervention groups, supplementation did not improve cognitive performance. | No reported, significant decline in cognition observed in participants randomized to the low-dose intervention. |
| VITACOG trial Smith et al., 2010; de Jäger et al., 2012 | n = 168; 85 intervention, 83 placebo | 76–77 | Patients > 70 years with evident MCI (MCI = TICS–M 17–29/30) | Placebo vs. oral TriBeB Plus® containing 0.8 mg folic acid, 0.5 mg B12, 20 mg B6 | 2 | Episodic memory, semantic memory, global cognition (MMSE) | Plasma [Hcy], Rate of brain atrophy | Sevenfold less regional atrophy in intervention group vs. placebo group. However, this effect was significant only in those with elevated Hcy above the median (i.e., > 11.3 µmol/L). Individuals with baseline plasma Hcy above the median, cognitive decline was prevented across all domains assessed. | VITACOG was not powered to detect any effect of plasma [Hcy] on cognition. |

(Continued)

TABLE 1 | Continued

| Reference | Sample size | Mean participant age (years) | Selection criteria | Intervention | Duration of intervention (years) | Cognitive domains assessed | Additional Study Endpoints | Study Conclusion | Study Limitations |
|--|---|------------------------------|---|--|----------------------------------|---|--|---|---|
| McMahon et al., 2006 | n = 253; 127 intervention, 126 placebo | 73–74 | Patients with plasma [Hcy] > 13 μmol/L but normal creatinine concentrations | Placebo vs. 500 μg B12, 1000 μg folic acid, 10 mg B6 | 2.2 | Short-term memory, Learning capacity, Verbal fluency, Semantic fluency, Information processing, Reasoning ability | Plasma [Hcy], Plasma [creatinine] | Despite normalization of plasma [Hcy], B vitamin supplementation did not improve cognitive performance. In fact, information processing speed was improved in those randomized to the placebo arm. | No reported, significant decline in cognition observed in participants randomized to the placebo arm. |
| FACIT Trial Durga et al., 2007 | n = 818; 405 intervention, 413 placebo | 60 | Patients with plasma [Hcy] > 13 and <26 μmol/L | Placebo vs. 800 μg folic acid | 3 | Memory, Sensorimotor speed, Complex speed, Information processing, Verbal fluency | N/A | Folic acid supplementation improved subdomains of cognitive function that typically decline with age. | N/A |
| Vitaltops Trial Study Group, 2010 | n = 8164; 4089 intervention, 4075 placebo | 62–63 | Patients with recent stroke or TIA | Placebo vs. 0.5 mg B12, 2 mg folic acid, 25 mg B6 | 3.2 | MMSE | Plasma [Hcy], stroke, MI, vascular death | Despite normalization of plasma [Hcy], B vitamin supplementation did not improve global cognitive performance. | Baseline cognitive scores not provided. |
| NORVIT Trial Bonna et al., 2006 | n = 3749; 937 folic acid, B12, B6, 935 folic acid and B12, 934 plus B6, 943 placebo | 62–63 | Patients ages 30–85 years exhibiting acute MI 7 days before randomization. | Placebo vs. 40 mg B6 vs. 0.8 mg folic acid, 0.4 mg B12 vs. 0.8 mg folic acid, 0.4 mg B12, 40 mg B16 (latter referred to as "combination therapy) | 3.3 | None | Death, MI, unstable angina pectoris requiring hospitalization, coronary revascularization with percutaneous coronary intervention or coronary artery bypass grafting, and stroke | Despite normalization of plasma [Hcy], B vitamin supplementation did not reduce the risk of recurrent CVD following acute MI. A harmful effect from combined B vitamin supplementation was suggested. | Cognition was not assessed. |
| SU.FOL.O.M3 Trial Andreeva et al., 2011 | n = 1748; 884 intervention, 864 control (no supplementation) | 61 | Patients with vascular disease | Placebo vs. 20 μg B12, 560 μg CH3-THF, 3 mg B6 ± 600 mg 2:1 ratio of eicosapentaenoic and docosahexaenoic acid | 4.5 | TICS-M | Plasma [Hcy] | Despite normalization of plasma [Hcy], B vitamin supplementation did not improve global cognitive performance. However, a possible effect of B vitamin supplementation on cognitive function in those with history of stroke was noted. | Baseline cognitive scores not provided. |

(Continued)

TABLE 1 | Continued

| Reference | Sample size | Mean participant age (years) | Selection criteria | Intervention | Duration of intervention (years) | Cognitive domains assessed | Additional Study Endpoints | Study Conclusion | Study Limitations |
|--|--|------------------------------|---|--|----------------------------------|-----------------------------|--|--|---|
| HOPE-2 Trial Lonn et al., 2006 | n = 5522; 2758 intervention, 2764 placebo | 69 | Patients with history of vascular disease or with history of diabetes and atherosclerosis | Placebo vs. 1 mg B12, 2.5 mg folic acid, 50 mg B6 | 4.8 | MMSE | Plasma [Hcy], death resulting from CVD, MI or stroke | Despite normalization of plasma [Hcy], B vitamin supplementation did not improve global cognitive performance. | No reported, significant decline in cognition observed in participants randomized to the placebo arm. |
| WAFACS Trial *Sub-study of HOPE-2 Trial* Kang et al., 2008 | n = 2009; 1002 intervention, 1007 placebo | 71 | Patients with CVD or 3 or more coronary risk factors | Placebo vs. 1 mg B12, 2.5 mg folic acid, 50 mg B6 | 6 | TICS | N/A | B vitamin supplementation failed to delay cognitive decline among patients with CVD or CVD risk factors. | No reported, significant decline in cognition observed in participants randomized to the placebo arm. |
| Study of the Effectiveness of Additional Interventions, Reductions in Cholesterol and Homocysteine (SEARCH) Collaborative Group et al., 2010 | n = 12064; 6033 interventions, 6031 placebo | 64 | Patients with history of MI | Placebo + 20 mg or 80 mg simvastatin vs. 1 mg B12, 2 mg folic acid + 20mg or 80 mg simvastatin | 7.1 | TICS-M, verbal fluency test | Plasma [Hcy], major coronary events, stroke, noncoronary revascularizations, death attributable to vascular or non-vascular causes | Despite normalization of plasma [Hcy], B vitamin supplementation did not improve global cognitive performance. | Baseline cognitive scores not provided. |

CH3-THF, methyltetrahydrofolate; CHF, congestive heart failure; CVD, cardiovascular disease; Hcy, homocysteine; MI, myocardial infarction; MMSE, mini-mental state examination; TIA, transient ischemic attack; TICS, telephone interview for cognitive status; TICS-M, modified telephone interview for cognitive status.

normal cognition generally decline only by ~ 0.1 points on MMSE each year (Hainsworth et al., 2016; Smith and Refsum, 2016). Thus, duration of the intervention must be sufficient to observe cognitive decline, especially if MMSE is to be used as an assessment tool. Three trials represented in **Table 1** and seven of the nine trials examined by Wald et al. (2010) in their meta-analysis were of short duration (< 12 months) and therefore too short to identify an effect on cognition. Assessment tools must also be sensitive enough to detect subtle changes over the course of the trial. Collectively, age range of the trial cohort, duration of the intervention, and assessment tools will dictate whether the trial design is sufficient to detect an effect. Finally, the appropriateness of the intervention and whether the chosen cohort is likely to respond to that intervention must be considered. Supplied daily doses of B vitamins should be sufficient to lower plasma homocysteine concentrations by at least 20% (Hainsworth et al., 2016). For example, 1 trial addressed in **Table 1** prescribed doses of folic acid (0.2 mg) and vitamin B12 (1 μg) that were too low to influence plasma homocysteine (McMahon et al., 2006; Hainsworth et al., 2016). Furthermore, the baseline B vitamin status of each potential participant must be considered at the time of enrollment. This relates back to a cardinal principle of nutrition in which the relationship between vitamin status and a given outcome follows a sigmoidal curve. For example, if a participant exhibited low levels of vitamin B6, additional B6 intake would likely be beneficial, with the opposite being true if the participant's B6 intake were already high. Additionally, when at the plateau phase (i.e., adequate B6 intake), additional B6 intake will likely have no effect. Despite having critical implications for clinical trials, this principle is often overlooked. Consideration of the participant's B vitamin status at the time of enrollment would therefore aid in determining whether they are likely to respond to intervention. As such, trial enrollment should only be open to those with insufficient B vitamin status or elevated plasma homocysteine concentration at baseline. Together, these considerations suggest the conclusion that homocysteine-lowering by B vitamin supplementation has no effect on cognition is premature.

Beneficial Effects of B Vitamin Supplementation

As previously mentioned, results from three trials (FACIT, WAFACS, and VITACOG) do support a beneficial effect of B vitamin supplementation on cognition. The FACIT trial showed significant effects of B vitamins on cognition in participants with high plasma homocysteine, while the WAFACS trial showed similarly significant effects in those with inadequate B vitamin status (Durga et al., 2007; Kang et al., 2008). Furthermore, the VITACOG trial revealed strong effects of B vitamins on both rates of brain atrophy and cognition in individuals with mild cognitive impairment (MCI; Douaud et al., 2013). Further data analysis revealed the sevenfold reduction in regional brain atrophy to be significant only in those with plasma homocysteine concentrations above the median ($> 11.3 \mu\text{mol/L}$) (Douaud et al., 2013). Results of the VITACOG trial thereby imply a threshold effect of plasma homocysteine on measures of brain atrophy

and cognition. A threshold effect of plasma homocysteine is further supported by results of the OPTIMA study in which only plasma homocysteine concentrations $> 11 \mu\text{mol/L}$ were associated with an increased rate of atrophy of the medial temporal lobe (Clarke et al., 1998). The threshold concept is further supported by a study showing a plasma homocysteine concentration-dependent increase in the rate of cognitive decline in AD patients (Oulhaj et al., 2010). Jointly, these studies suggest the threshold for effect of plasma homocysteine lies between 10 and 11 μM , which may explain why studies conducted in countries that employ mandatory folic acid fortification do not find associations between plasma homocysteine and cognition. Retrospective analysis of the VITACOG data revealed that the protective effect of B vitamin supplementation on both brain atrophy and cognition only occurred in those participants with adequate omega-3 fatty acid status (Jerneren et al., 2015). Additionally, the beneficial effect of B vitamin supplementation on brain atrophy was observed only in participants not routinely taking aspirin (Smith et al., 2010). Omega-3 fatty acid and aspirin statuses may therefore contribute to the failure of B vitamin trials.

In all, given the challenges faced by previous trials, further B vitamin supplementation trials are needed. New trials will be most successful if they prescribe a full combination supplement (B6, B12, and folic acid) at high dose (i.e., dosage sufficient to reduce plasma homocysteine by 20%) to at-risk age participants with elevated plasma homocysteine or inadequate B vitamin status at baseline, adequate omega-3 fatty acid status at baseline, and who do not routinely take aspirin.

CONCLUSION

With the number of people aged over 60 expected to increase worldwide by 1.25 billion by 2050, accounting for 22% of the world's population, it is crucial to understand the causes of dementia and develop treatments (Prince et al., 2015). Current clinical and preclinical data provide strong evidence that HHcy is a key risk factor for VCID. With B vitamin supplementation being an inexpensive and safe therapeutic possibility, it would seem that treatment of HHcy-induced VCID would allow for some progress in lowering the number of dementia patients. Unfortunately, the mechanisms through which HHcy induces cognitive impairment remain unclear, with several different mechanisms proposed. In addition, clinical trials aimed at lowering homocysteine levels via B vitamin supplementation have also been lacking in their study design and ability to properly test the hypothesis that lowering homocysteine can slow and/or prevent cognitive decline. Future studies involving preclinical animal models and properly designed clinical trials will be necessary in order to effectively treat HHcy-induced VCID and lower the incidence of dementia.

AUTHOR CONTRIBUTIONS

BP and EW each wrote 50% of the manuscript. DW edited for content, checked for accuracy, and provided guidance in the preparation of the content.

FUNDING

This work was supported by National Institutes of Health grants RO1NS079637 and RO1NS097722 to DW, and

REFERENCES

- Abraham, J. M., and Cho, L. (2010). The homocysteine hypothesis: still relevant to the prevention and treatment of cardiovascular disease? *Cleve. Clin. J. Med.* 77, 911–918. doi: 10.3949/ccjm.77a.10036
- Andreeva, V. A., Kesse-Guyot, E., Barberger-Gateau, P., Fezeu, L., Hercberg, S., and Galan, P. (2011). Cognitive function after supplementation with B vitamins and long-chain omega-3 fatty acids: ancillary findings from the SU.FOL.OM3 randomized trial. *Am. J. Clin. Nutr.* 94, 278–286. doi: 10.3945/ajcn.110.006320
- Baumbach, G. L., Sigmund, C. D., Bottiglieri, T., and Lentz, S. R. (2002). Structure of cerebral arterioles in cystathionine beta-synthase-deficient mice. *Circ. Res.* 91, 931–937. doi: 10.1161/01.RES.0000041408.64867.1D
- Beydoun, M. A., Beydoun, H. A., Gamaldo, A. A., Teel, A., Zonderman, A. B., and Wang, Y. (2014). Epidemiologic studies of modifiable factors associated with cognition and dementia: systematic review and meta-analysis. *BMC Public Health* 14:643. doi: 10.1186/1471-2458-14-643
- Bonaa, K. H., Njolstad, I., Ueland, P. M., Schirmer, H., Tverdal, A., Steigen, T., et al. (2006). Homocysteine lowering and cardiovascular events after acute myocardial infarction. *N. Engl. J. Med.* 354, 1578–1588. doi: 10.1056/NEJMoa055227
- Bostom, A. G., Rosenberg, I. H., Silbershatz, H., Jacques, P. F., Selhub, J., D'Agostino, R. B., et al. (1999). Nonfasting plasma total homocysteine levels and stroke incidence in elderly persons: the Framingham Study. *Ann. Intern. Med.* 131, 352–355. doi: 10.7326/0003-4819-131-5-199909070-00006
- Bowler, J. V., Munoz, D. G., Merskey, H., and Hachinski, V. (1998). Fallacies in the pathological confirmation of the diagnosis of Alzheimer's disease. *J. Neurol. Neurosurg. Psychiatry* 64, 18–24. doi: 10.1136/jnnp.64.1.18
- Candelario-Jalil, E., Thompson, J., Taheri, S., Grossetete, M., Adair, J. C., Edmonds, E., et al. (2011). Matrix metalloproteinases are associated with increased blood-brain barrier opening in vascular cognitive impairment. *Stroke* 42, 1345–1350. doi: 10.1161/STROKEAHA.110.600825
- Canobbio, I., Abubaker, A. A., Visconte, C., Torti, M., and Pula, G. (2015). Role of amyloid peptides in vascular dysfunction and platelet dysregulation in Alzheimer's disease. *Front. Cell. Neurosci.* 9:65. doi: 10.3389/fncel.2015.00065
- Chen, Z., Karaplis, A. C., Ackerman, S. L., Pogribny, I. P., Melnyk, S., Lussier-Cacan, S., et al. (2001). Mice deficient in methylenetetrahydrofolate reductase exhibit hyperhomocysteinemia and decreased methylation capacity, with neuropathology and aortic lipid deposition. *Hum. Mol. Genet.* 10, 433–443. doi: 10.1093/hmg/10.5.433
- Clarke, R., Bennett, D., Parish, S., Lewington, S., Skeaff, M., Eussen, S. J., et al. (2014). Effects of homocysteine lowering with B vitamins on cognitive aging: meta-analysis of 11 trials with cognitive data on 22,000 individuals. *Am. J. Clin. Nutr.* 100, 657–666. doi: 10.3945/ajcn.113.076349
- Clarke, R., Harrison, G., Richards, S., and Vital Trial Collaborative Group (2003). Effect of vitamins and aspirin on markers of platelet activation, oxidative stress and homocysteine in people at high risk of dementia. *J. Intern. Med.* 254, 67–75.
- Clarke, R., Smith, A. D., Jobst, K. A., Refsum, H., Sutton, L., and Ueland, P. M. (1998). Folate, vitamin B12, and serum total homocysteine levels in confirmed Alzheimer disease. *Arch. Neurol.* 55, 1449–1455. doi: 10.1001/archneur.55.11.1449
- Clayton, P. T., Smith, I., Harding, B., Hyland, K., Leonard, J. V., and Leeming, R. J. (1986). Subacute combined degeneration of the cord, dementia and parkinsonism due to an inborn error of folate metabolism. *J. Neurol. Neurosurg. Psychiatry* 49, 920–927. doi: 10.1136/jnnp.49.8.920
- Crider, K. S., Bailey, L. B., and Berry, R. J. (2011). Folic acid food fortification—its history, effect, concerns, and future directions. *Nutrients* 3, 370–384. doi: 10.3390/nu3030370
- Cuenod, M., Do, K. Q., and Streit, P. (1990). Homocysteic acid as an endogenous excitatory amino acid. *Trends Pharmacol. Sci.* 11, 477–478. doi: 10.1016/0165-6147(90)90137-W
- fellowship F31NS092202 to EW. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.
- de Jager, C. A., Oulhaj, A., Jacoby, R., Refsum, H., and Smith, A. D. (2012). Cognitive and clinical outcomes of homocysteine-lowering B-vitamin treatment in mild cognitive impairment: a randomized controlled trial. *Int. J. Geriatr. Psychiatry* 27, 592–600. doi: 10.1002/gps.2758
- Douaud, G., Refsum, H., de Jager, C. A., Jacoby, R., Nichols, T. E., Smith, S. M., et al. (2013). Preventing Alzheimer's disease-related gray matter atrophy by B-vitamin treatment. *Proc. Natl. Acad. Sci. U.S.A.* 110, 9523–9528. doi: 10.1073/pnas.1301816110
- Durga, J., van Boxtel, M. P., Schouten, E. G., Kok, F. J., Jolles, J., Katan, M. B., et al. (2007). Effect of 3-year folic acid supplementation on cognitive function in older adults in the FACIT trial: a randomised, double blind, controlled trial. *Lancet* 369, 208–216. doi: 10.1016/S0140-6736(07)60109-3
- Eikelboom, J. W., Lonn, E., Genest, J. Jr., Hankey, G., and Yusuf, S. (1999). Homocyst(e)ine and cardiovascular disease: a critical review of the epidemiologic evidence. *Ann. Intern. Med.* 131, 363–375. doi: 10.7326/0003-4819-131-5-199909070-00008
- Eussen, S. J., de Groot, L. C., Joosten, L. W., Bloo, R. J., Clarke, R., Ueland, P. M., et al. (2006). Effect of oral vitamin B-12 with or without folic acid on cognitive function in older people with mild vitamin B-12 deficiency: a randomized, placebo-controlled trial. *Am. J. Clin. Nutr.* 84, 361–370. doi: 10.1093/ajcn/84.1.361
- Firbank, M. J., Narayan, S. K., Saxby, B. K., Ford, G. A., and O'Brien, J. T. (2010). Homocysteine is associated with hippocampal and white matter atrophy in older subjects with mild hypertension. *Int. Psychogeriatr.* 22, 804–811. doi: 10.1017/S1041610210000499
- Ford, A. H., and Almeida, O. P. (2012). Effect of homocysteine lowering treatment on cognitive function: a systematic review and meta-analysis of randomized controlled trials. *J. Alzheimers Dis.* 29, 133–149. doi: 10.3233/JAD-2012-111739
- Galis, Z. S., Muszynski, M., Sukhova, G. K., Simon-Morrissey, E., Unemori, E. N., Lark, M. W., et al. (1994). Cytokine-stimulated human vascular smooth muscle cells synthesize a complement of enzymes required for extracellular matrix digestion. *Circ. Res.* 75, 181–189. doi: 10.1161/01.RES.75.1.181
- Gibson, J. B., Carson, N. A., and Neill, D. W. (1964). Pathological findings in homocystinuria. *J. Clin. Pathol.* 17, 427–437. doi: 10.1136/jcp.17.4.427
- Graham, I. M., Daly, L. E., Refsum, H. M., Robinson, K., Brattstrom, L. E., Ueland, P. M., et al. (1997). Plasma homocysteine as a risk factor for vascular disease. The European Concerted Action Project. *JAMA* 277, 1775–1781. doi: 10.1001/jama.1997.03540460039030
- Hainsworth, A. H., Yeo, N. E., Weekman, E. M., and Wilcock, D. M. (2016). Homocysteine, hyperhomocysteinemia and vascular contributions to cognitive impairment and dementia (VCID). *Biochim. Biophys. Acta* 1862, 1008–1017. doi: 10.1016/j.bbadis.2015.11.015
- Ho, P. I., Ashline, D., Dhitavat, S., Ortiz, D., Collins, S. C., Shea, T. B., et al. (2003). Folate deprivation induces neurodegeneration: roles of oxidative stress and increased homocysteine. *Neurobiol. Dis.* 14, 32–42. doi: 10.1016/S0969-9961(03)00070-6
- Hyland, K., Smith, I., Bottiglieri, T., Perry, J., Wendel, U., Clayton, P. T., et al. (1988). Demyelination and decreased S-adenosylmethionine in 5,10-methylenetetrahydrofolate reductase deficiency. *Neurology* 38, 459–462. doi: 10.1212/WNL.38.3.459
- Jakubowski, H. (1999). Protein homocysteinylation: possible mechanism underlying pathological consequences of elevated homocysteine levels. *FASEB J.* 13, 2277–2283. doi: 10.1096/fasebj.13.15.2277
- Jakubowski, H., Zhang, L., Bardeguet, A., and Aviv, A. (2000). Homocysteine thiolactone and protein homocysteinylation in human endothelial cells: implications for atherosclerosis. *Circ. Res.* 87, 45–51. doi: 10.1161/01.RES.87.1.45
- Janota, C., Lemere, C. A., and Brito, M. A. (2016). Dissecting the contribution of vascular alterations and aging to Alzheimer's disease. *Mol. Neurobiol.* 53, 3793–3811. doi: 10.1007/s12035-015-9319-7

- Jellinger, K. A., and Attems, J. (2010). Is there pure vascular dementia in old age? *J. Neurol. Sci.* 299, 150–154. doi: 10.1016/j.jns.2010.08.038
- Jerneren, F., Elshorbagy, A. K., Oulhaj, A., Smith, S. M., Refsum, H., and Smith, A. D. (2015). Brain atrophy in cognitively impaired elderly: the importance of long-chain omega-3 fatty acids and B vitamin status in a randomized controlled trial. *Am. J. Clin. Nutr.* 102, 215–221. doi: 10.3945/ajcn.114.103283
- Jiang, X., Yang, F., Tan, H., Liao, D., Bryan, R. M. Jr., Randhawa, J. K., et al. (2005). Hyperhomocysteinemia impairs endothelial function and eNOS activity via PKC activation. *Arterioscler. Thromb. Vasc. Biol.* 25, 2515–2521. doi: 10.1161/01.ATV.0000189559.87328.e4
- Joosten, E., van den Berg, A., Riezler, R., Naurath, H. J., Lindenbaum, J., Stabler, S. P., et al. (1993). Metabolic evidence that deficiencies of vitamin B-12 (cobalamin), folate, and vitamin B-6 occur commonly in elderly people. *Am. J. Clin. Nutr.* 58, 468–476. doi: 10.1093/ajcn/58.4.468
- Kamath, A. F., Chauhan, A. K., Kisucka, J., Dole, V. S., Loscalzo, J., Handy, D. E., et al. (2006). Elevated levels of homocysteine compromise blood-brain barrier integrity in mice. *Blood* 107, 591–593. doi: 10.1182/blood-2005-06-2506
- Kang, J. H., Cook, N., Manson, J., Buring, J. E., Albert, C. M., and Grodstein, F. (2008). A trial of B vitamins and cognitive function among women at high risk of cardiovascular disease. *Am. J. Clin. Nutr.* 88, 1602–1610. doi: 10.3945/ajcn.2008.26404
- Kim, J. M., Lee, H., and Chang, N. (2002). Hyperhomocysteinemia due to short-term folate deprivation is related to electron microscopic changes in the rat brain. *J. Nutr.* 132, 3418–3421. doi: 10.1093/jn/132.11.3418
- Klein, T., and Bischoff, R. (2011). Physiology and pathophysiology of matrix metalloproteinases. *Amino Acids* 41, 271–290. doi: 10.1007/s00726-010-0689-x
- Langa, K. M., Foster, N. L., and Larson, E. B. (2004). Mixed dementia: emerging concepts and therapeutic implications. *JAMA* 292, 2901–2908. doi: 10.1001/jama.292.23.2901
- Lewerin, C., Matousek, M., Steen, G., Johansson, B., Steen, B., and Nilsson-Ehle, H. (2005). Significant correlations of plasma homocysteine and serum methylmalonic acid with movement and cognitive performance in elderly subjects but no improvement from short-term vitamin therapy: a placebo-controlled randomized study. *Am. J. Clin. Nutr.* 81, 1155–1162. doi: 10.1093/ajcn/81.5.1155
- Locasale, J. W. (2013). Serine, glycine and one-carbon units: cancer metabolism in full circle. *Nat. Rev. Cancer* 13, 572–583. doi: 10.1038/nrc3557
- Lonn, E., Yusuf, S., Arnold, M. J., Sheridan, P., Pogue, J., Micks, M., et al. (2006). Homocysteine lowering with folic acid and B vitamins in vascular disease. *N. Engl. J. Med.* 354, 1567–1577. doi: 10.1056/NEJMoa060900
- Maron, B. A., and Loscalzo, J. (2009). The treatment of hyperhomocysteinemia. *Annu. Rev. Med.* 60, 39–54. doi: 10.1146/annurev.med.60.041807.123308
- Mattson, M. P., and Shea, T. B. (2003). Folate and homocysteine metabolism in neural plasticity and neurodegenerative disorders. *Trends Neurosci.* 26, 137–146. doi: 10.1016/S0166-2236(03)00032-8
- McCully, K. S. (1969). Vascular pathology of homocysteinemia: implications for the pathogenesis of arteriosclerosis. *Am. J. Pathol.* 56, 111–128.
- McCully, K. S. (2007). Homocysteine, vitamins, and vascular disease prevention. *Am. J. Clin. Nutr.* 86, 1563S–1568S. doi: 10.1093/ajcn/86.5.1563S
- McMahon, J. A., Green, T. J., Skeaff, C. M., Knight, R. G., Mann, J. I., and Williams, S. M. (2006). A controlled trial of homocysteine lowering and cognitive performance. *N. Engl. J. Med.* 354, 2764–2772. doi: 10.1056/NEJMoa054025
- Michaluk, P., Kolodziej, L., Mioduszevska, B., Wilczynski, G. M., Dzwonek, J., Jaworski, J., et al. (2007). Beta-dystroglycan as a target for MMP-9, in response to enhanced neuronal activity. *J. Biol. Chem.* 282, 16036–16041. doi: 10.1074/jbc.M700641200
- Miller, J. W., Green, R., Mungas, D. M., Reed, B. R., and Jagust, W. J. (2002). Homocysteine, vitamin B6, and vascular disease in AD patients. *Neurology* 58, 1471–1475. doi: 10.1212/WNL.58.10.1471
- Moll, S., and Varga, E. A. (2015). Homocysteine and MTHFR mutations. *Circulation* 132, e6–e9. doi: 10.1161/CIRCULATIONAHA.114.013311
- Neves, M. F., Endemann, D., Amiri, F., Viridis, A., Pu, Q., Rozen, R., et al. (2004). Small artery mechanics in hyperhomocysteinemic mice: effects of angiotensin II. *J. Hypertens.* 22, 959–966. doi: 10.1097/00004872-200405000-00018
- Olney, J. W., Price, M. T., Salles, K. S., Labruyere, J., Ryerson, R., Mahan, K., et al. (1987). L-homocysteic acid: an endogenous excitotoxic ligand of the NMDA receptor. *Brain Res. Bull.* 19, 597–602. doi: 10.1016/0361-9230(87)90077-3
- Oulhaj, A., Refsum, H., Beaumont, H., Williams, J., King, E., Jacoby, R., et al. (2010). Homocysteine as a predictor of cognitive decline in Alzheimer's disease. *Int. J. Geriatr. Psychiatry* 25, 82–90. doi: 10.1002/gps.2303
- Peng, H. Y., Man, C. F., Xu, J., and Fan, Y. (2015). Elevated homocysteine levels and risk of cardiovascular and all-cause mortality: a meta-analysis of prospective studies. *J. Zhejiang Univ. Sci. B* 16, 78–86. doi: 10.1631/jzus.B1400183
- Perla-Kajan, J., Twardowski, T., and Jakubowski, H. (2007). Mechanisms of homocysteine toxicity in humans. *Amino Acids* 32, 561–572. doi: 10.1007/s00726-006-0432-9
- Pirchl, M., Ullrich, C., and Humpel, C. (2010). Differential effects of short- and long-term hyperhomocysteinemia on cholinergic neurons, spatial memory and microbleedings in vivo in rats. *Eur. J. Neurosci.* 32, 1516–1527. doi: 10.1111/j.1460-9568.2010.07434.x
- Poloni, S., Sperb-Ludwig, F., Borsatto, T., Weber Hoss, G., Doriqui, M. J. R., Embirucu, E. K., et al. (2018). CBS mutations are good predictors for B6-responsiveness: a study based on the analysis of 35 Brazilian Classical Homocystinuria patients. *Mol. Genet. Genomic Med.* 6, 160–170. doi: 10.1002/mgg3.342
- Prince, M., Wimo, A., Guerchet, M., Ali, G., Wu, Y., and Prina, M. (2015). *The Global Impact of Dementia: An Analysis of Prevalence, Incidence, Cost and Trends*. London: Alzheimer's Disease International,
- Rasic-Markovic, A., Stanojlovic, O., Hrcic, D., Krstic, D., Colovic, M., Susic, V., et al. (2009). The activity of erythrocyte and brain Na⁺/K⁺ and Mg²⁺-ATPases in rats subjected to acute homocysteine and homocysteine thiolactone administration. *Mol. Cell. Biochem.* 327, 39–45. doi: 10.1007/s11010-009-0040-6
- Rimm, E. B., Willett, W. C., Hu, F. B., Sampson, L., Colditz, G. A., Manson, J. E., et al. (1998). Folate and vitamin B6 from diet and supplements in relation to risk of coronary heart disease among women. *JAMA* 279, 359–364. doi: 10.1001/jama.279.5.359
- Robert, K., Pages, C., Ledru, A., Delabar, J., Caboche, J., and Janel, N. (2005). Regulation of extracellular signal-regulated kinase by homocysteine in hippocampus. *Neuroscience* 133, 925–935. doi: 10.1016/j.neuroscience.2005.03.034
- Sacharow, S. J., Picker, J. D., and Levy, H. L. (1993). "Homocystinuria caused by cystathionine beta-synthase deficiency," in *GeneReviews(R)*, eds M. P. Adam, H. H. Ardinger, R. A. Pagon, S. E. Wallace, L. J. H. Bean, K. Stephens, et al. (Seattle, WA: University of Washington).
- Schneider, J. A., and Bennett, D. A. (2010). Where vascular meets neurodegenerative disease. *Stroke* 41(Suppl. 10), S144–S146. doi: 10.1161/STROKEAHA.110.598326
- Selhub, J. (1999). Homocysteine metabolism. *Annu. Rev. Nutr.* 19, 217–246. doi: 10.1146/annurev.nutr.19.1.217
- Selhub, J. (2006). The many facets of hyperhomocysteinemia: studies from the Framingham cohorts. *J. Nutr.* 136(Suppl. 6), 1726S–1730S. doi: 10.1093/jn/136.6.1726S
- Shastri, S., and Tyagi, S. C. (2004). Homocysteine induces metalloproteinase and shedding of beta-1 integrin in microvessel endothelial cells. *J. Cell. Biochem.* 93, 207–213. doi: 10.1002/jcb.20137
- Simard, M., and Nedergaard, M. (2004). The neurobiology of glia in the context of water and ion homeostasis. *Neuroscience* 129, 877–896. doi: 10.1016/j.neuroscience.2004.09.053
- Smith, A. D., and Refsum, H. (2016). Homocysteine, B vitamins, and cognitive impairment. *Annu. Rev. Nutr.* 36, 211–239. doi: 10.1146/annurev-nutr-071715-050947
- Smith, A. D., Smith, S. M., de Jager, C. A., Whitbread, P., Johnston, C., Agacinski, G., et al. (2010). Homocysteine-lowering by B vitamins slows the rate of accelerated brain atrophy in mild cognitive impairment: a randomized controlled trial. *PLoS One* 5:e12244. doi: 10.1371/journal.pone.0012244
- Stott, D. J., MacIntosh, G., Lowe, G. D., Rumley, A., McMahon, A. D., Langhorne, P., et al. (2005). Randomized controlled trial of homocysteine-lowering vitamin treatment in elderly patients with vascular disease. *Am. J. Clin. Nutr.* 82, 1320–1326. doi: 10.1093/ajcn/82.6.1320
- Study of the Effectiveness of Additional Reductions in Cholesterol and Homocysteine (SEARCH) Collaborative Group, Armitage, J. M., Bowman, L., Clarke, R. J., Wallendszus, K., Bulbulia, R., et al. (2010). Effects of homocysteine-lowering with folic acid plus vitamin B12 vs placebo on mortality

- and major morbidity in myocardial infarction survivors: a randomized trial. *JAMA* 303, 2486–2494. doi: 10.1001/jama.2010.840
- Sudduth, T. L., Powell, D. K., Smith, C. D., Greenstein, A., and Wilcock, D. M. (2013). Induction of hyperhomocysteinemia models vascular dementia by induction of cerebral microhemorrhages and neuroinflammation. *J. Cereb. Blood Flow Metab.* 33, 708–715. doi: 10.1038/jcbfm.2013.1
- Sudduth, T. L., Weekman, E. M., Price, B. R., Gooch, J. L., Woolums, A., Norris, C. M., et al. (2017). Time-course of glial changes in the hyperhomocysteinemia model of vascular cognitive impairment and dementia (VCID). *Neuroscience* 341, 42–51. doi: 10.1016/j.neuroscience.2016.11.024
- Sunden, S. L., Renduchintala, M. S., Park, E. I., Miklasz, S. D., and Garrow, T. A. (1997). Betaine-homocysteine methyltransferase expression in porcine and human tissues and chromosomal localization of the human gene. *Arch. Biochem. Biophys.* 345, 171–174. doi: 10.1006/abbi.1997.0246
- Surtees, R., Leonard, J., and Austin, S. (1991). Association of demyelination with deficiency of cerebrospinal-fluid S-adenosylmethionine in inborn errors of methyl-transfer pathway. *Lancet* 338, 1550–1554. doi: 10.1016/0140-6736(91)92373-A
- Toole, J. F., Malinow, M. R., Chambless, L. E., Spence, J. D., Pettigrew, L. C., Howard, V. J., et al. (2004). Lowering homocysteine in patients with ischemic stroke to prevent recurrent stroke, myocardial infarction, and death: the Vitamin Intervention for Stroke Prevention (VISP) randomized controlled trial. *JAMA* 291, 565–575. doi: 10.1001/jama.291.5.565
- Troen, A. M., Shea-Budgell, M., Shukitt-Hale, B., Smith, D. E., Selhub, J., and Rosenberg, I. H. (2008). B-vitamin deficiency causes hyperhomocysteinemia and vascular cognitive impairment in mice. *Proc. Natl. Acad. Sci. U.S.A.* 105, 12474–12479. doi: 10.1073/pnas.0805350105
- Vecil, G. G., Larsen, P. H., Corley, S. M., Herx, L. M., Besson, A., Goodyer, C. G., et al. (2000). Interleukin-1 is a key regulator of matrix metalloproteinase-9 expression in human neurons in culture and following mouse brain trauma in vivo. *J. Neurosci. Res.* 61, 212–224. doi: 10.1002/1097-4547(20000715)61:2<212::AID-JNRI12>3.0.CO;2-9
- Vemuri, P., and Knopman, D. S. (2016). The role of cerebrovascular disease when there is concomitant Alzheimer disease. *Biochim. Biophys. Acta* 1862, 952–956. doi: 10.1016/j.bbdis.2015.09.013
- Vermeer, S. E., van Dijk, E. J., Koudstaal, P. J., Oudkerk, M., Hofman, A., Clarke, R., et al. (2002). Homocysteine, silent brain infarcts, and white matter lesions: the Rotterdam Scan Study. *Ann. Neurol.* 51, 285–289. doi: 10.1002/ana.10111
- Vitapops Trial Study Group (2010). B vitamins in patients with recent transient ischaemic attack or stroke in the VITamins TO Prevent Stroke (VITATOPS) trial: a randomised, double-blind, parallel, placebo-controlled trial. *Lancet Neurol.* 9, 855–865. doi: 10.1016/S1474-4422(10)70187-3
- Wald, D. S., Kasturiratne, A., and Simmonds, M. (2010). Effect of folic acid, with or without other B vitamins, on cognitive decline: meta-analysis of randomized trials. *Am. J. Med.* 123, 522–527.e2. doi: 10.1016/j.amjmed.2010.01.017
- Watanabe, M., Osada, J., Aratani, Y., Kluckman, K., Reddick, R., Malinow, M. R., et al. (1995). Mice deficient in cystathionine beta-synthase: animal models for mild and severe homocyst(e)inemia. *Proc. Natl. Acad. Sci. U.S.A.* 92, 1585–1589. doi: 10.1073/pnas.92.5.1585
- Weiss, N., Heydrick, S. J., Postea, O., Keller, C., Keaney, J. F. Jr., and Loscalzo, J. (2003). Influence of hyperhomocysteinemia on the cellular redox state—impact on homocysteine-induced endothelial dysfunction. *Clin. Chem. Lab. Med.* 41, 1455–1461. doi: 10.1515/CCLM.2003.223
- Welch, G. N., and Loscalzo, J. (1998). Homocysteine and atherothrombosis. *N. Engl. J. Med.* 338, 1042–1050. doi: 10.1056/NEJM199804093381507
- Yeh, Y. C., Huang, M. F., Hwang, S. J., Tsai, J. C., Liu, T. L., Hsiao, S. M., et al. (2016). Association of homocysteine level and vascular burden and cognitive function in middle-aged and older adults with chronic kidney disease. *Int. J. Geriatr. Psychiatry* 31, 723–730. doi: 10.1002/gps.4383
- Zekry, D., Hauw, J. J., and Gold, G. (2002). Mixed dementia: epidemiology, diagnosis, and treatment. *J. Am. Geriatr. Soc.* 50, 1431–1438. doi: 10.1046/j.1532-5415.2002.50367.x
- Zou, C. G., and Banerjee, R. (2005). Homocysteine and redox signaling. *Antioxid. Redox Signal.* 7, 547–559. doi: 10.1089/ars.2005.7.547

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Price, Wilcock and Weekman. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.