

University of Kentucky

UKnowledge

Plant and Soil Sciences Faculty Publications

Plant and Soil Sciences

12-11-2018

Occurrence of Alkaloids in Grass Seeds Symbiotic With Vertically-Transmitted *Epichloë* Fungal Endophytes and Its Relationship With Antioxidants

Pedro E. Gundel

Universidad de Buenos Aires, Argentina

Charlotte E. Seal

Royal Botanic Gardens, UK

Fernando Biganzoli

Universidad de Buenos Aires, Argentina

Marco A. Molina-Montenegro

Universidad de Talca, Chile

Beatriz R. Vázquez-de-Aldana

Instituto de Recursos Naturales y Agrobiología de Salamanca, Spain

Follow this and additional works at: https://uknowledge.uky.edu/pss_facpub



Part of the [Ecology and Evolutionary Biology Commons](#), and the [Plant Sciences Commons](#)

See next page for additional authors

Right click to open a feedback form in a new tab to let us know how this document benefits you.

Repository Citation

Gundel, Pedro E.; Seal, Charlotte E.; Biganzoli, Fernando; Molina-Montenegro, Marco A.; Vázquez-de-Aldana, Beatriz R.; Zabalgoceazcoa, Iñigo; Bush, Lowell P.; Martínez-Ghersa, María A.; and Ghersa, Claudio M., "Occurrence of Alkaloids in Grass Seeds Symbiotic With Vertically-Transmitted *Epichloë* Fungal Endophytes and Its Relationship With Antioxidants" (2018). *Plant and Soil Sciences Faculty Publications*. 138.

https://uknowledge.uky.edu/pss_facpub/138

This Article is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in Plant and Soil Sciences Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Occurrence of Alkaloids in Grass Seeds Symbiotic With Vertically-Transmitted *Epichloë* Fungal Endophytes and Its Relationship With Antioxidants

Digital Object Identifier (DOI)

<https://doi.org/10.3389/fevo.2018.00211>

Notes/Citation Information

Published in *Frontiers in Ecology and Evolution*, v. 6, article 211, p. 1-7.

Copyright © 2018 Gundel, Seal, Biganzoli, Molina-Montenegro, Vázquez-de-Aldana, Zabalgogezcoa, Bush, Martínez-Ghersa and Ghersa.

This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](#). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Authors

Pedro E. Gundel, Charlotte E. Seal, Fernando Biganzoli, Marco A. Molina-Montenegro, Beatriz R. Vázquez-de-Aldana, Iñigo Zabalgogezcoa, Lowell P. Bush, María A. Martínez-Ghersa, and Claudio M. Ghersa



Occurrence of Alkaloids in Grass Seeds Symbiotic With Vertically-Transmitted *Epichloë* Fungal Endophytes and Its Relationship With Antioxidants

Pedro E. Gundel^{1*}, Charlotte E. Seal², Fernando Biganzoli³, Marco A. Molina-Montenegro^{4,5}, Beatriz R. Vázquez-de-Aldana⁶, Iñigo Zabalgoceazcoa⁶, Lowell P. Bush⁷, María A. Martínez-Ghersa¹ and Claudio M. Ghersa¹

¹IFEVA, Facultad de Agronomía, Universidad de Buenos Aires, CONICET, Buenos Aires, Argentina, ²Department of Comparative Plant and Fungal Biology, Royal Botanic Gardens, Kew, United Kingdom, ³Departamento de Métodos Cuantitativos y Sistemas de Información, Facultad de Agronomía, Universidad de Buenos Aires, Buenos Aires, Argentina, ⁴CEMF, Instituto de Ciencias Biológicas, Universidad de Talca, Talca, Chile, ⁵CEAZA, Universidad Católica del Norte, Coquimbo, Chile, ⁶Instituto de Recursos Naturales y Agrobiología de Salamanca (IRNASA-CSIC), Salamanca, Spain, ⁷Department of Plant & Soil Sciences, University of Kentucky, Lexington, KY, United States

OPEN ACCESS

Edited by:

Margot Schulz,
Universität Bonn, Germany

Reviewed by:

Benjamin Fuchs,
Freie Universität Berlin, Germany
Chunjie Li,
Lanzhou University, China

*Correspondence:

Pedro E. Gundel
gundel@agro.uba.ar

Specialty section:

This article was submitted to
Chemical Ecology,
a section of the journal
Frontiers in Ecology and Evolution

Received: 08 October 2018

Accepted: 27 November 2018

Published: 11 December 2018

Citation:

Gundel PE, Seal CE, Biganzoli F, Molina-Montenegro MA, Vázquez-de-Aldana BR, Zabalgoceazcoa I, Bush LP, Martínez-Ghersa MA and Ghersa CM (2018) Occurrence of Alkaloids in Grass Seeds Symbiotic With Vertically-Transmitted *Epichloë* Fungal Endophytes and Its Relationship With Antioxidants. *Front. Ecol. Evol.* 6:211. doi: 10.3389/fevo.2018.00211

Host organisms can acquire new functional traits through symbiosis. Seed-transmitted *Epichloë* fungal endophytes are known to protect host plants against herbivores and increase tolerance to abiotic stresses by alkaloids and antioxidants, respectively (currencies of mutualism). Whereas, alkaloids are fungal products with demonstrated effects at plant vegetative stage, few studies have focused on alkaloids in seeds. We assessed the occurrence of fungal alkaloids and determined their concentrations in seeds of two host grasses, *Festuca rubra* and *Lolium multiflorum*. Then, we sought for a relationship with the antioxidants tocochromanols and glutathione, which are involved in the control of oxidative stress. Different alkaloids were detected depending on the species and plant genotype. Most notably, loline alkaloids were not detected in *F. rubra* seeds, whereas ergovaline and peramine were absent in *L. multiflorum*. In *F. rubra*, ergovaline concentration was dependent on the maternal line in interaction with the production year, diminishing in seeds after 1 year of storage. The exposure of *L. multiflorum* plants to ozone had no effect on the seed concentration of lolines. There was a significant positive relationship between the concentrations of ergovaline and tocochromanols in both species, and between ergovaline concentration and E_{GSSG/2GSH} (glutathione half-cell reduction potential) in RAB maternal line of *F. rubra*. These results suggest that alkaloid and antioxidants have a close association in seeds of host grasses, and that the alkaloid bioactivity could be related with the antioxidant capacity to control stress. This has important implications for the ecology of partner species, thus supporting its consideration for further research.

Keywords: grass-endophyte symbiosis, secondary compounds, defensive mutualism, currency of mutualism, seed quality

INTRODUCTION

Host organisms can acquire new functional traits through symbiosis (Kiers and West, 2015). In the symbiosis between cool-season grasses (Subfamily Poöideae) and *Epichloë* fungal endophytes (Clavicipitaceae) (reviewed by Clay and Scharndl, 2002), the high prevalence of symbiotic individuals in populations has been ascribed to both an endophyte-mediated resistance to herbivory by alkaloids and tolerance to abiotic stress by antioxidants, known as the “currencies of the mutualism” (Scharndl et al., 2007; Hamilton and Bauerle, 2012). Nonetheless, the effects of fungal endophytes on host plant fitness can vary depending to partners’ species and genotypes, and as a result of the ecological context (Clay and Scharndl, 2002; Saikkonen et al., 2013).

Seeds are critical to the grass-endophyte symbiosis as fungus carriers from one generation to the next (Gundel et al., 2011a, 2017). The seed is the plant structure with the greatest fungus/plant biomass ratio which stores fungal alkaloids (TePaske et al., 1993; Justus et al., 1997; Vázquez-de-Aldana et al., 2003). The profile and concentration of alkaloids varies greatly responding to factors such as species identities, plant tissue/stage, and growth conditions (Bush et al., 1997; Faeth et al., 2002; Rasmussen et al., 2007; Mace et al., 2014). Production of alkaloids can be triggered by herbivory (Sullivan et al., 2007; Fuchs et al., 2017a), although it remains unknown whether alkaloids accumulate in seeds as a consequence (Gundel et al., 2017). Endophyte presence in seeds is deemed to deter granivorous animals (Uchitel et al., 2011), an effect that may be extended to protect seedlings from herbivores (Stewart, 1985; Czarnoleski et al., 2010). However, the endophyte-conferred resistance in seedlings may be ineffective (Lewis and Clements, 1986) because of a very low alkaloid level (Dymock et al., 1989) and/or because of fungus viability loss. Since the alkaloids are present in the seed, their loss rate could be independent of the endophyte viability (Stewart, 1985), but dependent on seed quality parameters.

A potentiated antioxidant system is proposed as a mechanism by which symbiotic plants usually exhibit higher tolerance to abiotic stress (White and Torres, 2010; Hamilton et al., 2012). The presence of endophytes in seeds may be related to a higher antioxidant capacity. For example, endophytic seeds of *F. rubra* (RAB maternal line) had a higher concentration of the lipid-soluble antioxidants α -tocopherol and α -tocotrienol than non-endophytic seeds (Gundel et al., 2012), that may contribute to stabilize membrane functionality (Falk and Munné-Bosch, 2010). In contrast, the glutathione half-cell reduction potential ($E_{GSSG/2GSH}$), a marker of the cellular redox state that shifts toward positive values with a loss of seed viability (Kranmer et al., 2010; Seal et al., 2010), was more positive in endophytic compared to non-endophytic seeds of *L. multiflorum* (Gundel et al., 2015). However, the presence of an endophyte does not always contribute toward viability. Whereas endophytes enhance the survival of germinating seeds in *L. multiflorum* and *F. rubra* (Gundel et al., 2006, 2011b), they had a negative effect on seed viability under both cool-dry and wet-warm storage conditions (Gundel et al., 2012, 2015). Therefore, endophyte effects on seeds may differ depending

on the plant species, seed quality, seed storage conditions and antioxidant.

With growing evidence beginning to elucidate the ecological functions of alkaloids and antioxidants in the grass-endophyte symbiosis independently, it is striking that any relationship between these two compounds is yet to be explored. Here, we first present data on the concentration of different fungal alkaloids in seeds of two grass species, *F. rubra* and *L. multiflorum*. With the aim to have within-species variation in the profile and level of alkaloids, we analyzed seeds produced by different plants and years in the case of *F. rubra*, and produced in atmospheres with high and low concentration of ozone in the case of *L. multiflorum*. Ozone is a stress factor associated to tropospheric contamination that depending on the intensity, can cause growth losses or induced resistance in plants by activating the antioxidant system (Menéndez et al., 2009). Then, we explored the existence of relationships between alkaloids and antioxidants. A relationship between these currencies of mutualism could have significant ecological consequences: a negative relationship elicited by different factors (i.e., antioxidants to abiotic stress and alkaloids to herbivory) would mean that the occurrence of a given factor (e.g., stress factor) would render the plant susceptible to another factor (e.g., herbivory) and *vice-versa*. Alternatively, a positive relationship would indicate that a lower concentration of antioxidants would imply a lower concentration of alkaloids and consequently seeds may be more prone to herbivory.

MATERIALS AND METHODS

Seed Material

Festuca rubra

We used half-sib seeds produced by clones of three individual plants (RAB, SAN, and PEN) infected by *Epichloë festucae*, the common fungal endophyte of fine fescues (*Zabalgogea* et al., 2006; Dirihan et al., 2016). The original plants were collected from three sites separated by at least 40 km, in dehesa grasslands of Salamanca, Spain. *Epichloë* fungal endophytes have shown to affect different growth attributes and biochemical variables of those maternal lines (*Zabalgogea* et al., 2006; Vázquez-de-Aldana et al., 2013). Each plant was split into three ramets, randomly transplanted in a field plot on a research farm (IRNASA-CSIC, Salamanca, Spain; Gundel et al., 2011b). The seeds were produced in 3 years: 2009, 2010, and 2011, and stored for 24, 12, and 1 month, respectively, under dry and cold (10°C) conditions. In 2012, seed lots produced by each maternal line and ramet ($n = 2$; since a ramet was lost in two of the maternal lines, we decided to use just 2 ramets each) in each year, were processed for alkaloid determination.

Lolium multiflorum

The original population was collected in a semi-natural grassland (Pampa Region, Argentina) where populations usually present high frequency of endophyte-infected plants (Gundel et al., 2009). In 2011, *L. multiflorum* plants with and without *Epichloë occaltans* were exposed to a high concentration of tropospheric ozone (Gundel et al., 2015). At pre-anthesis, 64 plants growing

independently in 2 L pots were placed in any of the 8 open-top chambers with either high (≈ 120 ppb) or low (< 10 ppb) levels of ozone for 5 days (2 h at noon). Seeds produced by all the plants within an open-top chamber were pooled and stored in dry and cold conditions for 3 months. In 2012, seed lots produced by all the plants within a chamber were pooled ($n = 4$ for each ozone treatment) and were processed for alkaloid determination.

Alkaloid Analysis

Concentration of the ergopetine alkaloid, ergovaline (Chemical name: 12'-Hydroxy-2'-methyl-5'alpha-(1-methylethyl)-ergotaman-3',6',18-trione) was quantified by HPLC following a modification of the methods described by Hill et al. (1993) and Yue et al. (2000). A 1.0 g ground seed sample was extracted in 20 ml of CHCl_3 and 1 ml of 0.5 mM NaOH for 2 h. A solution ($10 \mu\text{g mL}^{-1}$) of ergotamine ditartrate (Sigma-Aldrich) was added as internal standard. The mixture was vacuum-filtered through Whatman n° 2 filter paper and a 10 ml aliquot of filtrate was passed through a 500 mg Ergosil (Analtech; Newark, USA) solid-phase column preconditioned with CHCl_3 . Plant pigments were removed with 5 ml of chloroform:acetone (1:3). The sample was eluted with 2 ml of methanol and vacuum concentrated, redissolved in 1 ml of methanol, and filtered through a $0.22 \mu\text{m}$ nylon filter. Ergovaline quantification in extracts was performed in a HPLC system (Waters 2690) with an Xterra MS C18 Waters column (4.6×100 mm) and a fluorescence detector (Waters 2475) $\lambda_{\text{exc}} = 250$ nm and $\lambda_{\text{em}} = 420$ nm. The mobile phase was acetonitrile and 0.01M ammonium acetate with a gradient flow of 0.8 mL min^{-1} .

Peramine (Chemical name: 2-[3-(2-methyl-1-oxopyrrolo[1,2-a]pyrazin-3-yl)propyl]guanidine) was determined using the HPLC method described by Barker et al. (1993) and Yue et al. (2000). A freeze-dried and ground sample (100 mg) was extracted in 3 ml of 30% isopropanol for 30 min at 90°C . The mixture was centrifuged and the extract was passed through a preconditioned Varian Bond Elut carboxylic acid (CBA) column packed with 100 mg of adsorbent. After a wash of the column with 1–2 ml of methanol, peramine was eluted with 1 ml of 5% formic acid in 80% aqueous methanol. The extract was filtered through a $0.22 \mu\text{m}$ nylon filter and chromatographed in a Waters 2690 system with a Nova Pak C18 Waters column (3.9×150 mm). The isocratic mobile phase consisted of 18% (v/v) acetonitrile in a guanidine carbonate (10 mM)-formic acid buffer. Detection was performed with a Photodiode Array Detector (Waters 2996) at 280 nm. For more details see Vázquez-de-Aldana et al. (2010).

Pyrrolizidine alkaloids (NANL: N-acetilnorloline, NFL: N-formylloline, and NAL: N-acetylloine) [Chemical name: (6r,7r,7as)-n-methylhexahydro-1h-1,6-epoxyppyrolizin-7-amine] were extracted from powdered plant material with ethanol:methylene chloride (4:1, v/v) containing internal standard quinoline and sodium bicarbonate following protocol in Helander et al. (2016). Individual alkaloids were resolved and quantified by GC equipped with FID detector. Chromatographic conditions were $15 \text{ m} \times 0.53 \text{ mm DB5}$ column with initial oven temperature of 70°C increased to 160°C at $45^\circ\text{C min}^{-1}$, held for 5 min and increased to 290 at 45 min^{-1} and held for 7 min.

Antioxidant Analysis

In searching for relationships with the alkaloids, we focussed on total tocochromanols (that comprises several independent compounds of tocopherols and tocotrienols) and the glutathione half-cell reduction potential ($E_{\text{GSSG}/2\text{GSH}}$) that were found to be higher in concentration (tocochromanols) or more negative ($E_{\text{GSSG}/2\text{GSH}}$) in the presence of seed endophytes (Gundel et al., 2012, 2015). Published values were taken from our earlier works which analyzed the same seed lots (Gundel et al., 2012, 2015).

Data Analysis

We built generalized least squares models (gls function, nlme package, Pinheiro et al., 2017) in R (R. Core Team, 2017) to test the effects of the different variation factors on the concentration of alkaloids in seeds. In the case of *F. rubra*, the alkaloid was ergovaline while the factors were maternal line (PEN and RAB; SAN was not included in the analysis since we did not detect ergovaline) and production year (2009, 2010, and 2011). In the case of *L. multiflorum*, the alkaloids were “total lolines” and derivatives (NANL, NFL, and NAL) as affected by the maternal exposure to ozone. When necessary, we also modeled variance heterogeneity of errors with the option weights. Significance was tested with Type II Likelihood Ratio Test (car package; Fox and Weisberg, 2011). Finally, we tested the relationship between the alkaloids (ergovaline and total lolines) with the total tocochromanols and the $E_{\text{GSSG}/2\text{GSH}}$ in the seeds.

RESULTS

Festuca rubra

Loline alkaloids were not detected in any of the assessed *F. rubra* seed. The alkaloid peramine was only present in seeds of the maternal line PEN produced in 2010 ($3.12 \pm 0.01 \mu\text{g.g}^{-1}$) and 2011 ($5.68 \pm 0.01 \mu\text{g.g}^{-1}$; **Table 1**).

Ergovaline was not detected in seeds from the maternal line SAN in any of the production years (**Table 1**). The concentration of ergovaline in seeds of the other two maternal lines was found dependent on production year ($X_1^2 = 302.870$, $P < 0.001$). The concentration in seeds of RAB was 15-fold higher in comparison to that of PEN with a mean difference across production years of $0.61 \mu\text{g.g}^{-1}$ (**Table 1**).

Ergovaline concentration was related to tocochromanol concentrations but varied between maternal lines ($X_1^2 = 20.492$, $P < 0.001$). For the RAB lineage, a significant positive relationship between the alkaloid and tocochromanol concentrations was apparent, although this relationship was not related with the production year ($X_1^2 = 0.003$, $P = 0.956$; **Figure 1A**). The relationship between alkaloid and tocochromanol concentration was not significant in PEN.

Independently of the year ($X_1^2 = 2.065$, $P = 0.151$) but influenced by the maternal line ($X_1^2 = 37.944$, $P < 0.001$), the concentration of ergovaline was negatively related to $E_{\text{GSSG}/2\text{GSH}}$ for RAB but positively related for PEN (**Figure 1B**).

Lolium multiflorum

Neither ergovaline nor peramine were detected in any of the *L. multiflorum* seed lots (**Table 1**).

TABLE 1 | Concentration of fungal alkaloids in endophyte-symbiotic seeds of *Festuca rubra* and *Lolium multiflorum* as affected by “maternal line and production year”, and “maternal exposure to ozone,” respectively.

Species	Variation factor		Alkaloid		
	Maternal line	Production year	Ergovaline ($\mu\text{g}\cdot\text{g}^{-1}$ DM)	Lolines ($\mu\text{g}\cdot\text{g}^{-1}$ DM)	Peramine ($\mu\text{g}\cdot\text{g}^{-1}$ DM)
<i>Festuca rubra</i>	SAN	2009	n.d.	n.d.	n.d.
		2010	n.d.	n.d.	n.d.
		2011	n.d.	n.d.	n.d.
	PEN	2009	0.08 (0.00)	n.d.	n.d.
		2010	0.01 (0.00)	n.d.	3.12 (0.01)
		2011	0.04 (0.00)	n.d.	5.68 (0.01)
	RAB	2009	0.43 (0.01)	n.d.	n.d.
		2010	0.80 (0.02)	n.d.	n.d.
		2011	0.72 (0.02)	n.d.	n.d.
	Maternal ozone exposure				
<i>Lolium multiflorum</i>	high (≈ 120 ppb)		n.d.	1440.75 (226.00)	n.d.
	low (< 10 ppb)		n.d.	1426.50 (171.74)	n.d.

Values are mean (SE); $n = 2$, and $n = 4$, for *F. rubra* and *L. multiflorum*, respectively.

The exposure of mother plants to ozone did not affect either the concentration of total lolines in the seed ($X_1^2 = 0.003$, $P = 0.961$; **Table 1**) nor that of the lolines individually (NANL: $X_1^2 = 0.820$, $P = 0.365$, NFL: $X_1^2 = 0.046$, $P = 0.831$, and NAL: $X_1^2 = 0.001$, $P = 0.974$). The average concentration of each individual loline alkaloid in seeds produced by mother plants treated and non-treated with ozone was 254.87 ± 20.84 , 1147.12 ± 113.43 , and 31.625 ± 3.49 (mean \pm SE; $\mu\text{g}\cdot\text{g}^{-1}$ DM) for NANL, NFL, and NAL, respectively.

A significant positive relationship between the concentration of lolines and tocochromanols ($X_1^2 = 11.693$, $P = 0.001$) was independent of mother plants exposure to ozone ($X_1^2 = 2.773$, $P = 0.096$; **Figure 1C**). There was no relationship between loline concentration and $E_{\text{GSSG}/2\text{GSH}}$ ($X_1^2 = 0.133$, $P = 0.715$; **Figure 1D**).


DISCUSSION

Here we showed that, despite variability accounted for by genotype and species identity and the environmental conditions of seed production, seeds are carriers of fungal alkaloids, and the concentration may be related with the antioxidants system of the seeds. Because antioxidants are related to seed quality and functionality (Kranner et al., 2010; Seal et al., 2010), they may also be involved in the control of endophyte longevity (Gundel et al., 2012, 2015) and alkaloids bioactivity. Although more studies are certainly required, the two currencies of mutualism (Scharndl et al., 2007; Hamilton and Bauerle, 2012) appear as fundamental but interactive players for the ecological fate of the symbiotic unit (i.e., the seed). By joining host grass life cycle, fungal endophytes multiply and disperse through the seed (Gundel et al., 2011a). Thus, seeds are a crucial structure to protect (Gundel et al., 2017).

The alkaloid ergovaline is mainly known due to its toxicity on mammals (Bush et al., 1997; Saikkonen et al., 2013), although it can also be effective in deterring herbivorous insects (Potter

et al., 2008). The concentration of the alkaloid ergovaline varies greatly among plant species, genotypes and tissues (Scharndl et al., 2013; Mace et al., 2014). A great variation was previously shown between plants ($0\text{--}0.47 \mu\text{g}\cdot\text{g}^{-1}$) in stems and leaves of *F. rubra* (Vázquez-de-Aldana et al., 2007, 2010). In our study, the concentrations of ergovaline in seeds were similar to those found in plant vegetative tissues in Italian ecotypes (Jensen et al., 2007) but lower than those from Switzerland (Leuchtmann et al., 2000). An ergovaline concentration of $0.40 \mu\text{g}\cdot\text{g}^{-1}$ is considered a critical threshold in cattle diets above which symptoms of toxicosis can be observed (Bony and Delatour, 2001). Although much lower than the concentration found in seeds of *Schedonorus arundinaceus* ($1.1\text{--}4.4 \mu\text{g}\cdot\text{g}^{-1}$; TePaske et al., 1993), the seeds produced by the RAB plants exhibited seed alkaloid concentrations higher than the toxicosis threshold (from 0.41 to $0.83 \mu\text{g}\cdot\text{g}^{-1}$), being probably effective in deterring small rodents and herbivorous insects. Additionally, the observed variation among plants in alkaloid profile and concentration would indicate potential for populations to respond to selection pressures exerted by herbivory.

A similar variation has been observed in plant species associated with loline-producing endophytes. Concentration of lolines in *L. multiflorum* seeds showed high variability ($880\text{--}1,922 \mu\text{g}\cdot\text{g}^{-1}$) but it was on average significantly higher ($\approx 1430.63 \mu\text{g}\cdot\text{g}^{-1}$) compared to the concentration found in dry seed of the same species previously (TePaske et al., 1993) ($\approx 52 \mu\text{g}\cdot\text{g}^{-1}$). However, the concentrations of lolines we found are slightly lower to that reported for *F. pratensis* dry seeds ($\approx 1,801 \mu\text{g}\cdot\text{g}^{-1}$) by Justus et al. (1997). The fungal endophytes of these two last grass species, *E. occultans* and *E. uncinatum*, are well-known as loline-producing endophytes and the latter one is known for its high production level (Scharndl et al., 2007). Since lolines are insect deterrents and non-toxic for vertebrates, forage species infected with those endophytes such as *L. multiflorum*, *L. rigidum*, and *F. pratensis*, are naturally protected against



agricultural plagues. Lolines are also found in seeds of *S. arundinaceus* infected by *E. coenophiala* (Jackson et al., 1984). The existence of strains producing lolines and/or peramine but not ergovaline or lolitrem-B (that are toxic for livestock) is the foundation of the use of the “safe-endophyte technology” in forage breeding (Gundel et al., 2013). This technology relies on the quality of the seeds since not only the fungus but also its alkaloids determine its successful application in agriculture.

Interestingly, even though the gene required for the peramine (perA) is found in *E. occultaans*, the alkaloid remains undetected in tissue and now in seeds, of *Lolium multiflorum* (Bastías et al., 2017). Peramine is a unique and potent insecticide fungal alkaloid (Bush et al., 1997; Scharld et al., 2013). Here, it was only detected in seeds produced by the *F. rubra* PEN but with variation

among years. There was a clear relationship between the level of peramine and the year (2009: $0 \mu\text{g}\cdot\text{g}^{-1}$, 2010: $3.12 \mu\text{g}\cdot\text{g}^{-1}$, and 2011: $5.68 \mu\text{g}\cdot\text{g}^{-1}$, corresponding to 24, 12, and 1 month of storage) suggesting a susceptibility to denaturalize with time. It was found in *L. perenne* that the level of peramine follows the growing cycle of the host grass, showing a minimum ($< 1.0 \mu\text{g}\cdot\text{g}^{-1}$) in winter and a maximum ($4\text{--}8 \mu\text{g}\cdot\text{g}^{-1}$) in summer (Fuchs et al., 2017b). With a suggested threshold of $2 \mu\text{g}\cdot\text{g}^{-1}$ for peramine to be effective against herbivores (see references in Fuchs et al., 2017b), the concentration observed in seeds of PEN *F. rubra* line has the potential to be preserved for up to 1 year of storage.

Unlike the alkaloids, for which the genetic information for their synthesis is encoded in the endophyte genome

(Schardl et al., 2013), both the plant and the fungus can, in principle, produce molecules with antioxidant capacity (Huang et al., 2007). However, no precise information exists regarding the direct contribution of the fungus toward the antioxidant activity of plants (Hamilton et al., 2012) and their seeds (Gundel et al., 2012, 2015). Our results support the hypothesis that there is a positive relationship between both these two groups of secondary compounds. It was particularly evident for ergovaline in *F. rubra* RAB maternal line, as the concentration of alkaloid was related to a higher antioxidant capacity with regard to both tocochromanols and $E_{GSSG}/2GSH$, whereas in *L. multiflorum* seeds, a positive relationship was only found between loline alkaloids and tocochromanols.

The functionality of both tocochromanols and glutathione in the seed has implications for the ecology of the species. Previous studies that show a protective role of endophytes on seeds and seedlings, highlight alkaloids as the compounds responsible for this fact. For example, endophyte-infected seedlings of *L. perenne* were resistant to the weevil *Listronotus bonariensis* even though the fungus was dead, proving that alkaloids remain bioactive long after the demise of the fungus (Stewart, 1985). Nonetheless, this bioactivity could be broken if alkaloids are target of oxidative processes associated with aging and deteriorating conditions (Walters, 1998). We suggest that a lower antioxidant capacity is

associated with a lower concentration of alkaloids. As seed quality parameters decline, the potential for the endophytes to protect the seeds is likely to diminish.

AUTHOR CONTRIBUTIONS

PG, MAM-G, and CMG designed the research. PG, FB and MAM-M analyzed the data and made the figures. BRVdA, IZ, LPB, and CS performed the analyses for alkaloid and antioxidant quantification. PG, CS, BRVdA, and MAM-M wrote the manuscript. PG edited and submitted the manuscript. All authors have approved the final version of the manuscript.

FUNDING

This article was supported by the projects PII20150126 (Fondecyt-Chile) and AGL2016-76035-C21R (Spanish, MICINN).

ACKNOWLEDGMENTS

Authors thank Andrea C. Ueno for his technical assistance of preparing seeds for their analyses.

REFERENCES

- Barker, D. J., Davies, E., Lane, G. A., Latch, G. C. M., Nott, H. M., and Tapper, B. A. (1993). "Effect of water deficit on alkaloid concentrations in perennial ryegrass endophyte associations," in *Proceedings of the 2nd International Symposium on Acremonium/Grass Interactions*, eds D. E. Hume, G. C. M. Latch, and H. S. Easton (Palmerston North), 67–71.
- Bastias, D. A., Ueno, A. C., Machado-Asseff, C. R., Alvarez, A. E., Young, C. A., and Gundel, P. E. (2017). Metabolism or behavior: explaining the performance of aphids on alkaloid-producing fungal endophytes in annual ryegrass (*Lolium multiflorum*). *Oecologia* 185, 245–256. doi: 10.1007/s00442-017-3940-2
- Bony, S., and Delatour, P. (2001). "Relevance and impact of grass endophyte toxins in Europe," in *The Grassland Conference*, eds V. H. Paul and P. D. Dapprich (Soest).
- Bush, L. P., Wilkinson, H. H., and Schardl, C. L. (1997). Bioprotective alkaloids of grass-fungal endophyte symbioses. *Plant Physiol.* 114, 1–7. doi: 10.1104/pp.114.1.1
- Clay, K., and Schardl, C. P. (2002). Evolutionary origins and ecological consequences of endophyte symbiosis with grasses. *Am. Nat.* 160, 99–127. doi: 10.1086/342161
- Czarnoleski, M., Olejniczak, P., Mikołajczak, P., Lembicz, M., and Kozłowski, J. (2010). Fungal endophytes protect grass seedlings against herbivory and allow economical seed production. *Evol. Ecol. Res.* 12, 1–9.
- Dirihan, S., Helander, M., Väre, H., Gundel, P. E., Garibaldi, L. A., Irisarri, J. G. N., et al. (2016). Geographic variation in *Festuca rubra* L. ploidy levels and systemic fungal endophyte frequencies. *PLoS ONE* 11:e0166264. doi: 10.1371/journal.pone.0166264
- Dymock, J. J., Latch, G. C. M., and Tapper, B. A. (1989). "Novel combinations of endophytes in ryegrasses and fescues and their effects on Argentine stem weevil (*Listronotus bonariensis*) feeding," in *Proc Of 5th Australasian Conf. Grassland Invertebrate Ecology*, ed P. P. Stahle Melbourne, VIC 28–34.
- Faeth, S. H., Bush, L. P., and Sullivan, T. J. (2002). Peramine alkaloid variation in *Neotyphodium*-infected Arizona Fescue: effects of endophyte and host genotype and environment. *J. Chem. Ecol.* 28, 1511–1525. doi: 10.1023/A:1019916227153
- Falk, J., and Munné-Bosch, S. (2010). Tocochromanol functions in plants: antioxidation and beyond. *J. Exp. Bot.* 61, 1549–1566. doi: 10.1093/jxb/erq030
- Fox, J., and Weisberg, S. (2011). *An {R} Companion to Applied Regression, 2nd Edn*. Thousand Oaks, CA: Sage. Available online at: <http://socserv.socsci.mcmaster.ca/jfox/Books/Companion>
- Fuchs, B., Krischke, M., Mueller, M. J., and Krauss, J. (2017a). Herbivore-specific induction of defence metabolites in a grass–endophyte association. *Funct. Ecol.* 31, 318–324. doi: 10.1111/1365-2435.12755
- Fuchs, B., Krischke, M., Mueller, M. J., and Krauss, J. (2017b). Plant age and seasonal timing determine endophyte growth and alkaloid biosynthesis. *Fungal Ecol.* 29, 52–58. doi: 10.1016/j.funeco.2017.06.003
- Gundel, P. E., Garibaldi, L. A., Tognetti, P. M., Aragón, R., Ghersa, C. M., and Omacini, M. (2009). Imperfect vertical transmission of the endophyte *Neotyphodium* in exotic grasses in grasslands of the Flooding Pampa. *Microb. Ecol.* 57, 740–748. doi: 10.1007/s00248-008-9447-y
- Gundel, P. E., Hamilton, C. E., Seal, C. E., Helander, M., Martínez-Ghersa, M. A., Ghersa, C. M., et al. (2012). Antioxidants in *Festuca rubra* L. seeds affected by the fungal symbiont *Epichloë festucae*. *Symbiosis* 58, 73–80. doi: 10.1007/s13199-012-0194-y
- Gundel, P. E., Maseda, P. H., Vila-Aiub, M. M., Ghersa, C. M., and Benech-Arnold, R. L. (2006). *Neotyphodium* fungi effects on *Lolium multiflorum* seed germination in relation to water availability. *Ann. Bot. Lond.* 97, 571–577. doi: 10.1093/aob/mcl004
- Gundel, P. E., Pérez, L. I., Helander, M., and Saikkonen, K. (2013). Symbiotically modified organisms: nontoxic fungal endophytes in grasses. *Trends Plant Sci.* 18, 420–427. doi: 10.1016/j.tplants.2013.03.003
- Gundel, P. E., Rudgers, J. A., and Ghersa, C. M. (2011a). Incorporating the process of vertical transmission into understanding of host-symbiont dynamics. *Oikos* 120, 1121–1128.
- Gundel, P. E., Rudgers, J. A., and Whitney, K. D. (2017). Vertically transmitted symbionts as mechanisms of transgenerational effects. *Am. J. Bot.* 104, 787–792. doi: 10.3732/ajb.1700036
- Gundel, P. E., Sorzoli, N., Ueno, A. C., Ghersa, C. M., Seal, C. E., Bastias, D. A., et al. (2015). Impact of ozone on the viability and antioxidant content of grass seeds is affected by a vertically transmitted symbiotic fungus. *Environ. Exp. Bot.* 113, 40–46. doi: 10.1016/j.envexpbot.2015.01.001

- Gundel, P. E., Zabalgoagezcoa, I., and Vázquez-de-Aldana, B. R. (2011b). Interaction between plant genotype and the symbiosis with *Epichloë* fungal endophytes in seeds of red fescue (*Festuca rubra*). *Crop Pasture Sci.* 62, 1010–1016. doi: 10.1071/CP11300
- Hamilton, C. E., and Bauerle, T. L. (2012). A new currency for mutualism: *Neotyphodium* antioxidants and host drought response. *Fungal Divers.* 54, 39–49. doi: 10.1007/s13225-012-0156-y
- Hamilton, C. E., Gundel, P. E., Helander, M., and Saikkonen, K. (2012). Endophytic mediation of reactive oxygen species and antioxidant activity in plants: a review. *Fungal Divers.* 54, 1–10. doi: 10.1007/s13225-012-0158-9
- Helander, M., Phillips, T., Faeth, S. H., Bush, L. P., and McCulley, R. (2016). Alkaloid quantities in endophyte-infected tall fescue are affected by the plant-fungus combination and environment. *J. Chem. Ecol.* 42, 118–126. doi: 10.1007/s10886-016-0667-1
- Hill, N. S., Rottinghaus, G. E., Agee, C. S., and Schultz, L. M. (1993). Simplified sample preparation for HPLC analysis of ergovaline in tall fescue. *Crop Sci.* 33, 331–333. doi: 10.2135/cropsci1993.0011183X003300020024x
- Huang, W.-Y., Cai, Y.-Z., Xing, J., Corke, H., and Sun, M. (2007). A potential antioxidant resource: endophytic fungi from medicinal plants. *Econ. Bot.* 61, 14–30. doi: 10.1663/0013-0001(2007)61[14:APAREF]2.0.CO;2
- Jackson, J. A. Jr., Hemken, R. W., Boling, J. A., Harmon, R. J., Buckner, R. C., et al. (1984). Loline alkaloids in tall fescue hay and seed and their relationship to summer fescue toxicosis in cattle. *J. Dairy Sci.* 67, 104–109. doi: 10.3168/jds.S0022-0302(84)81272-2
- Jensen, A. M. D., Mikkelsen, L., and Roulund, N. (2007). Variation in genetic markers and ergovaline production in endophyte (*Neotyphodium*)-infected fescue species collected in Italy, Spain, and Denmark. *Crop Sci.* 47, 139–147. doi: 10.2135/cropsci2005.10.0352
- Justus, M., Witte, L., and Hartmann, T. (1997). Levels and tissue distribution of loline alkaloids in endophyte-infected *Festuca pratensis*. *Phytochemistry* 44, 51–57. doi: 10.1016/S0031-9422(96)00535-3
- Kiers, E. T., and West, S. A. (2015). Evolving new organisms via symbiosis. *Science* 348, 392–394. doi: 10.1126/science.aaa9605
- Kranner, I., Minibayeva, F. V., Beckett, R. P., and Seal, C. E. (2010). What is stress? Concepts, definitions and applications in seed science. *New Phytol.* 188, 655–673. doi: 10.1111/j.1469-8137.2010.03461.x
- Leuchtmann, A., Schmidt, D., and Bush, L. P. (2000). Different levels of protective alkaloids in grasses with stroma-forming and seed-transmitted *Epichloë/Neotyphodium* endophytes. *J. Chem. Ecol.* 26, 1025–1036. doi: 10.1023/A:1005489032025
- Lewis, G. C., and Clements, R. O. (1986). A survey of ryegrass endophyte (*Acremonium loliae*) in the U.K. and its apparent ineffectuality on a seedling pest. *J. Agric. Sci.* 107, 633–638. doi: 10.1017/S002185960006980X
- Mace, W. J., Lunn, K. L., Kaur, N., and Lloyd-West, C. M. (2014). Variation in the expression of ergot alkaloids between individual tillers of perennial ryegrass. *Front. Chem.* 2, 1–6. doi: 10.3389/fchem.2014.00107
- Menéndez, A. I., A. M. Romero, A. M. Folcia, and Martínez-Ghersa, M. A. (2009). Getting the interactions right: will higher O₃ levels interfere with induced defense to aphid feeding? *Basic Appl. Ecol.* 10, 255–264. doi: 10.1016/j.baee.2008.03.010
- Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., and Core Team (2017). *nlme: Linear and Nonlinear Mixed Effects Models. R package version 3.1-131*. Available online at: <https://CRAN.R-project.org/package=nlme>
- Potter, D. A., Stokes, J. T., Redmond, C. T., Schardl, C. L., and Panaccione, D. G. (2008). Contribution of ergot alkaloids to suppression of a grass-feeding caterpillar assessed with gene knockout endophytes in perennial ryegrass. *Entomol. Exp. Appl.* 126, 138–147. doi: 10.1111/j.1570-7458.2007.00650.x
- Rasmussen, S., Parsons, A. J., Bassett, S., Christensen, M. J., Hume, D. E., Johnson, L. J., et al. (2007). High nitrogen supply and carbohydrate content reduce fungal endophyte and alkaloid concentration in *Lolium perenne*. *New Phytol.* 173, 787–797. doi: 10.1111/j.1469-8137.2006.01960.x
- R. Core Team (2017). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna. Available online at: <https://www.R-project.org/>
- Saikkonen, K., Gundel, P. E., and Helander, M. (2013). Chemical ecology mediated by fungal endophytes in grasses. *J. Chem. Ecol.* 39, 962–968. doi: 10.1007/s10886-013-0310-3
- Schardl, C. L., Grossman, R. B., Nagabhyru, P., Faulkner, J. R., and Mallik, U. P. (2007). Loline alkaloids: currencies of mutualism. *Phytochemistry* 68, 980–996. doi: 10.1016/j.phytochem.2007.01.010
- Schardl, C. L., Young, C. A., Pan, J., Florea, S., Takach, J. E., Panaccione, D. G., et al. (2013). Currencies of mutualisms: sources of alkaloid genes in vertically transmitted *Epichloae*. *Toxins* 5, 1064–1088. doi: 10.3390/toxins5061064
- Seal, C. E., Zammit, R., Scott, P., Flowers, T. J., and Kranner, I. (2010). Glutathione half-cell reduction potential and α -tocopherol as viability markers during the prolonged storage of *Suaeda maritima* seeds. *Seed Sci. Res.* 20, 47–53. doi: 10.1017/S0960258509990250
- Stewart, A. V. (1985). Perennial ryegrass seedling resistance to Argentine stem weevil. *N. Zeal. J. Agr. Res.* 28, 403–407. doi: 10.1080/00288233.1985.10430445
- Sullivan, T. J., Rodstrom, J., Vandop, J., Librizzi, J., Graham, C., Schardl, C. L., et al. (2007). Symbiont-mediated changes in *Lolium arundinaceum* inducible defenses: Evidence from changes in gene expression and leaf composition expression and leaf composition. *New Phytol.* 176, 673–679. doi: 10.1111/j.1469-8137.2007.02201.x
- TePaske, M. R., Powell, R. G., and Clement, S. L. (1993). Analyses of selected endophyte-infected grasses for the presence of loline-type and ergot-type alkaloids. *J. Agr. Food Chem.* 41, 2299–2303. doi: 10.1021/jf00036a015
- Uchitel, A., Omacini, M., and Chaneton, E. J. (2011). Inherited fungal symbionts enhance establishment of an invasive annual grass across successional habitats. *Oecologia* 165, 465–475. doi: 10.1007/s00442-010-1740-z
- Vázquez-de-Aldana, B. R., García-Ciudad, A., García-Criado, B., Vicente-Tavera, S., and Zabalgoagezcoa, I. (2013). Fungal endophyte (*Epichloë festucae*) alters the nutrient content of *Festuca rubra* regardless of water availability. *PLoS ONE* 8:e84539. doi: 10.1371/journal.pone.0084539
- Vázquez-de-Aldana, B. R., Zabalgoagezcoa, I., García-Ciudad, A., and García-Criado, B. (2003). Ergovaline occurrence in grasses infected by fungal endophytes of semi-arid pastures in Spain. *J. Sci. Food Agric.* 83, 347–353. doi: 10.1002/jsfa.1319
- Vázquez-de-Aldana, B. R., Zabalgoagezcoa, I., García-Ciudad, A., and García-Criado, B. (2007). Fungal alkaloids in populations of endophyte-infected *Festuca rubra* subsp. *pruinosa*. *Grass Forage Sci.* 62, 364–371. doi: 10.1111/j.1365-2494.2007.00591.x
- Vázquez-de-Aldana, B. R., Zabalgoagezcoa, I., Rubio-de-Casas, R., García-Ciudad, A., and García-Criado, B. (2010). Relationships between the genetic distance of *Epichloë festucae* isolates and the ergovaline and peramine contents of their *Festuca rubra* hosts. *Ann. Appl. Biol.* 156, 51–61. doi: 10.1111/j.1744-7348.2009.00360.x
- Walters, C. (1998). Understanding the mechanisms and kinetics of seed aging. *Seed Sci. Res.* 8, 223–244. doi: 10.1017/S096025850000413X
- White, J. F. Jr., and Torres, M. S. (2010). Is plant endophyte-mediated defensive mutualism the result of oxidative stress protection? *Physiol. Plantarum* 138, 440–446. doi: 10.1111/j.1399-3054.2009.01332.x
- Yue, Q., Jonson-Cicalese, J., Gianfagna, T. J., and Meyer, W. A. (2000). Alkaloid production and chinch bug resistance in endophyte-inoculated chewings and strong creeping red fescues. *J. Chem. Ecol.* 26, 279–292. doi: 10.1023/A:1005410115940
- Zabalgoagezcoa, I., García-Ciudad, A., Vázquez-de-Aldana, B. R., and García-Criado, B. (2006). Effects of the infection by the fungal endophyte *Epichloë festucae* in the growth and nutrient content of *Festuca rubra*. *Eur. J. Agron.* 24, 374–384. doi: 10.1016/j.eja.2006.01.003

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Gundel, Seal, Biganzoli, Molina-Montenegro, Vázquez-de-Aldana, Zabalgoagezcoa, Bush, Martínez-Ghersa and Ghersa. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.