



2021

## Hearing Health Healthcare Disparities in Appalachia

Matthew L. Bush

*University of Kentucky*, [matthew.bush@uky.edu](mailto:matthew.bush@uky.edu)

Follow this and additional works at: <https://uknowledge.uky.edu/jah>



Part of the [Medicine and Health Sciences Commons](#)

### Recommended Citation

Bush M. Hearing Health Healthcare Disparities in Appalachia. *J Appalach Health* 2021;3(4):5–10. DOI: <https://doi.org/10.13023/jah.0304.02>

Copyright © 2021 Matthew L. Bush

This Commentary is brought to you for free and open access by the College of Public Health at the University of Kentucky.

---

## Hearing Health Healthcare Disparities in Appalachia

### Abstract

Hearing loss is a global public health issue with disproportionate negative impacts on those who live in rural regions, such as Appalachia. This commentary provides an overview of hearing health and healthcare disparities in rural regions along with discussion of the significance of recent research findings which highlight the incidence of hearing loss and the shortage of hearing specialists in Appalachia.

### Keywords

Appalachia, hearing loss, audiologists, rural health

### Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

### Cover Page Footnote

No competing financial or editorial interests were reported by the authors of this editorial.

Hearing loss is a major global public health problem. According to the 2021 World Report on Hearing, approximately 1.5 billion people, including adults and children, worldwide have hearing loss.<sup>1</sup> It currently represents the third most prevalent chronic health condition among adults within the U.S.<sup>2</sup> Children with hearing loss, either of congenital or acquired causes, are at risk for delayed language development.<sup>3</sup> Undiagnosed and untreated hearing loss causes a negative impact on the social, occupational, cognitive, and emotional well-being of those affected.<sup>4,5</sup> There is a significant global financial impact, as well, with nearly \$980 billion in yearly economic losses due to hearing loss.<sup>1</sup> Unfortunately, as is the case with many chronic diseases, the global trends in hearing loss indicate that the greatest prevalence and burden of disease occurs in regions with the least accessible healthcare. The research presented in this issue of the *Journal of Appalachian Health* by Pudrith et al.<sup>6</sup> illustrates that trend by focusing our attention on the hearing health and healthcare status of the Appalachian region of the U.S., which has suffered from pervasive and persistent health disparities.

Hearing loss is medical condition that involves multi-specialty hearing healthcare specialists such as otolaryngologists and audiologists and may be managed with a variety of medical, surgical, and technological treatments. The lack of hearing healthcare access, utilization, and affordability continues to play a major role in preventing timely diagnosis and treatment of hearing loss. The inaccessibility and under-utilization of hearing healthcare is amplified in vulnerable populations, such as rural adults. Approximately 20% of the US population reside in rural areas and rural Americans tend to be poorer, less educated, suffer higher rates of disease and disability, and are more likely to be uninsured than their urban counterparts.<sup>7,8</sup> Rural adults face a greater risk of death from major diseases; however, these rural residents may have the most limited access to and poorest utilization of healthcare.<sup>9</sup> Systemic factors such as lack of public transportation, lack of telecommunications access, fewer community services, and a shortage of healthcare providers contribute to sub-optimal health among rural Americans.<sup>8</sup> Hearing loss is more prevalent among adults living in rural communities compared with those in urban settings.<sup>10,11</sup> Awareness of benefits from diagnosis of hearing loss and treatment is low among the rural population and among healthcare professionals.<sup>12,13</sup> Screening for hearing loss among adults is uncommon in the primary care setting and there is poor adherence to recommended treatment.<sup>14,15</sup> There is also a critical shortage of hearing healthcare specialists within rural areas.<sup>16</sup> While these features are common in much of rural America, rural Appalachia possesses an extreme

version of these health disparity and care inequity characteristics.<sup>8</sup> Appalachia is an expansive and diverse region encompassing over 25 million people throughout 13 states.<sup>17</sup> Across the lifespan, rural Appalachian adults with hearing loss have faced delayed diagnosis and treatment, lack of treatment, and poorer adherence/utilization of evidence-based treatments.<sup>18,19</sup>

Against this backdrop of hearing loss significance and long-standing rural hearing healthcare disparities, this manuscript presents novel findings regarding hearing loss prevalence and audiological disparities in Appalachia. Much of what we know about rural hearing loss prevalence and accessibility to hearing healthcare specialists in rural regions is based on work outside the United States or in non-Appalachian regions of the U.S. This manuscript utilizes a combination of population databases to conduct their univariate and multivariate analysis comparing Appalachian regions with non-Appalachian regions. The authors have quantified the higher incidence of subjective hearing loss among Appalachian residents and through their multiple linear regression analyses have presented data that demonstrates that Appalachian residence, apart from rural residence, is independently associated with a higher incidence of hearing loss. Furthermore, their pairwise linear regression data provides us with a deeper understanding of the disparity at hand by quantifying that relationship between rurality, prevalence of hearing loss, and the shortage of audiologists. While we neither understand the causes underlying this health disparity nor the solution to this problem, this study gives us clearly articulated evidence that helps us understand this disparity better. This data pushes us to consider how specialty care can be delivered to vulnerable populations in remote locations. Considerable innovation is needed to address rural and Appalachian hearing health and healthcare disparities. New models and methods for care delivery are needed to explore how to deliver specialty healthcare when there are no or very few specialists nearby. Research related to hearing loss may be less familiar for the readers of this journal; however, I encourage readers to consider the importance of work such as this to define and describe a complex health disparity issue within a resource-limited region such as Appalachia.

## REFERENCES

1. World Report on Hearing. 2021. World Health Organization. <https://www.who.int/activities/highlighting-priorities-for-ear-and-hearing-care> Accessed 10/4/2021.
2. Lin FR, Niparko JK, Ferrucci L. Hearing loss prevalence in the United States. *Arch Intern Med*. 2011 Nov 14;171(20):1851–2. PMID: 22083573 PMCID: PMC3564588
3. Erenberg A, Lemons J, Sia C, Trunkel D, Ziring P. Newborn and infant hearing loss: detection and intervention. American Academy of Pediatrics. Task Force on Newborn and Infant Hearing, 1998– 1999. *Pediatrics*. 1999;103(2):527–30.
4. Jung D, Bhattacharyya N. Association of Hearing Loss with Decreased Employment and Income Among Adults in the United States. *Ann of Otol Rhinol Laryngol* 2012; 121(12): 771–5. PMID: 23342548
5. Li CM, Zhang X, Hoffman HJ, Cotch MF, Theman CL, Wilson MR. Hearing Impairment Associated with Depression in US Adults, National Health and Nutrition Examination Survey 2005–2010. *JAMA Otolaryngol Head Neck Surg*. 2014; 140(4): 293–302. PMID: 24604103 PMCID: PMC4102382
6. Pudrith C, Grider E, Kitner B. Perceived Hearing Loss and Availability of Audiologists in Appalachia. *J Appalach Health* 2021;3(4):29–44. DOI: <https://doi.org/10.13023/jah.0304.04>
7. US Congress, Office of Technology Assessment. Healthcare in Rural America. Washington DC: Government Printing Office, 1990. (Office of Technology Assessment Pub. No. OTA-H-434.)
8. Friedell G, Rubio A, Maretzki A, Garland B, Brown P, Crane M, Hickman P. Community cancer control in a rural, underserved population: the Appalachia Leadership Initiative on Cancer Project. *Journal of Health Care for the Poor and Underserved*. 2010;12:5–19.
9. About Rural Health. Rural Health. Centers for Disease Control and Prevention. <https://www.cdc.gov/ruralhealth/about.html>. Accessed 10/4/2021.
10. Merchant JA, Stromquist AM, Kelly KM, et al. Chronic disease and injury in an agricultural county: The Keokuk County Rural Health Cohort Study. *J Rural Health*. 2002 Fall; 18(4):521–35. PMID: 12380895
11. Cruickshanks KJ, Wiley TL, Tweed TS, Klein BE, et al. Prevalence of hearing loss in older adults in Beaver Dam, Wisconsin. The Epidemiology of Hearing Loss Study. *Am J Epidemiol*. 1998 Nov 1; 148(9):879–86. PMID: 9801018

12. Bush M, Alexander D, Noblitt B, Lester C, Shinn J. Pediatric Hearing Healthcare in Kentucky's Appalachian Primary Care Setting. *J Community Health*. 2015. 40(4):762–8. PMID: 25672888 PMCID: PMC4490996
13. Bush M, Hardin B, Rayle C, Lester C, Studts C, Shinn J. Rural Barriers to Early Diagnosis and Treatment of Infant Hearing Loss in Appalachia. *Otology & Neurotology* 2015. 36(1): 93–8. PMID: 25325844 PMCID: PMC4268139
14. Cohen SM, Labadie RF, Haynes DS. Primary care approach to hearing loss: the hidden disability. *Ear Nose Throat J*. 2005; 84(1):26, 29–31, 44. PMID: 15724769
15. Chou R, Dana T, Bougatsos C, Fleming C, Beil T. Screening adults aged 50 years or older for hearing loss: a review of the evidence for the US Preventive Services Task Force. *Ann Intern Med*. 2011; 154(5):347–55. PMID: 21357912
16. Sebastian S. American Academy of Audiology Director of Membership. Personal communication. July 15, 2015.
17. About The Appalachian Region. Appalachian Regional Commission. <https://www.arc.gov/about-the-appalachian-region/> Accessed 10/4/21.
18. Chan S, Hixon B, Adkins M, Shinn JB, Bush ML. Rurality and determinants of hearing healthcare in adult hearing aid recipients. *The Laryngoscope*. 2017;127(10):2362–7.
19. Hixon B, Chan S, Adkins M, Shinn JB, Bush ML. Timing and Impact of Hearing Healthcare in Adult Cochlear Implant Recipients. *Otology & Neurotology*. 2016;37(9):1320–4.