University of Kentucky
UKnowledge

Biosystems and Agricultural Engineering
Faculty Publications

Biosystems and Agricultural Engineering

7-9-2013

The Use of Co-Culturing in Solid Substrate Cultivation and Possible Solutions to Scientific Challenges

Wanying Yao
University of Kentucky

Sue E. Nokes
University of Kentucky, sue.nokes@uky.edu

Follow this and additional works at: https://uknowledge.uky.edu/bae_facpub

Part of the [Bioresource and Agricultural Engineering Commons](#), and the [Biotechnology Commons](#)
Right click to open a feedback form in a new tab to let us know how this document benefits you.

Repository Citation

Yao, Wanying and Nokes, Sue E., "The Use of Co-Culturing in Solid Substrate Cultivation and Possible Solutions to Scientific Challenges" (2013). *Biosystems and Agricultural Engineering Faculty Publications*. 169.
https://uknowledge.uky.edu/bae_facpub/169

This Article is brought to you for free and open access by the Biosystems and Agricultural Engineering at UKnowledge. It has been accepted for inclusion in Biosystems and Agricultural Engineering Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

The Use of Co-Culturing in Solid Substrate Cultivation and Possible Solutions to Scientific Challenges

Digital Object Identifier (DOI)

<https://doi.org/10.1002/bbb.1389>

Notes/Citation Information

Published in *Biofuels, Bioproducts and Biorefining*, v. 7, issue 4, p. 361-372.

Copyright © 2013 Society of Chemical Industry and John Wiley & Sons, Ltd

This is the peer reviewed version of the following article: Yao, W., & Nokes, S. E. (2013). The use of co-culturing in solid substrate cultivation and possible solutions to scientific challenges. *Biofuels, Bioproducts and Biorefining*, 7(4), 361-372. doi: 10.1002/bbb.1389, which has been published in final form at <https://doi.org/10.1002/bbb.1389>. This article may be used for non-commercial purposes in accordance with [Wiley Terms and Conditions for Use of Self-Archived Versions](#).

Perspective: The Use of Co-culturing in Solid Substrate Cultivation and Possible Solutions to Scientific Challenges

*Wanying Yao^a, *Sue E. Nokes^a*

^aDepartment of Biosystem and Agricultural Engineering, University of Kentucky
128 C.E. Barnhart Building
Lexington, KY, US
E-mail: Julia.yao@uky.edu; snokes@uky.edu

Abstract

This perspective systematically summarizes the use of solid substrate co-cultures in agriculture, food, plant and industrial biotechnology applications. The summarization is organized by organism, i.e. fungus, bacteria, yeast and then co-cultivation of either two or three organisms. Generally, in solid substrate co-culture, the organisms synergistically penetrate and degrade the solid substrate thereby increasing product yield and productivity over a monoculture. Efforts to increase co-culture performance include optimizing process parameters (pH, temperature, moisture and oxygen demand) and defining the acceptable types of substrate. Scientific challenges exist in understanding the interactions between microbial strains, such as viability, suite of products, and bio-transformations. The perspective details possible solutions to these challenges and highlights future research directions for co-cultures using either solid or liquid fermentation.

Key words: solid substrate cultivation, co-cultural microorganisms, lignocellulosic biomass, bioproducts and biofuels

1. Background

Currently, industrial demand for bioproducts is predominately being met using submerged fermentation (SmF) processes, generally employing modified strains of microorganisms.

However the cost of production in SmF systems is prohibitive for many commodity chemicals. Production cost can be reduced by deploying alternative methods such as solid substrate cultivation (SSC) systems.¹⁻³

Microbial consortia consisting of two or more microorganisms are responsible for many bio-transformations in natural environments.⁴ Interactions between the organisms can be triggered either through direct cell-to-cell communications or by signal substances that can bind to cell-surface proteins in the fermentation broth.⁵ In other words a stable co-culture could be maintained by the interactions of syntrophic relationships, competition for substrates as well as growth promoters or inhibitors such as antibiotics.^{5, 6} For example, one of the syntrophic interactions in co-cultures is the widely distributed phenomenon of interspecies hydrogen transfer between sulfate reducers such as *Syntrophobacter* sp. and methanogens, where carbon dioxide is reduced into methane by obtaining the energy from hydrogen.^{7, 8}

Currently, co-culture SmF fermentations (mostly positive interactions) are widely used in the production of antibiotics, enzymes, fermented food, composting and bioconversion of wastewater sludge.⁹ Although not as widely applied as co-cultures in SmF, the advantages of co-cultures may be even more pronounced in SSC, because the colonization, penetration, and degradation of the solid substrate depends on a symbiotic association where each species can synergistically function with the other specie(s) in a substrate niche via a specific set of enzymes.¹⁰

2. Fungal co-culture under SSC

Filamentous fungi typically grow in symbiotic associations on solid substrates, such as wood, seeds, stems, roots and leaves of plants.¹¹ Similar to natural microbial processes, cultivation of mixed species of fungi can demonstrate more effective utilization of the substrate, increased productivity of the desired metabolites owing to comprehensive enzyme mixtures and higher resistance to contamination.^{10, 12} In addition, fungal solid substrate cultivation is a low-cost fermentation process particularly suited to the needs in developing pilot scale processes, or producing high volume, low cost commodity chemicals.¹³

2.1 Production of lignocellulolytic enzymes and enzymes mixtures

The composition of lignocellulosic waste/biomass is complex and therefore requires multiple enzymes for decomposition.¹⁰ Ligninolytic, cellulolytic, pectinolytic and xylanolytic enzymes are the main hydrolytic enzyme systems that either pretreat biomass or generate reducing sugars. Ligninolytic enzymes (lignin peroxidase, manganese peroxidase and laccase), directly and specifically attack lignin and oxidize it, often rendering the lignin water soluble, which enable the utilization of this complex biopolymer for the production of fine chemicals.¹⁴ A cellulolytic enzyme system hydrolyzes lignocellulosic biomass synergistically using endoglucanase, exoglucanase and β -glucosidase.¹⁵ Xylanase complements the cellulolytic enzyme system as xylanase is needed to elicit complete and efficient hydrolysis of the lignocellulosic biomass, which contains an appreciable amount of hemicellulose in both hardwood and herbaceous biomass – composed of xylan predominately.¹⁶

The association of two or more fungi results in higher enzyme production, which by acting synergistically, effectively degrades various biomass or agricultural wastes. Among the cellulolytic fungi, genera like *Trichoderma*, *Aspergillus*, *Penicillium* and *Fusarium* are regarded as cellulase producers.¹⁷ The feasibility and performance of mixed cultures (either of different species or genera) using industrial and agricultural carbon wastes were investigated and enhanced by optimizing incubation time, carbon sources and initial pH of the fermentation medium¹⁸⁻²¹ and are specified below.

Aspergillus is by far the most efficient producer of β -glucosidase among the microorganisms investigated in terms of high, active and stable enzymatic properties maintained at the existence of inhibitors and broad pH range.^{9, 22}

Strains of *Trichoderma* can accumulate high activities of endo and exo-glucanase, but are deficient in β -glucosidase whereas some strains of *Aspergillus* have high β -glucosidase activity.²³⁻²⁵ Therefore, strains of both *Trichoderma* and *Aspergillus* can be successfully cultivated together in SSC to produce multi-complex cellulase systems as reported in Table 1 (see Supplement for detailed discussion of studies).^{16, 19, 20, 26, 27} Besides the improvement in the activity of individual enzyme system, co-culturing two or more fungal strains using SSC can also produce multiple enzyme complexes.^{10, 28-30}

As evidenced in Table 1, fungal co-cultures regulated the complete utilization of the complex biomass by producing a variety of different enzymes complexes. Even for the same species, different strain combinations can correspond to different enzyme complexes. The informed selection of compatible microorganisms in mixed fungal systems can correspond to the desired enzymatic levels and compositions in order to accomplish different objectives. For a selection of an optimum fungal system as related to strain compatibility, a direct target is the high yield of enzyme activities. Besides the activity of enzyme, enzyme volumetric productivities (the amount of enzyme produced per m³ bioreactor volume per unit time) can be used as a criteria to evaluate the performance of co-cultures as compared to mono culture.

Table 1 List of fungal co-cultures for hydrolytic enzymes

Product	Substrate	Increased amount as comparisons to monoculture	Co-culturing strains	References
Xylanase and laccase	Mixed wood pulp	Not specified	Mutant <i>Penicillium oxalicum</i> SAUE-3.510 and <i>Pleurotus ostreatus</i> MTCC 1804 (ratio not specified)	26
Mixture of Laccase, and glucoamylase or β -glucosidase	Wheat bran	1.5-fold increase in β -glucosidase over that of <i>T. aurantiacus</i> ; 2.5-fold increase in laccase over that of <i>A. niger</i> , A; 1.68-fold increase in glucoamylase over that of <i>T. aurantiacus</i>	<i>T. aurantiacus</i> with <i>A. niger</i> A or B (ratio not specified)	27
Laccase and pectinase	Wheat bran, oats straw and beetroot press	8.4-fold increase in laccase over the mono culture	<i>A. niger</i> and <i>Fusarium moniliforme</i> ; <i>Trametes versicolor</i> and <i>A. niger</i>	10
Xylanase and cellulase	Sugarcane bagasse	35 - 45% increase in xylanase and 20-142% increase in cellulase compared to single cultures of either partner	<i>T. reesei</i> with either <i>A. niger</i> or <i>A. phoenicis</i> (ratio 3:2)	28
Cellulase complex	Waste paper	Up to 2.4-fold increase in cellulase over mono culture of <i>A. niger</i> ; 3.7-fold increase in cellulase over <i>T. viride</i>	<i>A. niger</i> and <i>T. viride</i> (ratio 1:1)	29
β -glucosidase	Defined medium	More than 10-fold rise than that of <i>A. niger</i> or <i>A. oryzae</i>	<i>A. niger</i> and <i>A. oryzae</i> (ratio 3:1)	9

cellulase (β -glucosidase and endocellulase), xylanase	Soybean hulls supplemented with wheat bran	65% increase in β -glucosidase and 67.3% increase in endocellulase (100 IU/g) compared to that of <i>T. reesei</i> (60 IU/g)	<i>T. reesei</i> and <i>A. oryzae</i> (ratio 1:1)	16
Cellulase, hemicellulase, glucoamylase, pectinase, and acidic proteinase	Bran mash	At least 65% activity increase for all the tested enzymes compared to each mono culture.	Species of <i>A. niger</i> (ratio 1:4)	30
Cellulase complex	Sugar cane bagasse	1.4-fold increase in β -glucosidase and 2.9-fold increase in endoglucanase compared to that of <i>T. reesei</i> LM-UC4	<i>T. reesei</i> LM-UC4 and <i>A. phoenicu</i> (ratio 1:1)	19
Cellulase complex	Bagasse	4.3-fold increase in β -glucosidase; 7.3-fold increase in endoglucanase over single culture of <i>T. reesei</i> in flask culture	<i>T. reesei</i> LM-UC4 and <i>A. phoenicis</i> QM329 (ratio 2:1)	20
Cellulase and hemicellulase	Tomato pomace	Up to 1.8-fold increase in cellulose conversion over mono culture of <i>T. reesei</i>	<i>Trichoderma</i> and <i>Sporotrichum</i> sp	21

2.2 Production of tannase and gallic acid

Tannin acyl hydrolase, commonly called tannase, has been widely used in the food and pharmaceutical industries. Due to its hydrolysis of gallotannin to gallic acid, this enzyme is mainly commercially applied to the synthesis of an antifolate and antibacterial drug trimethoprim.³¹

The following microorganisms; Fungi-*Aspergillus sp.*, *Penicillium sp.*, *Rhizopus sp.*, Yeast-*Candida sp.* and bacteria - *Bacillus sp.* have been used as inoculum for tannase and gallic acid production from tannin-rich substrates using SSC.^{32, 33} Various fungal combinations have been investigated successfully, as summarized in Table 2 (see Supplement for detailed discussion of studies).

Table 2 List of fungal co-cultures for tannase and gallic acid production

Product	Substrate	Increased amount over mono-culture	Selected co-culturing strains	References
Tannase and gallic acid	Grape waste	33.3% to 37.7% increase in enzyme volumetric productivity (U/g/min) over single cultures of either partner	<i>P. chrysogenum</i> and <i>T. viride</i> (ratio 1:1)	34
Tannase	Tamarind seed powder	35.1% to 57.4% increase in enzyme volumetric productivity (U/g/day) over single cultures of either partner	<i>R. oryzae</i> (MTCC 1987) and <i>A. foetidus</i> (MTCC 3557) (ratio 1:1)	35
Tannase	Fruits of Terminalia chebula (myrobalan) and Caesalpinia digyna cover	Not specified	<i>R. oryzae</i> (NRRL 21498) and <i>A. foetidus</i> (MTCC 3557) (ratio 1:1)	36
Gallic acid	Fruits of Terminalia chebula (containing about 32% tannin) and Caesalpinia digyna cover powder (45% tannin)	29.8% increase in enzyme activity over <i>R. oryzae</i> ; 13.4% increase in enzyme activity over <i>A. foetidus</i>	<i>R. oryzae</i> (NRRL 21498) and <i>A. foetidus</i> (MTCC 3557) (ratio 1:1)	37

2.3 Other bioproducts produced from fungal co-cultures

Fungal co-cultures can produce unique enzymatic complex synergistically transforming raw materials or agricultural residues into high protein biomass or secondary metabolites ³⁸, which is attractive in comparison with monoculture as demonstrated in Table 3 (see Supplement for detailed discussion of studies).

Other than fungal mixed cultures, co-cultivation of yeast, bacteria and fungi have also been used to enhance the nutrient availability for animal feed under SSC, which will be discussed in section 3 of this perspective.

Table 3 List of fungal co-cultures for producing other value-added products

Product	Substrate	Benefits or increased amount over mono cultures	Co-culturing strains	References
Protein enrichment for animal feeds	Cassava lees	Increase of protein from 1.42% to 16.08% and 18.54%	<i>A. niger</i> and <i>Geotrichum sp</i> <i>A. niger</i> and <i>Penicillium sp.</i> (ratio not specified)	39
	Rapeseed meal	Decrease of glucosinolates by 90.71%	<i>A. oryzae</i> and <i>T. viride</i> (ratio 1:1)	40
	Sweet potato residue	10.2% to 24.1% increase in protein content	<i>Aspergillus sp.</i> and <i>Rhizopus sp.</i> (ratio 1:1)	41
Lovastatin	Red mold rice	Up to 64.7% increment in lovastatin yield over the mono culture of <i>Monascus ruber</i> in literature ⁴²	<i>Monascus purpureus</i> and <i>Monascus ruber</i> (ratio 1:1)	43
Decolorization of textile dye	Coomassie brilliant blue, bromophenol blue and malachite green	-15.6% to -21.8% degradation efficiency	<i>Trichoderma sp.</i> and <i>A. flavus</i> (ratio not specified)	44

3. Bacterial co-cultures

SSC has been used as a principal technology to produce biological pest control agents for crops, to prepare a fertilizer or to improve the quality and flavor of the product in traditional food fermentation in which bacterial co-cultures provided the biotransformation. The corresponding literature is listed in Table 4 (see Supplement for detailed discussion of studies).

Table 4 List of bacterial co-cultures on SSC

Product	Substrate	Advantages or increased amount over single culture	Co-culturing strains	References
Biopesticide	Kitchen waste	Decreased oil content thus facilitating biopesticide production	<i>Bacillus thuringiensis</i> and <i>Bacillus subtilis</i> (ratio 3:1-1:3)	45
Bioactive compounds	Carrot pomace	10-fold increase in mucilage and 6.2-fold increase in fibrinolytic enzyme activity	<i>Bacillus subtilis</i> and <i>Leuconostoc mesenteroides</i>	46
Vitamin B ₁₂	Agriculture crop residues (sugar-cane bagasse, wheat straw, rice straw, bean straw and cotton stalks)	At least 20% increase in vitamin B ₁₂ over mono cultures	<i>Bacillus firmus</i> AZ-78B and <i>Streptomyces halstedii</i> AZ- 8A (ratio 2:1)	47
Fertilizer	Corn stover, soybean meal, bran	Shortened fermentation period from 72 h to 52 h and increased spore formation rate from 50 % to 90%	<i>Bacillus mucilaginosus</i> and <i>Bacillus megaterium</i> (ratio 2:1)	48
Lactic acid	Pine needles	At least 72% increase in lactic acid yield over mono Lactobacilli cultures	<i>Lactobacilli delbrueckii</i> (NCIM2025) and <i>Lactobacilli pentosus</i> (NCIM 2912) (ratio not specified)	49
Vinegar	Grains and bran	Improved product flavor and quality	<i>Acetobacter rancens huniang 1.01</i> , <i>Acetobacter pasteurianus</i> As 1.41, two <i>Gluconobaceter</i> isolates (ratio 2:1:3:2:2)	50

4. Co-cultures of bacteria, yeast and fungi

4.1 Co-cultures for protein enrichment and animal feed production

Microbial SSC conversion of agricultural byproducts or residues into cell protein has received much attention because it requires low level technology, increases protein availability in animal feed allowing direct use of the fermented product for feeding purposes as well as solving pollution and waste disposal problems.^{38, 51} Co-culturing two or more strains are favored for protein enrichment because the synergism of the strains improves the digestibility and nutrient value of agriculture residues designed for animal feed products. *Enterococcus faecium* and *Lactobacillus casei* are important bacteria for the health of the lower gastrointestinal tract. They assist in pH modulation thereby creating an optimum environment for endogenous enzymes to process feed efficiently. A further benefit is that these bacteria produce bacteriocins that inhibit the proliferation of *Escherichia*, *Salmonella* and *Clostridia*. *Bacillus subtilis* are aerobes that produce a variety of enzymes with good protease, amylase, lipase, esterase and some xylanase and cellulase activity. The enzymes produced by the *Bacillus* strains also help to improve feed efficiency and conversion. Additions of fungi, such as *Aspergillus* and *Rhizopus*, increase the accessibility of the lignocelluloses or starch materials (Table 3). Yeasts, notably *S. cerevisiae*, are important probiotic organisms that help modulate the pH of the caecum and colon, as well as stimulate the growth of beneficial fiber and lactic acid-utilizing bacteria.

Processing parameters, such as moisture, pH of the substrate, inoculum size, temperature, and nitrogen source have been optimized to achieve the best synergistic effect from the microorganisms as illustrated in Table 5 (see Supplement for detailed discussion of studies).

Besides optimizing the process conditions for co-cultures to improve protein content, evaluation of the solid state fermented feed on animal performance is a frequent topic in the literature. Published reports have demonstrated the positive performances of SSC on nutrient digestibility by increasing the amount of water-soluble protein content as well as a reduction in pathogens by the accumulation of acetic acid.^{52, 53}

Table 5 List of co-cultures of bacteria, yeast and fungi for protein enrichment on SSC

Substrate	Protein yield/ enhancement in comparisons with raw materials	Co-culturing strains	References
Rapeseed meal wheat bran, corn flour and rice bran with inorganic salts	12.8% crude protein	<i>Geotrichum candidum</i> and <i>Bacillus subtilis</i>	54
Rapeseed meal, wheat bran and brown sugar	36.6% crude protein	<i>Lactobacillus fermentum</i> , <i>Enterococcus faecium</i> , <i>S. cerevisiae</i> and <i>Bacillus subtilis</i> (ratio 1:1:1:1).	53
Soybean meal, wheat bran	28.5%, two-times higher than the raw material	<i>A. niger</i> , <i>T. viride</i> , <i>S. cerevisiae</i> , and <i>Candida utilis</i> (ratio 1:1:1:3)	55
Soybean meal, corn gluten meal, rapeseed meal and peanut meal	31% with 9.2% of water soluble protein	<i>Lactobacillus fermentum</i> , <i>S. cerevisiae</i> and <i>Bacillus subtilis</i> (ratio 1:1:1)	52
Sweet potato	Increased by 101% in amino acid composition over raw material	<i>R. oligosporus</i> and <i>Candida arborea</i>	56

4.2 Co-cultures for ethanol or alcoholic drink

Yeasts and bacteria co-cultures for ethanol fermentations, especially for co-fermenting glucose and xylose, have been widely investigated in submerged liquid cultivation⁵⁷⁻⁵⁹ while only a few reports on SSC co-cultures for ethanol have been published. Sree⁶⁰ used a thermotolerant *S. cerevisiae* strain and a local isolate of an amylolytic *Bacillus* sp. to produce fuel ethanol from sweet sorghum and sweet potato using SSC. More ethanol (42.3% increase) was produced at 37 °C by the co-culture than by the thermotolerant yeast alone, due to the increased availability of reducing sugars from *Bacillus* sp.. A sequential co-culture of fungi and yeast was used to produce a traditional alcoholic drink from steamed cereals using SSC.⁶¹ The process involves two steps using fungus (generally, *A. kawachii* or *A. oryzae*) for saccharifying the feedstock and a yeast (usually, *S. cerevisiae*) for ethanol production.⁶² Seventy percent of total carbohydrates were converted to ethanol after 50 days of cultivation.

5. Future aspects for solid substrate, liquid or submerged co-cultures

5.1 Considerations and techniques on establishment of co-culture system

The criteria considered to select the co-culture systems include: 1) high compatibility between co-culture partners with little or no metabolic inhibitions; 2) compromise of the differentiation in culturing parameters to the mutual culture conditions; 3) maintaining the culture stability so no outcompeting occurs (except sequential co culture); 4) improved or even complete utilization of complex substrate cocktail; and 5) improved product yield and rate as compared to mono culture. These criteria characterize an efficient and feasible co-culture system, which assures process success. Further optimization of the co-culture (including optimal ratio of co-culture partners, moisture content, temperature, time, pH, substrate, etc) will need to be deployed to improve the process efficiency with respect to product target, system stability and productivity. However, efficient screening and analytical techniques to predict the co-culture performance are not yet developed.

5.1.1 Screening of co-culture system - Selectivity and compatibility

When selecting microbial species for a co-culture system, the first step is to choose individual strains with the desired properties, then test their compatibility and study their co-fermentation

performance. Screening technologies for identifying more desirable strains for co-culture systems are still under development.⁵ Usually the screening tests are performed on the petri dish to observe the organisms' compatibility or inhibitions, but this can only be regarded as prescreening due to the limited growth environment. Other requirements for a stable co-culture would still need to be verified, such as metabolic interactions (i.e., growth promoter or inhibitors) and mutually adjustable cultivation conditions.⁶ Candidates identified from pre-screenings would need to be scaled up to eliminate uncertainty. Microbial growth rates, product tolerances, productivity and product yield of co-cultures also need to be quantified for selecting the optimum combination of microorganisms.

5.1.2 Evaluation of the stability of co-cultures

Culture stability is another trait vital to co-culture systems' success, yet one for which criteria have not yet been well established. Usually, the co-culture system is considered stable when the ratio of the live cells of the two organisms remains constant within the limits of experimental error, demonstrating that neither organism is outcompeting the other. The determination can be accomplished by the enumeration of the two microbial populations using the standard technique of a solid growth medium inoculated with serial dilutions of the broth cultures. However, this simple technique is only adapted to bacterial populations that can be differentiated on petri dish. Some researchers defined the criteria for structurally stable bacterial communities as one in which all the members persist over more than 20 times subculturing.⁶ But the adaptability of this criteria still needs to be explored and verified.^{6, 63} Nowadays, analytical techniques from molecular biology like denaturing gradient gel electrophoresis (DGGE) are used to investigate the dynamics of microorganism populations and fermentation characteristics of in vitro co-cultures.⁶⁴

Additionally, besides screening for the optimum combination of co-culture, selection of co-cultures which exhibit desired properties is another prospective method. Despite the advent of molecular biology and rational screening techniques, classical mutation still plays a major and irreplaceable role in improving industrial mono-cultures, and can also be an option for co-culture selection. To guarantee culture compatibility, cells or fungal spores can be blended together when exposed to the mutagenic agents instead of mutating the mono-culture separately. The

difficulty of using classical mutation on a co-culture is in determining which micro-organism was significantly improved; i.e. separating the effects of the mutation(s) on the two different organisms.

5.2 Solutions to determining the mechanism of the co-culture system

A viable co-culture system will exhibit positive interactions between the two or more partners. However, the interactions between microorganisms in co-culture environments may not always correspond to desirable consequences.⁵ Therefore, knowing the interactions between associated strains in a co-culture system is essential. Very little research has been conducted in this area, mainly because of the difficulties involved in working with complex systems containing multiple microorganisms. Therefore, co-culture fermentations offer significant opportunities for future research.

5.2.1 Current understanding on the co-cultural mechanism

As evidenced by the cases illustrated above, the advantage of co-culture is evident in SSC because the slow growth of colonization allows better equilibration of populations and may degrade the substrate better in symbiotic associations as each species having its own niche for growth and substrate degradation. It is understandable that strains with different enzyme complexes can be combined to produce a suite of enzymes capable of achieving synergistic degradation of complicated substrates. But in the case of mutual products enhancement of the SSC co-culture, the underlying mechanisms have not been well elucidated in the literature, such as how the metabolism is reinforced on the proteomics and genetic level; how intense the cooperation and mutual interdependence is. One possible assumption is that one organism utilizes the metabolites (e.g. precursors of vitamins or certain amino acids) excreted from a partner organism and profits from this support even if it could synthesize the respective compound on its own. In this case, the microbes under such conditions may display more refined types of interaction with others.⁶⁵ This kind of job-sharing among co cultures ascribes to every single organism only a limited task it has to fulfill and makes the whole process more efficient via saving biosynthetic energy.

5.2.2 Kinetic model

In theory, a kinetic model can be used instead of experimentation to find optimum operating conditions which will maximize profitability, by accurately simulating the dynamic properties of the system. It is therefore usually believed to be advantageous to develop an appropriate kinetic model to characterize the desired co-culture systems. However, due to the difficulty in separating the kinetic parameters of each strain in a co-culture system, few modeling studies have been done to describe co-culture system dynamics. One possible solution to resolving the dynamic analysis of each strain is to use a molecular technique to observe population dynamics, such as DGGE, but this technique would increase experimental expense. Another solution is to regard the co-culture system as a mono culture, which ignores the details of the interactions between co-cultures thus avoiding the difficulty in differentiating each strain in the co-cultures. A logistic model has been used to closely predict product formation rate, and the authors state the model was successful by treating a co-culture system as mono culture.⁶⁶

5.2.3 Metabolic network model

Another research tool which would allow better understanding of the mechanism of the co-culture system is metabolic network modeling. A metabolic network model can be used for predicting metabolic pathways, explaining phenotypic behavior or designing ideal microbial strains for industrial production.⁶⁷ To deal with the difficulty of discerning behavioral interactions between individual microbes, the manner in which products are exchanged between species can be inferred biochemically and genetically by developing a mutualism model for co-cultured strains which have sequenced genomes. For a sequenced genome, a metabolic network model can be constructed from the biochemical and physiological information available. A combined metabolic model could be developed on the basis of the genome-scale reconstructed metabolic network model for each individual strain in a co-culture system, which may illustrate the potential metabolic interactions between the two strains.⁵ Some publications have proven that reconstructed metabolic networks and stoichiometric models can also serve to capture growth parameters and community composition of simple bacterial communities besides predicting metabolic fluxes and growth phenotypes of single organisms.^{68, 69}

5.3 Scale up of the co-culture fermentation process

5.3.1 Cultivation mode investigation in SSC

The selection of fermentation configuration depends on the microbes used in the co-culture. Product yield improvement, inhibitor alleviation and efficient substrate utilization can be enhanced in liquid co-cultures by employing fed-batch or continuous modes.⁷⁰ The related techniques which work in a mono culture or liquid co-cultures, such as feeding strategy, feedback/feedforward control and scale up modeling, can be extended to the SSC co-cultivations if well-verified yield performance, compatibility and stability have been determined. However, limited research is available focused on control of cultivation modes of the SSC co-culture systems due to lack of empirical data. Continuous flushing systems for SSC has been illustrated for cellulose conversion using *Clostridium thermocellum* by Dharmagadda et al.⁷¹. Chromatography columns fitted with porous polyethylene discs at the bottom were periodically flushed with culture medium and the solid substrate was retained. The flushed cultures exhibited higher cellulose conversion (25%-47%) than non-flushed SSC (<8%). Other empirical approaches to designing the SSC process, like tray bioreactor (Koji-type processes)⁷² or deep bed bioreactors⁷³, and integrating these with downstream processing would be beneficial to support the investigation of the cultivation modes of SSC.

Another future research area will be to extend metabolic modeling to dynamic situations where SSC is performed in various cultivation modes, and the model can then assist with process control. The classical flux balance analysis method has been successfully extended for such dynamic situations in liquid cultures.⁷⁴ Dynamic flux balance models can be gained by integrating stoichiometric equations for intracellular metabolism with dynamic mass balances on main substrates and products under the assumption of rapid intracellular changes. The cellular growth rate and substrate uptake kinetics provide the connection between the intracellular and extracellular models. This connection can be used to systematically interpret the regulatory effects such as inhibition by growth by-products.⁷⁵

5.3.2 Downstream processing

Generally the design and cost of the downstream processing is dependent on process-related contaminants, including byproducts generated and remains of the host cells (proteins, nucleic acids) or the cell culture (components of the culture media used). Although in many cases, the

performances of the co-cultures with respect to product yields are doubtless higher than single culture, the insufficient record of process-related inhibitors and contaminants which accumulate during cultivation overshadowed the influences on the product recovery in the downstream process or affected the entire process economics. Robust and cost-effective purification processes need to be developed or traditional processes modified in order to improve process efficiency and economics for co-cultures.

5.4 Strategy of co-culture – an alternative to co-ferment the hydrolysates for biofuel production

One of the bottlenecks in the field of lignocellulosic biofuels is the limited utilization of the sugars in the hydrolysates by the microorganisms, which significantly influences the fermentation efficiency and processing costs. The lack of industrially robust microbes for co-fermentation of glucose and xylose has been a major technical barrier. Researchers have been trying to genetically modify microorganisms by creating pathways for xylose fermentation. Some genetically modified microorganisms have been engineered to ferment glucose and xylose for high ethanol yields (55-57 g/L), such as recombinant *E. coli*, *Zymomonas mobilis* (*Z. mobilis*), and *S. cerevisiae*.^{58, 76, 77} One remaining challenge is to control the sugar consumption rates so that one sugar does not end up being partially unconsumed even if the organism could consume the two or more sugars simultaneously.

Another approach is to co-culture microorganisms which vary in their sugar consumption for biofuel production. This appears to have advantages over single culture fermentations since co-culturing provides an opportunity to adjust microbial populations to the changing sugar concentrations. Additionally, the co-culture can achieve simultaneous sugar conversion of C5 and C6 sugars, maximizing substrate utilization rate and increasing ethanol yield. Some research focus has been placed on the utilization of co-culture for ethanol production by co-fermenting glucose and xylose from various biomass hydrolysates. The yeast genus *Saccharomyces* or respiratory-deficient mutant is preferably used as the glucose-fermenting strain with a xylose-fermenting strain such as *Pichia stipitis* used due to the compatible fermentation conditions of pH, temperature and oxygen. Other co-cultures (*E. coli*, *S. cerevisiae* or *Z. mobilis*) were also shown to simultaneously consume xylose and glucose with either independent or overlapping pathways.^{59, 78, 79}

However, due to the complexity and variety of the hydrolysates' composition (a function of the biomass itself and its corresponding hydrolysis methods), limited research has been conducted to improve the sugar utilization using co-cultures. For example, the sugars released from sugar beet pulp by enzymatic hydrolysis were mainly arabinose and glucose (15-18 %) instead of xylose and glucose.⁸⁰ To develop robust co-culture systems for biomass conversion, a systematic study of the substrate-selective uptake range of each single strain needs to be explored. Understanding the strain compatibility and stability of the co-culture system will allow us to control and predict process performance. The metabolic pathways of each strain could then be further optimized for co-culturing to optimize yield and productivity of a particular product.

5 Conclusion

The examples of solid state co-culture processes in this perspective demonstrate the importance of this technique in the production of animal feed, enzymes, fertilizer and the process of bioremediation. Moreover, increasing demand for using renewable resources as feedstock for fermentative biofuel production, combined with advances in biotechnology and innovative process developments invokes further interest in solid state co-culture. Much of the interest is generated because SSC offers the opportunity to efficiently use inexpensive substrates by allowing microorganisms to act synergistically via a suite of enzymes specifically adapted for effectiveness in this solid niche. However, further progress in the co-culture technique will require a deep understanding of the interactions between associated strains in a co-culture system as well as developing efficient evaluation tools to monitor the co-culture's activity.

Acknowledgments

The authors gratefully acknowledge the financial support of the United States Department of Agriculture National Institute for Food and Agriculture Biomass Research and Development Initiative Grant # 2011-10006-30363. The investigation reported in this paper (No. 13-05-042) is a part of a project of the Kentucky Agricultural Experiment Station and is published with the approval of the director.

References

1. Pandey A, Soccol CR and Mitchell D, New developments in solid state fermentation: I-bioprocesses and products. *Process Biochemistry* **35**: 1153-69 (2000).
2. Raghavarao K, Ranganathan TV and Karanth NG, Some engineering aspects of solid-state fermentation. *Biochemical Engineering Journal* **13**: 127-35 (2003).
3. Chinn MS, Nokes SE and Strobel HJ, Screening of thermophilic anaerobic bacteria for solid substrate cultivation on lignocellulosic substrates. *Biotechnology Progress* **22**: 53-59 (2006).
4. Peng X, Sun J, Iserentant D, Michiels C and Verachtert H, Flocculation and coflocculation of bacteria by yeasts. *Applied Microbiology and Biotechnology* **55**: 777-81 (2001).
5. Chen YL, Development and application of co-culture for ethanol production by co-fermentation of glucose and xylose: A systematic review. *Journal of Industrial Microbiology & Biotechnology* **38**: 581-97 (2011).
6. Maki M, Leung KT and Qin WS, The prospects of cellulase-producing bacteria for the bioconversion of lignocellulosic biomass. *International Journal of Biological Sciences* **5**: 500-16 (2009).
7. Sakai S, Imachi H, Sekiguchi Y, Ohashi A, Harada H and Kamagata Y, Isolation of key methanogens for global methane emission from rice paddy fields: A novel isolate affiliated with the clone cluster rice cluster i. *Applied and Environmental Microbiology* **73**: 4326-31 (2007).
8. Stams AJM, Plugge CM, de Bok FAM, van Houten B, Lens P, Dijkman H and Weijma J, Metabolic interactions in methanogenic and sulfate-reducing bioreactors. *Water Science and Technology* **52**: 13-20 (2005).
9. Raza F, Raza NA, Hameed U, Ikram ul H and Mariam I, Solid state fermentation for the production of b-glucosidase by co-culture of *Aspergillus niger* and *R. oryzae*. *Pakistan Journal of Botany* **43**: 75-83 (2011).
10. Stoilova I and Krastanov A, Overproduction of laccase and pectinase by microbial associations in solid substrate fermentation. *Applied Biochemistry and Biotechnology* **149**: 45-51 (2008).
11. Castillo MR, Gutierrezcorrea M, Linden JC and Tengerdy RP, Mixed culture solid substrate fermentation for cellulolytic enzyme-production. *Biotechnology Letters* **16**: 967-72 (1994).

12. Ahamed A and Vermette P, Enhanced enzyme production from mixed cultures of *Trichoderma reesei* rut-c30 and *Aspergillus niger* lma grown as fed batch in a stirred tank bioreactor. *Biochemical Engineering Journal* **42**: 41-46 (2008).
13. Dashtban M, Schraft H and Qin WS, Fungal bioconversion of lignocellulosic residues; opportunities & perspectives. *International Journal of Biological Sciences* **5**: 578-95 (2009).
14. Maciel MJM, Silva ACE and Ribeiro HCT, Industrial and biotechnological applications of ligninolytic enzymes of the basidiomycota: A review. *Electronic Journal of Biotechnology* **13** (2010).
15. Holker U, Hofer M and Lenz J, Biotechnological advantages of laboratory-scale solid-state fermentation with fungi. *Applied Microbiology and Biotechnology* **64**: 175-86 (2004).
16. Brijwani K, Oberoi HS and Vadlani PV, Production of a cellulolytic enzyme system in mixed-culture solid-state fermentation of soybean hulls supplemented with wheat bran. *Process Biochemistry* **45**: 120-28 (2010).
17. Jahangeer S, Khan N, Sohail M, Shahzad S, Ahmad A and Khan SA, Screening and characterization of fungal cellulases isolated from the native environmental source. *Pakistan Journal of Botany* **37**: 739-48 (2005).
18. Juwaied AA, Al-amiry A and Anaam U, Optimization of cellulase production by *Aspergillus niger* and *Trichoderma viride* using sugar cane waste. *Journal of Yeast and Fungal Research* **2**: 19-23 (2011).
19. GutierrezCorrea M and Tengerdy RP, Production of cellulase on sugar cane bagasse by fungal mixed culture solid substrate fermentation. *Biotechnology Letters* **19**: 665-67 (1997).
20. Duenas R, Tengerdy RP and Gutierrezcorrea M, Cellulase production by mixed fungi in solid-substrate fermentation of bagasse. *World Journal of Microbiology & Biotechnology* **11**: 333-37 (1995).
21. Carvalheiro F, Roseiro JC and Collaco MTA, Biological conversion of tomato pomace by pure and mixed fungal cultures. *Process Biochemistry* **29**: 601-05 (1994).
22. Dan S, Marton I, Dekel M, Bravdo BA, He SM, Withers SG and Shoseyov O, Cloning, expression, characterization, and nucleophile identification of family 3, *Aspergillus niger* beta-glucosidase. *Journal of Biological Chemistry* **275**: 4973-80 (2000).
23. Wang C, Wu G, Chen C and Chen S, High production of β -glucosidase by *Aspergillus niger* on corncob. *Applied Biochemistry and Biotechnology* **168**: 58-67 (2012).
24. Kumar R, Singh S and Singh OV, Bioconversion of lignocellulosic biomass: Biochemical and molecular perspectives. *Journal of Industrial Microbiology & Biotechnology* **35**: 377-91 (2008).

25. Oberoi HS, Babbar N, Dhaliwal SS, Kaur S, Vadlani PV, Bhargav VK and Patil RT, Enhanced oil recovery by pre-treatment of mustard seeds using crude enzyme extract obtained from mixed-culture solid-state fermentation of kinnow (citrus reticulata) waste and wheat bran. *Food and Bioprocess Technology* **5**: 759-67 (2012).
26. Dwivedi P, Vivekanand V, Pareek N, Sharma A and Singh RP, Bleach enhancement of mixed wood pulp by xylanase-laccase concoction derived through co-culture strategy. *Applied Biochemistry and Biotechnology* **160**: 255-68 (2010).
27. Stoilova IS, Gargove SA and Krastanov A, Production of enzymes by mixed culture from micelial fungi in solid-state fermentation. *Biotechnol&Biotechnol* **19**: 103-08 (2005).
28. Gutierrez-Correa M and Tengerdy RP, Xylanase production by fungal mixed culture solid substrate fermentation on sugar cane bagasse. *Biotechnology Letters* **20**: 45-47 (1998).
29. Juwaied AA, Adnan S and Al-amieriy A, Production cellulase by different co-culture of *Aspergillus niger* and *Tricoderma viride* from waste paper. *Journal of Yeast and Fungal Research* **1**: 108-11 (2010).
30. Wang XJ, Bai JG and Liang YX, Optimization of multienzyme production by two mixed strains in solid-state fermentation. *Applied Microbiology and Biotechnology* **73**: 533-40 (2006).
31. Paranthaman R, Vidyalakshmi R, Muruges S and Singaravadivel K, Effects of fungal co-culture for the biosynthesis of tannase and gallic acid from grape wastes under solid state fermentation. *Global Journal of Biotechnology & Biochemistry* **4**: 29-36 (2009).
32. Jana A, Maity C, Halder SK, Mondal KC, Pati BR and Das Mohapatra PK, Tannase production by *Penicillium purpurogenum* paf6 in solid state fermentation of tannin-rich plant residues following ovat and rsm. *Applied Biochemistry and Biotechnology* **167**: 1254-69 (2012).
33. Rodriguez-Duran LV, Contreras-Esquivel JC, Rodriguez R, Prado-Barragan LA and Aguilar CN, Optimization of tannase production by *Aspergillus niger* in solid-state packed-bed bioreactor. *Journal of Microbiology and Biotechnology* **21**: 960-67 (2011).
34. Paranthaman R, Vidyalakshmi R, Murgesh S and Singaravadivel K, Effects of fungal co-culture for the biosynthesis of tannase and gallic acid from grape wastes under solid state fermentation. *Global Journal of Biotechnology& Biochemistry* **4**: 29-36 (2009).
35. Perinbam K, Kanimozhi S, Bharathy K and Ramnath V, Production of enzyme tannase by solid state fermentation using tamarind seed powder. *Biosciences, Biotechnology Research Asia* **6**: 209-14 (2009).
36. Mukherjee G and Banerjee R, Effects of temperature, pH and additives on the activity of tannase produced by a co-culture of *Rhizopus oryzae* and *Aspergillus foetidus*. *World Journal of Microbiology & Biotechnology* **22**: 207-12 (2006).

37. Banerjee R, Mukherjee G and Patra KC, Microbial transformation of tannin-rich substrate to gallic acid through co-culture method. *Bioresource Technology* **96**: 949-53 (2005).
38. Yalemtesfa B, Alemu T and Santhanam A, Solid substrate fermentation and conversion of orange waste in to fungal biomass using *Aspergillus niger* ka-06 and *Chaetomium spp* kc-06. *African Journal of Microbiology Research* **4**: 1275-81 (2010).
39. Ge R, Li H, Zhang H, Li b and Su Z, Screening of the strains for producing protein feeds by solid-state fermentation from cassava lees. *Shengwu Jishu* **20**: 82-84 (2010).
40. Gu XH, Zhang YG, Zhao GM, Zhang R, Zhang F and Zhu JH, Improvement of feeding value of rapeseed meal by mix solid state fermentation *Animal Husbandry and Feed Science* **2**: 12-17 (2010).
41. Yang SS, Protein enrichment of sweet-potato residue with coculture of amylolytic fungi by solid-state fermentation. *Biotechnology Advances* **11**: 495-505 (1993).
42. Xu BJ, Wang QJ, Jia XQ and Sung CK, Enhanced lovastatin production by solid state fermentation of *Monascus ruber*. *Biotechnology and Bioprocess Engineering* **10**: 78-84 (2005).
43. Panda BP, Javed S and Ali M, Optimization of fermentation parameters for higher lovastatin production in red mold rice through co-culture of *Monascus purpureus* and *Monascus ruber*. *Food and Bioprocess Technology* **3**: 373-78 (2010).
44. Ramalingam, Saraswathy N, Shanmugapriya S, Shakthipriyadarshini S, Sadasivam S and Shanmugaprasanth M, Decolorization of textile dyes by *Aspergillus tamarii*, mixed fungal culture and *Penicillium purpurogenum*. *Journal of Scientific & Industrial Research* **69**: 151-53 (2010).
45. Gong A, Qiu L, Wang D, Zhou L, Zhu Y, Zhang Z and Liu Q, Method for producing *Bacillus thuringiensis* (bt) biopesticide by co-culturing bt and *Bacillus subtilis* (bs) with kitchen waste via solid fermentation. China(2011).
46. Jung H and Lee S, Production of carrot pomace fortified with mucilage, fibrinolytic enzyme and probiotics by solid-state fermentation using the mixed culture of *Bacillus subtilis* and *Leuconostoc mesenteroides*. *Journal of Food Science and Nutrition* **14**: 335-42 (2009).
47. Atta H, Arafa R, Salem M and El-Meleigy M, Microbiological studies on the production of vitamin b12 from two mixed cultures under solid state fermentation condition. *Journal of Applied Sciences Research* **4**: 1463-77 (2008).
48. Tian G, Zhang L and Li G, Method for preparing composite biological fertilizer by combined solid fermentation of double bacterium China patent CN 101585722 (2009).
49. Ghosh MK and Ghosh UK, Utilization of pine needles as bed material in solid state fermentation for production of lactic acid by lactobacillus strains. *Bioresources* **6**: 1556-75 (2011).

50. Song C, Zhang Q and Li L, Composite acetic acid bacteria, culture method thereof, and method for producing acetic acid through solid-state fermentation. China patent CN 101875908 (2009).
51. Pandey A, Selvakumar P, Soccol CR and Nigam P, Solid state fermentation for the production of industrial enzymes. *Current Science* **77**: 149-62 (1999).
52. Hu JK, Lu WQ, Wang CL, Zhu RH and Qiao JY, Characteristics of solid-state fermented feed and its effects on performance and nutrient digestibility in growing-finishing pigs. *Asian-Australasian Journal of Animal Sciences* **21**: 1635-41 (2008).
53. Chiang G, Lu WQ, Piao XS, Hu JK, Gong LM and Thacker PA, Effects of feeding solid-state fermented rapeseed meal on performance, nutrient digestibility, intestinal ecology and intestinal morphology of broiler chickens. *Asian-Australasian Journal of Animal Sciences* **23**: 263-71 (2010).
54. Gu B, Ma H and Liu B, Preparation of bio-feedstuff with rich peptides from rapeseed meal by mixed solid fermentation. *Journal of the Chinese Cereals and Oils Association* **26** (2011).
55. Pan T, Zhang D, Zhao C and Li K, Study on microbial protein production by mixed culture solid-state fermentation on soybean waste. *Fermentation and Bioindustrial Chemistry* **16** (2004).
56. Wang P, Huang S and Yu B, Production of protein-enriched feed by direct bioconversion from sweet potato with mixed microbial culture. *Food and Feed Chemistry* **17** (2000).
57. Fu N, Peiris P, Markham J and Bavor J, A novel co-culture process with *Zymomonas mobilis* and *pichia stipitis* for efficient ethanol production on glucose/xylose mixtures. *Enzyme and Microbial Technology* **45**: 210-17 (2009).
58. Eliasson A, Christensson C, Wahlbom CF and Hahn-Hagerdal B, Anaerobic xylose fermentation by recombinant *Saccharomyces cerevisiae* carrying *xyl1*, *xyl2*, and *xks1* in mineral medium chemostat cultures. *Applied and Environmental Microbiology* **66**: 3381-86 (2000).
59. Okuda N, Ninomiya K, Katakura Y and Shioya S, Strategies for reducing supplemental medium cost in bioethanol production from waste house wood hydrolysate by ethanologenic *Escherichia coli*: Inoculum size increase and coculture with *Saccharomyces cerevisiae*. *Journal of Bioscience and Bioengineering* **105**: 90-96 (2008).
60. Sree NK, Sridhar M, Rao LV and Pandey A, Ethanol production in solid substrate fermentation using thermotolerant yeast. *Process Biochemistry* **34**: 115-19 (1999).
61. Fukuda H, Kizaki Y, Tsukihashi T and Wakabayashi S, Shochu brewing characteristics and properties of a trichothecin-resistant shochu yeast mutant. *Biotechnology Letters* **23**: 2009-13 (2001).

62. Murado MA, Pastrana L, Vazquez JA, Miron J and Gonzalez MP, Alcoholic chestnut fermentation in mixed culture. Compatibility criteria between *Aspergillus oryzae* and *Saccharomyces cerevisiae* strains. *Bioresource Technology* **99**: 7255-63 (2008).
63. Kato S, Haruta S, Cui ZJ, Ishii M and Igarashi Y, Network relationships of bacteria in a stable mixed culture. *Microbial Ecology* **56**: 403-11 (2008).
64. Sun YZ, Mao SY, Yao W and WY. Z, The dynamics of microorganism populations and fermentation characters of co-cultures of rumen fungi and cellulolytic bacteria on different substrates. *Wei Sheng Wu Xue Bao* **46**: 422-26 (2006).
65. Schink B, Syntrophic associations in methanogenic degradation. *Progress in Molecular and Subcellular Biology* **41**: 1-19 (2006).
66. Manikandan K and Viruthagiri T, Optimization of C/N ratio of the medium and fermentation conditions tapioca starch using co – culture of *Aspergillus niger*. *International Journal of ChemTech Research* **2**: 947-55 (2010).
67. Cho BK, Charusanti P, Herrgard MJ and Palsson BO, Microbial regulatory and metabolic networks. *Current Opinion in Biotechnology* **18**: 360-64 (2007).
68. Stolyar S, Van Dien S, Hillesland KL, Pinel N, Lie TJ, Leigh JA and Stahl DA, Metabolic modeling of a mutualistic microbial community. *Molecular Systems Biology* **3** (2007).
69. Pramanik J and Keasling JD, Stoichiometric model of *Escherichia coli* metabolism: Incorporation of growth-rate dependent biomass composition and mechanistic energy requirements. *Biotechnology and Bioengineering* **56**: 398-421 (1997).
70. Laplace JM, Delgenes JP, Moletta R and Navarro JM, Cofermentation of glucose and xylose to ethanol by a respiratory-deficient mutant of *Saccharomyces cerevisiae* co-cultivated with a xylose-fermenting yeast. *Journal of Fermentation and Bioengineering* **75**: 207-12 (1993).
71. Dharmagadda VSS, Nokes SE, Strobel HJ and Flythe MD, Investigation of the metabolic inhibition observed in solid-substrate cultivation of *Clostridium thermocellum* on cellulose. *Bioresource Technology* **101**: 6039-44 (2010).
72. Durand A, Bioreactor designs for solid state fermentation. *Biochemical Engineering Journal* **13**: 113-25 (2003).
73. Brijwani K, Vadlani PV, Hohn KL and Maier DE, Experimental and theoretical analysis of a novel deep-bed solid-state bioreactor for cellulolytic enzymes production. *Biochemical Engineering Journal* **58-59**: 110-23 (2011).
74. Mahadevan R, Edwards JS and Doyle FJ, Dynamic flux balance analysis of diauxic growth in *Escherichia coli*. *Biophysical Journal* **83**: 1331-40 (2002).

75. Lisha KP and Debasis S (2011) Dynamic flux balance modeling of fed-batch co-culture fermentation of glucose/xylose mixture for improved bioethanol in *3rd International Conference on Chemical, Biological and Environmental Engineering*, LACSIT Press, Singapore.
76. Qureshi N, Dien BS, Nichols NN, Saha BC and Cotta MA, Genetically engineered *Escherichia coli* for ethanol production from xylose - substrate and product inhibition and kinetic parameters. *Food and Bioproducts Processing* **84**: 114-22 (2006).
77. Zhang M, Eddy C, Deanda K, Finkenstein M and Picataggio S, Metabolic engineering of a pentose metabolism pathway in ethanologenic *Zymomonas mobilis*. *Science* **267**: 240-43 (1995).
78. Eiteman M, Lee S and Altman E, A co-fermentation strategy to consume sugar mixtures effectively. *Journal of Biological Engineering* **2**: 1-8 (2008).
79. Patle S and Lal B, Ethanol production from hydrolysed agricultural wastes using mixed culture of *Zymomonas mobilis* and *Candida tropicalis*. *Biotechnology Letters* **29**: 1839-43 (2007).
80. Rorick R, Nahar N and Pryor S (2009) Enzymatic hydrolysis and fermentation of sugar beet pulp in *American Society of Agriculture and Biological Engineering* Nevada.
81. Duff SJB, Cooper DG and Fuller OM, Cellulase and beta-glucosidase production by mixed culture of *Trichoderma reesei* rut c30 and *Aspergillus phoenicis*. *Biotechnology Letters* **7**: 185-90 (1985).
82. Mukherjee G and Banerjee R, Biosynthesis of tannase and gallic acid from tannin rich substrates by *Rhizopus oryzae* and *Aspergillus foetidus*. *Journal of Basic Microbiology* **44**: 42-48 (2004).
83. Nigam P, Process selection for protein-enrichment - fermentation of the sugar-industry by-products molasses and sugar-beet pulp. *Process Biochemistry* **29**: 337-42 (1994).
84. Demain AL, Pharmaceutically active secondary metabolites of microorganisms. *Applied Microbiology and Biotechnology* **52**: 455-63 (1999).
85. Manzoni M and Rollini N, Biosynthesis and biotechnological production of statins by filamentous fungi and application of these cholesterol-lowering drugs. *Applied Microbiology and Biotechnology* **58**: 555-64 (2002).
86. Chang YN, Huang JC, Lee CC, Shih IL and Tzeng YM, Use of response surface methodology to optimize culture medium for production of lovastatin by *Monascus ruber*. *Enzyme and Microbial Technology* **30**: 889-94 (2002).
87. Chang YN, Lin YC, Lee CC, Liu BL and Tzeng YM, Effect of rice-glycerol complex medium on the production of lovastatin by *Monascus ruber*. *Folia Microbiologica* **47**: 677-84 (2002).

Supplement

Discussion of the studies in Table 1

Co cultures for cellulase system

Various mixed cultures of *Aspergillus* species were tested by Raza.⁹ Among each of the two genera combinations of *A. niger*, *A. awamori* and *A. oryzae*, the co- culture of *A. niger* and *A. oryzae* at a ratio of 3:1 gave the highest production rate of β -glucosidase, 2975 ± 5.3 U/g/min at pH 5.0 among all the tested combinations, which was more than ten-fold higher productivity than similar monocultures.

Brijwani et al.¹⁶ demonstrated that co-culturing of *T. reesei* with *A. oryzae* in 1:1 ratio boosted the production of β -glucosidase activity by 65% in SSC of soybean hulls supplemented with wheat bran, when compared to a monoculture of *T. reesei*. The co-culture system also synergistically produced 100 IU/g endocellulase while only 60 IU/g and 68 IU/g were obtained with a monoculture of *T. reesei* and *A. oryzae*, respectively. A mixed culture of *T. reesei* and *A. phoenicis* in SSC using bagasse was investigated by Duenas *et al.*²⁰ to produce a cellulase with an increased activity of β -glucosidase (See Table 1). The highest cellulase and β -glucosidase by co-culture obtained were 18.7 and 38.6 IU/g, respectively, representing approximately 3- and 6-fold increase over the activities attained in single-culture SSC. The same co-culture system was adopted by Gutierrez-Correa and Tengerdy¹⁹ and up to 2.7-, 3.8- and 2.3-fold increase of activities of cellulase, endoglucanase and β -glucosidase respectively over that from monocultures. In addition, the reported volumetric productivity rates of cellulase production using co-culture SSC by Duenas *et al.*²⁰ and Gutierrez-Correa and Tengerdy¹⁹ were $280 \text{ IU.L}^{-1}.\text{h}^{-1}$ and $210 \text{ IU.L}^{-1}.\text{h}^{-1}$ with no washed dilution, respectively, significantly higher than that obtained in mixed submerged fermentations; $12.5 \text{ IU.L}^{-1}.\text{h}^{-1}$ by *T. reesei* and *A. phoenicis*⁸¹, which demonstrated an advantage of SSC over SmF in terms of concentrated activities. Juwaied²⁹ tested co-cultures of *A. niger* and *T. viride* at various coculture initiation times. The study found

that the cellulase obtained from the continuously mixed cultures had maximum activity (about 2.4 U/ml) as compared to their pure cultures (about 0.6-1.0 U/ml).

Co cultures for multiple enzyme complexes

T. reesei was co-cultured with either *A. niger* or *A. phoenicis* in SSC on sugar cane bagasse and this resulted in higher xylanase activities than their corresponding monocultures. When soymeal was used as the nitrogen supplement at a C:N ratio of 10:1, activity levels of xylanase (2,600 – 2,800 IU/g dry biomass) and cellulase (14 – 15 IU/g dry biomass), corresponding to 35 - 45% and 20 - 142% higher than single cultures of either partner, were obtained by both mixed-culture systems after 72 hours of fermentation.²⁸ Another study produced both xylanase and laccase by mutant *Penicillium oxalicum* SAUE-3.510 cultured with *Pleurotus ostreatus* MTCC 1804 under SSC and the collected enzyme mixture (xylanase/laccase, 22:1) of 8 IU/g bleached the mixed wood pulp with a notable decrease of 21% in kappa number and increase of 8% in brightness as compared to xylanase alone.²⁶ Wang³⁰ selected the co-culture of two strains of *A. niger* (in a proportion of 1:4) and optimized the medium with 54.5% (w/w dry basis) water and demonstrated that the activities of cellulase, hemicellulase, glucoamylase, pectinase, and acidic protease acted together to provide a 65% increase in reducing sugars released as compared to each monoculture. Stoilova²⁷ cultivated *Thermoascus aurantiacus* with *A. niger* strains A and B respectively in SSC. A synergistic effect was observed between *Thermoascus aurantiacus* and *A. niger* – A. Higher activity of β -glucosidase (500 IU/g) was produced, surpassing the activity from the monoculture of *A. niger* - A by a factor of 2.5. The synergism between the species *Thermoascus aurantiacus* and *A. niger* - B led to an increased biosynthesis of laccase (31IU/g) and glucoamylase (308 IU/g), which was an activity 1.5 times higher than that of the *T. aurantiacus* monoculture. Two microbial fungal associations, *A. niger* & *Fusarium moniliforme* and *Trametes versicolor* & *A. niger*, were investigated in a study by Stoilova and Krastanov.¹⁰ The synergism between *A. niger* and *Fusarium moniliforme* increased the biosynthesis of α -amylase and pectinase, necessary for fungal growth. *Trametes versicolor* and *A. niger* proved to be compatible partners to produce high activity of laccase (97,600 IU/g dry weight substrate), surpassing the enzymatic activity level in the monoculture by a factor of 8.4.

Discussion of the studies in Table 2

Co-culture of *R. oryzae* and *A. foetidus* is one of the most widely investigated microbial combinations for bio-converting tannin-rich substrate to gallic acid. When either strain, *R. oryzae* or *A. foetidus* was grown separately under SSC on powdered fruits of *Terminalia chebula* and *Caesalpinia digyna* pod, *A. foetidus* showed a marginally higher yield of tannase activity (36.4 U/ml in 72 h) than *R. oryzae* (31.8 U/ml in 60 h), however *A. foetidus* required a longer incubation period to produce this amount of enzyme activity.⁸² A higher amount of tannase (41.3 U/ml) was produced in a shorter time period (48 h) by these two fungal strains together than monoculture under SSC using the same substrate.³⁷ Mukherjee³⁶ investigated the effects of temperature and pH on the activity of purified tannase produced by co-culturing *R. oryzae* and *A. foetidus* on the fruits of *Terminalia chebula* and *teri* pod with the culture conditions of 40 °C and 5.0 respectively.

Other fungal combinations for tannase production were investigated successfully. Among the co-cultures of : *A. niger* and *P. chrysogenum*; *P. chrysogenum* and *T. viride*; *A. niger* and *T. viride*; *A. niger* & *P. chrysogenum* & *T. viride* using grape's peel as a substrate, the optimal combination was *P. chrysogenum* and *T. viride* which produced enzyme activity of 84±2 U/g/min, which was higher than the respective mono cultures (61- 63 U/g/min).³⁴

Discussion of the studies in Table 3

Mixed fungal cultures generate diverse enzymes permitting transformation of raw materials or agricultural residues into high protein biomass with interesting amino acid compositions. Combinations of amylolytic *Aspergillus sp.* and *Rhizopus sp.* on sweet potato residue showed higher protein enrichment (32.4% protein) than mono cultures of fungus (26% -29% protein) and amylolytic yeast (17-18.4% protein).⁴¹ For agricultural residues and lignocellulose materials containing fiber and other anti-nutrients, high levels of enzymatic production during SSC, especially of cellulolytic enzymes, enhanced the digestibility of the materials designed for animal feed products and furthered the interest in SSC of mixed cultures.^{10, 83} Co-cultures like *Aspergillus sp.* and *T. viride* using rapeseed meal or *Aspergillus sp.* and *Penicillium sp.* using food waste like cassava lees (byproduct in the process of alcohol production by utilizing cassava) demonstrated the ability to use fiber (decreased by 19-22 %) thus stimulating the accumulation of nitrogenous nutrients.^{39, 40}

As discussed above, co-culture of fungi on SSC helps in maximizing the utilization of substrate; moreover co-cultures provide support for secondary metabolite production. A typical example is fermented rice (red mold rice) by mold, which has been applied as a functional food to reduce the serum total cholesterol and triglycerides due to presence of lovastatin in the fermented rice.^{84, 85} Different *Monascus* species such as *M. ruber*, *M. purpureus*, *M. anka*, and *M. pilosus* are utilized in China to produce red mold rice. Co-cultures of *M. purpureus* MTCC 369 and *M. ruber* MTCC 1880 under SSC were reported to yield 2.83 mg/g lovastatin in red mold rice under optimized process conditions.⁴³ Up to 64.7% increment in lovastatin yield over the mono culture of *M. ruber* reported by Xu⁴². By volume, this yield is equivalent to about 1.4 g/L losvatin, which represents about 8.8~10.5-fold increase over the yield of lovastatin generated by mono culture under optimized submerged liquid conditions in other papers.^{86, 87}

Not all fungal co-cultures demonstrated better performances than monoculture however. The degradative ability of *A. tamari*, *P. purpurogenum* and mixed culture (*Trichoderma sp.*, and *A. flavus*) on various synthetic dyes were described by Ramalingam.⁴⁴ The results showed that each monoculture completely decolorized coomassie brilliant blue after 20 days while mixed cultures decolorized coomassie brilliant blue, bromophenol blue and malachite green with a 15.6%-21.8% decrease in degradation efficiency. The intrinsic reasons were not investigated in the study. In addition, lack of comparison with the monoculture of *Trichoderma sp.* and *A. flavus* in this study did not allow differentiation between monoculture effects and synergistic effects of the co-culture.

Discussion of the studies in Table 4

Solid substrate bacterial co cultures have been reported to produce biological pest control agents or produce a fertilizer. Gong⁴⁵ inoculated 10-30% (by mass) *Bacillus thuringiensis* and *Bacillus subtilis* into kitchen waste containing 45-60% water for pesticide production. The main advantage of using the probiotic *Bacillus subtilis* lies in its ability to effectively degrade oils in kitchen waste. Atta⁴⁷ selected two microbial isolates for their high yield of vitamin B₁₂; a bacteria (*Bacillus firmus* AZ-78B) and an actinomycete (*Streptomyces halstedii*, AZ- 8A) from among 140 microbial isolates from soil. The co-culture of these two microorganisms produced 37.7 µg /ml of vitamin B₁₂ from agricultural crop residues which translates to more than a 20%

increase over mono-cultures. Tian⁴⁸ used *Bacillus mucilaginosus* and *Bacillus megaterium* to produce an efficient composite bacterial fertilizer by adjusting the ratio of inoculum to 2:1 and controlling the cultivation conditions at various stages of production. The process improved with the co-culture by demonstrating an accelerated strain propagation, short fermentation period of 52 h (traditionally 72-100 h) and high spore formation rate of 90% (50-70% for mono-cultures) thus significantly lowering the production cost. In other applications, *Bacillus subtilis* HA and *Leuconostoc mesenteroides* were co-cultivated on carrot pomace by SSC to increase production of the bioactive compounds, such as 24% (g/g) of mucilage content (2.31% by *Bacillus subtilis* alone), fibrinolytic enzyme of 104.9 U/g (16.95 U/g by *Bacillus subtilis*) and probiotics. The fermented carrot pomace could be utilized as a valuable ingredient for functional foods or in the cosmetic industries.⁴⁶

Another interesting prospect is that co-cultivation of bacteria may be an effective method to overcome the recalcitrance of biomass. Ghosh⁴⁹ utilized pine needles as feedstock in lactic acid production in SSC using *L. delbrueckii* (NCIM2025) and *L. pentosus* (NCIM 2912). They saw a greater than 72% increase in the yield of lactic acid over mono cultures (8.72 to 26.15 g/L).

In traditional food fermentation, co-cultures of bacteria played a vital role in improving the quality and flavor of the final product. Song⁵⁰ screened five types of composite acetic acid bacteria from fermented grains of Shanxi mature vinegar. The composite acetic acid bacteria culture solution was inoculated into fermented grains and bran using SSC. The mixed culture improved acetic acid production rate, shortened fermentation time, and increased gluconic acid content in the suite of metabolites. The quality of the obtained table vinegar was also improved.

Discussion of the studies in Table 5

Research on co-cultures of bacteria, yeast and fungi for protein enrichment and animal feed mainly involves identification of suitable process parameters. Gu⁵⁴ reported that the combination of *Geotrichum candidum* and *Bacillus subtilis* resulted in the highest protein enrichment (12.81%) among the other combinations of *A. oryzae*, *Geotrichum candidum*, *Candida tropicalis*, *Saccharomyces cerevisiae* (*S. cerevisiae*) and *Bacillus subtilis* grown on rapeseed meal and salts (CaCl₂ 0.1% and NH₄Cl 1.5%). The study by Chiang⁵³ showed protein enrichment of up to

36.6% by a mixed culture of *Lactobacillus fermentum*, *Enterococcus faecium*, *Saccharomyces cerevisiae* and *Bacillus subtilis* (30-day culture) on rapeseed meal supplemented with wheat bran and brown sugar. Pan⁵⁵ enriched crude protein from soybean meal by optimizing cultivation conditions which were as follows: inoculum volume *A. niger*: *T. viride*: *S. cerevisiae*: *Candida utilis* = 1:1:1:3, initial pH value of 5.5 and cultivation temperature of 32 °C. Under the optimum conditions, the amount of crude protein could be as high as 28.5% on a dry basis, which was approximately twice that of the raw material. Hu⁵² designed a substrate medium containing soybean meal, corn gluten meal, rapeseed meal and peanut meal for co-culturing *Lactobacillus fermentum*, *S. cerevisiae* and *Bacillus subtilis*. The crude protein increased to 31% with 9.2% of water soluble protein after a 30-day culture. A mixed culture of *R. oligosporus* and *Candida arborea* using SSC in a shallow dish (length× width × height = 290mm×190mm×55mm) enriched protein from sweet potato to 22.1% when fermented at 28°C for 72 h. Using a solid-substrate bioreactor (length× width × height = 500mm×350mm×250mm) for 48 h at 28°C, 15.9% crude protein was obtained. The total amount of amino acids were increased by 101%, compared with that of raw material.⁵⁶