2-17-2021

CERT_L Reduces C16 Ceramide, Amyloid- β Levels, and Inflammation in a Model of Alzheimer's Disease

Simone M. Crivelli

University of Kentucky, s.crivelli@uky.edu

Qian Luo

Maastricht University, the Netherlands

Jo A. A. Stevens

Maastricht University, the Netherlands

Caterina Giovagnoni

Maastricht University, the Netherlands

Daan van Kruining

Maastricht University, the Netherlands

Follow this and additional works at: https://uknowledge.uky.edu/physiology_facpub

Part of the [Internal Medicine Commons](#), [Medical Physiology Commons](#), [Neurology Commons](#), and the [Neuroscience Commons](#)

See next page for additional authors

[Right click to open a feedback form in a new tab to let us know how this document benefits you.](#)

Repository Citation

Crivelli, Simone M.; Luo, Qian; Stevens, Jo A. A.; Giovagnoni, Caterina; van Kruining, Daan; Bode, Gerard; den Hoedt, Sandra; Hobo, Barbara; Scheithauer, Anna-Lena; Walter, Jochen; Mulder, Monique T.; Exley, Christopher; Mold, Matthew; Mielke, Michelle M.; De Vries, Helga E.; Wouters, Kristiaan; van den Hove, Daniel L. A.; Berkes, Dusan; Ledesma, María Dolores; Verhaagen, Joost; Losen, Mario; Bieberich, Erhard; and Martinez-Martinez, Pilar, "CERT_L Reduces C16 Ceramide, Amyloid- β Levels, and Inflammation in a Model of Alzheimer's Disease" (2021). *Physiology Faculty Publications*. 169.
https://uknowledge.uky.edu/physiology_facpub/169

This Article is brought to you for free and open access by the Physiology at UKnowledge. It has been accepted for inclusion in Physiology Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

CERT_L Reduces C16 Ceramide, Amyloid- β Levels, and Inflammation in a Model of Alzheimer's Disease

Digital Object Identifier (DOI)

<https://doi.org/10.1186/s13195-021-00780-0>

Notes/Citation Information

Published in *Alzheimer's Research & Therapy*, v. 13, issue 1, article no. 45.

© The Author(s) 2021

This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <https://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<https://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Authors

Simone M. Crivelli, Qian Luo, Jo A. A. Stevens, Caterina Giovagnoni, Daan van Kruining, Gerard Bode, Sandra den Hoedt, Barbara Hobo, Anna-Lena Scheithauer, Jochen Walter, Monique T. Mulder, Christopher Exley, Matthew Mold, Michelle M. Mielke, Helga E. De Vries, Kristiaan Wouters, Daniel L. A. van den Hove, Dusan Berkes, María Dolores Ledesma, Joost Verhaagen, Mario Losen, Erhard Bieberich, and Pilar Martinez-Martinez

RESEARCH

Open Access

CERT_L reduces C16 ceramide, amyloid- β levels, and inflammation in a model of Alzheimer's disease

Simone M. Crivelli^{1,2,3†}, Qian Luo^{1†}, Jo A.A. Stevens¹, Caterina Giovagnoni¹, Daan van Kruining¹, Gerard Bode¹, Sandra den Hoedt⁴, Barbara Hobo⁵, Anna-Lena Scheithauer⁶, Jochen Walter⁶, Monique T. Mulder⁴, Christopher Exley⁷, Matthew Mold⁷, Michelle M. Mielke⁸, Helga E. De Vries⁹, Kristiaan Wouters^{10,11}, Daniel L. A. van den Hove¹², Dusan Berkes¹³, María Dolores Ledesma¹⁴, Joost Verhaagen⁵, Mario Losen¹, Erhard Bieberich^{2,3} and Pilar Martinez-Martinez^{1*}

Abstract

Background: Dysregulation of ceramide and sphingomyelin levels have been suggested to contribute to the pathogenesis of Alzheimer's disease (AD). Ceramide transfer proteins (CERTs) are ceramide carriers which are crucial for ceramide and sphingomyelin balance in cells. Extracellular forms of CERTs co-localize with amyloid- β (A β) plaques in AD brains. To date, the significance of these observations for the pathophysiology of AD remains uncertain.

Methods: A plasmid expressing CERT_L, the long isoform of CERTs, was used to study the interaction of CERT_L with amyloid precursor protein (APP) by co-immunoprecipitation and immunofluorescence in HEK cells. The recombinant CERT_L protein was employed to study interaction of CERT_L with amyloid- β (A β), A β aggregation process in presence of CERT_L, and the resulting changes in A β toxicity in neuroblastoma cells. CERT_L was overexpressed in neurons by adeno-associated virus (AAV) in a mouse model of familial AD (5xFAD). Ten weeks after transduction, animals were challenged with behavior tests for memory, anxiety, and locomotion. At week 12, brains were investigated for sphingolipid levels by mass spectrometry, plaques, and neuroinflammation by immunohistochemistry, gene expression, and/or immunoassay.

Results: Here, we report that CERT_L binds to APP, modifies A β aggregation, and reduces A β neurotoxicity in vitro. Furthermore, we show that intracortical injection of AAV, mediating the expression of CERT_L, decreases levels of ceramide d18:1/16:0 and increases sphingomyelin levels in the brain of male 5xFAD mice. CERT_L in vivo over-expression has a mild effect on animal locomotion, decreases A β formation, and modulates microglia by decreasing their pro-inflammatory phenotype.

(Continued on next page)

* Correspondence: p.martinez@maastrichtuniversity.nl

[†]Simone M. Crivelli and Qian Luo contributed equally to this work.

¹Department of Psychiatry and Neuropsychology, School for Mental Health and Neuroscience, Maastricht University, Universiteitssingel 50, 6229 ER Maastricht, the Netherlands

Full list of author information is available at the end of the article

© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

(Continued from previous page)

Conclusion: Our results demonstrate a crucial role of CERT_L in regulating ceramide levels in the brain, in amyloid plaque formation and neuroinflammation, thereby opening research avenues for therapeutic targets of AD and other neurodegenerative diseases.

Keywords: Ceramide, Sphingomyelin, Ceramide transporter protein (CERT), Adeno-associated virus (AAV), Alzheimer's disease (AD), 5xFAD, Amyloid- β plaques, Neuroinflammation, Microglia

Background

Key pathological features of Alzheimer's disease (AD) are aggregates of amyloid- β peptides (A β) and neurofibrillary tangles (NFTs), and neurodegeneration, together with blood-brain barrier (BBB) dysfunction, neuroinflammation, and lipid disbalance. To date, the molecular mechanism underlying neurodegeneration in AD remains unclear. Elucidation of the dysregulated biological mechanisms that lead to the onset and progression of AD is critical to identify new treatment strategies [1–5].

Sphingolipids (SLs) are waxy lipids formed by a sphingosine backbone, important for the cell membrane architecture and for the function of transmembrane proteins. Furthermore, SLs such as ceramides (Cer) and sphingosine-1-phosphate (S1P) are potent second messengers that regulate various important cellular processes, including cell growth and apoptosis [6–9]. Cer are formed by two metabolic pathways: *de novo* synthesis initiated with the precursor palmitoyl-CoA, or catabolism of complex SLs such as sphingomyelin (SM) [10–12]. In the cell membrane, SLs are typically organized in microdomains, called lipid rafts, characterized by specific SL species composition.

Several studies have analyzed lipid composition in AD brain tissue, reporting an increase of Cer species [13–16]. Lipid rafts enriched in Cer, isolated postmortem from frontal cortex tissue of AD patients, showed a reduction of SM levels compared to those isolated from the control brains [17]. Moreover, in AD brain tissue, SM levels were reduced in brain regions particularly vulnerable to A β plaque formation [18–20]. Strong evidence links A β pathology to SL homeostasis. The enzymes β -secretase and γ -secretase, which cleave the amyloid precursor protein (APP) to generate A β , are stabilized and have an increased half-life in Cer enriched membranes, thus increasing A β biogenesis [21, 22]. In turn, A β can stimulate Cer production by directly activating the phosphodiesterase enzyme sphingomyelinase which converts SM to Cer [18, 19].

Ceramide transfer proteins (CERTs) contain a steroidogenic acute regulatory protein (StAR)-related lipid transfer (START) domain that confers the ability to transport Cer intracellularly between the endoplasmic reticulum (ER) and the Golgi [23]. CERTs are found in at least two isoforms, which differ for the presence of a

26-amino acid serine-rich domain [24]. CERTs are expressed in the central nervous system, and they are crucial in embryogenesis and brain development [24–26]. When CERTs' activity is blocked pharmacologically or genetically, by compromising the START domain of the protein, SM production decreases significantly [27, 28]. CERT_L can be secreted extracellularly and was found to partially co-localize with serum amyloid P component (SAP) and with amyloid plaques in AD brain [29, 30]. Besides, CERTs are also potent activators of the classical complement pathway that plays an active role in AD pathogenesis [31]. To date, the significance of these observations for the pathophysiology of AD remains uncertain.

In the current study, we investigated the interaction of CERT_L with APP and A β *in vitro*. Next, we explored the effect of CERT_L overexpression on SL composition, amyloid formation, and inflammation *in vivo* using adeno-associated virus (AAV)-mediated gene delivery in the 5xFAD mouse model [32]. Our findings showed that an increase of CERT_L modulated SL levels by reducing specific Cer and elevating SM. Notably, CERT_L also affected amyloid plaque formation and brain inflammation, supporting the idea that enzymes and transporters of the SL pathways are at the core of the pathophysiological changes observed in AD.

Material and methods

CERT_L interaction with APP/A β

Immunoprecipitation (IP)

Wild-type HEK293 and transgenic HEK293 cells that stably overexpress human APP695 isoform (NP_958817.1) [33] were cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with fetal bovine serum (FBS), penicillin/streptomycin (Pen/Strep), and L-glutamine. Stable transfected HEK-APP were maintained in G418 selective medium. Prior to the experiment, cells were seeded in 25-cm² flasks and maintained in serum-free DMEM for 24 h. For the homogenization, cells were washed two times with phosphate-buffered saline (PBS), collected in lysis buffer (25 mM Tris HCl pH 7.5/150 mM NaCl, 0.5% Triton X-100 and protease inhibitors), and centrifuged at 20,000g for 30 min, and the resultant supernatants collected for the Bradford protein analysis. Protein extracts (100 μ g) from HEK or HEK-APP were used

for immunoprecipitation experiments. Pull down of endogenous CERT_L and APP was performed with 1 µg mAb anti CERT_L (3A1-C1) [29] and anti-Aβ mAb 6E10 (Covance), respectively by 1-h incubation at room temperature. mAb anti-syntaxin 6335 (clone 3D10, Abcam) was used as an isotype control. Next, anti-mouse secondary antibodies (Eurogentec) were used to pull down the immune complex. Thereafter, samples were centrifuged at 20,000g for 30 min. Pellets were washed three times in 50 µL PBS and boiled in reducing sample buffer containing mercaptoethanol to solubilize immunocomplexes. Then, the proteins were separated on a Tris-HCl 4–15% gradient gel (Bio-Rad) and blotted on nitrocellulose membrane (Millipore). Next, the membranes were probed with anti-Aβ/APP (6E10) or rabbit pAb anti-CERTs (epitope 1–50 of human CERTs, Bethyl Laboratories) antibodies. After 3 washes, membranes were incubated with donkey anti-mouse IRDye 680 and goat anti-rabbit IRDye 800 (Rockland Immunochemicals) and scanned using the Odyssey infrared imaging system (LI-COR Biosciences).

Neuronal culture and immunofluorescent staining

Primary neurons were cultured from 5xFAD neonates P0 as described, with modifications [34]. After the cortical area was dissected, the tissue was digested in 0.25% trypsin in Hank's Balanced Salt Solution (HBSS, Corning) for 15 min. Trypsin activity was stopped with plating medium, DMEM (Gibco, Invitrogen) containing 10% FBS and N2 supplement. Then, the digested tissue was passed through a cell strainer, spun down, and cells resuspended in plating medium. Cells were seeded onto poly-D-lysine-coated coverslips and cultured at 37 °C in a 5% CO₂ atmosphere. After 4 h, the plating medium was replaced with Neurobasal medium supplemented with B27 supplement, Pen/Strep, and 0.5 mM L-glutamine and kept on for 10–14 days. Every other day, supplemented Neurobasal medium was partially replaced.

Neurons were fixed with 4% PFA in PBS (Thermo Scientific) at 4 °C for 10 min, permeabilized with 0.25% Triton-X in PBS for 5 min, washed three times with PBS, and incubated with 3% BSA for 30 min. Cells were stained with rabbit polyclonal anti-CERTs (epitope 300–350 of human CERTs, Bethyl Laboratories), goat anti-MAP-2 (D-19) (Santa Cruz Biotechnology), and 6E10 anti APP/ Aβ [29]. The following secondary antibodies conjugated to fluorophores were used for detection: anti-mouse IgG Alexa 647, anti-rabbit IgG cy3, and anti-goat IgG Alexa 488. Fluorescence microscopy was performed using Eclipse Ti2-E inverted microscope system (Nikon). Images were processed using Nikon NIS-Elements software equipped with a 3D deconvolution program.

Microscale thermophoresis binding analysis

Microscale thermophoresis (MST) analysis was performed in the Monolith NT.155 instrument (Nanotemper). In brief, 20 nM of NT647 labeled CERT was incubated for 20 min at room temperature in the dark with different concentrations of either Aβ_{1–42} (rPeptide Athens) (3–100,000 nM) or control 17 kDa Lama antibody fragment (H6) (1–35,000 nM) in PBS Tween20 (0.01%). Afterward, 3–5 µL of the samples were loaded into glass capillaries (Monolith NT Capillaries, Cat#K002), and the thermophoresis analysis was performed (LED 40.51%, IR laser 80%). Statistical analysis was performed with Origin8.5 Software.

Aβ aggregation assay and cell-based toxicity assay

Transmission electron microscopy (TEM)

Aβ_{1–42}, purchased as the lyophilized salt (Bachem), was dissolved in 0.01 M NaOH in ultrapure water to give an Aβ_{1–42} stock solution of *ca* 0.20 mM, which was used immediately to prepare each of the treatments. The remaining peptide stock solution was frozen at –20 °C until required. Under these highly alkaline conditions, the peptide is fully dissolved and exists only as monomers [35]. Treatments containing Aβ_{1–42} and/or CERT_L [29] were prepared in 0.20-µm filtered modified Krebs-Henseleit (KH) medium (118.5 mM NaCl, 4.8 mM KCl, 1.2 mM MgSO₄, 1.4 mM CaCl₂, 11.0 mM glucose), buffered in 100 mM PIPES at pH 7.4, including 0.05% w/v sodium azide to inhibit microbial growth. Samples were incubated at 37 °C until their specified time points prior to being prepared onto TEM grids. For replicate samples, Aβ_{1–42} was thawed thoroughly immediately before use and then vortexed briefly. The stock solution was centrifuged at 15,000 rpm for 5 min, and 2.0 µL was then taken ready for concentration determination by absorbance at 280 nm, utilizing a NanoDrop 1000 spectrophotometer (Thermo). The concentration of Aβ_{1–42} was calculated with the Beer-Lambert law and the extinction coefficient 1390 M^{–1} cm^{–1}. CERT_L concentrations were determined in the same manner with a value of 107,925 M^{–1} cm^{–1} taken as the extinction coefficient using the 72 kDa recombinant CERT_L sequence (hCERT_L, 1875 bp NP_005704.1).

All samples for TEM were prepared via a modified TEM staining protocol [36]. Pre-coated S162 200 mesh formvar/carbon-coated copper grids (Agar Scientific) were inserted into 20.0 µL of the sample beaded onto paraffin film for 60 s, then wicked, passed through ultrapure water, re-wicked, and placed into 30 µL 2% uranyl acetate (in 70% ethanol), for 30 s. Following staining with uranyl acetate, grids were removed, wicked, passed through ultra-pure water, re-wicked, and placed into 30 µL 30% ethanol for 30 s. Grids were finally re-wicked following this step, covered and allowed to dry for up to 24 h, prior to analysis via TEM.

Samples for TEM were viewed on a JEOL 1230 transmission electron microscope operated at 100.0 kV (spot size 1), equipped with a Megaview III digital camera from Soft Imaging Systems (SIS). Images were obtained on the iTEM universal TEM imaging platform software.

Aggregation assay

A β_{1-42} was purchased from Anaspec. The peptide was solubilized in sterilized PBS, 0.1% trifluoroacetic acid (TFA) at the concentration 2 mM and frozen in aliquots at -80°C . Aliquots were diluted at the final concentration of 20 μM in a total volume of 400 μL containing 1 or 2.5 μM of affinity-purified recombinant CERT_L [29]. Samples were kept under rotarod shaking for 1, 2, 4, 8, 12, and 24 h at 37°C before adding 5 μL to 95 μL of thioflavin T (ThT) concentrated 20 μM , dispensed in 96-well optical plate and measuring fluorescent excitation at 450 nm and emission 486 nm in Victor X3 plate reader (Perkin-Elmer). Two A β antibodies against epitope 1–16 (6E10, Biolegend) and 17–24 (4G8, Biolegend) respectively were used to antagonize aggregation at a concentration of 0.1 mg/ml.

Toxicity assay

SH-SY5Y cells were seeded on a 96-well plate at a density of 3×10^4 cells per well in 1:1 DMEM:F12 with phenol red, 4 mM glutamine, 200 U/ml penicillin, 200 U/ml streptomycin, MEM non-essential amino acids (100 $\times$; Gibco), and 10% FBS and incubated at 37°C for 24 h, reaching up to 100% confluency, with 5% CO₂. After 24 h, the medium was removed and replaced with 100 μL /well medium without phenol-red, containing 2% FBS and with 10 μM A β_{1-42} oligomers, and/or 1 μM CERT_L, or alone to control wells and incubated for 24 h. Ten microliters of MTT (4 mg/ml) was added to each well and incubated at 37°C for 3 h. MTT solution was decanted and the formazan was extracted with 100 μL of 4:1 DMSO:EtOH. Plates were read at 570 nm, with a reference filter at 690 nm.

Generation of adeno-associated virus

Human collagen type IV alpha 3 binding protein cDNA sequence (hCERT_L, 1875 bp NP_005704.1) was cloned into the plasmid AAV-6P-SEW a kind gift of Prof. S. Kugler, Department of Neurology, University of Göttingen. The transgene expression was controlled by a human synapsin-1 promoter (hSYN, 480 bp), and an internal ribosome entry site (IRES 566 bp) enabled the co-expression of EGFP [37]. The plasmid expressing exclusively EGFP was used as a control (pAAV-EGFP). The AAV-CERT_L plasmid was sequenced by GATC Biotech laboratories and both AAVs plasmids were tested in vitro. AAVs particles were produced as explained previously [38]. In brief, the transfer plasmids pAAV-EGFP

or pAAV-CERT_L were used to produce AAV2 particles. Eight 15-cm petri dishes each containing 1.25×10^7 HEK 293 T cells in DMEM containing 10% fetal calf serum (FCS) and 1% Pen/Strep (all GIBCO-Invitrogen Corp., New York, NY, USA) were prepared 1 day before transfection. The medium was refreshed 1 h prior to transfection to Iscove's modified Eagle medium (IMEM) containing 10% FCS, 1% Pen/Strep, and 1% Glutamine. Transfer plasmids were co-transfected using polyethylenimine (PEI, MV25000; Polysciences Inc.) in a ratio of 1:3 with the pAAV-EGFP or pAAV-CERT_L resulting in a total amount of 50 μg of plasmid DNA per plate. The day after transfection, the medium was replaced with fresh IMEM with 10% FCS, 1% PS, and 1% glutamine. Two days later (3 days post-transfection), cells were harvested in Dulbecco-PBS (D-PBS, GIBCO) and lysed with 3 freeze-thaw cycles. Genomic DNA was digested by adding 10 $\mu\text{g}/\text{ml}$ DNaseI (Roche Diagnostics GmbH) into the lysate and incubated for 1 h at 37°C . The crude lysate was cleared by centrifugation at 4000 rpm for 30 min. The virus was purified from the crude lysate using the iodixanol gradient method, diluted in D-PBS/5% sucrose and concentrated using an Amicon 100 kDa MWCO Ultra-15 device (Millipore). All AAV vectors were stored at -80°C until use. Titers (genomic copies/ml) were determined by quantitative PCR on viral DNA primers directed against the EGFP portion (Forward: GTCTATATCATGGCCGACAA; Reverse: CTTGAAGTTCACCTTGATGC). The AAV particles produced with pAAV-CERT_L are referred to in this paper as AAV-CERT_L while the particles produced with pAAV-EGFP are named AAV-control.

Animals

In this study, male mice were used. To investigate transduction efficiency over time, we employed 24 C57BL/6 wild-type (WT) animals. B6/SJL WT and 5xFAD animals were obtained from the Jackson Laboratory and bred in house using 5xFAD $\times$ non-carriers. This breeding strategy may breed out the retinal degeneration allele Pde6brd1 from the original strain. The Jackson Lab has observed a less robust amyloid phenotype in this strain. The 5xFAD model carries 5 familial AD mutations, three of them in the human APP transgene (Swedish, Florida, and London), and two in the human presenilin-1 (PS1) transgene (M146L and L286V mutations). These mutations lead to an increase in A β peptide production [32]. Animals were individually housed under a 12 h light/dark cycle in individually ventilated cages. One week before behavioral tests, animals were adjusted to a reversed day-night cycle. Food and water were provided ad libitum throughout the study. All experiments were approved by the Animal Welfare Committee of Maastricht University (project number

DEC2013-056 and DEC2015-002) and followed the laws, rules, and guidelines of the Netherlands.

Stereotactic injection

The animals underwent bilateral stereotactic injections. Mice were placed in a stereotactic head frame, and after midline incision of the skin, two holes were drilled in the skull in the appropriate location using bregma and lambda as references. The layer V of the motor-sensory frontal cortex was targeted; this was verified by light microscopy to observe the dye. Coordinates were determined as follows: anterior-posterior [AP] 0.06, mediolateral [ML] ± 0.15 , and dorsoventral [DV] -0.1 [39]. The AAVs were injected at the dose of 1.12×10^8 transducing unit (t.u.) in the anesthetized mice at a rate of $0.2 \mu\text{L}/\text{min}$ with a final volume of $1 \mu\text{L}$ for each side.

Behavioral procedures

The *open field* (OF) task was performed as described elsewhere [40]. Briefly, locomotion activity was assessed in a square divided into 4 equal arenas. At the start of a trial, the animals were placed in the center of each arena. The total distance traveled was measured under low light conditions by a video camera connected to a video tracking system (Ethovision Pro, Noldus).

The *Y-maze spontaneous alternation* (AYM) test was conducted to assess spatial working memory. Mice were placed randomly in one of the three arms of the Y-maze and were left free to explore the arena for 6 min. The number of arm entries and the number of triads were recorded in order to calculate the percentage of alternations to measure working memory.

The *elevated zero-maze* (EZM) was used to measure anxiety. It consists of a circular runway which is divided equally into two opposite open and two opposite enclosed arms. The mice were placed into one of the open arms and allowed to explore the maze over a period of 5 min. The total and relative duration (in %) and distance traveled in the open and enclosed arms were measured in the dark via an infrared video camera connected to a video tracking system (Ethovision Pro, Noldus). Percentage of time spent in the open arms was corrected for latency to first closed arm entry.

The *Y-maze spatial memory* test (SYM) was performed using the same arena as described in AYM above. One arm of the arena was closed by a removable blockade placed in front of it. The mice were placed in one of the open arms, which was randomized over the groups, and allowed to explore the 2 open arms of the maze for 5 min (pre-test). Afterward, the animal was taken from the arena and put back into its home cage. Five hours later, the mouse was placed back into its corresponding start arm of the arena, now with all three arms accessible (post-test). The previously blocked arm

was termed the “novel arm”. Memory was evaluated by calculating the amount of time spent in the novel arm corrected for the latency to move from the start arm to another arm and the amount of time the animal spent in the center of the maze [41].

Immunofluorescence staining

Mice were sacrificed by intracardial perfusion using Tyrode's solution for the first minute, followed by fixation solution 4% paraformaldehyde (PFA) for 10 min under deep sodium pentobarbital anesthesia (150 mg/kg). The brains were removed and post-fixed overnight in 4% PFA fixation solution and subsequently moved every 24 h in a buffer containing a gradually higher sucrose percentage: 10% and 20% sucrose in 0.1 M PBS. Afterwards, brains were quickly frozen using CO_2 and dissected into $16\text{-}\mu\text{m}$ -thick sagittal sections using a cryostat (at -25°C ; Leica). All series of sections were subsequently stored at -80°C until further processing. For the CERTs and neuron colocalization stain, we incubated the antibodies separately to reduce the antibody-antigen interaction. Before the antibodies incubation, the slice sections were fixed with acetone 10 min and blocked with 0.3% H_2O_2 for 1 h. The sections were incubated with a monoclonal NeuN primary antibody (1:50, chemicon international Inc., Temecula, CA, USA) overnight at 4°C . Sections were washed 3 times with Tris-buffered saline (TBS), TBS with 0.2% TritonX-100, and TBS. Subsequently, streptavidin Alexa 594 (1:500) applied for 1 h at room temperature. Then, rabbit polyclonal anti-CERTs (epitope 300–350, Bethyl Laboratories) diluted 1:250 was used to detect CERTs. After overnight incubation, and the corresponding secondary antibody Alexa Fluor-647 (1:100) was applied for 1 h at RT. The slices were mounted and stored in 4°C before taking pictures. Next immunofluorescence co-labeling was performed with either rabbit IgG anti-Iba1 (Wako Pure Chemical Corporation) or mouse IgG anti-glial fibrillary acidic protein (GFAP) combined with human IgG anti-A β [29]. Subsequently, the corresponding anti-rabbit or anti-mouse and anti-human secondary antibodies conjugated to Alexa Fluor-594 or 488 (Jackson ImmunoResearch Laboratories) were added for 2 h. Washes were performed 3 times for 10 min in TBS, TBS with 0.2% TritonX-100, and TBS, respectively in between the antibody incubation steps. Densitometric analysis of the stainings was performed on sagittal brain sections at different lateral depth (6–9 sections per animal) with ImageJ. Microglia ramification and sphericity were analyzed as described [42].

Sphingolipid analysis

High pressure liquid chromatography-tandem mass spectrometry (HPLC-MS/MS)

Neuro-2a (N2a) were maintained and prepared for HPLC-MS/MS as described in [Supplementary methods](#).

Powder aliquots of cortex, hippocampus, and cerebellum tissue or Neuro-2a (N2a) cell pellet were homogenized in PBS at the concentration of 10 μ L/mg. Then, 50 μ L of the brain preparation (or 25 μ L of plasma) were used to measure Cer, sphinganine (SPA), sphingosine (SPH), and sphingosine-1-phosphate (S1P) as previously described [43, 44]. Briefly, brain preparation (or plasma) was spiked with internal standards mixture prior to undergoing extraction. Data acquisition was done using select ion monitor (SRM) after chromatographic separation and electrospray ionization on the Thermo TSQ Quantum Ultra mass spectrometer (West Palm Beach) coupled with a Waters Acquity UPLC system (Milford) for Cer, sphinganine (SPA), sphingosine (SPH), and sphingosine-1-phosphate (S1P). SM data acquisition was done using multiple reaction monitoring after chromatographic separation and electrospray ionization on the Sciex Qtrap 5500 quadrupole mass spectrometer (AB Sciex Inc., Thornhill, Ontario, Canada) coupled with a Shimadzu HPLC system (Shimadzu, Kyoto, Japan). Concentrations of each analyte were calculated against each corresponding calibration curve and corrected for internal standard concentrations.

Protein extraction

Mice were terminally anesthetized with sodium pentobarbital, perfused, and the brain removed and dissected into the cortex, hippocampus, and cerebellum. Each brain region was then powdered in iron mortar partly emerged in liquid nitrogen, and aliquoted. Frozen tissue from dissected brains was sonicated in about 15 volumes (w/v) of TBS with PhosSTOP and protein inhibitors (Roche). Samples were centrifuged, and the TBS-soluble fraction was aliquoted prior to freezing in liquid nitrogen and stored at -80°C in aliquots. The pellet was resuspended by sonication for 10 s in about 15 volumes of TBS containing 1% Triton-X 100 (TBS-T) and protease inhibitor cocktail. Samples were centrifuged, and the TBS-T-soluble fraction and frozen in aliquots as described for the TBS fraction. The pellet was resuspended in 70% formic acid to 150 mg/ml based on tissue weight, and mixed by rotation at room temperature for 2 h. Samples were centrifuged, and the formic acid-soluble fraction was neutralized (with 20 volumes of 1 M Tris base) and frozen in aliquots at -80°C . Total protein content in the TBS and TBS-T extractions was determined with Bio-Rad DC (Life Science Group) protein assay following the manufacturer's instructions.

A β immunoassay and Western blot

Immunoassay

Microplates Microton/F-shape REF 655092 (Greiner) were coated with 1 μ g/mL of human 3D6 [29] overnight

at room temperature in coating buffer (sodium carbonate pH = 9.6 0.05 M NaCO_3 in MQ water). After washing plates (washing buffer 0.05% Tween- 20 in PBS) were blocked with 4% not fat dry milk and incubated with brain homogenates or with A β to generate the standard curve. Next plates were washed, incubated with 50 ng/mL of biotinylated human 20C2 [29], and washed again. Finally, plates were incubated with streptavidin-HRP (Jackson ImmunoResearch Laboratories) diluted 1:8000 and developed using 3,3',5,5'-Tetramethylbenzidine (TMB). The absorption was measured at 450 nm within 30 min of stopping the reaction with 2 M H_2SO_4 using the Perkin Elmer 2030 manager system.

Western blot

TBS and TBS-T samples corresponding to 40 μ g of total protein we loaded onto a precast TGX 4–16% gel (Bio-Rad). Then, samples were transferred to PDVF membranes, blocked with Odyssey blocking buffer (LI-COR Bioscience) and probed with 1 μ g/mL mAb anti-A β (6E10, Covance). After incubation with donkey anti-mouse IRdye680 (Rockland Immunochemicals) diluted 1:1000 in Odyssey blocking buffer, the membrane was scanned and analyzed with the Odyssey imager. The intensities of the APP bands detected at 100 kDa were measured with the Odyssey imager.

Reverse transcription polymerase chain reaction (RT-PCR)

Total RNA was isolated from 20 to 50 mg of cortex using Trizol reagent (Invitrogen). One microgram of total RNA was treated with DNase I and transcribed into cDNA (Superscript III, Invitrogen). PCR was performed in duplicate with SensiMix™ SYBR® Low-ROX Kit (Bioline) using the set of primers reported in supplementary Table 1. Fold changes of expression relative to control were determined after normalization to GAPDH and Actin. Fold change was calculated by the comparative Ct method [45].

Statistical analysis

Statistical analysis was performed using the Statistical Package for the Social Sciences (SPSS 23.0 SPSS Inc., Chicago, IL, USA). All graphs were designed in GraphPad Prism (version 5, GraphPad Software, San Diego, CA, USA). All data shown are expressed as mean $\pm$ standard error of the mean (SEM). Behavioral tasks, SLs, RT-PCR, and cytokines data were analyzed with two-way analysis of variances (ANOVA) with AAV treatment and genotype as independent factors. Least significant difference (LSD) was used for post hoc testing. Fluorescence amyloid- β aggregation assays were analyzed with repeated measure ANOVA or ANOVA followed by Dunnett's multiple comparisons test. Comparison of mean values from two groups was performed by an

unpaired two-tailed Student's *t* test or by Mann–Whitney *U* test for non-parametric testing. *P* values were considered as significant if $p \leq 0.05$ and marked with (*). Results were marked with (**) if $p \leq 0.01$ or (***) if $p \leq 0.001$.

Results

CERT_L directly affects A β aggregation and toxicity in vitro

It has been demonstrated that specific forms of CERTs can be released extracellularly or found membrane bound [30]. We have shown that CERTs can be found in proximity to A β plaques in AD brain, where they partially co-localize with SAP and with amyloid fibrils [29]. Here, we tested whether the long isoform of CERT, CERT_L, interacts with A β and its precursor protein APP. Cellular extracts of HEK-APP, expressing endogenous CERT_L, and stably expressing APP, were incubated with (mouse) monoclonal antibodies (mAbs) against CERT_L, APP, or syntaxin 6 (used as a negative control) for immunoprecipitation (IP). Pull-down was performed with anti-mouse antibodies for all 3 conditions. Western blot analysis of the samples showed efficient direct IP of CERT_L and APP by their respective antibodies. APP was detected as a band of ~100 kDa whereas CERT_L was detected as a band of ~200 kDa (Fig. 1a, left panel). As a result of co-immunoprecipitation, these two respective bands were also found when either CERT_L or APP were pulled down, but not when syntaxin 6 was used instead (Fig. 1a, right panel). The results, representative of three independent experiments, showed an interaction between CERT_L and APP/A β confirming our previous pull down of APP with CERT in the brain lysate of AD animals manifesting severe A β pathology [29]. The two proteins showed partial colocalization in primary neurons of 5xFAD mice at the plasma membrane and in the perinuclear region (Fig. 1b). This was also observed in brain sections of 5xFAD immunolabeled for CERTs and APP/A β (Supplementary Figure 1A). As shown by microscale thermophoresis (MST) analysis, CERT_L also bound to A β_{1-42} ($K_d = 2.5 \mu\text{M}$) (Fig. 1c) and by Western blot (Supplementary Figure 1B and C). Having shown that CERT_L directly binds to APP and A β , we next tested whether binding of CERT_L to A β_{1-42} could directly influence the spontaneous fibrillization of A β_{1-42} by thioflavin T (ThT) fluorescence spectrometry and TEM. In the absence of CERT_L, ThT fluorescence of A β_{1-42} peaked after 10–12 h indicating amyloid formation. With the addition of CERT_L (2.5 μM) A β_{1-42} maximum ThT fluorescence reduced of about 75% around 24 h. In contrast, at 1 μM concentration, CERT_L was ineffective at preventing A β_{1-42} fibrillization (Fig. 1d). Fibrillization was also blocked by A β antibodies (Supplementary Figure 1D).

In addition to the ThT fluorescence assay, we studied A β aggregation in the presence of CERT_L by TEM. The TEM images showed that amyloid-like fibrils were observed in both conditions with and without CERT_L in combination with A β . However, the amyloid structure identified in the presence of CERT_L was not linear in shape and the fibril width varied significantly (Fig. 1e, f) when compared to A β_{1-42} alone. The aggregates observed with CERT_L plus A β_{1-42} could be precursors to amyloid fibril formation, but not classic straight amyloid fibril formation as previously reported [46].

Furthermore, using SHSY-5Y cells, we examined the effect of CERT_L on cell viability when A β_{1-42} was present. The addition of oligomeric A β_{1-42} to the culture medium of SHSY-5Y cells resulted in a 38% decrease in viability after 24 h, as measured by MTT reduction relative to control conditions ($p < 0.01$). Interestingly, the simultaneous addition of CERT_L significantly ameliorated the toxic effect of A β_{1-42} (Fig. 1g). Our results indicate that CERT_L forms complexes with APP and A β . The interaction of CERT_L with A β affected spontaneous aggregation and toxicity of the peptide.

As aforementioned, CERTs are important regulators of cellular Cer and SM balance. For this reason, we investigated the effect of modulating CERT_L levels on SL composition in vitro. After 48-h cell transfection with pcDNA3.1 driving expression of CERT_L, SM d18:1/16:0 was significantly increased while Cer were unchanged (Fig. 1h). Furthermore, CERT_L overexpression did not affect cell viability (data not shown).

Ceramide species are increased in 5xFAD compared to WT animals depending on brain region and acyl chain length

The 5xFAD model carries 5 familial AD mutations. These mutations lead to a rapid increase of A β peptide production. By 6 weeks of age, mice display elevated levels of A β , amyloid deposits, and age-dependent amyloid pathology accompanied by increase of inflammatory marker levels in the CNS [47–50]. However, it is unknown if this model shows also an increase of Cer level in the brain as it has been reported in the brains of AD patients [13–16]. Sphingolipid species were determined with HPLC-MS/MS in the cortex, hippocampus, and cerebellum of 5xFAD and wild-type (WT) male mice at 25–26 weeks of age (Supplementary Table 1 reports complete analysis). The analysis showed a significant elevation of Cer d18:1/16:0 levels in the cortex and of sphinganine (SPA), S1P, Cer d18:1/16:0, Cer d18:1/18:1, Cer 18:d1/20:0, and Cer d18:1/22:0 in the hippocampus of the 5xFAD animals compared to WT mice (Fig. 2a, b). In the cerebellum, only

Fig. 1 CERT_L binds directly to APP and Aβ and reduces Aβ aggregation and toxicity in vitro. **a** Protein interaction detected using co-IP of APP and CERT_L in HEK-APP. The total cell lysate of HEK-APP cells (L) lane 1 and the total cell lysate of HEK-APP cells IP using APP/Aβ (lane 2), CERT_L (lane 3), and syntaxin (isotype control) (lane 4) antibodies were analyzed by Western blot. APP (1) and CERTs proteins were detected. The isotype control syntaxin protein was negative. Molecular weight markers are indicated (kDa). **b** Immunofluorescent staining showing co-localization of CERTs and APP/Aβ in primary neurons isolated from 5xFAD brains. Neuronal marker MAP 2 was used to immuno-label neuronal cells. DAPI was used for nuclei staining. Scale bar 5 μm. **c** CERT_L and Aβ₁₋₄₂ interaction was measured by microscale thermophoresis. The dissociation constant (K_d) calculated was $2.5 \pm 0.3 \mu M$. **d** Percentage of Thioflavin T (ThT) fluorescence intensity to detect Aβ₁₋₄₂ 20 μM aggregation in the absence or presence of recombinant CERT_L 1 or 2.5 μM at different time points. Each data point represents the percentage of mean fluorescent intensity of three wells (repeated measures ANOVA; Dunnett's multiple comparisons test $**p < 0.01$, $***p < 0.001$). **e, f** TEM analysis of 2 μM Aβ₁₋₄₂ aggregation in the absence and presence of 0.1 μM CERT_L showed a different aggregation pattern quantified by Aβ width (Student's *t* test $***p < 0.001$). **g** Measurement of cell metabolic activity of SH-SY5Y by MTT assay in cells incubated with medium alone (control) or medium containing 10 μM Aβ₁₋₄₂, Aβ₁₋₄₂, and 1 μM CERT_L, or CERT_L alone for 24 h. Graph bar expressed as means $\pm$ S.E.M % of control $N = 5-10$ (one-way ANOVA, Bonferroni correction $*p < 0.05$, $**p < 0.01$). **h** SM d18:1/16:0, SM d18:1/18:1, SM d18:1/18:0, SM d18:1/24:1, and SM d18:1/24:0 measured by HPLC-MS/MS in N2a cells after 48-h transfection with vector control or pcDNA-CERT_L. Graph bar expressed as means $\pm$ S.E.M % of control $N = 4/\text{group}$ (one-way ANOVA, Holm-Sidak's multiple comparisons test $***p < 0.001$)

S1P levels were found to be significantly higher in the 5xFAD animals compared to controls. These results indicate that at that specific age and disease stage of

the animals, the cortex and the hippocampus are more susceptible to increase of Cer levels while the cerebellum is less affected. This suggests that the

increase of Cer levels correlate with amyloid burden. In line, the cortex and hippocampus are the areas where the plaques are first reported to appear in this model [32]. Quantification of CERT levels in the cortex by immunoassay showed a significant reduction of the protein concentration in AD brains compared to controls at 25–26 weeks of age [51]. CERTs reduction was not observed in cerebellum (Fig. 2d). Our data suggest that SL metabolism is shifted towards increased production of Cer at different degrees depending on the brain region. Furthermore, CERT concentration was reduced in the cortex of 5xFAD mice compared to WT.

AAV-mediated neuronal expression of CERT_L in mouse brain

To test the hypothesis that increasing CERT_L levels in the cortex counteract the SL dysbalance in 5xFAD animals and Aβ formation, we generated AAV2 particles carrying the CERT_L cDNA sequence controlled by the neuron-specific synapsin promoter (Fig. 3a). The AAV-CERT_L was tested in vitro on cortical rat primary cell culture, proving to effectively transduce neurons (Supplementary Figure 2A). Next, 8-week-old WT animals underwent stereotactic surgery and AAV-CERT_L or AAV-control were injected in the layer V of the motor cortex (Supplementary Figure 2B). The transduction

efficiency was evaluated 1, 2, 6, and 12 weeks post-injection by immunohistochemistry (Supplementary Figure 1C-D) and by RT-PCR (Supplementary Figure 1E). The AAV-CERT_L was shown to effectively transduce neurons and expressed the CERT_L protein for at least 12 weeks. The AAVs were then injected in 5xFAD mice in a similar fashion. Mice underwent stereotactic surgery for CNS administration of AAV particles at 12–13 weeks of age and were monitored for 12 weeks. Layer V of the frontal cortex was targeted since the A β accumulation is the most severe there [32] and CERT levels are reduced in this mouse model. After 12 weeks CERT overexpression was confirmed by immunofluorescence (Fig. 3b) and by Western blot analysis (Fig. 3c, d). The immunolabeling of CERTs, with neuronal marker NeuN, showed colocalization in the layer V of the motor cortex. Relative quantification of CERTs levels in cortex homogenates by Western blot illustrated a significant increase of CERTs levels in AAV-CERT_L-treated groups (Fig. 3d).

5xFAD and WT mice injected with AAV-CERT_L and AAV-WT did not show behavioral abnormalities

At week 10 post-injection, mice were challenged with a behavioral test battery in the following sequence: open field (OF) for assessing locomotion activity, Y-maze spontaneous alternation (AYM), and SYM for assessing spatial memory and elevated zero-maze (EZM) for examining anxiety (Fig. 4a–e). It has been reported that 5xFAD mice exhibit changes in hippocampus-dependent spatial working memory by 16 to 24 weeks of age [32]. However, in our hands, no difference was found in the performance of 5xFAD compared to WT in locomotion (Fig. 4b), memory (Fig. 4c, e), and anxiety (Fig. 4d). Our data showed that the WT animals treated with the control virus performed above the 50% chance level in the AYM (Fig. 4c). Furthermore, in the OF, the AAV-CERT_L attenuated 5xFAD hyperactivity, even though 5xFAD treated with AAV-control did not perform significantly different from WT animals (Fig. 4b, two-way

Fig. 3 AAV-mediated neuronal expression of CERT_L in mouse brain. **a** The recombinant genomes of the two AAV-2 vectors. → Abbreviations: From left to right, ITR, inverted terminal repeats; hSYN1, human synapsin 1 gene promoter; CERT_L, cDNA sequence coding for ceramide transfer protein long isoform (hCERT_L, 1875 bp NP_005704.1); IRES, internal ribosome sequence for translation initiation; EGFP, cDNA coding for enhanced green fluorescent protein (GFP); WPRE, woodchuck hepatitis virus posttranscriptional control element; bGH, bovine growth hormone gene-derived polyadenylation site; TB, synthetic transcription blocker. **b** Representative images of immunofluorescent staining of cortical brain area from 5xFAD animals treated with AAV-control or AAV-CERT_L. Section was co-stain for CERTs protein (green), neuronal marker NeuN (red). Scale bar 200 μm and 50 μm. **c** Western blot showing band intensities of CERTs and GAPDH. **d** Relative quantification of CERTs levels normalized to GAPDH in cortical protein extract from WT and 5xFAD animals treated with AAV-control or AAV-CERT_L. Bars represent mean one representative experiment with 5–7 samples per group (Mann-Whitney test, **p* < 0.05)

ANOVA, interaction $F = 4.170$, $p = 0.0463$). Overall, these data suggest that no detrimental behavioral effects in the animals were observed after CERT_L overexpression.

AAV-CERT_L decreased Cer d18:1/16:0 and increased SM levels in the cortex

One of the main functions of CERTs is to shuttle Cer from the ER to the Golgi [23]. It has been reported that toxic increase of Cer species in the muscle can be attenuated by overexpressing hCERT cDNA [52]. HPLC-MS/MS analysis of brain cortex tissue revealed a significant reduction of Cer d18:1/16:0 level due to CERT_L overexpression ($p < 0.05$). This effect was not observed in WT animals but only in 5xFAD animals where Cer d18:1/16:0 level was significantly elevated ($p < 0.01$). This transport of Cer to the Golgi is crucial for the de novo synthesis of more complex SLs such as SM. Previous data from in vitro experiments reported that blocking CERTs function SM levels would significantly decrease [27]. Hence, if CERTs activity is enhanced, we expected an increase in SM levels. As reported above, we found that in vitro overexpression of CERT_L increased the levels of certain species of SM. SL analysis of the cortex showed that the levels of most of SM species (SM d18:1/16:0 $p < 0.001$, SM d18:1/18:0 $p < 0.001$, SM d18:1/18:1 $p < 0.01$, SM d18:1/20:0 $p < 0.05$, SM d18:1/22:0 $p < 0.05$, and SM d18:1/24:1 $p < 0.01$) were increased in AAV-CERT_L treated animals (Fig. 5). The only SM species not found significantly elevated was the SM d18:1/24:0, whose precursor Cer d18:1/24:0 has been reported to be poorly transferred by CERTs [53].

These data suggest that CERT_L overexpression is effective in reducing Cer increase by intensifying the transfer to the Golgi, which leads to an increase of SM. While the Cer attenuation was restricted to a pathological increase of Cer d18:1/16:0 level, SM elevation was consistent in all AAV-CERT_L-treated animals. This shift in SL composition did not affect apoptosis markers in the cortex quantified by RT-PCR (Supplementary Figure 6A).

AAV-CERT_L reduces A β levels by decreasing APP cleavage in the cortex of 5xFAD mice

Since our previous data indicated that CERT_L could be released to the extracellular milieu and directly affects A β aggregation and toxicity in vitro, we investigated the effect of CERT_L overexpression on A β deposition. (Fig. 6a). Statistical analysis showed no significant difference in plaque load between the 5xFAD groups at 24–26 weeks of age (Fig. 6b). However, the percentage of small plaques size (10–25 μ m) was reduced ($p < 0.05$) in AAV-CERT_L-treated 5xFAD brains (Fig. 6c). Furthermore, A β quantification of brain homogenate in TBS soluble and TBS-T soluble fraction showed that A β

levels were reduced in samples treated with AAV-CERT_L ($p = 0.04$ and $p = 0.03$, respectively), while in the formic acid soluble fraction no change was observed ($p = 0.29$) (Fig. 6d). Since it has been reported that APP cleavage can be affected by Cer composition [21, 22], we investigated if the reduction of Cer and increase of SM levels mediated by AAV-CERT_L is associated with altered processing of APP. The ratio A β /FL-APP was decreased in CERT_L overexpressing mice implying a reduction of A β biogenesis or increased clearance of A β ($p < 0.01$). Since the CTF β is the product of β -secretase cleavage of APP and the immediate precursor of A β formation [54], the ratio of CTF β /FL-APP bands' intensities was used to assess APP processing by β - and γ -secretase. The CTF β /FL-APP was found an increase in AAV-CERT_L treated 5xFAD animals ($p < 0.05$) (Fig. 6e). Meanwhile, the ratio A β /CTF β was reduced in brains of CERT_L overexpressing mice ($p < 0.01$) (Fig. 6e and Supplementary Figure 5). These results indicate that AAV-CERT_L affects the proteolytic processing of APP by β - and/or γ -secretase.

These results show that a specific balance of Cer to SM and/or interaction of CERT_L with APP is critical for APP cleavage and A β biogenesis.

Four weeks of administration of CERTs inhibitor (HPA-12) exacerbates Cer and A β pathology in AD transgenic mice

To test if efficient Cer trafficking from the ER to the Golgi is vital in the regulation of the Cer levels and A β formation, we administered the CERTs inhibitor N-(3-hydroxy-1-hydroxymethyl-3-phenylpropyl) dodecanamide (HPA-12) for 4 weeks to AD transgenic mice. As expected Cer d18:1/16:0, Cer d18:1/20:0, Cer d18:1/22:0, and Cer d18:1/24:1 levels were found an increase in the brain (Fig. 7a). Moreover, A β levels were increased by 117% in the TBS soluble fraction ($p < 0.05$) and by 47% in the TBS-T soluble fraction of brain homogenates ($*p < 0.05$) of CERTs inhibitor-treated AD animals. No significant changes were found in the FA insoluble fraction (Fig. 7b).

These data suggest that efficient Cer transfer from ER to *trans*-Golgi is critical to control Cer levels and thereafter APP processing. Pharmacological interference with CERT activity augments Cer levels and increases substantially A β biogenesis and/or fibrillization.

AAV-CERT_L reduces microglia immunoreactivity as shown by Iba1 labeling and CD86 expression

We have observed that CERTs are associated with A β plaques in AD brain where microglia cells are engaged [29]. Thus, we investigated if microglia cells were affected by CERT_L overexpression in the cortex.

Fig. 4 No behavioral abnormalities 10 weeks after injection of AAV-CERT_L. **a** The effects of CERT_L over-expression were investigated in 30 5xFAD and 30 WT males. Mice were bilaterally injected at 12–13 weeks of age with AAV-CERT_L or AAV-control particles at the dose 1.12×10^8 transducing unit (t.u.). Starting at week 22 of age, animals were challenged with the following behavioral tests: open field (OF) for locomotion activity, alternate Y-maze (AYM) and spatial Y-maze (SYM) for spatial memory, and elevated zero-maze (EZM) for anxiety. **b** Locomotion expressed as distance traveled in OF task. **c** The graph shows the results of the working memory in the AYM task as a percentage of correct alternation in the first four triads. Percentage were compared to 50% chance levels (one sample *t* test $**p < 0.01$). **d** Anxiety was assessed, measuring the percentage of time spent in the closed arm in EZM. **e** Memory was measured in SYM expressed as a percentage of time spent in the novel arm. Bars represent the means $\pm$ S.E.M per group $N = 10$ –20 (two-way ANOVA, interaction effect $F = 4.170$ $p = 0.0463$, LSD, $*p < 0.05$)

To achieve this aim, brain sections were analyzed for Iba1 reactivity. AAV-CERT_L-treated brains had a decreased immunoreactivity to Iba1 (percentage of area $p < 0.001$) (Fig. 8a–c). Iba1 is considered a constitutive marker for microglia, which is highly increased in 5xFAD animals compared to WT. [55, 56] Furthermore, it has been shown to be important for membrane ruffling, a process crucial for macrophage and microglia motility and chemotaxis [57]. We further characterized microglia based on ramifications and sphericity. The analysis showed that microglia of

AAV-CERT_L-injected mice had longer ramifications and lower sphericity index (Fig. 8d, e). To investigate if AAV-CERT_L affected other microglia membrane markers, we analyzed cortex tissue by RT-PCR. We quantified the CD86 membrane marker, which is an indicator for microglia pro-inflammatory polarization. Statistical analysis showed that CD86 was increased in AAV-control-treated 5xFAD animals compared to AAV-control-treated WT and that AAV-CERT_L specifically decreased CD86 in the 5xFAD group (Fig. 8f).

Fig. 5 AAV-CERT_L reduces Cer d18:1/16:0 and increases sphingomyelin species in the cortex. Sphingolipids levels were measured in the cortex by HPLC-MS/MS. Ceramides were classified based on acyl chain number of carbons (Cer d18:1/16:0, Cer d18:1/18:0, Cer d18:1/18:1, Cer d18:1/20:0, Cer d18:1/22:0, and Cer d18:1/24:1) as well as sphingomyelin (SM d18:1/16:0, SM d18:1/18:0, SM d18:1/18:1, SM d18:1/20:0, SM d18:1/22:0, and SM d18:1/24:1). Ceramide levels were expressed as ng/mg tissue, while sphingomyelins were expressed as pmol/mg tissue. Bars represent the mean $\pm$ S.E.M per group $N = 5-12$ (two-way ANOVA, LSD, significant effects, * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$)

Next, we analyzed cytokines levels in brain homogenate using multiplex technology. The assay did not reveal any significant effect of AAV-CERT_L treatment. However, we observed a significant increment of IL-1 β ($p < 0.01$) and a significant reduction of IL-4 ($p < 0.05$) when comparing WT to 5xFAD animals in AAV-control treated groups (Supplementary Figure 6A).

These results indicate overall that AAV-CERT_L decreases pro-inflammatory processes in microglia.

AAV-CERT_L does not change the immunoreactivity of astrocytes as shown by GFAP labeling in the cortex of 5xFAD mice

Reactive astrocytes have been described as a pathological hallmark that generally occurs in response to neurodegeneration in AD [58]. To determine the extent of astrogliosis in the 5xFAD mouse brain treated with AAV-CERT_L compared to control, we performed immunofluorescence GFAP labeling on sagittal brain sections (Fig. 8g). Densitometric and

particle analysis performed with ImageJ showed a 47% reduction in astrocyte immunoreactivity, which was not statistically significant (Fig. 6h). However, as previously reported, we observed a 3-fold increase of GFAP immunoreactivity in the 5xFAD animals compared to WT controls [32, 59].

Discussion

In this work, we provide evidence suggesting that CERT_L plays an important role in characteristic processes of AD by affecting A β production and aggregation, neuroinflammation, and SL disbalance typical of AD. Our data show that the pathological increase of Cer d18:1/16:0 can be reduced to normal levels by upregulating CERT_L. The reduction of Cer levels also proved to be effective in attenuating A β formation. Furthermore, CERT_L overexpression in neurons revealed that CERT_L can downregulate membrane markers indicating a pro-inflammatory status in microglia.

Fig. 6 Neuronal increase of CERT_L reduces Aβ by decreasing APP cleavage. **a** Representative photomicrographs of sagittal brain sections imaging the motor sensory cortex (M1 and M2) stained for nuclei in blue and Aβ plaques in green. All photomicrographs were exposed and processed identically. Scale bar represents 200 and 50 μm (from right to left). **b** Immunofluorescent quantification of plaques measured by the percentage of area, plaque counts/mm². **c** Frequency distribution of plaques based on size (10–25 μm) (error bars represent ± SEM of 4–6 animals per experimental condition, ANOVA, Bonferroni correction, significant effects, **p* < 0.05; ***p* < 0.01). **d** Aβ quantification in three extraction buffers, BS, TBS-T, and formic acid (FA) by ELISA showed that Aβ was significantly reduced in the soluble fractions in the cortex but not in the insoluble fraction (Student's *t* test **p* < 0.05). **e** Western blot analysis of TBS cortex homogenate stained with 6E10 antibody showed that ratios of amyloid Aβ/FL-APP and Aβ/CTFβ are reduced while CTFβ/FL-APP is increased in AAV-CERT_L-treated animals while CTFβ/FL-APP is increased. Error bars represent ± SEM of 5 animals per experimental condition (Student's *t* test **p* < 0.05; ***p* < 0.01) (full length amyloid precursor protein = FL-APP; amyloid-β peptide = Aβ; C-terminal fragment β = CTFβ). Western blot membranes are shown in Supplementary Figure 5

It has previously been reported that APP can interact with extracellular matrix proteins like collagen I [60, 61]. This interaction is thought to be important in neuronal cell to cell adhesion with APP functioning as an anchor [62]. CERT_L is known to be crucial in stabilizing the basal membrane [63]. We previously reported that CERTs can be found in close location to plaques in AD brains, where they co-localize with amyloid fibrils [29]. In line with this, we found that APP can interact with CERT_L and form complexes that can be immunoprecipitated. Also Aβ can bind to a variety of biomolecules, including lipids, proteins, and proteoglycans [64]. Here, we show that CERT_L binds directly to Aβ peptides and that this interaction affects Aβ fibrillization by organizing Aβ into less neurotoxic aggregates.

Manipulation of CNS SL metabolism can be challenging. Removal or addition of genes encoding for enzymes and transporters in the SL pathway can be

deleterious for brain function [65, 66]. A previous report where human acid sphingomyelinase was increased in the brain of rodents and primates showed that motor function could be severely affected [67]. In contrast, in our study, behavioral testing at 9–10 weeks post AAV particle injection revealed no detectable side effects when comparing AAV-CERT_L- to AAV-control-treated animals in various memory- and anxiety-related behavioral tasks. Of note though, we found a significant reduction of locomotor activity in the 5xFAD due to AAV-CERT_L treatment. It has been reported that 5xFAD animals exhibit a hyperactive behavior compared to control animals even though this specific behavior is not well understood and translation to human symptoms is unclear [68]. However, the question whether AAV-CERT_L could improve memory or other behavioral deficits observed in the 5xFAD model could not be answered by our data. No significant impairments in spatial memory

(See figure on previous page.) **Fig. 8** AAV-CERT_L reduces microglia reactivity to Iba1 and CD86 expression levels but has no significant effect on GFAP immunoreactivity in the cortex of 5xFAD mice. **a** Representative photomicrograph of Iba1 staining in the cortical motor sensory region of 5xFAD animals treated with AAV-control or AAV-CERT_L (scale bar 50 μ m). **b** Densitometric analysis of Iba1 staining represented as a percentage of the area (AAV-control $n = 6$ and AAV-CERT_L $n = 4$ for WT and 5xFAD groups). **c** Densitometric analysis of Iba1 staining represented as number of positive Iba1 cells/mm² (AAV-control $n = 6$ and AAV-CERT_L $n = 4$ for 5xFAD groups). **d** Length of microglia ramification and sphericity per cell in AAV-control or AAV-CERT_L. Morphological analysis was performed on 3–5 pictures/group. **e** Illustrations of the microglia morphological analysis applied to a fluorescent photomicrograph captured with $\times 60$ objective with a single cell cropped to show details. Scale bar = 20. **f** Analysis of gene expression of membrane markers CD86 (4–5 number of animals per group). **g** Representative photomicrographs of GFAP staining in the cortical motor sensory region of 5xFAD animals treated with AAV-control or AAV-CERT_L (scale bar 50 μ m). **h** Densitometric analysis of GFAP staining represented as a percentage of the area (AAV-control $n = 6$ and AAV-CERT_L $n = 4$ for WT and 5xFAD groups). Bars represent the mean $\pm$ S.E.M per group (two-way ANOVA, LSD, significant effects, * $p < 0.05$; ** $p < 0.01$)

were detected in the AYM and the SYM tests. In the original 5xFAD line (on a hybrid B6SJL background and carrying the Pde6b gene), Jawhar et al. reported deficits of working memory, assessed by a cross-maze test, to appear at 6 months [69]. In contrast, the behavioral phenotype of the 5xFAD line on a C57BL/6J background and without the Pde6b gene, are much less clear. Richard et al. reported in this last line of 5xFAD spatial memory deficits were only visible after 7 months of age [70]. Others reported spatial memory impairment at an earlier age when using the water maze but with a very small difference between WT and 5xFAD [71, 72]. In the present study, none of the groups tested in the SYM test performed above chance level indicating lack of recognition in all groups examined. However, in the AYM, the WT group treated with AAV-control performed above chance level (mean of 68%), which in the AYM is considered to be 50%. As such, future studies should additionally assess shorter inter-trial intervals in this respect.

Cer generation is abnormal in AD, causing an increase in Cer formation [15, 73]. Furthermore, an accumulating body of evidence consistently reported a global rise in Cer levels in specific brain regions of AD patients [13, 74, 75]. In agreement with these observations, we found that 5xFAD transgenic mice at 6 months of age also showed an increase of Cer d18:1/16:0 in the cortex and of several Cer species in the hippocampus (Cer d18:1/16:0, Cer d18:1/18:1, Cer d18:1/20:0 and Cer d18:1/22:0). Previously, other studies showed increased brain Cer levels in different transgenic AD models (APP, PS1, and PS1-APP mutated mice) [76]. This suggests that there is a causal relationship between amyloid pathology and Cer imbalance.

In this study, we report that increasing Cer trafficking from the ER to the Golgi by overexpressing CERT_L reversed the pathological increase of Cer in the cortex. A similar effect of CERTs overexpression in Cer elevation state has been reported in a lipotoxic mouse model where muscle cells in overload Cer status were rescued by increasing the expression of hCERT [52]. In AD brains, we observed that CERT_L reduced 11% of the total Cer content restoring it close to normal levels. In

particular, Cer d18:1/16:0 was the most affected species by AAV-CERT_L being reduced up to 35%. These results demonstrate the importance of physiological ER-to-Golgi Cer traffic in preserving the physiological balance of Cer levels in AD pathology.

The soluble A β forms were reduced in the CERT_L-treated animals, whereas the insoluble A β forms were not altered by the treatment. This reduction in the soluble forms could be explained in two ways: (i) the 11% reduction of total Cer due to CERT_L overexpression and (ii) CERT_L interaction with A β . As aforementioned, the amyloidogenic cleavage of APP is favored resulting in more A β formation in Cer enriched conditions [21, 22]. In our study, the total Cer reduction may have affected the secretases activity in the opposite way. We found that the A β /APP ratio, which describes the APP processing to form A β , was lower in AAV-CERT_L animals implying that lesser APP is processed to generate the A β peptide. Importantly, APP processing takes place in different cell compartments not only on the cell surface [77] and the Cer shift is not confined to the ER compartment but can affect the whole cell [78, 79]. This conclusion was further confirmed by pharmacological inhibition of CERT Cer transfer activity with CERT inhibitor HPA-12. Recently, the pharmacokinetics of HPA-12 was described, and it was proven that the compound reaches the brain intact [25]. Here, we found that after 4 weeks, the treatment of HPA-12 increases Cer and A β levels.

It is now thought that one of the crucial processes in the development and exacerbation of AD is neuroinflammation [80]. Our lab demonstrated that CERT_L interacts with SAP which belongs to the pentraxin family of the innate immune system [29]. Additionally, we reported that CERT_L can activate the complement system [31]. Here, we found that AAV-CERT_L influenced microglia activation even though CERT_L was specifically expressed in neurons under the control of the synapsin promoter. It has been consistently reported that 5xFAD microglia are polarized towards a more pro-inflammatory status, in response to the extensive plaque formation. Consequently, the Iba1 microglia marker is highly expressed in AD models [55, 56]. Our

findings suggest that CERT_L could play a role in the cross-talk between neurons and microglia. Interestingly, neuronal-derived CERT_L activity is exerted only when there is an inflammatory reaction ongoing by reducing membrane markers for the pro/inflammatory status of microglia. Nevertheless, it remains unclear by which mechanism AAV-CERT_L decreased Iba1- and CD86-positive cells. Here, we propose two hypotheses. First, the reduction of Iba1 and CD86 is a direct action of CERT_L on microglia activation status once secreted by neurons. It is known that forms of CERT_L can be released in the extracellular space [30]. The second, Iba1 and CD86 are decreased because of a modified by shifts in Cer and SM composition or other indirect effects like reduction of Aβ levels. In the CNS, there is an extensive cross-talk ongoing between neurons and microglia, which takes advantage of lipid vesicles.

Furthermore, toxic Aβ has also been reported among the content of exosome and reduction of exosome secretion was correlated to Aβ reduction [81, 82].

Similarly, to microglia, also astrocyte activation was reduced by AAV-CERT_L, even though not significantly. During inflammation astrocytes are enriched in Cer. They seem to produce the pro-apoptotic Cer d18:1/16:0 [83, 84]. Further, reactive astrocytes release extracellular vesicle enriched in Cer that carry Aβ peptides [84]. These specific extracellular vesicles isolated from brains of 5xFAD mice showed to be particularly toxic for neurons.

Limitations

The limitations of this study are three folded. First, we did not detect memory difference between WT and 5xFAD. This could be due to our in-house breeding

approach, which is explained in material and methods section. We adopted a breeding scheme where the Pde6b gene, which is associated to retinal degeneration, was bred out of the genetic background of the 5xFAD transgenic mice. The Pde6b gene is thought to be crucial for early detection of memory impairments. The second limitation is connected to the first. Since no memory impairment was detected, it is unclear if the AAV-CERT_L would protect from memory decline. For this purpose, animals could be tested at older age and the AAV-CERT_L could be injected in the hippocampus. This follow-up study would be of interest also because the hippocampus, as we show for the first time in this study, is particularly affected by the Cer pathology in 5xFAD males. The third limitation is the effect of AAV-CERT_L on astrocytes. While we had sufficient statistical power to detect the microglia changes, astrocytes denoted a similar trend, which was not statistically significant. These three limitations set the ground for future studies.

Conclusion

In conclusion, by increasing CERT_L expression in neuronal cells, we were able to increase SM production and reduce Cer d18:1/16:0 especially in the CNS. Next, after proving that CERT_L binds and modifies A β aggregation in vitro, we observed that administration of AAV-CERT_L in AD animals reduced A β production by at least 2 mechanisms: by altering SL composition and by direct interaction with APP in 5xFAD animals. Moreover, we reported a new immune role of CERT_L. AAV-CERT_L decreased membrane markers important for the pro/inflammatory status of microglia. Overall, our experiments are the first to demonstrate that an increase of CERT_L modulates SL levels and affects amyloid plaque formation and brain inflammation in AD (see the model in Fig. 9). These data open research pathways for therapeutic targets of AD and related neurodegenerative diseases.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13195-021-00780-0>.

Additional file 1: Supplementary methods.

Additional file 2: Supplementary Tables.

Additional file 3: Supplementary Figures.

Abbreviations

AD: Alzheimer's disease; A β : Amyloid- β peptides; NFTs: Neurofibrillary tangles; BBB: Blood-brain barrier; SLs: Sphingolipids; Cer: Ceramide; S1P: Sphingosine-1-phosphate; SM: Sphingomyelin; CERTs: Ceramide transfer proteins; CERT_L: Ceramide transfer proteins long form; START: Steroidogenic acute regulatory protein (StAR)-related lipid transfer; ER: Endoplasmic reticulum; AAV: Adeno-associated virus; IP: Immunoprecipitation; APP: Amyloid precursor protein; FL-APP: Full length amyloid precursor protein; CTFB: C-terminal fragment- β ; MST: Microscale thermophoresis; TEM: Transmission

electron microscopy; ThT: Thioflavin T; OF: Open field; AYM: Y-maze spontaneous alternation; EZM: Elevated zero-maze; SYM: Y-maze spatial memory test; HPLC-MS/MS: High pressure liquid chromatography-tandem mass spectrometry; (mAbs): Monoclonal antibodies; (HPA-12): N-(3-Hydroxy-1-hydroxymethyl-3-phenylpropyl) dodecanamide

Acknowledgements

We thank Prof. S. Kugler Department of Neurology, University of Gottingen, for the kind gift of adeno-associated virus plasmid. We thank Geertjan van Zonneveld for the professional approach and dedication to complete the illustrations of this research paper.

Authors' contributions

PMM and ML conceived the project. PMM, ML, SMC, QL, CG, JS, DVK, and GB designed and performed the animal behavior, biochemical, and molecular biological experiments with critical input from ML, DVH, HEV, MDL, and PMM. JV and BH generated the adeno-associated virus particles. CH and MM performed the electron microscopy. MM, MTM, and SVH performed lipid mass spectrometry analyses. ALS and JW analyzed APP processing and contributed important reagents and cell models. DB synthesized and provided the CERTs inhibitor. BE helped to perform and interpreted the cytokine measurements in the brain homogenate. WK helped experiment design and data interpretation of neuroinflammation. The manuscript was written by SMC and PMM. All authors contributed by critically revising the manuscript. The author(s) read and approved the final manuscript.

Funding

This work was supported by grants to NMdW, SdH, MTM, JW, AR, PMM, JV, and HEV from ZonMw Memorabel program (projectnr: 733050105). PMM is also supported by the International Foundation for Alzheimer Research (ISAO) (projectnr: 14545). SMC received a travel grant support from Alzheimer Netherlands foundation (AN) to spend a month in the laboratory of EB, University of Kentucky, Lexington KY, USA. Aspects of this work were supported by the grants NIH R01AG034389, R01NS095215, and R56AG064234; VA I01BX003643 to EB. MMM was supported by R01AG049704.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

All experiments were approved by the Animal Welfare Committee of Maastricht University (project number DEC2013-056 and DEC2015-002) and followed the laws, rules, and guidelines of the Netherlands.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Psychiatry and Neuropsychology, School for Mental Health and Neuroscience, Maastricht University, Universiteitssingel 50, 6229 ER Maastricht, the Netherlands. ²Department of Physiology, University of Kentucky College of Medicine, Lexington, KY, USA. ³Veterans Affairs Medical Center, Lexington, KY 40502, USA. ⁴Department of Internal Medicine, Laboratory Vascular Medicine, Erasmus MC University Medical Center, Rotterdam, the Netherlands. ⁵Laboratory for Neuroregeneration, Netherlands institute for Neuroscience, Amsterdam, the Netherlands. ⁶Department of Neurology, University Hospital Bonn, University of Bonn, Bonn, Germany. ⁷The Birchall Centre, Lennard-Jones Laboratories, Keele University, Staffordshire, UK. ⁸Division of Epidemiology, Department of Health Science Research, and Department of Neurology, Mayo Clinic Rochester, Rochester, MN, USA. ⁹Department of Molecular Cell Biology and Immunology, Amsterdam Neuroscience, Amsterdam UMC, Amsterdam, the Netherlands. ¹⁰Department of Internal Medicine, Maastricht University Medical Centre, Maastricht, the Netherlands. ¹¹Cardiovascular Research Institute Maastricht (CARIM), Maastricht, the Netherlands. ¹²Department of Psychiatry, Psychosomatics and Psychotherapy, University of Wuerzburg, Wuerzburg,

Germany. ¹³Department of Organic Chemistry, Slovak University of Technology, Radlinského 9, 81237 Bratislava, Slovak Republic. ¹⁴Department of Molecular Neuropathology, Centro de Biología Molecular "Severo Ochoa" (CSIC-UAM), Madrid, Spain.

Received: 10 July 2020 Accepted: 31 January 2021

Published online: 17 February 2021

References

1. Rogers J, Morrison JH. Quantitative morphology and regional and laminar distributions of senile plaques in Alzheimer's disease. *J Neurosci*. 1985;5(10):2801–8.
2. Kawasaki H, Murayama S, Tomonaga M, Izumiya N, Shimada H. Neurofibrillary tangles in human upper cervical ganglia. Morphological study with immunohistochemistry and electron microscopy. *Acta Neuropathol*. 1987;75(2):156–9.
3. Wisniewski HM, Kozlowski PB. Evidence for blood-brain barrier changes in senile dementia of the Alzheimer type (SDAT). *Ann N Y Acad Sci*. 1982;396:119–29.
4. Krabbe G, Halle A, Matyash V, Rinnenthal JL, Eom GD, Bernhardt U, et al. Functional impairment of microglia coincides with Beta-amyloid deposition in mice with Alzheimer-like pathology. *PLoS One*. 2013;8(4):e60921.
5. Crivelli SM, Giovagnoni C, Visseren L, Scheithauer AL, de Wit N, den Hoedt S, et al. Sphingolipids in Alzheimer's disease, how can we target them? *Adv Drug Deliv Rev*. 2020.
6. Hannun YA, Obeid LM. Principles of bioactive lipid signalling: lessons from sphingolipids. *Nat Rev Mol Cell Biol*. 2008;9(2):139–50.
7. Zhang Y, Li X, Becker KA, Gulbins E. Ceramide-enriched membrane domains—structure and function. *Biochim Biophys Acta*. 2009;1788(1):178–83.
8. Schwarz A, Futerman AH. Distinct roles for ceramide and glucosylceramide at different stages of neuronal growth. *J Neurosci*. 1997;17(9):2929–38.
9. Zhang H, Desai NN, Olivera A, Seki T, Brooker G, Spiegel S. Sphingosine-1-phosphate, a novel lipid, involved in cellular proliferation. *J Cell Biol*. 1991;114(1):155–67.
10. Bartke N, Hannun YA. Bioactive sphingolipids: metabolism and function. *J Lipid Res*. 2009;50(Suppl):S91–6.
11. Perry DK, Hannun YA. The role of ceramide in cell signaling. *Biochim Biophys Acta*. 1998;1436(1–2):233–43.
12. van Meer G, Holthuis JC. Sphingolipid transport in eukaryotic cells. *Biochim Biophys Acta*. 2000;1486(1):145–70.
13. Cutler RG, Kelly J, Storie K, Pedersen WA, Tammara A, Hatanpaa K, et al. Involvement of oxidative stress-induced abnormalities in ceramide and cholesterol metabolism in brain aging and Alzheimer's disease. *Proc Natl Acad Sci U S A*. 2004;101(7):2070–5.
14. Mielke MM, Bandaru VV, Haughey NJ, Xia J, Fried LP, Yasar S, et al. Serum ceramides increase the risk of Alzheimer disease: the Women's Health and Aging Study II. *Neurology*. 2012;79(7):633–41.
15. He X, Huang Y, Li B, Gong CX, Schuchman EH. Deregulation of sphingolipid metabolism in Alzheimer's disease. *Neurobiol Aging*. 2010;31(3):398–408.
16. de Wit NM, Snkhchyan H, den Hoedt S, Wattimena D, de Vos R, Mulder MT, et al. Altered sphingolipid balance in capillary cerebral amyloid angiopathy. *J Alzheimers Dis*. 2016.
17. Fabelo N, Martín V, Marin R, Moreno D, Ferrer I, Díaz M. Altered lipid composition in cortical lipid rafts occurs at early stages of sporadic Alzheimer's disease and facilitates APP/BACE1 interactions. *Neurobiol Aging*. 2014;35(8):1801–12.
18. Lee JT, Xu J, Lee JM, Ku G, Han X, Yang DI, et al. Amyloid-beta peptide induces oligodendrocyte death by activating the neutral sphingomyelinase-ceramide pathway. *J Cell Biol*. 2004;164(1):123–31.
19. Malaplate-Armand C, Florent-Bechard S, Youssef I, Koziel V, Sponne I, Kriem B, et al. Soluble oligomers of amyloid-beta peptide induce neuronal apoptosis by activating a cPLA2-dependent sphingomyelinase-ceramide pathway. *Neurobiol Dis*. 2006;23(1):178–89.
20. Isobe C, Abe T, Terayama Y. Levels of reduced and oxidized coenzyme Q-10 and 8-hydroxy-2'-deoxyguanosine in the CSF of patients with Alzheimer's disease demonstrate that mitochondrial oxidative damage and/or oxidative DNA damage contributes to the neurodegenerative process. *J Neurol*. 2010;257(3):399–404.
21. Pugliese L, Ellis BC, Saunders AJ, Kovacs DM. Ceramide stabilizes beta-site amyloid precursor protein-cleaving enzyme 1 and promotes amyloid beta-peptide biogenesis. *J Biol Chem*. 2003;278(22):19777–83.
22. Takasugi N, Sasaki T, Shinohara M, Iwatsubo T, Tomita T. Synthetic ceramide analogues increase amyloid-beta 42 production by modulating gamma-secretase activity. *Biochem Biophys Res Commun*. 2015;457(2):194–9.
23. Hanada K, Kumagai K, Yasuda S, Miura Y, Kawano M, Fukasawa M, et al. Molecular machinery for non-vesicular trafficking of ceramide. *Nature*. 2003;426(6968):803–9.
24. Mencarelli C, Losen M, Hammels C, De Vry J, Hesselink MK, Steinbusch HW, et al. The ceramide transporter and the Goodpasture antigen binding protein: one protein—one function? *J Neurochem*. 2010;113(6):1369–86.
25. Crivelli SM, Paulus A, Markus J, Bauwens M, Berkes D, De Vries HE, et al. Synthesis, radiosynthesis, and preliminary in vitro and in vivo evaluation of the fluorinated ceramide trafficking inhibitor (HPA-12) for brain applications. *J Alzheimers Dis*. 2017;60(3):783–94.
26. Wang X, Rao RP, Kosakowska-Cholody T, Masood MA, Southon E, Zhang H, et al. Mitochondrial degeneration and not apoptosis is the primary cause of embryonic lethality in ceramide transfer protein mutant mice. *J Cell Biol*. 2009;184(1):143–58.
27. Yasuda S, Kitagawa H, Ueno M, Ishitani H, Fukasawa M, Nishijima M, et al. A novel inhibitor of ceramide trafficking from the endoplasmic reticulum to the site of sphingomyelin synthesis. *J Biol Chem*. 2001;276(47):43994–4002.
28. Yamaji T, Hanada K. Establishment of HeLa cell mutants deficient in sphingolipid-related genes using TALENs. *PLoS One*. 2014;9(2):e88124.
29. Mencarelli C, Bode GH, Losen M, Kulharia M, Molenaar PC, Veerhuis R, et al. Goodpasture antigen-binding protein/ceramide transporter binds to human serum amyloid P-component and is present in brain amyloid plaques. *J Biol Chem*. 2012;287(18):14897–911.
30. Revert F, Ventura I, Martinez-Martinez P, Granero-Molto F, Revert-Ros F, Macias J, et al. Goodpasture antigen-binding protein is a soluble exportable protein that interacts with type IV collagen. Identification of novel membrane-bound isoforms. *J Biol Chem*. 2008;283(44):30246–55.
31. Bode GH, Losen M, Buurman WA, Veerhuis R, Molenaar PC, Steinbusch HW, et al. Complement activation by ceramide transporter proteins. *J Immunol*. 2014;192(3):1154–61.
32. Oakley H, Cole SL, Logan S, Maus E, Shao P, Craft J, et al. Intraneuronal β -amyloid aggregates, neurodegeneration, and neuron loss in transgenic mice with five familial Alzheimer's disease mutations: potential factors in amyloid plaque formation. *J Neurosci*. 2006;26(40):10129–40.
33. Flührer R, Multhaup G, Schlicksupp A, Okochi M, Takeda M, Lammich S, et al. Identification of a beta-secretase activity, which truncates amyloid beta-peptide after its presenilin-dependent generation. *J Biol Chem*. 2003;278(8):5531–8.
34. Wang G, Silva J, Dasgupta S, Bieberich E. Long-chain ceramide is elevated in presenilin 1 (PS1M146V) mouse brain and induces apoptosis in PS1 astrocytes. *Glia*. 2008;56(4):449–56.
35. House E, Collingwood J, Khan A, Korchazkina O, Berthon G, Exley C. Aluminium, iron, zinc and copper influence the in vitro formation of amyloid fibrils of A beta (42) in a manner which may have consequences for metal chelation therapy in Alzheimer's disease. *J Alzheimers Dis*. 2004;6(3):291–301.
36. Mold M, Ouro-Gnao L, Wieckowski BM, Exley C. Copper prevents amyloid-beta (1–42) from forming amyloid fibrils under near-physiological conditions in vitro. *Sci Rep*. 2013;3:1256.
37. Kugler S, Lingor P, Scholl U, Zolotukhin S, Bahr M. Differential transgene expression in brain cells in vivo and in vitro from AAV-2 vectors with small transcriptional control units. *Virology*. 2003;311(1):89–95.
38. Verhaagen J, Hobo B, Ehler EME, Eggers R, Korecka JA, Hoyng SA, et al. Small scale production of recombinant adeno-associated viral vectors for gene delivery to the nervous system. *Methods Mol Biol*. 2018;1715:3–17.
39. Franklin KB, Paxinos G, editors. The mouse brain in stereotaxic coordinates. San Diego: Academic Press; 1997.
40. Prickaerts J, Moechars D, Cryns K, Lenaerts I, van Craenendonck H, Goris I, et al. Transgenic mice overexpressing glycogen synthase kinase 3beta: a putative model of hyperactivity and mania. *J Neurosci*. 2006;26(35):9022–9.
41. Sierksma AS, Prickaerts J, Chouliaras L, Rostamian S, Delbroek L, Rutten BP, et al. Behavioral and neurobiological effects of prenatal stress exposure in male and female APPswe/PS1dE9 mice. *Neurobiol Aging*. 2013;34(1):319–37.
42. Young K, Morrison H. Quantifying microglia morphology from photomicrographs of immunohistochemistry prepared tissue using ImageJ. *J Vis Exp*. 2018;136.
43. Blachnio-Zabielska AU, Persson XM, Koutsari C, Zabielski P, Jensen MD. A liquid chromatography/tandem mass spectrometry method for measuring the in vivo incorporation of plasma free fatty acids into intramyocellular ceramides in humans. *Rapid Commun Mass Spectrom*. 2012;26(9):1134–40.

44. Hoogendoorn A, den Hoedt S, Hartman EMJ, Krabbendam-Peters I, Te Lintel HM, van der Zee L, et al. Variation in coronary atherosclerosis severity related to a distinct LDL (low-density lipoprotein) profile: findings from a familial hypercholesterolemia pig model. *Arterioscler Thromb Vasc Biol*. 2019;39(11):2338–52.
45. Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative C(T) method. *Nat Protoc*. 2008;3(6):1101–8.
46. Gras SL, Waddington LJ, Goldie KN. Transmission electron microscopy of amyloid fibrils. *Methods Mol Biol*. 2011;752:197–214.
47. Oakley H, Cole SL, Logan S, Maus E, Shao P, Craft J, et al. Intraneuronal beta-amyloid aggregates, neurodegeneration, and neuron loss in transgenic mice with five familial Alzheimer's disease mutations: potential factors in amyloid plaque formation. *J Neurosci*. 2006;26(40):10129–40.
48. Ohno M, Cole SL, Yasvoina M, Zhao J, Citron M, Berry R, et al. BACE1 gene deletion prevents neuron loss and memory deficits in 5XFAD APP/PS1 transgenic mice. *Neurobiol Dis*. 2007;26(1):134–45.
49. Kimura R, Devi L, Ohno M. Partial reduction of BACE1 improves synaptic plasticity, recent and remote memories in Alzheimer's disease transgenic mice. *J Neurochem*. 2010;113(1):248–61.
50. Fukumoto H, Cheung BS, Hyman BT, Irizarry MC. Beta-secretase protein and activity are increased in the neocortex in Alzheimer disease. *Arch Neurol*. 2002;59(9):1381–9.
51. Matarin M, Salih DA, Yasvoina M, Cummings DM, Guelfi S, Liu W, et al. A genome-wide gene-expression analysis and database in transgenic mice during development of amyloid or tau pathology. *Cell Rep*. 2015;10(4):633–44.
52. Bandet CL, Mahfouz R, Veret J, Sotiropoulos A, Poirier M, Giussani P, et al. Ceramide transporter CERT is involved in muscle insulin signaling defects under lipotoxic conditions. *Diabetes*. 2018;67(7):1258–71.
53. Kumagai K, Yasuda S, Okemoto K, Nishijima M, Kobayashi S, Hanada K. CERT mediates intermembrane transfer of various molecular species of ceramides. *J Biol Chem*. 2005;280(8):6488–95.
54. De Strooper B, Annaert W. Proteolytic processing and cell biological functions of the amyloid precursor protein. *J Cell Sci*. 2000;113(Pt 11):1857–70.
55. Girard SD, Jacquet M, Baranger K, Migliorati M, Escoffier G, Bernard A, et al. Onset of hippocampus-dependent memory impairments in 5XFAD transgenic mouse model of Alzheimer's disease. *Hippocampus*. 2014;24(7):762–72.
56. Savage JC, Jay T, Goduni E, Quigley C, Mariani MM, Malm T, et al. Nuclear receptors license phagocytosis by trem2+ myeloid cells in mouse models of Alzheimer's disease. *J Neurosci*. 2015;35(16):6532–43.
57. Kanazawa H, Ohsawa K, Sasaki Y, Kohsaka S, Imai Y. Macrophage/microglia-specific protein Iba1 enhances membrane ruffling and Rac activation via phospholipase C-gamma -dependent pathway. *J Biol Chem*. 2002;277(22):20026–32.
58. Olabarria M, Noristani HN, Verkhratsky A, Rodriguez JJ. Concomitant astroglial atrophy and astrogliosis in a triple transgenic animal model of Alzheimer's disease. *Glia*. 2010;58(7):831–8.
59. Dinkins MB, Enasko J, Hernandez C, Wang G, Kong J, Helwa I, et al. Neutral sphingomyelinase-2 deficiency ameliorates Alzheimer's disease pathology and improves cognition in the 5XFAD mouse. *J Neurosci*. 2016;36(33):8653–67.
60. Behr D, Hesse L, Masters CL, Multhaup G. Regulation of amyloid protein precursor (APP) binding to collagen and mapping of the binding sites on APP and collagen type I. *J Biol Chem*. 1996;271(3):1613–20.
61. Multhaup G. Identification and regulation of the high affinity binding site of the Alzheimer's disease amyloid protein precursor (APP) to glycosaminoglycans. *Biochimie*. 1994;76(3–4):304–11.
62. Breen KC, Bruce M, Anderton BH. Beta amyloid precursor protein mediates neuronal cell-cell and cell-surface adhesion. *J Neurosci Res*. 1991;28(1):90–100.
63. Turner N, Mason PJ, Brown R, Fox M, Povey S, Rees A, et al. Molecular cloning of the human Goodpasture antigen demonstrates it to be the alpha 3 chain of type IV collagen. *J Clin Invest*. 1992;89(2):592–601.
64. Verdier Y, Zarandi M, Penke B. Amyloid beta-peptide interactions with neuronal and glial cell plasma membrane: binding sites and implications for Alzheimer's disease. *J Pept Sci*. 2004;10(5):229–48.
65. Ebel P, Vom Dorp K, Petrasch-Parwez E, Zlomuzica A, Kinugawa K, Mariani J, et al. Inactivation of ceramide synthase 6 in mice results in an altered sphingolipid metabolism and behavioral abnormalities. *J Biol Chem*. 2013;288(29):21433–47.
66. Imgrund S, Hartmann D, Farwanah H, Eckhardt M, Sandhoff R, Degen J, et al. Adult ceramide synthase 2 (CERS2)-deficient mice exhibit myelin sheath defects, cerebellar degeneration, and hepatocarcinomas. *J Biol Chem*. 2009;284(48):33549–60.
67. Salegio EA, Samaranch L, Jenkins RW, Clarke CJ, Lamarre C, Beyer J, et al. Safety study of adeno-associated virus serotype 2-mediated human acid sphingomyelinase expression in the nonhuman primate brain. *Hum Gene Ther*. 2012;23(8):891–902.
68. Flanigan TJ, Xue Y, Kishan Rao S, Dhanushkodi A, McDonald MP. Abnormal vibrissa-related behavior and loss of barrel field inhibitory neurons in 5XFAD transgenics. *Genes Brain Behav*. 2014;13(5):488–500.
69. Jawhar S, Trawicka A, Jenneckens C, Bayer TA, Wirths O. Motor deficits, neuron loss, and reduced anxiety coinciding with axonal degeneration and intraneuronal Abeta aggregation in the 5XFAD mouse model of Alzheimer's disease. *Neurobiol Aging*. 2012;33(1):196 e29–40.
70. Richard BC, Kurdakova A, Baches S, Bayer TA, Weggen S, Wirths O. Gene dosage dependent aggravation of the neurological phenotype in the 5XFAD mouse model of Alzheimer's disease. *J Alzheimers Dis*. 2015;45(4):1223–36.
71. Wu D, Tang X, Gu LH, Li XL, Qi XY, Bai F, et al. LINGO-1 antibody ameliorates myelin impairment and spatial memory deficits in the early stage of 5XFAD mice. *CNS Neurosci Ther*. 2018;24(5):381–93.
72. Gu L, Wu D, Tang X, Qi X, Li X, Bai F, et al. Myelin changes at the early stage of 5XFAD mice. *Brain Res Bull*. 2018;137:285–93.
73. Katsel P, Li C, Haroutunian V. Gene expression alterations in the sphingolipid metabolism pathways during progression of dementia and Alzheimer's disease: a shift toward ceramide accumulation at the earliest recognizable stages of Alzheimer's disease? *Neurochem Res*. 2007;32(4–5):845–56.
74. Couttas TA, Kain N, Tran C, Chatterton Z, Kwok JB, Don AS. Age-dependent changes to sphingolipid balance in the human hippocampus are gender-specific and may sensitize to neurodegeneration. *J Alzheimers Dis*. 2018;63(2):503–14.
75. Han X, D MH, McKeel DW, Jr., Kelley J, Morris JC. Substantial sulfatide deficiency and ceramide elevation in very early Alzheimer's disease: potential role in disease pathogenesis. *J Neurochem*. 2002;82(4):809–818.
76. Barrier L, Fauconneau B, Noel A, Ingrand S. Ceramide and related-sphingolipid levels are not altered in disease-associated brain regions of APP and APP/PS1 mouse models of Alzheimer's disease: relationship with the lack of neurodegeneration? *Int J Alzheimers Dis*. 2011;2011:920958.
77. Thinakaran G, Koo EH. Amyloid precursor protein trafficking, processing, and function. *J Biol Chem*. 2008;283(44):29615–9.
78. Lopez-Montero I, Rodriguez N, Cribier S, Pohl A, Velez M, Devaux PF. Rapid transbilayer movement of ceramides in phospholipid vesicles and in human erythrocytes. *J Biol Chem*. 2005;280(27):25811–9.
79. Silva L, de Almeida RF, Fedorov A, Matos AP, Prieto M. Ceramide-platform formation and -induced biophysical changes in a fluid phospholipid membrane. *Mol Membr Biol*. 2006;23(2):137–48.
80. Heppner FL, Ransohoff RM, Becher B. Immune attack: the role of inflammation in Alzheimer disease. *Nat Rev Neurosci*. 2015;16(6):358–72.
81. Wang G, Dinkins M, He Q, Zhu G, Poirier C, Campbell A, et al. Astrocytes secrete exosomes enriched with proapoptotic ceramide and prostate apoptosis response 4 (PAR-4): potential mechanism of apoptosis induction in Alzheimer disease (AD). *J Biol Chem*. 2012;287(25):21384–95.
82. Dinkins MB, Dasgupta S, Wang G, Zhu G, Bieberich E. Exosome reduction in vivo is associated with lower amyloid plaque load in the 5XFAD mouse model of Alzheimer's disease. *Neurobiol Aging*. 2014;35(8):1792–800.
83. de Wit NM, den Hoedt S, Martinez-Martinez P, Rozemuller AJ, Mulder MT, de Vries HE. Astrocytic ceramide as possible indicator of neuroinflammation. *J Neuroinflammation*. 2019;16(1):48.
84. Elsherbini A, Kirov AS, Dinkins MB, Wang G, Qin H, Zhu Z, et al. Association of Abeta with ceramide-enriched astrosomes mediates Abeta neurotoxicity. *Acta Neuropathol Commun*. 2020;8(1):60.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.