



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

22nd International Grassland Congress

Evaluation of a Drought Tolerance Native Grass: *Cleistogenes songorica* for the Turf Use Purpose

Xinyong Li
Lanzhou University, China

Xue Wei
Lanzhou University, China

Yanrong Wang
Lanzhou University, China

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/22/1-6/5>

The 22nd International Grassland Congress (Revitalising Grasslands to Sustain Our Communities) took place in Sydney, Australia from September 15 through September 19, 2013.

Proceedings Editors: David L. Michalk, Geoffrey D. Millar, Warwick B. Badgery, and Kim M.

Broadfoot

Publisher: New South Wales Department of Primary Industry, Kite St., Orange New South Wales, Australia

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Evaluation of a drought tolerance native grass: *Cleistogenes songorica* for the turf use purpose

Xinyong Li, Xue Wei and Yanrong Wang

State Key Laboratory of Grassland Agro-ecosystems, College of Pastoral Agriculture Science and Technology, Lanzhou University, People's Republic of China

Contact email: yrwang@lzu.edu.cn

Keywords: Native grass, turfgrass, drought tolerance, *Cleistogenes songorica*.

Introduction

Water deficit is one of the most important factors to restrict growth of turf grass, especially in northwest China where water available for landscape irrigation is increasingly limited. Use of drought-tolerant turfgrass species or cultivars is one of the strategies used to reduce water utilization and irrigation requirement (Nielsen and Stewart 1990). Recent study showed that regionally adapted native grass species are worth investigating as suitable alternatives to the conventional turfgrasses in many applications (Mark *et al.* 2011), and several native grass species are suitable for low-maintenance turfgrass use has also been reported (Mintenko *et al.* 2002). Awnless cleistogenes (*Cleistogenes songorica*) is a drought tolerance perennial grass native to the northwest desert grasslands of China. It grows well with a mean annual rainfall of 100 to 200 mm and is tolerant to the winter temperatures as low as -40°C . A series of studies related to the domestication of this species have been reported by the writers including seed germination

ecology (Yu *et al.* 2004), seedling establishment techniques (Tai *et al.* 2008) and seed production (Wei 2010). This paper reports the performance of *C. songorica* as a turf use grass under drought conditions.

Material and methods

The turf performance test of *C. songorica* was conducted in the Hexi Corridor of Gansu Province, China ($38^{\circ}24' \text{N}$, $100^{\circ}29' \text{E}$, a.s.l. 1450 m) from 2008 to 2010. Two varieties of tall fescue: Houndog 5 and Water Saver were used as controls. The sowing density of three varieties was 15,000 plants/m². The experiment was performed in split plots based on a completely randomized blocks design with four replications. The plots were irrigated at 10 day intervals in 2008 and 2009, and 20 day intervals in 2010. The characteristics measured included: plant density, coverage, green period, and growth rate in 20 days after cutting each year. In 2010, leaf relative water content (RWC) was also determined according to Barrs and Weatherley (1962).




Figure 1. Performances comparison between awnless cleistogenes and two tall fescue varieties in 2008, 2009 and 2010. **A:** coverage, **B:** green period, **C:** plant height, **D:** RWC. Different letters indicate significant differences between test materials at 5% level in LSD test.

Results

Compared with two varieties of tall fescue, *C. songorica* showed a higher coverage ($P<0.05$), particularly under the drought stress in 2010 (Fig. 1A). Although the green period of *C. songorica* was shorter in 2008 and 2009, it was more consistent over the three years of observations compared with the tall fescues (Fig. 1B). By 2010, the coverage and green period of *C. songorica* were 84% and 223d, compared with 78% and 225d for Water Saver, and 54% and 210d for Houndog5, respectively. After cutting, regrowth of *C. songorica* was much slower ($P<0.05$) than that of two fescue varieties (Fig. 1C), indicating low management inputs needed. The RWC of *C. songorica* was much higher ($P<0.05$) under drought stress. For example, after 20 days water stress RWC of *C. songorica* was 60%, compared with 45% and 30% for Water Saver and Houndog5, respectively (Fig. 1D).

Conclusion

It is concluded that *C. songorica* can be used as a turfgrass or groundcover grass which has advantages of drought tolerance and low management inputs.

Acknowledgments

This study was financially supported by the Natural Science Foundation of China (30771532)

References

- Barrs HD, Weatherley PE (1962) A re-examination of the relative turgidity technique for estimating deficit in leaves. *Australian Journal of Biological Sciences* **15**, 413-428.
- Mark S, Michelle B, Steve W, Holly Z (2011) The performance of native and non-native turfgrass monocultures and native turfgrass polycultures: an ecological approach to sustainable lawns. *Ecological Engineering* **37**, 1095-1103.
- Mintenko AS, Smith SR and Cattanic DJ (2002) Turfgrass evaluation of native grasses for the Northern Great Plains Region. *Crop Science* **42**, 2018-2024.
- Tai JH, Wang YR, Chen G (2008) Responses of Seed Germination, Seedling Emergence and Seedling Growth in *Cleistogenes songorica* to Soil Water Content. *Acta Prataculturae Sinica* **17**, 105-110.
- Yu XJ, Wang YR, Zeng YJ, Su D (2004) Effect of temperature and osmotic potential on seed germination of *Cleistogenes songorica* and *Plantago lessingii*. *Acta Ecologica Sinica* **24**, 883-887.