



8-1963

Determine Liming Needs on Tobacco Land This Fall

Jones H. Smiley
University of Kentucky

Harold F. Miller
University of Kentucky

Right click to open a feedback form in a new tab to let us know how this document benefits you.

Follow this and additional works at: https://uknowledge.uky.edu/pss_notes

 Part of the [Agronomy and Crop Sciences Commons](#)

Repository Citation

Smiley, Jones H. and Miller, Harold F., "Determine Liming Needs on Tobacco Land This Fall" (1963). *Agronomy Notes*. 149.
https://uknowledge.uky.edu/pss_notes/149

This Report is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in Agronomy Notes by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

AGRONOMY NOTES

SOILS • CROPS

Prepared by Department of Agronomy, University of Kentucky Cooperative Extension Service

Number 3

August, 1963

DETERMINE LIMING NEEDS ON TOBACCO LAND THIS FALL

There have been far too many reports of manganese toxicity in tobacco this season for a condition which is so easily corrected.

Manganese is one of the minor elements necessary for plant growth, but it is needed in very small amounts. When large amounts are taken up by a tobacco plant, light green or yellowish areas appear between the larger veins of the leaf. In severe cases the plant may be stunted or even die.

The availability of manganese is closely associated with soil acidity. As the soil becomes more acid the amount of manganese in the soil solution is increased to a point where it becomes toxic to plants. Fertilization practices on burley tobacco frequently cause the soil to become more acid, which results in a need for ground limestone to sweeten the soil.

Since not all tobacco fields will need lime, a good representative soil sample should be taken from each tobacco field as soon as the crop is harvested to determine whether lime is needed. If soil test results show that limestone is needed, it should be applied and disked in before the cover crop is seeded this fall. When this is done the limestone will have the winter months to be reducing soil acidity before next year's crop is planted. For this reason it is important that tobacco ground be tested immediately following harvest.

900-8-63
Filing Code 1

Jones H. Smiley
Harold F. Miller

(To simplify information in this publication, trade names of some products are used. No endorsement is intended, nor is criticism implied of similar products not named.)

Cooperative Extension Work in Agriculture and Home Economics: College of Agriculture and Home Economics, University of Kentucky, Lexington, and the United States Department of Agriculture, cooperating. William A. Seay, Director. Issued in furtherance of the Acts of May 8 and June 30, 1914.