



Copper Status of Free Ranging Cattle: What's Hidden Behind? A Pilot Study at the Gilgel Gibe Catchment, Ethiopia

Veronique Dermauw
Ghent University, Belgium

Thomas Van Hecke
Ghent University, Belgium

Kechero Yisehak
Jimma University, Ethiopia

Gijs Du Laing
Ghent University, Belgium

Eric Van Ranst
Ghent University, Belgium

See next page for additional authors

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/22/1-10/12>

The 22nd International Grassland Congress (Revitalising Grasslands to Sustain Our Communities) took place in Sydney, Australia from September 15 through September 19, 2013.

Proceedings Editors: David L. Michalk, Geoffrey D. Millar, Warwick B. Badgery, and Kim M.

Broadfoot

Publisher: New South Wales Department of Primary Industry, Kite St., Orange New South Wales, Australia

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Presenter Information

Veronique Dermauw, Thomas Van Hecke, Kechero Yisehak, Gijs Du Laing, Eric Van Ranst, Luc Duchateau, and Geert P. J. Janssens

Copper status of free ranging cattle: what's hidden behind? A pilot study at the Gilgel Gibe catchment, Ethiopia

Veronique Dermauw^{A*}, Thomas Van Hecke^{A*}, Kechero Yisehak^B, Gijs Du Laing^C, Eric Van Ranst^D, Luc Duchateau^E and Geert P J Janssens^A

^A Department of Nutrition, Genetics and Ethology, Faculty of Veterinary Medicine, Ghent University, Heidestraat 19, B-9820, Merelbeke, Belgium

^B Department of Animal Sciences, Jimma University College of Agriculture and Veterinary Medicine, PO Box 307, Jimma, Ethiopia

^C Department of Applied Analytical and Physical Chemistry, Ghent University, Coupure Links 653, B-9000 Gent, Belgium

^D Department of Geology and Soil Science, Ghent University, Krijgslaan 281/S8, B-9000 Gent, Belgium

^E Department of Comparative Physiology and Biometrics, Ghent University, Salisburylaan 133, B-9820 Merelbeke, Belgium.

* Both authors contributed equally to this work

Contact email: veronique.dermauw@ugent.be, thomas.vanhecke@ugent.be

Keywords: Mineral, soil, humid tropics, zebu.

Introduction

Copper (Cu) deficiency is known to be a major risk for cattle health and production. The widespread problem is due to low absorption rates in the rumen, low Cu concentrations in the natural diet and the presence of dietary Cu antagonists, such as S, Mo and Fe (Suttle 2010). Cattle in the tropics are even more prone to shortage of this mineral, given that the animals are often extensively kept and largely dependent on natural pasture for mineral supply (McDowell and Arthington, 2005). More specifically, in Ethiopia, Cu deficiency was previously described in zebu (*Bos indicus*) cattle by several authors (*e. g.* Dermauw *et al.*). Similar to other minerals, Cu is part of the soil-plant-animal chain, with many factors influencing Cu concentrations at every level. In our study, the overall goal was to investigate the possible influence of certain environment and management factors on dietary concentrations of Cu and antagonists and Cu status in free ranging cattle.

Methods

This study was conducted in the Gilgel Gibe catchment, south-western Ethiopia. In three elevation zones (1= 1700-1800m above sea level (a.s.l.), 2= 1800-2000m a.s.l., 3= 2200m a.s.l.), 19 herds (<10 to >70 animals per herd) of free ranging zebu (*Bos indicus*) cattle were randomly selected within a radius of 35 km around the town of Jimma, and observed during one day. One cow per herd was closely observed, the body condition score (BCS) was recorded and a blood sample was obtained. The soil type (Nitrisols, Acrisols and Ferralsols (soil association NAF) or Planosols and Vertisols (soil association PV) (Van Ranst *et al.* 2011)) was identified based upon landscape position and visual inspection of the soil. Management practices were evaluated based upon grazing strategy (communal versus individual), supplementation with crop residues (as % of dietary observations) and the total distance covered (during

one grazing day). The dietary plant composition was estimated through observation of ingested plant species every 10 minutes. One ingested plant per observation was equal to 1 point, several ingested species per observation were allocated 1 point divided by the number recorded. A sample of all ingested plants was collected. To estimate the dietary mineral concentrations, the ingested proportions of each plant were multiplied by mineral content and summed. If no sample could be obtained due to small plant size, the mineral concentration of the known percentage of ingested plants was extrapolated to 100%. Finally, the number of ingested plant species (NIPS) was recorded. Plasma was obtained through centrifugation at 1500 × g for 10 min. Plant samples were oven dried at 65 °C for 72 hours. All samples were digested using microwave destruction with 10 ml HNO₃ (plasma: open, plants: closed vessels) followed by filtration and mineral analysis through inductively coupled plasma optical emission spectrometry & mass spectrometry (ICP-OES & -MS) (Vista MPX radial, Varian, Palo Alto, USA; Elan DRC-e, Perkin Elmer, Sunnyvale, CA, USA, respectively). Statistical analysis was performed using SAS Version 9.3 (SAS Institute Inc., Cary, NC, USA). To determine the association between categorical variables, a Fisher's Exact test was used. To determine the effect of the physical environment and management on mineral concentrations in ingested diet and status, a fixed effects model with one categorical or one continuous covariate at a time was fitted to the different response variables. Hypotheses were tested using the F-test at the 5% significance level.

Results

Out of the 19 observed animals, 12 were severely Cu deficient (mean: 0.48 ± 0.19 (SD) mg/l), while estimated dietary Cu concentrations were below requirements for all herds (5.8 ± 1.4 mg/kg DM). Soil type and elevation were significantly associated ($P < 0.001$), with the typical occur-

rence of Planosol/Vertisol associations (PV) at lower elevation. Soil type did not affect the amount crop residues offered by the farmers ($P>0.05$), but significantly influenced herd type (more communal grazing for longer distances on PV soils). Focusing on the impact of this soil type on diet, NIPS (10.7 vs. 17.2 ± 1.3 (SEM)), dietary concentrations of Cu (4.7 vs. 6.4 ± 0.3 mg/kg DM) and Fe (765 vs. 1672 ± 183 mg/kg DM) were lower on PV soils whereas concentrations of Mo (1.47 vs. 0.96 ± 0.10 mg/kg DM) were higher than on NAF soils (all $P<0.05$). Dietary concentrations of S tended to be lower on PV soils (0.15 vs. 0.19 ± 0.01 % DM, $P = 0.082$). Animal Cu status, however, was not affected by soil type. Concerning the impact of management practices on diet and status, dietary Mo concentrations were higher in communal grazing herds (1.4 vs. 1.0 ± 0.1 mg/kg DM) whereas the Cu:Mo ratio was higher in individual grazing herds (6.8 vs. 4.0 ± 1.0) (both $P<0.050$), but Cu status was not affected ($P>0.050$). The amounts of crop residues offered to grazing animals had a significant negative effect on dietary Zn concentrations ($P = 0.033$) and tended to negatively affect dietary Mo and S concentrations ($P = 0.064$; $P = 0.063$ respectively). Animal mineral status was not affected by this, but BCS was positively affected ($P = 0.010$). Herding distance tended to negatively affect Cu:Mo ratio ($P = 0.082$), but did not affect Cu status nor of Cu antagonists.

Conclusion

Zebu cattle grazing in the study area were under high risk for Cu deficiency, potentially aggravated due to Fe, Mo and S overload. Lower dietary Cu concentrations were ingested

on PV soils, which coincided with higher dietary Mo and lower Fe concentrations. Herd type, crop residue supplementation and herding distance had or tended to have an effect on dietary concentrations of Cu antagonists. Despite the unaffected Cu status, these results accentuate the impact of both environment and management on dietary concentrations of Cu and antagonists. The the impact of these factors may encourage farmers to take informed choices, which might improve the copper status of their animals.

Acknowledgements

This study was funded by Grant SB091348 from IWT-Vlaanderen. Furthermore, we would like to thank the IUC-JU programme of VLIR-UOS for logistical support and Ria Van Hulle for performing mineral analyses.

References

- Dermauw V, Yisehak K, Belay D, Van Hecke T, Du Laing G, Duchateau L, Janssens GPJ (2013). Mineral deficiency status of ranging zebu (*Bos indicus*) cattle around the Gilgel Gibe catchment, Ethiopia. *Tropical Animal Health and Production* **45**, 1139-1147
- McDowell LR, Arthington JD (2005). Minerals for grazing ruminants in tropical regions. 4th Edition. (University of Florida, IFAS, Gainesville, FL, USA)
- Suttle NF (2010). Mineral nutrition of livestock. 4th Edition. (CABI, Wallingford, UK)
- Van Ranst E, Dumon M, Tolossa AR, Cornelis, JT, Stoops G, Vandenberghe RE, Deckers J (2011). Revisiting ferrolysis processes in the formation of Planosols for rationalizing the soils with stagnic properties in WRB. *Geoderma* **163**, 265–274.