



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

22nd International Grassland Congress

Challenges and Solutions for Forage Conservation for Small and Large Enterprises

M. Goto
Mie University, Japan

H. Kawamoto
NARO, Japan

H. Matsuyama
NARO, Japan

Ryuichi Uegaki
NARO, Japan

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/22/1-11/1>

The 22nd International Grassland Congress (Revitalising Grasslands to Sustain Our Communities) took place in Sydney, Australia from September 15 through September 19, 2013.

Proceedings Editors: David L. Michalk, Geoffrey D. Millar, Warwick B. Badgery, and Kim M.

Broadfoot

Publisher: New South Wales Department of Primary Industry, Kite St., Orange New South Wales, Australia

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Challenges and solutions for forage conservation for small and large enterprises

M Goto ^A, H Kawamoto ^B, H Matsuyama ^C and R Uegaki ^D

^A Graduate school of Bioresources, Mie University, Tsu, Japan 514-8507, www.bio.mie-u.ac.jp

^B Tohoku Agricultural Research Center, NARO, Morioka, Japan 020-0198, www.naro.affrc.go.jp

^C Institute of Livestock and Grassland Science, NARO, Nasushiobara, Japan 328-2793
www.naro.affrc.go.jp

^D Institute of Livestock and Grassland Science, NARO, Nasushiobara, Japan 328-2793,
www.naro.affrc.go.jp

Contact email: goto@bio.mie-u.ac.jp

Abstract. Forage conservation, particularly silage making, is one of the major technologies used as an interface between forage production and animal production, and the advanced technology of biological, chemical, and enzymatic additives for making silage has contributed significantly to the development of livestock production. However, the increasing demand for meat and dairy products, severe environmental deterioration induced by livestock production, and the critical risks to human health associated with mycotoxin contamination of forage crops remain to be addressed. Here we review the extant literature regarding treatment with various silage additives in relation to new paddy field forage production systems, mycotoxin contamination of forage crops, and methane and ammonia production from livestock farming.

Keywords: Paste ammonia, conservation, methane, mycotoxins, rice whole crop, silage additive.

Introduction

The consumption of meat and dairy products continues to increase worldwide, with estimates of the increased demand ranging from 229 and 580 million tons of meat and dairy products in the years 1999/2001 to 465 and 1043 million tons in the year 2050, respectively (UN Food and Agriculture Organization 2006). Particularly in Asian countries, which are home to over half of the world's population and where paddy fields occupy over 11% of the world's total cultivated area (Maclean *et al.* 2002), the demand for livestock products is increasing. Forage and livestock production systems are thus focused on establishing the most effective use of paddy fields, especially surplus fields. For example, rice paddy fields occupy 54% of the total cultivated land area in Japan, and 46% of the paddy field is not planted due to reduced rice consumption and less suitability of other crops that are associated with humid soil conditions. This reduced use of land for rice farming is very serious in view of not only the economics of rice farming but also that of land management, including the control of water, and thus the establishment of forage rice crops is increasingly important, especially in Asian countries.

According to the UN Food and Agriculture Organization (FAO 2006) and the Intergovernmental Panel on Climate Change (2007), worldwide the livestock industry accounted for 18% of the greenhouse gas emissions equivalent to carbon dioxide, and 23% of the methane (CH₄) emission from anthropogenic activities and sources. Severe environmental deterioration induced by nitrogen imbalances from livestock farming have

created ammonia pollution on the ground and in the atmosphere, acid rain, and nitrate contamination of underground water. The control of nitrogen (N) pollution and CH₄ emissions produced by livestock farming is another economic concern, since only approx. 50% of total N ingested by ruminants is retained, and the enteric CH₄ emission, a byproduct of mainly fibrous forage digestion by rumen micro-organisms, reached 3%–12% of the loss of dietary gross energy (GE), as calculated based on a value of 55 J/g CH₄. The simultaneous increase of the use and/or efficiency of dietary N and carbohydrate sources is another important goal toward directly reducing their environmental impact and enhancing sustainable development.

In addition, there are serious and valid concerns about the safety of forage. Among the many hygienic challenges involved in forage and livestock production, the existence of mycotoxins is a major problem that can damage the health of not only animals but also humans. Since these toxic compounds often contaminate feedstuff including forage crop silages and many different forage plant species (Scudamore and Livesey 1998; Miller 2008), the difficulty of controlling the generation of mycotoxins at pre- and post-harvest in livestock farming remains.

Since forage conservation, particularly silage making, is the main interface between forage production and animal production, further advances in the technology involving biological, chemical, and enzymatic additives are needed to evaluate how these additives can contribute to improvements in forage and livestock

farming, especially in light of the newly developed forage production systems in Asia, the safety of meat and dairy products, and the environmental impact of livestock farming. In this paper, we review the treatment of various silage additives in relation to the new forage production system for paddy fields, mycotoxin contamination of forages, and methane and ammonia production from livestock farming.

Silage production fields in Asian countries

When forage rice is to be fed to cattle, the entire rice plant including panicles, leaves and stems is generally harvested at the yellow ripening stage and conditioned in whole crop silage (WCS). The introduction of round balers with an attached crawler-type driving system successfully improved harvesting operations in wet soil conditions, which had not been achieved for forage production before, and it has contributed to the promotion of rice WCS production and use in Japan (Urakawa and Yoshimura 2003). The development of attachable short-length cutting equipment has resulted in the further improvement of fermentation quality (Kawamoto *et al.* 2007; Urakawa 2009) and has enabled a new conservation system for ensiling total mixed ration (TMR) (Shito *et al.* 2006). The livestock production system in Japan is moving toward regional cooperative farming, where TMR contractor network with rice crop farmers to supply regional self-sufficient forages.

However at the present, the rice WCS system has several problems involving yield, nutrition and feeding management, and fermentation quality. Although a new forage-type cultivar achieved an almost twofold yield (24.1 dry matter [DM] t/ha and 12.7 total digestible nutrients [TDN] t/ha) compared to common grain-type cultivars (Kato 2009), the rice-winter crop rotation system was implemented in order to increase the annual forage yield (Iijima *et al.* 2005; Kamoshita *et al.* 2007). The new cultivar "Tachisuzuka" is characterized by a high straw yield and high sugar content, with a much lower grain:straw ratio (0.08–0.14) and higher sugar content (11.5% DM) compared to the ratio 0.69 and 1.7% DM of the common cultivar "Kusanohoshi" (Matsushita *et al.* 2011). A higher sugar content in a cultivar is advantageous for achieving preferential fermentation profiles similar to those of forage corn and others, by reducing the influence of soil contamination (undesirable microorganisms) and lower ensiling density of round bales, which is commonly observed with rice WCS production. The lower ratio of grain:straw in the cultivar Tachisuzuka contributes to the reduction of considerable energy losses and difficulty in the feeding management of rice WCS. Shinde (2010) reported that the DM percentage of undigested rice grains was 23% of the amount of the grain ingested as harvested at the dough-ripe stage and 43% of that with advancing maturation to the yellow-ripe stage.

A number of chemical and biological additives have been proposed to improve the microbial profiles, fermentation quality, aerobic stability, and energy loss of forage crops including rice WCS, alfalfa (*Medicago sativa* L.), orchardgrass (*Dactylis glomerata* L.), and

other grasses and legumes. Treatment with combinations of homofermentative and heterofermentative bacterial inoculants has been extensively demonstrated to alter the effectiveness of microbial inoculation, since they have shown no interactions and exhibited different beneficial effects on the resulting silage (Hu *et al.* 2009). The shifting of fermentation toward lactic acid with homofermentative lactic acid bacteria improved the fermentation profiles, and that toward acetic acid with heterofermentative lactic acid bacteria improved the aerobic stability and thereby both the performance of the silage and the acceptance of the silage by animals.

The potential of using commercialized lactic acid bacteria inoculants such as *Lactobacillus plantarum*, *Lactobacillus fermentum*, *Pediococcus pentosaceus*, *Enterococcus faecium*, and *Lactobacillus buchneri* has also been described in the literature (Adesogan *et al.* 2003; Hu *et al.* 2009; Cao *et al.* 2011). A commercial lactic acid bacteria inoculant – a strain of *L. plantarum* developed for rice whole crops from a national research project – is also used for ensiling rice forage crops, and strain-dependent effect inoculation of *L. plantarum* on the fermentation quality of rice WCS were observed (Tohno *et al.* 2012).

Fermentation factors such as carbohydrate assimilation, pH range, temperature allowing growth, and growth regulation of undesirable microorganisms are important, and the diversity of epiphytic lactic acid bacteria (fermented juice of epiphytic lactic acid bacteria; FJLB) was observed to affect the anaerobic fermentation of forage crops including rice whole crop, alfalfa, Italian ryegrass (*Lolium multiflorum* L.), orchardgrass, and some tropical grasses (Cao *et al.* 2002; Yahaya *et al.* 2004; Hiraoka *et al.* 2006). A simple screening of effective microflora by using FJLB, which can be obtained by prefermenting fresh rice leaf and stem fractions for 2–3 days at about 30°C in water containing 1–2% sugars (w/v) and a few drops of acetic acid (vinegar) satisfactorily comprised over 2.0×10^8 colony-forming units per milliliter (cfu/ml) of the compositional percentage of 55% *L. lactis* subsp. *lactis*, 17% *L. fermentum*, 16% *L. plantarum*, and 11% *Weissella confusa* (Hiraoka *et al.* 2006). The diverse members of *L. lactis* subsp. *lactis*, *L. fermentum*, *L. plantarum*, and *W. confusa* respond to various environmental and material conditions. Further research using FJLB preparations of mixed and balanced inoculants of homofermentative and heterofermentative lactic acid bacteria is of interest, and is expected to contribute to preferential fermentation quality without reducing the advantages of ease of use and cost-effectiveness.

Antifungal and anti-yeast activities of chemical additives strongly improved the stability of silage when it was exposed to air (Kung *et al.* 2004). Propionic acid-based additives and commercial preservatives (additives containing a mixture of ammonium formate, propionate, ethyl benzoate, and benzoate) did not affect fermentation at low rates but altered the aerobic stability of forage crops including corn, barley, wheat grains and TMR, and, in some circumstances when treated with microbial inoculants, the silage was preferentially fermented

(Adesogan *et al.* 2003; Kung *et al.* 2004; Tyrolová and Výborná 2011). Chemical additives, in combination with enzymatic treatment, certainly improved fermentation quality in the silage (Chungheng *et al.* 2008; Sun *et al.* 2009), while chemical treatment including urea and sodium bicarbonate alone promoted the *Enterococcus* dominant microbial flora and induced preferential fermentation in whole crop wheat silage (Bal and Bal 2012).

The amino acid fermentation byproduct (AFB), which is the mother-liquid residues of amino acid-producing bacteria and is normally used as an N supplement for ruminants, has been recognized as a very useful silage additive with beneficial effects on the fermentation, aerobic stability, and digestibility of silages of Italian ryegrass, rice whole crop, barley whole crop (*Hordeum vulgare* L.), and corn (*Zea mays*) (Yimiti *et al.* 2006a; 2006b, Okiyama *et al.* 2008, Yamamoto *et al.* 2008, Yimamu *et al.* 2008). The addition of AFB at the ensiling of rice whole crop increased the digestibility of rice grains of lactating dairy cattle, which was advantageous for the feeding management because the byproduct possess the ability to assist the microbes to access morphological and anatomical tissues of forage plants. The *in situ* rumen degradability of DM and NDF of AFB-treated silages was also increased by averages of 7% and 6.5% compared to untreated silages and in some circumstances the milk production was increased after the treatment of rice WCS (Fang *et al.* 2006; Okiyama *et al.* 2008).

AFB additives consistently enhanced the growth of *L. lactis* subsp. *lactis*, *P. acidilactici*, *Leuconostoc mesenteroides* subsp. *mesenteroides*, and *L. plantarum*, to levels higher than the growth of *Clostridium butyricum* and the growth of yeast isolated from Italian ryegrass silage (Yimiti *et al.* 2006). Aerobic stability after the silo was opened was also in evidence based on lowered lactic acid consumption and lowered pH rising in AFB-added corn silage, although the colony numbers of yeast and fungi were increased, with no significant difference between the untreated and treated silages (Yimamu *et al.* 2008). AFB consistently contains higher concentrations of ammonia N, free amino acids, reducing sugars and carbohydrates and various minerals, showing that concentrations of total N, potassium, calcium, magnesium, and total cations were related to the improvement of both lactic acid fermentation and rumen degradability (Yamamoto *et al.* 2008). Significant improvements in NDF degradability might therefore be expected with the commonly used agricultural byproducts.

Hygienic measures in forage production and conservation

Aflatoxin (AFT), fumonisin (FUM), trichothecium mycotoxins (TRCs) such as deoxynivalenol (DON), nivalenol (NIV), and T-2 toxin (T2) and zearalenon (ZEN), the secondary metabolites produced by fungi, are the major mycotoxins detected in forage crops and silage including corn, sorghum (*Sorghum bicolor* Moench), barley, wheat (*Triticum aestivum* L.), rice plants, and

food-processed byproducts including bran, gluten, and corn distillers' dried grains with solubles (Scudamore and Livesey 1998; González *et al.* 2008; Miller 2008). Approximately 2–6.2% of the AFT B₁ ingested was carried over to milk (Fink-Gremmels 2008), and some was metabolized in the rumen and detected as AFT M₁ in the milk; this toxicity is almost the same as that of AFT B₁ (Trucksess *et al.* 1983; Allcroft *et al.* 1968). DON was completely metabolized in the rumen and detected as the de-epoxides in the milk (Miyazaki 2007; Fink-Gremmels 2008). The ZEN was converted into α -zearalenol in the rumen (Kiessling *et al.* 1984), and the ZEN and metabolite in the milk was only detected at low concentrations of 0.06–0.08%.

Several studies have described the occurrence of liver cancer and other toxic mutations in domestic animals and humans due to AFT (Park and Stoloff 1989), liver and renal toxicities in domestic animals and humans and equine leukoencephalopathy due to FUM (Voss *et al.* 2007), alimentary and immune-toxic injuries due to TRCs (Miyazaki 2007), and inactivation of female hormones due to ZEN (Donga *et al.* 2010). Thus the concentrations in feed are required to be <10 μ g/kg for AFT, <100 mg/kg for FUM (outside Japan), <5 mg/kg for DON (in Japan, EU, USA *etc.*) and <100 mg/kg for T2.

Toxic compounds are widely spread throughout Northern, Southeast, and Southern Asia, Oceania, Northern, Central, and Southern Europe, and the Mediterranean littoral (Binder *et al.* 2007). AFT and related compounds (AFTB₁, B₂, G₁, G₂) produced by *Aspergillus flavus* and *A. parasiticus* are detected particularly in the tropical zones, whereas FUM and related compounds (FUM B₁, B₂, B₃) produced mainly by *Fusarium fujikuroi* and *F. proliferatum* are detected especially in corn in temperate zones. T2 produced by *Fusarium* fungi is present primarily in Northern Europe and rarely in the Oceania and Asia. ZEN, produced by *Fusarium* fungi such as *F. graminearum*, is spread widely from the cold to the tropical zones.

Plant breeding has been conducted to obtain new resistant varieties and to deepen the understanding of mechanisms of toxin generation related to climate conditions and plant–pathogen interactions (Oerke *et al.* 2010; Woloshuk and Shim 2013). However, the control of mycotoxins is a complex undertaking in light of the generation behavior of each mycotoxin; for example, as reported for FUM, a higher toxic concentration was observed in plants showing slight or no symptoms of *Fusarium* head blight, while the opposite was also observed (Oerke *et al.* 2010). Another difficulty is that very little information has been obtained about AFT generation behavior during the pre- and post-harvesting periods and the interactions of AFT generation and environmental conditions. At present, it is thus practical to harvest properly at the growing stage before starting the generation of mycotoxins, to ensile forage immediately after harvest, and to maintain the hygienic conditions of silos and animal housing (Magan 2007).

Using as an example FUM in corn, the yellow-ripe stage is suitable for harvesting, since FUM mostly begins

to increase in some of the kernels and ears at the yellow-ripe stage (Warfield and Gilchrist 1999; Ariño *et al.* 2009; Uegaki *et al.* 2012a). Uegaki *et al.* (2012b) reported that there were no changes in the FUM concentration after and during the ensiling of contaminated forage crops. Commercial adsorbent might be also practically useful as a method for protecting animals against mycotoxicosis; a mixed adsorbent of zeolites and hydrated sodium calcium aluminosilicates was found to be near-perfect for preventing aflatoxicosis, but no adsorbent possesses complete effectiveness against all of the various types of mycotoxins (Alexander *et al.* 2001). The microbial and/or relative enzymatic degradation of mycotoxins collected by the *Nocardioides* sp. in wheat fields was capable of degrading DON to 3-epi-deoxynivalenol and its intermediates after prolonged incubation (Ikunaga *et al.* 2011), and microbial and enzymatic technology is expected to eventually provide detoxifying systems for toxic fungi in field soil and their secondary metabolite products during ensiling.

Environmental load related to forage production and utilization

Fecal and urinary N excretion per unit of dietary protein ingestion is influenced by low dietary protein quality alone, and in combination with the feeding of low-quality forage. Similarly, the enteric CH₄ emission per unit of organic matter (OM) and dry matter intake (DMI) is extensively influenced by the higher composition and lower digestibility of the fibrous components of feeds. Ominski *et al.* (2006) reported that when four formulations of chopped alfalfa-grass silage ranging in NDF content from 46.4 to 60.8% (DM basis) were fed to steers during the overwintering period, the enteric losses were lower at 27.1 L/kg DMI with a silage of 60.8% NDF than at that of 23.7 L/kg DMI with a silage of 46.4% NDF.

A similar trend was observed when CH₄ emissions were considered as a percent of GE. Therefore, treatment with chemical, biological, and/or enzymatic additives for silage making has been considered worthwhile to reduce the influence of agronomic factors including plant species and variety, fertilization, planting density, harvest time, and climate conditions, not only to improve fermentation and animal performance but also for reducing environmentally hazardous substances as ensiled forages.

Generally, forage conservation and hay or silage making have significant effects on the compositional percentage of non-protein N. It has been reported that the concentrations of non-protein N in silage accounted for 25–85% of the total N, higher than the 10–35% in the original materials (Muck 1988). Lowering the pH to near 4.0 or 4.5 immediately after ensiling is of great importance, since the protein degradation of forage crops is activated by plant proteolytic or silage bacterial activities at pH 5.5–6.0. Therefore, chemical additives, as mentioned above, are effective to control the protein degradation, particularly with lactic acid bacterial additives. The addition of cell-wall degrading enzymes was also effective to reduce ammonia production during

silage fermentation, similar to biological treatments (Nadeau *et al.* 2000; Zahiroddini *et al.* 2004; Dehghani *et al.* 2012).

Lactic acid bacterial additives including both homo- and hetero-fermentative lactic acid bacteria were observed to reduce the ammonia concentration in the silages of various forage crops by 9.4–41.1% (average of 22.7%) (Filya 2003; Kung *et al.* 2004; Hu *et al.* 2009; Chen *et al.* 2013). Since nowadays most silage bacterial additives are designed to protect the aerobic stability of the silage, their contribution to the environmental impact is more reliable and effective, although it is influenced by fermentation factors such as plant species, conservation period, pre-treatment of wilting, and year. Cao *et al.* (2002) demonstrated using alfalfa silage that the enhancement of lactic acid fermentation using FJLB significantly increased the *in vivo* DM and OM digestibility, probably as result of the acidic breakdown of pectin-rich alfalfa cell walls. The N retention was also higher at 13.9% of the total indigested N with FJLB-added silage compared to that at 2.1% with the control silage, accompanied by decreased NH₃ production during the ensiling. A similar trend was observed with higher urinary allantoin of cattle fed the FJLB-added silage, which was associated with higher bacterial growth in the rumen.

Methane is produced as a result of the microbial breakdown of forage cell walls in the rumen, reflected in the concentrations of NDF and hemicellulose. Ensiling alfalfa, compared with hay making, was assessed to decrease 32% and 28% of the enteric CH₄ production per unit of energy intake and per unit of DE, respectively, according to a mathematic and dynamic model of rumen digestion in which the fiber content was the primary definition factor (Benchaar *et al.* 2001). That decrease was also estimated to be higher compared to decreases obtained by changing the harvest time and the cutting length. However, anaerobic fermentation does not decrease the NDF and hemicellulose concentrations in all of the resulting silage of various forage crops (Nadeau *et al.* 2000; Zahiroddini *et al.* 2004; Xu *et al.* 2008).

Regarding cell wall constituents and digestibility, the mixture of cellulase and lactic acid bacteria additives is expected to be effective (Xu *et al.* 2008; Sun *et al.* 2009; Xing *et al.* 2009). Xing *et al.* (2009) reported that treatment with cellulolytic enzymes effectively decreased the NDF and ADF concentrations and improved the *in vitro* dry matter digestibility and *in vitro* NDF digestibility. However, those authors observe that the NDF degradation by cellulolytic enzyme activity during the ensiling was primarily the result in the lactic acid bacterial consumption of the enzyme-degradable NDF fraction, the subsequently increased NDF concentration, and the decreased digestibility of enzyme-undegradable NDF residues by rumen microorganisms.

The surfactant Tween 80, registered as a food additive, was reported to increase the water and enzyme absorption and cellulose degradation of grass hay when used in a buffered enzyme solution with a low concentration of 0.005% Tween 80 (Goto *et al.* 2003a). A considerable increase in forage degradability at *in vitro*

rumen digestion was also obtained by adding a surfactant at a low concentration, together with significant increases in the ratio of VFA:CH₄. The surfactant probably induces hydration and alters the surface of plant materials, in accord with the nonspecific enhancement of pure culture growth of *Streptococcus bovis*, *Prevotella ruminicola*, *Megashaera elsdinii*, *Selenomonas ruminantium*, *Butyrivibrio fibrisolvens*, and *Fibrobacta succinogenes*, which are fiber-, starch-, protein-, and metabolite-degrading bacteria in the rumen (Goto et al. 2003b; Kim et al. 2005). In addition, AFB, a promising agent for silage additives as described above, was revealed to possess the ability to assist the accessibility of morphological and anatomical tissues to cellulolytic enzymes, probably due to the increased ion strength of higher concentrations of ammonium sulfate and ammonium chloride. The current research suggests the importance of the further development and use of agents that aid microbial and enzymatic additives, in cross-relation to agricultural factors such as forage species and growth and maturation stages at harvest.

Moss et al. (1994) studied the enteric CH₄ emissions from sheep fed straw of three types (oat, wheat, and barley), and they found that treatment with ammonia (35 g/kg DM) and sodium hydroxide (45 g/kg DM) decreased the enteric CH₄ emissions per digestible OM by 20.5% and 12.5%, respectively. According to a very common observation following alkali treatment, the application to major forage crops must be considered. Additionally, nutrition and feeding management techniques such as supplementation with ionophores and fumaric and malic acids, plus lipid-rich diets with ingredients such as whole cotton seed, canola seed, sunflower, flax, coconut oil, and copra meal, are important to reduce methane emissions produced by livestock farming (Beauchemin et al. 2008).

Conclusion

The main purpose of ensiling is to preserve the nutritive and feeding values of a crop until it is fed to livestock. The present goals are to contribute to the expanding of forage and livestock production areas by increasing fermentation quality and to reduce both environmental deterioration and the risks to food safety. Ensiling technology including the development of new bacterial, enzymatic and chemical additives has already improved the fermentation and aerobic stability, and it has helped improve the nutrition and feeding management of silage. Increased effectiveness and efficiency of silage additives that have contributed to decreased environmental impact and improved livestock production have also been achieved, particularly with new silage additive candidates and supporting agents, as shown by AFB (mother-liquid residues of amino acid producing bacteria) and surfactant Tween 80. However, mycotoxin contamination in forage crops and silage remains a problem that must be addressed by plant breeding, cultivation, and conservation. Due to the possible effects of ensiling technology, especially silage additives, further microbiology research is needed to develop microbial and/or enzymatic detoxification techniques to

protect against food contamination and risks to animal and human health.

References

- Adesogan AT, Salawu MB, Ross AB, Davies DR, Brooks AE (2003) Effect of *Lactobacillus buchneri*, *Lactobacillus fermentum*, *Leuconostoc mesenteroides* inoculants, or a chemical additive on the fermentation, aerobic stability, and nutritive value of crimped wheat grains. *Journal of Dairy Science* **86**, 1789-1796.
- Alexander H, Stefan F, Othmar K, Hans D (2001) Mycotoxin detoxication of animal feed by different adsorbents. *Toxicology Letters* **122**, 179-188.
- Allcroft R, Roberts BA, Lloyd MK (1968) Excretion of aflatoxin in a lactating cow. *Food and Cosmetics Toxicology* **6**, 619-625.
- Ariño A, Herrera M, Juan T, Estopañan G, Carramiñana JJ, Rota C, Herrera A (2009) Influence of agricultural practices on the contamination of maize by fumonisin mycotoxins. *Journal Food Protection* **72**, 898-902.
- Bal EBB, Bal MA (2012) Effects of chemical additives and ensiling time on whole plant wheat silage microbial profiles inferred by phenotypic and 16S ribosomal DNA analysis. *World Journal of Microbiology and Technology* **28**, 767-776.
- Beauchemin KA, Kreuzer M, O'Mara F, McAllister TA (2008) Nutritional management for enteric methane abatement. *Australian Journal of Experimental Agriculture* **48**, 21-27.
- Benchaar C, Pomar C, Chiquette J (2001) Evaluation of dietary strategies to reduce methane production in ruminants: A modeling approach. *Canadian Journal of Animal Science* **81**, 563-574.
- Binder EM, Tanb LM, Chinb LJ, Handla J, Richardc J (2007) Worldwide occurrence of mycotoxins in commodities, feeds and feed ingredients. *Animal Feed Science Technology* **137**, 265-282.
- Cao LM, Goto M, Karita S, Yamamoto Y, Mizutani M, Deguchi Y, Urakawa S, Maekawa Y, Kawamoto Y, Masuko T (2002) Effect of fermented juice of epiphytic lactic acid bacteria on the fermentation quality of alfalfa (*Medicago sativa* L.) silage and its energy and nitrogen by dry cows. *Japanese Journal of Grassland Science* **48**, 227-235 [In Japanese with English abstract].
- Cao Y, Cai Y, Takahashi T, Yoshida N, Tohno M, Uegaki R, Nonaka K, Terada F (2011) Effect of lactic acid bacteria inoculant and beet pulp addition on fermentation characteristics and in vitro ruminal digestion of vegetable residue silage. *Journal of Dairy Science* **94**, 3902-3912.
- Chen MM, Liu QH, Xin GR, Zhang JG (2013) Characteristics of lactic acid bacteria isolates and their inoculating effects on the silage fermentation at high temperature. *Letters in Applied Microbiology* **56**, 71-78.
- Chuncheng X, Cai Y, Zhang J, Fukasawa M, Moriya N (2008) Ensiling and subsequent ruminal degradation characteristics of barley tea grounds treated with contrasting additives. *Animal Feed Science and Technology* **141**, 368-374.
- Dehghani MR, Weisbjerg MR, Hvelplund T, Kristensen NB (2012) Effect of enzyme addition to forage at ensiling on silage chemical composition and NDF degradation characteristics. *Livestock Science* **150**, 51-58.
- Donga M, Heb XJ, Tulayakule P, Lic JY, Donga KS, Manabec N, Nakayamab H, Kumagaia S (2010) The toxic effects and fate of intravenously administered zearalenone in goats. *Toxicon* **55**, 523-530.
- Fang J, Horiguchi K, Takahashi T, Goto M, Tanaka O (2006) Effect of the addition of amino acid by-product to unhulled rice and the treatment of germination on the digestibility

- and excretion of indigestible unhulled rice in cattle. *Japanese Journal of Grassland Science* **52**, 170-175 [In Japanese with English abstract].
- Filya I (2003) The effect of *Lactobacillus buchneri* and *Lactobacillus plantarum* on the fermentation, aerobic stability, and ruminal degradability of low dry matter corn and sorghum silages. *Journal of Dairy Science* **86**, 3575-3581.
- Fink-Gremmels J (2008) Mycotoxins in cattle feeds and carry-over to dairy milk. *Food Additives and Contaminations* **25**, 172-180.
- González PML, Alonso VA, Sager R, Morlaco MB, Magnoli CE, Astoreca AL, Rosa CAR, Chiacchiera SM, Dalcero AM, Cavaglieri LR (2008) Fungi and selected mycotoxins from pre- and postfermented corn silage. *Journal of Applied Microbiology* **104**, 1034-1041.
- Goto M, Bae H, Yahaya MS, Karita S, Kim W, Cheng KJ (2003a) Effects of surfactant Tween 80 on enzymatic accessibility and degradation of orchardgrass (*Dactylis glomerata* L.) at different growth stages. *Asian-Australasian Journal Animal Science* **16**, 672-676.
- Goto M, Bae H, Yahaya MS, Karita S, Kim W, Baah J, Sugawara K, Cheng KJ (2003b) Effects of surfactant Tween 80 on forage degradability and microbial growth on the in vitro rumen mixed and pure cultures. *Asian-Australasian Journal Animal Science* **16**, 672-676.
- Hiraoka H, Yamamoto Y, Urakawa S, Yamada H, Karita S, Goto M (2006) Effective fractions and micro biota of fermented juice of epiphytic lactic acid bacteria (FJLB) of rice plant (*Oryza sativa* L.). *Japanese Journal of Grassland Science* **51**, 379-384 [In Japanese with English abstract].
- Hu W, Schmidt RJ, McDonnell EE, Klingerman CM, Kung L Jr. (2009) The effect of *Lactobacillus buchneri* 40788 or *Lactobacillus plantarum* MTD-1 on the fermentation and aerobic stability of corn silages ensiled at two dry matter contents. *Journal of Dairy Science* **92**, 3907-3914.
- Iijima M, Asai T, Zegada-Lizararu W, Nakajima Y, Hamada Y (2005) Productivity and water source of intercropped wheat and rice in a direct-sown sequential cropping system: the effects of no-tillage and drought. *Plant Production Science* **8**, 368-374.
- Ikunaga Y, Sato I, Grond S, Numaziri N, Yoshida S, Yamaya H, Hiradate S, Hasegawa M, Toshima H, Koitabashi M, Ito M, Karlovsky P, Tsushima S (2011) *Nocardioides* sp. strain WSN05-2, isolated from a wheat field, degrades deoxynivalenol, producing the novel intermediate 3-epi-deoxynivalenol. *Applied Microbiology and Biotechnology* **89**, 419-427.
- Intergovernmental Panel on Climate Change (2007) IPCC Forth Assessment Report (AR4).
- Kamoshita A, Ishikawa M, Abe J, Imoto H (2007) Evaluation of water-saving rice-winter crop rotation system in a suburb of Tokyo. *Plant Production Science* **10**, 219-231.
- Kato H (2009) Development of forage and grain feed rice varieties in Japan. *Proceedings of the First International Conference of China-Japan on Feed Science Research*. Guangzhou, China. pp.29-34 [In Japanese].
- Kawamoto H, Otani R, Oshibe A, Yamaguchi H, Deguchi S, Tanaka O, Uozumi S, Watanabe H (2007) Ensilage of wilted whole crop rice (*Oryza sativa* L.) using a roll baler for chopped material: Silage quality in long-term storage. *Grassland Science* **53**, 85-90.
- Kiessling KH, Pettersson H, Sandholm K, Olsen M (1984) Metabolism of aflatoxin, ochratoxin, zearalenone, and three trichothecenes by intact rumen fluid, rumen protozoa, and rumen bacteria. *Applied and Environmental Microbiology* **47**, 1070-1073.
- Kim W, Gamo Y, Yahaya MS, Yimiti W, Goto M (2005) Pretreatment of surfactant Tween 80 and fibrolytic enzyme influencing volatile fatty acid and methane production of a total mixed ration by mixed rumen microorganisms at *in vitro*. *Grassland Science* **51**, 145-152.
- Kung LJ, Myers CL, Neylon JM, Taylor CC, Lazartic J, Mills JA, Whiter AG (2004) The effects of buffered acid-based additives alone or combined with microbial inoculation on the fermentation of high moisture corn and whole-crop barley. *Journal of Dairy Science* **87**, 1310-1316.
- Maclean JL, Dawe DC, Hardy B, Hettel GP (2002) Rice Almanac: Source Book for the Most Important Economic Activity on Earth, 3rd edn (CABI Publishing, Oxford).
- Magan N, Aldred D (2007) Post-harvest control strategies: Minimizing mycotoxins in the food chain. *International Journal of Food Microbiology* **119**, 131-139.
- Matsushita K, Iida S, Ideta O, Sunohara Y, Maeda H, Tamura Y, Kouno S, Takakuwa M (2011) "Tachisuzuka", a new rice cultivar with high straw yield and high sugar content for whole-crop silage use. *Breeding Science* **61**:86-92.
- Miller JD (2008) Mycotoxins in small and maize: Old problems, new challenges. *Food Additives and Contaminations* **25**, 219-230.
- Miyazaki S (2007) Implications of natural toxicants in feed safety. *Japanese Society of Grassland Science* **52**, 257-282 [In Japanese].
- Moss AR, Givens DI, Garnsworthy PC (1994) The effect of alkali treatment of cereal straws on digestibility and methane production by sheep. *Animal Feed Science and Technology* **49**, 245-259.
- Muck RE (1988) Factors influencing silage quality and their implications for management. *Dairy Science* **71**, 2992-3002.
- Nadeau EMG, Russell JR, Buxton DR (2000) Intake, digestibility, and composition of orchardgrass and alfalfa silages treated with cellulose, inoculant, and formic acid fed to lambs. *Journal of Animal Science* **78**, 2980-2989.
- Oerke EC, Meier A, Dehne HW, Sulyok M, Krska R, Steiner U (2010) Spatial variability of fusarium head blight pathogens and associated mycotoxins in wheat crops. *Plant Pathology* **59**, 671-682.
- Okiyama T, Yimamu A, Gamo Y, Yimiti W, Takeda M, Yokoi D, Takahashi T, Horiguchi K, Karita S, Goto M (2008) Effects of a monosodium glutamate fermentation byproduct additive on the fermentation quality and digestibility of whole crops and grains in the whole rice crop silage fed lactating Holstein cattle. *Japanese Journal of Grassland Science* **54**, 64-68 [In Japanese with English abstract].
- Ominski KH., Boadi DA, Wittenberg KM (2006) Enteric methane emissions from backgrounded cattle consuming all-forage diets. *Canadian Journal of Animal Science* **86**, 393-400.
- Park DL, Stoloff L (1989) Aflatoxin control – How a regulatory agency managed risk from an unavoidable natural toxicant in food and feed. *Regulatory Toxicology and Pharmacology* **9**, 109-130.
- Scudamore KA, Livesey CT (1998) Occurrence and significance of mycotoxins in forage crops and silage. *Journal Agriculture and Food Chemistry* **77**, 1-17.
- Shinde S (2010) Feeding of rice whole crop silage for lactating dairy cow. *Japanese Journal of Grassland Science* **55**, 365-372 [In Japanese].
- Shito H, Yamana N, Shibuya Y, Takahashi K (2006) Development of the roll baler for chopped materials. *Japan Agricultural Research Quarterly* **40**, 233-237.
- Sun ZH, Liu SM, Tayo GO, Tang SX, Tan ZL, Lin B, He ZX, Hang XF, Zhou ZS, Wang M (2009) Effects of cellulose or

- lactic acid bacteria on silage fermentation and *in vitro* gas production of several morphological fractions of maize stover. *Animal Feed Science and Technology* **152**, 219-231.
- Tohno M, Kobayashi H, Tajima K, Uegaki R (2012) Strain-dependent effects of inoculation of *Lactobacillus plantarum* subsp. *Plantarum* on fermentation quality of paddy rice (*Oryza sativa* L. subsp. *japonica*) silage. *FEMS Microbiology Letters*. **337**, 112-119.
- Trucksess MW, Richard JL, Stoloff L, McDonald JS, Brumley WC (1983) Absorption and distribution patterns of aflatoxicol and aflatoxins B₁ and M1 in blood and milk of cows given aflatoxin B₁. *American Journal of Veterinary Research* **44**, 1753-1756.
- Tyrolová Y, Výborná A (2011) The effects of wilting and biological and chemical additives on the fermentation process in field pea silage. *Czech Journal Animal Science* **56**, 427-432.
- Uegaki R, Kobayashi H, Tohno M, Tsukiboshi T (2012a) Identification of mycotoxin-producing *Fusarium* spp. isolated from corn and the changes in concentration of fumonisin during the cultivation period. *Grassland Science* **58**, 121-126.
- Uegaki R, Kobayashi H, Inoue H, Tohno M, Tsukiboshi T (2012b) Changes of fumonisin production in rice grain during ensiling. *Animal Science Journal* **84**, 48-53.
- UN Food and Agriculture Organization (2006) Livestock's long shadow. 1-26.
- Urakawa S, Yoshimura Y (2003) Development of the cutting roll baler for the rice whole crop silage. *Japanese Journal of Grassland Science* **49**, 43-48 [In Japanese with English abstract].
- Urakawa S (2009) Development of harvesting machine for forage crop: Mechanization technologies for rice whole-crop silage. *Proceedings of the First International Conference of China-Japan on Feed Science Research*. Guangzhou, China. pp.110-116 [In Japanese].
- Voss KA, Smith GW, Haschek MW (2007) Fumonisin: toxicokinetics, mechanism of action and toxicity. *Animal Feed Science Technology* **137**, 299-325.
- Wang F, Nishino N (2008) Resistance to aerobic stability, of total mixed ration silage: effect of ration formulation, air infiltration and storage period on fermentation characteristics and aerobic stability. *Journal of the Science of Food and Agriculture* **88**, 133-140.
- Warfield CY, Gilchrist DG (1999) Influence of kernel age on fumonisin B₁ production in maize by *Fusarium moniliforme*. *Applied and Environmental Microbiology* **65**, 2853-2856.
- Woloshuk CP, Shim WB (2013) Aflatoxins, fumonisins, and trichothecenes: a convergence of knowledge. *FEMS Microbiology Reviews* **37**, 94-109.
- Xing L, Chen LJ, Han LJ (2009) The effect of an inoculant and enzymes on fermentation and nutritive value of sorghum straw silages. *Bioresource Technology* **100**, 488-491.
- Xu C, Cai Y, Zhang J, Fukasawa M, Moriya N (2008) Ensiling and subsequent ruminal degradation characteristics of barley tea grounds treated with contrasting additives. *Animal Feed Science and Technology* **141**, 368-374.
- Yahaya MS, Goto M, Yimiti W, Smerjai B, Kawamoto Y (2004) Evaluation of fermentation quality of a tropical and temperate forage crops ensiled with additives of fermented juice of epiphytic lactic acid bacteria (FJLB). *Asian-Australasian Journal of Animal Science* **17**, 942-946.
- Yamamoto Y, Yimamu A, Yimiti W, Hiraoka H, Inui K, Takeda M, Yokoi D, Karita S, Goto M (2008) Effects of various amino acid fermentation byproducts on the fermentation quality and feed characteristics of Italian ryegrass (*Lolium multiflorum* Lam.) silage. *Japanese Journal of Grassland Science* **53**, 289-294 [In Japanese with English abstract].
- Yimamu A, Fukui H, Hiraoka H, Yamamoto Y, Ogawa A, Takeda M, Yokoi D, Karita S, Goto M (2008) Effects of mixture of Lysine and Threonine fermentation byproducts on the fermentation quality and aerobic stability of corn (*Zea mays* L.) silage. *Japanese Journal of Grassland Science* **54**, 1-6. [In Japanese with English abstract].
- Yimiti W, Gamo Y, Kim W, Hiraoka H, Takeda M, Horiguchi K, Takahashi T, Yoshida N, Goto M (2006a) Effects of monosodium glutamate fermentation byproduct on the fermentation quality and *in situ* rumen degradability of whole crop barley (*Hordeum vulgare* L.) silage. *Grassland Science* **52**, 17-22 [In Japanese with English abstract].
- Yimiti W, Hiraoka H, Yahaya MS, Takeda M, Kim W, Kariya S, Horiguchi K, Takahashi T, Goto M (2006b) Effects of phenylalanine fermentation byproduct and sugarcane molasses on fermentation quality and rumen degradation of whole crop barley (*Hordeum vulgare* L.) silage in situ. *Grassland Science* **52**, 73-79.
- Zahiroddini H, Baah J, Absalom W, McAllister TA (2004) Effect of an inoculant and hydrolytic enzymes on fermentation and nutritive value of whole crop barley silage. *Animal Feed Science and Technology* **117**, 317-330.