



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

22nd International Grassland Congress

---

## Characterization and Identification of Lactic Acid Bacteria Isolated from Silage Prepared in Lao PDR

Yimin Cai

*Japan International Research Center for Agricultural Sciences, Japan*

Huili Pang

*Zhengzhou University, China*

Viengsakoun Napasirth

*National University of Laos, Lao*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/22/1-11/24>

The 22nd International Grassland Congress (Revitalising Grasslands to Sustain Our Communities) took place in Sydney, Australia from September 15 through September 19, 2013.

Proceedings Editors: David L. Michalk, Geoffrey D. Millar, Warwick B. Badgery, and Kim M.

Broadfoot

Publisher: New South Wales Department of Primary Industry, Kite St., Orange New South Wales, Australia

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

# Characterization and identification of lactic acid bacteria isolated from silage prepared in Lao PDR

Yimin Cai<sup>A</sup>, Huili Pang<sup>B</sup> and Viengsakoun Napasirth<sup>C</sup>

<sup>A</sup> Japan International Research Center for Agricultural Sciences, Tsukuba, Japan, [www.jircas.affrc.go.jp](http://www.jircas.affrc.go.jp)

<sup>B</sup> Henan Provincial Key Laboratory of Ion Beam Bio-engineering, Zhengzhou University, Henan, People's Republic of China, <http://english.zzu.edu.cn>

<sup>C</sup> Department of Livestock and Fisheries, Faculty of Agriculture, National University of Laos, Vientiane, Lao PDR, [www.nuol.edu.la](http://www.nuol.edu.la)

Contact email: [cai@affrc.go.jp](mailto:cai@affrc.go.jp)

**Keywords:** Lactic acid bacteria, 16S rRNA, silage.

## Introduction

Silage is now the most common preserved cattle feed in many countries and lactic acid bacteria (LAB) play an important role during silage fermentation process. In order to establish an animal feed production system to cover the shortage of animal feed in the dry season of Lao PDR, silage fermentation technology is needed that has potential application in poor villages. Therefore, the characterization of LAB in various forage crops and their species identification requires further study. In the present study, the characterization of LAB species isolated from forage crops in Lao PDR was examined. In order to determine their taxonomic status, these strains were also studied by 16S rRNA sequence analysis.

## Materials and methods

Silage samples were prepared in Vientiane, Lao PDR using corn (*Zea mays*), paddy rice (*Oryza sativa*) straw and sugarcane (*Saccharum officinarum*) tops. LAB were isolated on MRS agar (Difco Laboratories, Michigan, USA). Gram stain, morphology, catalase activity, spore formation, motility, nitrate reduction and gas production from glucose were determined. Carbohydrate fermentation patterns were examined using AP 50 CH strips. The isomers of lactate formed from glucose were determined enzymatically with reagents obtained from Boehringer GmbH, Mannheim, Germany. The 16S rRNA sequence was analyzed as described by Cai *et al.* (1999; 2012).

## Results and Discussion

Seven representative strains of lactic acid bacteria were isolated from silage in Vientiane, Laos (Table 1). These strains were Gram-positive, catalase-negative bacteria. Strains LA1 and LA 46 were homofermentative rods that formed DL-lactic acid. Strain LA 1 could grow in low pH condition (3.5), but Strain LA 46 did not grow below pH 4.5. Strains

LA 3 and LA 8 were homofermentative cocci that formed L(+) lactic acid. Strains LA 22 and LA 37 were heterofermentative cocci that formed D(-) lactic acid. Strain LA 40 was homofermentative cocci that formed DL lactic acid, and could grow at pH 3.5. Based on 16S rRNA gene sequence analysis, these isolates were identified as *Lactobacillus plantarum*, *Enterococcus faecalis*, *Leuconostoc mesenteroides*, *Leuconostoc mesenteroides*, *Lactococcus lactis* subsp. *Lactis*, *Weissella confusa*, *Pediococcus pentosus*, *Lactobacillus brevis*, respectively.

In the present study, the isolates were Gram-positive and catalase-negative cocci or rod that produced lactate from glucose, formed approximately equal quantities of L(+), DL and D(-) lactic acid. These properties showed that these strains belong to the genus of LAB. However, as available phenotypic procedures are difficult to assign isolates to known species because it is difficult to differentiate easily and clearly between species of LAB (Cai 1999; Ennahar *et al.* 2003; Pang *et al.* 2011). Based on 16S rRNA gene sequence analysis, these isolates could be assigned to species level.

It is well established that LAB play an important role in silage fermentation and animal health. LAB are naturally present on forage crops, silage and animal intestines, therefore collection and identification of LAB isolated from silage is important. In the present study, *Lactobacillus plantarum* and *Weissella confusa* were the most dominant LAB in the silage. Isolation and identification of the LAB strains are key first steps for future research in which the relationship between LAB species and silage fermentation is being evaluated.

## Conclusions

Seven representative strains of lactic acid bacteria were isolated from silage in Vientiane, Lao PDR. These strains were Gram-positive, catalase-negative cocci or rod, they are homo or heterofermentative

**Table 1. Characteristics of lactic acid bacteria isolated from silage in Laos. Isolates were *Lactobacillus plantarum* LA 1 (1), *Enterococcus faecalis* LA 8 (2), *Leuconostoc pseudo-mensenteroides* LA 22 (3), *Lactococcus lactis* subsp. *lactis* LA 3 (4), *Weissella confusa* LA 37 (5), *Pediococcus pentosus* LA 40 (6), and *Lactobacillus brevis* LA 46 (7). All strains were Gram-positive and catalase-negative bacteria. +, positive; -, negative; w, weakly positive.**

Character	Isolate 1	Isolate 2	Isolate 3	Isolate 3	Isolate 5	Isolate 6	Isolate 7
Cell form	Rod	Cocci	Cocci	Cocci	Cocci	Cocci	Rod
Fermentation type	Homo	Homo	Hetero	Homo	Hetero	Homo	Hetero
Lactate isomer	DL	L(+)	D(-)	L(+)	D(-)	DL	DL
Growth at temperature							
15 °	+	+	+	+	+	+	+
45 °	+	+	w	+	+	+	+
Growth at pH							
3.5	+	-	-	-	-	+	-
4.0	+	-	-	-	-	+	-
4.5	+	+	-	+	w	+	w
Growth at MRS medium							
OD 620	2.2	0.8	0.7	0.7	0.9	1.7	1.7
Lactate production (g/L)	14.5	8.7	8.9	8.8	8.1	12.3	11.2
Final pH	3.6	4.7	4.6	4.7	4.6	3.8	3.7
Similarity of 16S rRNA sequence	99.6	99.7	99.2	99.5	99.8	99.7	99.1

type and formed L(+), DL or D(-) lactic acid. Based on 16S rRNA gene sequence analysis, these isolates were identified as species of *Lactobacillus*, *Enterococcus*, *Leuconostoc*, *Weissella*, *Lactococcus* and *pediococcus*, respectively. *Lactobacillus plantarum* and *Weissella confusa* were the mostly dominate members of the population in these silages. The relationship between LAB species and silage fermentation is currently in progress.

## References

- Cai Y (1999) Identification and characterization of *Enterococcus* species isolated from forage crops and their influence on silage fermentation. *Journal of Dairy Science* **82**, 2466–2471.
- Cai Y, Kumai S, Ogawa M, Benno Y, Nakase T (1999) Characterization and identification of *Pediococcus* species isolated from forage crops and their application for silage preparation. *Applied Environmental Microbiology* **65**, 2901-2906.
- Cai Y, Pang H, Kitahara M, Ohkuma M. (2012) *Lactobacillus nasuensis* sp. nov., a lactic acid bacterium isolated from silage. *International Journal of Systematic and Evolutionary Microbiology* **62**, 1140–1144.
- Ennahar S, Cai Y, Fujita Y (2003) Phylogenetic Diversity of Lactic Acid Bacteria Associated with Paddy Rice Silage as Determined by 16S Ribosomal DNA Analysis. *Applied Environmental Microbiology* **69**(1), 444-451.
- Pang H, Qin G, Tan Z, Li Z, Wang Y, Cai Y (2011) Natural populations of lactic acid bacteria associated with silage fermentation as determined by phenotype, 16S ribosomal RNA and *RecA* gene analysis. *Systematic and Applied Microbiology* **34**, 235-241.