



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

XXII International Grassland Congress

Grazing Termination Dates of Summer-Dormant Flecha Tall Fescue

Sindy M. Interrante

The Samuel Roberts Noble Foundation

Twain J. Butler

The Samuel Roberts Noble Foundation

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/22/1-12/10>

The XXII International Grassland Congress (Revitalising Grasslands to Sustain Our Communities) took place in Sydney, Australia from September 15 through September 19, 2013.

Proceedings Editors: David L. Michalk, Geoffrey D. Millar, Warwick B. Badgery, and Kim M.

Broadfoot

Publisher: New South Wales Department of Primary Industry, Kite St., Orange New South Wales, Australia

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Grazing termination dates of summer-dormant Flecha tall fescue

Sindy M Interrante and Twain J Butler

The Samuel Roberts Noble Foundation, www.noble.org 2510 Sam Noble Parkway, Ardmore OK 73401, USA

Contact email: tjbutler@noble.org

Keywords: Grazing, production, summer-dormant tall fescue.

Introduction

Summer-dormant, Mediterranean-type tall fescue (*Festuca arundinacea* Schreb.) has potential to replace summer-active, Continental-type tall fescue and traditional, annual small grain graze-out systems in the Southern Great Plains region of the USA. This region is characterized by severe water deficits accompanied by extreme heat in summer, and by relatively mild, rainy winters (Malinowski *et al.* 2009). Although the climate of the southern Great Plains is different from the Mediterranean climate, the temperature and precipitation patterns during summer are historically similar. However, in the past decade, the bimodal precipitation pattern with peaks in May and September has become highly unpredictable, resulting in delayed planting of small grains in autumn and lack of winter forage for grazing livestock (Malinowski *et al.* 2009). As a perennial forage crop, summer-dormant tall fescue provides a source of forage during the winter months when warm-season grasses are dormant, reducing pasture establishment costs, soil erosion, and the time and labor associated with annual forage systems (Kindiger and Conley 2002; Beck *et al.* 2008; Islam *et al.* 2011).

Replacing summer-active with summer-dormant cool-season grasses can provide some resilience in the forage systems to extreme seasonal precipitation and temperature patterns (Clark and Harris 2009; Malinowski *et al.* 2005). These conditions, combined with relatively mild winters, allow summer-dormant tall fescue to be better adapted and more persistent in the southern Great Plains than traditional, summer-active types of tall fescue and other cool-season perennial grasses (Hopkins and Bhamidimarri, 2009; Malinowski *et al.* 2009). However there are no best-management practices that address the timing of grazing cessation as it relates to summer-dormant tall fescue stand persistence and animal production. The objectives of this study were to determine the effects of four grazing termination dates on endophyte-free 'Flecha' summer-dormant tall fescue forage production, animal production (average daily gain [ADG], gain, and grazing days), and tall fescue persistence (% stand and root mass).

Methods

Paddocks (0.1 ha each) were planted on 19 September 2010 at 17 kg pure live seed /ha of monoculture summer-dormant tall fescue. Tall fescue was planted according to

guidelines outlined by Butler *et al.* (2008), whereby annual grass weeds were sprayed with glyphosate in the spring to prevent seed production, followed by a second application of glyphosate after rainfall to control emerged grass weeds prior to planting tall fescue in the autumn. Soil type was a Weatherford fine sandy loam (Fine-loamy, siliceous, active, thermic Ultic Haplustalfs). Prior to planting in September, 45 kg P₂O₅, 112 kg K₂O, and 112 kg N/ha were applied. Paddocks were not grazed during the establishment season to avoid the confounding effects of grazing on persistence.

In both grazing season (2011-12, 2012-13), paddocks were continuously stocked with a variable stocking rate using ewes (*Ovis aries*; 41 ± 3.4 kg initial body weight), and grazing began on all treatments on 11 January 2012 and 30 November 2012. In both seasons (2011-12, 2012-13), grazing ceased on the terminations date of 3 May, 16 May, 31 May, and 9 June. Stocking rates were adjusted every 28-d based on forage mass (FM) with put-and-takes ewes. Forage mass was measured every 28 d during the grazing season. After grazing termination and after plants entered dormancy in mid-summer (July), FM, root mass, and percent tall fescue stand measurements were taken to estimate storage organ mass and overall health/status of the paddocks prior to recovery/green-up the following autumn.

Percent stand measurements were determined by counting the live plants within 100 - 10 x 10 cm squares of a 1 m² quadrat. As late spring temperatures increase and soil moisture is reduced, summer-dormant tall fescue prioritizes photosynthate to increased root and stem base masses and to storage as nonstructural carbohydrates as opposed to new above-ground growth (Thornton *et al.* 2000). Severe defoliation prior to prolonged stress such as drought may limit the persistence of cool-season grasses. Nonstructural carbohydrate reserves are used for respiration during dormancy and for subsequent regrowth under favorable growing conditions (Richards 1993), therefore, fewer carbohydrate reserves may result in less vigorous regrowth and potential stand failure.

Results

Precipitation during the growing season from September through June was approximately 59% below average during the establishment season (2010-11), similar to the 30-yr average (900 mm) in the 2011-12 season, and 27%

below average during the 2012-13 season. In the first grazing season, FM decreased (both $P < 0.001$) as the grazing season progressed. However, at their respective times of termination, there were no differences in FM between the grazing termination date treatments (average 840 kg DM/ha). There were fewer grazing days per hectare ($P = 0.01$) on 16 May (4470 d/ha) than the other termination dates (average 5656 d/ha), which was probably due to the lesser FM on 16 May. There was a tendency ($P = 0.09$) towards greater FM in July associated with the earliest (3 May) grazing termination date (2580 kg DM/ha), however there was no difference ($P = 0.55$) in root mass (average 1150 g DM/m) between termination dates. At July dormancy, tall fescue stand was greatest ($P = 0.05$) for the earliest termination date (3 May; 79%) and least in 16 May and 6 June (44 and 47%, respectively), while stand on 31 May was intermediate (65%). When measured in November 2012 prior to grazing initiation in season 2, stand was reduced ($P = 0.04$) across all treatments to 57% compared to 69% in July.

Conclusion

After one year, data show a reduction in percent stand and a tendency towards less forage mass and root mass associated with grazing termination dates later than 3 May, which may indicate the need to cease grazing at this time of year in order to maintain stand persistence. This research will be conducted for an additional year with the anticipation of gaining more information on the grazing management of summer-dormant tall fescue in the Southern Great Plains.

References

- Beck PA, Gunter SA, Lusby KS, West CP, Watkins KB, Hubbell DS III. (2008) Animal performance and economic comparison of novel and toxic endophyte tall fescues to cool-season annuals. *Journal of Animal Science* **86**, 2043-2055.
- Butler TJ, Islam AM, Muir JP (2008) Establishment of cool-season perennial grasses in the southern Great Plains. *Forage and Grazinglands* doi:10.1094/FG-2008-0911-01-RS.
- Clark S, Harris C (2009) Summer dormancy in Australian perennial grasses: Historical background, a simulation study, and current research. *Crop Science* **49**, 2328–2334.
- Hopkins AA, Bhamidimarri S (2009) Breeding summer-dormant grasses for the United States. *Crop Science* **49**, 2359–2362.
- Islam AM, Biermacher JT, Reuter RR, Interrante SM, Hopkins AA, Cook BJ, Bouton JH, Butler JT (2011) Production and economics of grazing rye-annual ryegrass and tall fescue systems. *Agronomy Journal* **103**, 558-564.
- Kindiger B, Conley T (2002) Competition and survival of perennial cool-season grass forages seeded with winter wheat in the southern Great Plains. *Journal of Sustainable Agriculture* **21**, 27-45.
- Malinowski DP, Kigel J, Pinchak WE (2009) Water deficit, heat tolerance, and persistence of summer-dormant grasses in the U.S. Southern Plains. *Crop Science* **49**, 2363–2370.
- Malinowski DP, Zuo H, Kramp BA, Muir JP, Pinchak WE (2005) Obligatory summer-dormant cool-season perennial grasses for semiarid environments of the Southern Great Plains. *Agronomy Journal* **97**, 147–154.
- Richards JH (1993) Physiology of plants recovering from defoliation. pp. 85-94. In *Proceedings of the 17th international Grassland Congress*, Rockhampton, Australia. 8-12 Feb. 1993. Dunmore Press Ltd., Palmerston North, New Zealand.
- Thornton B, Millard P, Bausenwein U (2000) Reserve formation and recycling of carbon and nitrogen during regrowth of defoliated plants. pp. 85–99. In : G Lemaire et al. (eds.). *Grassland ecophysiology and grazing ecology*. CAB International, New York.