



An Efficient Sampling Protocol for Sagebrush/Grassland Monitoring

Larry L. Larson
Oregon State University

Mounir Louhaichi
International Center for Agricultural Research in the Dry Areas, Jordan

Patrick E. Clark
U.S. Department of Agriculture

Douglas E. Johnson
Oregon State University

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/22/1-13/9>

The XXII International Grassland Congress (Revitalising Grasslands to Sustain Our Communities) took place in Sydney, Australia from September 15 through September 19, 2013.

Proceedings Editors: David L. Michalk, Geoffrey D. Millar, Warwick B. Badgery, and Kim M.

Broadfoot

Publisher: New South Wales Department of Primary Industry, Kite St., Orange New South Wales, Australia

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

An efficient sampling protocol for sagebrush/grassland monitoring

Larry L Larson^A, Mounir Louhaichi^B, Patrick E Clark^C and Douglas E Johnson^D

^A Department of Rangeland Ecology and Management, Oregon State University, Oregon, USA

^B International Center for Agricultural Research in the Dry Areas (ICARDA), Amman, Jordan

^C USDA/ARS, Boise, Idaho, USA

^D Department of Rangeland Ecology and Management, Oregon State University, Oregon, USA

Contact email: llarson@eou.edu

Keywords: Random sampling, digital photography, GPS unit, protocol.

Introduction

Rangeland scientists and quantitative ecologists have developed numerous methods and monitoring techniques that can be used for vegetation sampling (Barbour *et al.* 1987). The methods used to position samples (transects, quadrats, lines, and points) vary and can be classed as selective, capricious, systematic, or random. One of the prerequisites for valid statistical inference is that samples are taken randomly. A random sampling procedure implies that all elements or units of the population being studied have an equal chance of being represented in the sample. It also implies that selection of an element or unit does not influence the chance of other units being sampled. Data that is collected using random sampling procedures can be used to compare attributes of different populations or sites such as vegetative cover, density, production, growth rates, etc. This paper suggests a random sampling protocol that can be easily applied in the field for sagebrush/grassland monitoring.

Material and Methods

Our protocols typically employ a monopod or pole with a mounting head which allows us to point a digital camera vertically downward from a fixed height (Louhaichi *et al.* 2010). Also attached next to the camera is a continuously recording (1 Hz) WAAS enabled GPS unit (Fig. 1). This combination of instruments allows us to take high-resolution, vertical images from between 1.5 m and 5.0 m above the sample plot (Booth *et al.* 2004). The higher the camera, the larger the ground coverage and coarser the resolution. Advances in digital imaging technologies have compensated for the loss of resolution and modern cameras have sensors that can capture images with a pixel count of 5784 x 3861, which translates to a ground pixel resolution of 1mm for an image covering 5.78 by 3.86 m. The height, species list, and reference photos are completed in field using a pre-determined sampling pattern for each ecological site (Fig. 2 and 3).

The data collection process can be divided into a series of steps in two operations: field (Fig. 4) and office (Fig. 5).

Results and Discussion

Since photographs can be taken quickly, many quadrats can be photographically sampled during a workday. It is not



Figure 1. Digital charting apparatus.



Figure 2. Landscape photo is taken (left) then reference photographs are taken of plants that are identifiable (right).

- From the Reference Location Sample in the Following Pattern with 10 Photos per Line




Figure 3. Sampling pattern used within an ecological site.




Figure 4. Diagram of a field protocol for collecting digital charting information that can be used for random sampling.




Figure 5. The office process involves positioning, rotating, and scaling images and saving them as geo-registered bitmaps.

unusual for us to sample 4 quadrats (plots) per minute once we are at a site and set up in the field, providing the area being sampled is level and samples are 2 to 4 paces apart. With low staff sampling, 60 images provide approximately 150 m² of ground that is photographed. If 25 samples of 0.25 m² are randomly chosen from the complete photographic record, we have randomly sampled less than 5% of the total area photographed. Once geo-referenced photographic samples are taken, they can be examined, read, and/or processed (Fig. 6).



Figure 6. Example results from digital sampling of photo quadrats on a low sagebrush/perennial grasses site.

Conclusions

The proposed digital charting protocol can be used to monitor perennial grass, forb, shrub, and weed cover and density as well as plant species composition change over years, the presence of distinctive plants, and mortality of specific perennial plants. It can also be used to evaluate the intensity of grazing, presence of insect pests, and the effectiveness of herbicide treatments. Because field sampling is so fast, many sites can be sampled in the same phenological stage, encouraging uniformity and consistency of collected information.

References

- Barbour MG, Burk JH, Pitts WD (1987) *Terrestrial Plant Ecology*. Chapter 9: Method of sampling the plant community. Menlo Park, CA, USA: Benjamin/Cummings Publishing Co.
- Booth DT, Cox SE, Louhaichi M, Johnson DE (2004) Lightweight camera stand for close-to-earth remote sensing. *Journal of Range Management* **57**, 675-678.
- Louhaichi M, Johnson MD, Woerz AL, Jasra AW, Johnson DE (2010) Digital charting technique for monitoring rangeland vegetation cover at local scale. *International Journal of Agriculture and Biology*, **12**, 406-410.