11-4-2016

Sign-Problem-Free Monte Carlo Simulation of Certain Frustrated Quantum Magnets

Fabien Alet

Université de Toulouse, France

Kedar Damle

Tata Institute of Fundamental Research, India

Sumiran Pujari

University of Kentucky, sumiran.pujari@uky.edu

Follow this and additional works at: https://uknowledge.uky.edu/physastron_facpub

Part of the [Physics Commons](#)

Right click to open a feedback form in a new tab to let us know how this document benefits you.

Repository Citation

Alet, Fabien; Damle, Kedar; and Pujari, Sumiran, "Sign-Problem-Free Monte Carlo Simulation of Certain Frustrated Quantum Magnets" (2016). *Physics and Astronomy Faculty Publications*. 405.

https://uknowledge.uky.edu/physastron_facpub/405

This Article is brought to you for free and open access by the Physics and Astronomy at UKnowledge. It has been accepted for inclusion in Physics and Astronomy Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Sign-Problem-Free Monte Carlo Simulation of Certain Frustrated Quantum Magnets

Digital Object Identifier (DOI)

<https://doi.org/10.1103/PhysRevLett.117.197203>

Notes/Citation Information

Published in *Physical Review Letters*, v. 117, issue 19, 197293, p. 1-7.

© 2016 American Physical Society

The copyright holder has granted the permission for posting the article here.

Sign-Problem-Free Monte Carlo Simulation of Certain Frustrated Quantum Magnets

Fabien Alet,¹ Kedar Damle,² and Sumiran Pujari³

¹Laboratoire de Physique Théorique, IRSAMC, Université de Toulouse, CNRS, 31062 Toulouse, France

²Department of Theoretical Physics, Tata Institute of Fundamental Research, Mumbai 400 005, India

³Department of Physics and Astronomy, University of Kentucky, Lexington, Kentucky 40506-0055, USA
(Received 21 January 2016; revised manuscript received 27 August 2016; published 4 November 2016)

We introduce a quantum Monte Carlo (QMC) method for efficient sign-problem-free simulations of a broad class of frustrated $S = 1/2$ antiferromagnets using the basis of spin eigenstates of clusters to avoid the severe sign problem faced by other QMC methods. We demonstrate the utility of the method in several cases with competing exchange interactions and flag important limitations as well as possible extensions of the method.

DOI: 10.1103/PhysRevLett.117.197203

Quantum Monte Carlo (QMC) simulations compute equilibrium properties of a many-body system by importance sampling of the canonical partition function $Z = \text{Tr} \exp(-H/T)$, where H is the many-body Hamiltonian and T is the temperature [1–3]. They have emerged as a major tool for the study of lattice Hamiltonians that either model low- T thermodynamic properties of interesting strongly correlated materials [4,5] or provide concrete realizations of novel phases in such condensed matter systems [6,7]. However, models of *geometrically frustrated magnets* [5], in which antiferromagnetic interactions compete with each other due to the geometry of the exchange pathways, have typically remained beyond the reach of QMC methods. This is due to the presence of a *sign problem*, whereby the weights assigned to individual Monte Carlo configurations are no longer strictly positive in the commonly used basis of eigenstates of S_r^z , the z component of each spin. In such cases, the average sign decreases exponentially with the system size and inverse temperature, leading to unmanageably large statistical errors in the estimation of physical quantities. A similar sign problem crops up in diverse settings ranging from QCD to strongly correlated metals, and a general solution is considered unlikely [8].

Limited progress has been possible in a few cases, for instance, in anisotropic systems in which the frustration affects only S_r^z (thereby allowing sign-free simulation in the z basis [9–17]) or when the sign problem of the full theory can be finessed at low T by working with a low-energy effective Hamiltonian which has no sign problem [18–22]. For some models, sign-free simulations are possible by virtue of specific symmetries of the Hamiltonian [23–29]. In other strongly correlated systems with a full-fledged sign problem, progress has been made in some cases by developing *improved estimators* for computing physical observables [30–33]. In principle, one could also try to find an alternate basis in which all configurations have positive weights. However, this has been possible only in a few

interesting cases [34,35], including some models of topologically ordered states of matter [36–38].

Synopsis.—In this Letter, we introduce a QMC method that works in the basis of spin eigenstates of clusters to simulate a large class of frustrated quantum magnets in a provably sign-free manner. We focus our discussion on systems in which the clusters in question are made up of two spin-1/2 moments ($\vec{S}_{Ir}$ and $\vec{S}_{IIr}$) located on *layers* I and II at *sites* r of a bipartite Bravais lattice in any spatial dimension (Fig. 1), with Hamiltonian

$$H_{\text{bilayer}} = \sum_{\langle r_a r_b \rangle} (\mathcal{J}_z S_{Ir_a}^z S_{Ir_b}^z + \mathcal{J}_\perp \vec{S}_{Ir_a}^\perp \cdot \vec{S}_{Ir_b}^\perp + \text{I} \leftrightarrow \text{II}) + \sum_{\langle r_a r_b \rangle} (\mathcal{K}_z S_{Ir_a}^z S_{IIr_b}^z + \mathcal{K}_\perp \vec{S}_{Ir_a}^\perp \cdot \vec{S}_{IIr_b}^\perp + \text{I} \leftrightarrow \text{II}) + \sum_r (\mathcal{D}_z S_{Ir}^z S_{IIr}^z + \mathcal{D}_\perp \vec{S}_{Ir}^\perp \cdot \vec{S}_{IIr}^\perp), \quad (1)$$

FIG. 1. Vertices that appear in the SSE operator string for H_{bilayer} , with corresponding weights in the canonical cluster basis. All other valid vertices are obtained by symmetry operations that exchange left and right, or upper and lower, legs (keeping the weight fixed). The constant C in the function $f(l_A, l_B, m_A, m_B) = C - J_z m_A m_B - \zeta(\Delta_z - \Delta_\perp)(m_A^2 + m_B^2) - \zeta \Delta_\perp [l_A(l_A + 1) + l_B(l_B + 1)]$ is chosen to ensure that $f \geq 0$. Bottom right: Lattice structure and exchange couplings of H_{bilayer} . Vertices and lattice structure for H_{mixed} are detailed in Supplemental Material [56].

where the nearest-neighbor links of this Bravais lattice have been denoted by $\langle r_a r_b \rangle$ to emphasize its bipartite nature and $\vec{S}_{I/IIr}^\perp$ represents the vector formed by the two transverse components (x and y) of these spins. Geometric frustration of the exchange interactions leads to a severe sign problem for other QMC methods whenever $\mathcal{D}_\perp \mathcal{K}_\perp \mathcal{J}_\perp > 0$. Our central result is that such frustration leads to no sign problems in our method whenever the interactions in H_{bilayer} are constrained to satisfy *at least one* of the following three conditions: (i) $\mathcal{K}_z = \mathcal{J}_z$; (ii) $\mathcal{K}_\perp = \mathcal{J}_\perp$; (iii) $\mathcal{K}_\perp = -\mathcal{J}_\perp$. *Fully frustrated bilayer* systems [39–44], which have infinitely many conserved quantities, represent a special case with (i) and (ii) both being satisfied. The method also works when the B sublattice only hosts a single spin-1/2 moment $\vec{S}_{r_b}$ that couples symmetrically to $\vec{S}_{I r_a}$ and $\vec{S}_{II r_a}$ on neighboring A sublattice sites [45–47]:

$$H_{\text{mixed}} = \sum_{\langle r_a r_b \rangle} (\mathcal{J}_z \mathcal{S}_{I r_a}^z \mathcal{S}_{r_b}^z + \mathcal{J}_\perp \vec{S}_{I r_a}^\perp \cdot \vec{S}_{r_b}^\perp + \text{I} \leftrightarrow \text{II}) \\ + \sum_{r_a} (\mathcal{D}_z \mathcal{S}_{I r_a}^z \mathcal{S}_{II r_a}^z + \mathcal{D}_\perp \vec{S}_{I r_a}^\perp \cdot \vec{S}_{II r_a}^\perp). \quad (2)$$

For $\mathcal{D}_\perp > 0$, the usual QMC method has a sign problem, which is no longer present in our QMC scheme. Our method is also expected to apply to other such models with infinitely many conserved quantities [48–54]. Additionally, (iii) includes interesting examples of frustrated bilayer magnets with full SU(2) symmetry and no extra conservation laws. Some of our results have been obtained independently in recent parallel work [55].

Key idea.—We use the stochastic series expansion (SSE) QMC framework [3] and work at each Bravais lattice site r in the basis $\{|l, m\rangle\}$ of simultaneous eigenstates of the total spin $\vec{L}_r^2$ and its z component L_r^z , with eigenvalues $l(l+1)$ and m , respectively. For H_{bilayer} , we define $\vec{L}_r = \vec{S}_{I r} + \vec{S}_{II r}$ on both sublattices. For H_{mixed} , this is modified on the B sublattice by defining $\vec{L}_{r_b} = \vec{S}_{r_b}$. We decompose the Hamiltonian into terms living on bonds $\langle r_a r_b \rangle$ of the bipartite Bravais lattice, with the terms proportional to $\mathcal{D}_z$ and $\mathcal{D}_\perp$ at each site r being shared equally among all bonds emanating from r :

$$H_{1\langle r_a r_b \rangle} = J_z L_{r_a}^z L_{r_b}^z + \zeta \Delta_z [(L_{r_a}^z)^2 + (L_{r_b}^z)^2] \\ + \zeta \Delta_\perp [(\vec{L}_{r_a}^\perp)^2 + (\vec{L}_{r_b}^\perp)^2] - C, \\ H_{2\langle r_a r_b \rangle}^\pm = \frac{J_\perp}{2} (L_{r_a}^\pm \cdot L_{r_b}^\mp), \\ H_{3\langle r_a r_b \rangle} = K_z N_{r_a}^z N_{r_b}^z, \\ H_{4\langle r_a r_b \rangle}^\pm = \frac{K_\perp}{2} (N_{r_a}^\pm \cdot N_{r_b}^\mp), \quad (3)$$

with $\vec{L}_r^\perp$ the vector made of transverse (x/y) components of $\vec{L}_r$, $\vec{N}_r = \vec{S}_{I r} - \vec{S}_{II r}$, $L_r^\pm = L_r^x \pm i L_r^y$, $N_r^\pm = N_r^x \pm i N_r^y$, ζ

the inverse coordination number of the bipartite lattice, C a constant introduced to ensure negativity of all matrix elements of the *diagonal* operator $H_{1\langle r_a r_b \rangle}$ in our chosen basis, and $\Delta_\mu = \mathcal{D}_\mu/2$, $J_\mu = (\mathcal{J}_\mu + \mathcal{K}_\mu)/2$, $K_\mu = (\mathcal{J}_\mu - \mathcal{K}_\mu)/2$ (for $\mu = z, \perp$). Using this decomposition, we have $H_{\text{bilayer}} = \sum_{\langle r_a r_b \rangle} H_{1\langle r_a r_b \rangle} + H_{3\langle r_a r_b \rangle} + H_{2\langle r_a r_b \rangle}^+ + H_{4\langle r_a r_b \rangle}^+ + H_{2\langle r_a r_b \rangle}^- + H_{4\langle r_a r_b \rangle}^-$. H_{mixed} , when decomposed in the same way, has analogs only of the $H_{1\langle r_a r_b \rangle}$ and $H_{2\langle r_a r_b \rangle}^\pm$ terms [56].

Working within the SSE framework with this decomposition of H_{bilayer} , one writes

$$Z = \sum_{n=0}^{\infty} \frac{1}{n! T^n} \sum_{\mathcal{S}_n} \langle \alpha_0 | (-H_n) | \alpha_{n-1} \rangle \langle \alpha_{n-1} | (-H_{n-1}) | \alpha_{n-2} \rangle \dots \\ \times \langle \alpha_1 | (-H_1) | \alpha_0 \rangle, \quad (4)$$

where the sum over *operator strings* $\mathcal{S}_n$ of length n is implemented by allowing each $|\alpha_j\rangle$ to range over the full basis of states and each H_j to range over all bond operators $H_{1\langle r_a r_b \rangle}$, $H_{2\langle r_a r_b \rangle}^\pm$, $H_{3\langle r_a r_b \rangle}$, and $H_{4\langle r_a r_b \rangle}^\pm$. Along with the factor of $1/(n! T^n)$, the product of matrix elements appearing in the summand serves as the Monte Carlo weight of each operator string, and the QMC simulation proceeds by performing an importance sampling of Z .

Proof of positive-weight property.—Clearly, the weight of an operator string *does not* depend on the choice of arbitrary phase factors attached to individual basis states, since these phase factors cancel in pairs in the product of matrix elements that sets the weight. Fixing these phases, we define the *canonical* cluster basis as follows: $|s\rangle = (|\uparrow_I \downarrow_{II}\rangle - |\downarrow_I \uparrow_{II}\rangle)/\sqrt{2}$, $|t_{+1}\rangle = |\uparrow_I \uparrow_{II}\rangle$, $|t_{-1}\rangle = |\downarrow_I \downarrow_{II}\rangle$, and $|t_0\rangle = (|\uparrow_I \downarrow_{II}\rangle + |\downarrow_I \uparrow_{II}\rangle)/\sqrt{2}$. Next, we classify the off-diagonal matrix elements contributing to the weight of an operator string into five types (Fig. 1): (i) *m-type processes* that hop one quantum of L^z along link $\langle r_a r_b \rangle$ between two neighboring triplet clusters, (ii) *l-type processes* that exchange the states $|s\rangle$ and $|t_0\rangle$ of two neighboring clusters, (iii) *p-type processes* that take neighboring singlet clusters and promote both to the $|t_0\rangle$ state or vice versa, (iv) *lm-type processes* that exchange singlet and triplet states of neighboring clusters and simultaneously hop one quantum of L^z , and (v) *pm-type processes* that take neighboring singlet clusters to states $|t_{\pm 1}\rangle$ and $|t_{\mp 1}\rangle$, respectively, or vice versa.

All processes of a given type have a fixed sign for the corresponding matrix elements between basis states (Fig. 1). Therefore, a positive weight is guaranteed if $\mathcal{N}_t$, the number of occurrences (in any string $\mathcal{S}_n$) of t -type processes, has even parity for each type t . These parities are constrained by the periodicity of the operator string $\mathcal{S}_n$; i.e., the starting state $|\alpha_0\rangle$ is recovered after the action of n operators. Since pair creation of the l quantum number must be balanced by

corresponding pair destruction processes, $\mathcal{N}_p + \mathcal{N}_{pm}$ must be even. Since a bipartite lattice has only loops of even length, the number of occurrences of processes that hop the m quantum number must be even, implying that $\mathcal{N}_m + \mathcal{N}_{pm} + \mathcal{N}_{lm}$ is even. By the same argument applied to the l quantum number, $\mathcal{N}_l + \mathcal{N}_{lm}$ must also be even.

Since $H_{2(r_a r_b)}^\pm$ gives rise only to m -type processes, while $H_{3(r_a r_b)}$ gives rise to l -type and p -type processes and $H_{4(r_a r_b)}^\pm$ gives rise to lm -type and pm -type processes, $K_\perp = 0$ ($K_z = 0$) implies $\mathcal{N}_{lm} = \mathcal{N}_{pm} = 0$ ($\mathcal{N}_l = \mathcal{N}_p = 0$). The periodicity constraints then imply that all nonzero $\mathcal{N}_i$ are even in both these cases. As a result, in both these cases, each $\mathcal{S}_n$ has positive weight in this QMC scheme *regardless of the sign of all nonzero couplings*. On the other hand, if $J_\perp = 0$, i.e., $\mathcal{N}_m = 0$, only pm -type processes can create or destroy pairs of $m = \pm 1$ states on neighboring sites, thus ensuring that $\mathcal{N}_{pm}$ is even. Along with the other periodicity constraints, this again implies that all nonzero $\mathcal{N}_i$ are even, yielding a sign-problem-free method whenever $J_\perp = 0$, *independent of the sign of other couplings*.

Thus, the Monte Carlo weight is positive for frustrated bilayer magnets with Hamiltonian H_{bilayer} when at least one of the following conditions is satisfied: (i) $K_z = \mathcal{J}_z$; (ii) $K_\perp = \mathcal{J}_\perp$; (iii) $K_\perp = -\mathcal{J}_\perp$. When (i) and (ii) are both satisfied, one obtains fully frustrated bilayer systems [39–44] with infinitely many conservation laws, which can have either SU(2) or U(1) symmetry. Additionally, (iii) also contains other examples of SU(2) symmetric frustrated magnets (e.g., for $\mathcal{J}_z = \mathcal{J}_\perp = 0$, $K_z = K_\perp > 0$, and $\Delta_z = \Delta_\perp < 0$) and no extra conservation laws. A similar argument establishes the absence of a sign problem for H_{mixed} [56].

Alternately, this positive-weight property can be made explicit by switching from the canonical cluster basis to a rotated basis obtained by attaching phase factors $e^{i\theta_{|lm\rangle} + i\eta_{|lm\rangle}}$ to the states $|l, m\rangle$. Here, the $\theta_{|lm\rangle}$ are r independent, while the $\eta_{|l,m\rangle}$ are 0 on the B sublattice and constant on the A sublattice. These phases are chosen in each of these three generic cases to ensure that every nonzero contribution to the weight in the rotated basis is explicitly positive. For instance, when $K_\perp = 0$, we set $\eta_{|t_\pm\rangle} = \theta_{|l,m\rangle} = 0$ (for all l, m) while independently choosing $\eta_{|s\rangle}$ to be 0 or π and $\eta_{|t_0\rangle}$ to be 0 or π depending on the signs of K_z and $J_\perp$. The other sign-free cases can be handled with slightly different choices for these phase factors [56]. The positive-weight property of H_{mixed} can also be made explicit in the same way [56].

Implementation.—A key advantage of this QMC approach is that the usual SSE framework [3] remains valid with no change in the construction of the diagonal update and one new feature in the construction of directed loop updates: *Three* different kinds of directed loop updates [57,58] are now possible, involving changes to the m quantum number, the l quantum number, or both. Additionally, to improve statistics, one can use parallel

FIG. 2. Temperature (T) dependence of the susceptibility and specific heat of H_{bilayer} with $\mathcal{D}_z = \mathcal{D}_\perp = 1$, $\mathcal{J}_\perp = K_\perp = 1$, $\mathcal{J}_z = 1 + K_z$, and $K_z = 1 - K_\perp$. Symbols display data for a sample with $L = 64$ unit cells, plotted for a variety of values of K_z . The inset shows the perfect agreement between QMC data (symbols) and exact diagonalization results (lines) for $L = 6$.

tempering [59] as well as an additional local update, which identifies worldlines that are only touched by diagonal vertices, and changes their state using Metropolis-type acceptance probabilities.

Benchmarks.—Our implementation, which incorporates all these updates, has been successfully benchmarked against exact diagonalization in spatial dimension $d = 1$ for all the sign-free cases, including the two special cases with infinitely many conserved quantities [56]. In Fig. 2 (inset), we illustrate this for a representative example, focusing on the susceptibility $\chi = \beta \langle (S^z)^2 \rangle / N$ and specific heat per spin $C_v = \frac{1}{N} (d\langle H \rangle / dT)$ for H_{bilayer} in $d = 1$, with $K_\perp = 0$ ($N = 2L$ is the number of spins 1/2 in a ladder of length L). Data in the main panel illustrate the power of the method, which allows us to access the thermodynamics of this frustrated ladder for fairly large L up to low T and for a range of values of K_z .

Numerical results in $d = 2$.—We now consider H_{bilayer} on a square Bravais lattice in the presence of an additional *magnetic field*, which our method can handle without a sign problem: $H = H_{\text{bilayer}} - h \sum_r (S_{Ir}^z + S_{IIr}^z)$ (the magnetic field modifies only weights of the diagonal vertices). The physics of the fully frustrated special case ($K_z = K_\perp = 0$) in a certain field range was argued [60] to be dominated at low temperatures by sublattice-ordered configurations in which two-spin clusters on one spontaneously chosen sublattice are in the triplet state $|t_{+1}\rangle$, while two-spin clusters on the other sublattice are in a singlet state $|s\rangle$, allowing a low-temperature mapping to the ordered phase of hard squares on the square lattice. This predicts that the system undergoes a temperature-driven phase transition in the 2d Ising universality class to a high-temperature phase in which sublattice symmetry is restored [60,61].

Nevertheless, this construction of negative-weight configurations *relies on the existence of loops* in the underlying bipartite Bravais lattice and leaves open the possibility that this sign problem could be controlled in 1d systems with open boundaries. In summary, our work has led to a solution of the sign problem for a large and interesting class of frustrated quantum magnets. Given the ubiquity of the sign problem in computational physics, we hope that the strategy outlined in this work can be adapted to other systems.

We thank S. Wessel for generously sharing with us some details of the recent parallel work Ref. [55] and for stimulating discussions regarding the sign problem in the general case. We gratefully acknowledge J. Richter for suggesting the study of the finite-temperature Ising transition in the square bilayer system in a field. One of us (K. D.) also gratefully acknowledges the hospitality of ECT* (Trento), ICTP (Trieste), and ICTS-TIFR (Bengaluru) during the writing of this Letter. This work was supported by the Indo-French Centre for the Promotion of Advanced Research (IFCPAR/CEFIPRA Project 4504-1) and performed using numerical resources from GENCI (Grants No. 2014-x2014050225 and No. 2015-x2015050225) and CALMIP. Our QMC code is based on the SSE code of the ALPS libraries [58,68].

-
- [1] *Strongly Correlated Systems Numerical Methods*, edited by A. Avella and F. Mancini, Springer Series in Solid State Sciences Vol. 176 (Springer, Berlin, 2013).
 - [2] W. Krauth, *Algorithms and Computations*, Oxford Master Series in Statistical Computational and Theoretical Physics (Oxford University, New York, 2006).
 - [3] A. W. Sandvik, Computational studies of quantum spin systems, *AIP Conf. Proc.* **1297**, 135 (2010).
 - [4] D. C. Johnston, R. K. Kremer, M. Troyer, X. Wang, A. Klümper, S. L. Bud'ko, A. F. Panchula, and P. C. Canfield, Thermodynamics of spin $S = 1/2$ antiferromagnetic uniform and alternating-exchange Heisenberg chains, *Phys. Rev. B* **61**, 9558 (2000).
 - [5] *Introduction to Frustrated Magnetism Materials, Experiments, Theory*, edited by C. Lacroix, P. Mendels, and F. Mila, Springer Series in Solid State Sciences Vol. 176 (Springer, Berlin, 2011).
 - [6] S. Wessel, in *Lecture Notes of the Autumn School Correlated Electrons 2013*, edited by E. Pavarini, E. Koch, and U. Schollwck (Forschungszentrum, Jülich, 2013).
 - [7] R. K. Kaul, R. G. Melko, and A. W. Sandvik, Bridging lattice-scale physics and continuum field theory with quantum Monte Carlo simulations, *Annu. Rev. Condens. Matter Phys.* **4**, 179 (2013).
 - [8] M. Troyer and U.-J. Wiese, Computational Complexity and Fundamental Limitations to Fermionic Quantum Monte Carlo Simulations, *Phys. Rev. Lett.* **94**, 170201 (2005).
 - [9] R. Moessner and S. L. Sondhi, Ising models of quantum frustration, *Phys. Rev. B* **63**, 224401 (2001).
 - [10] S. V. Isakov and R. Moessner, Interplay of quantum and thermal fluctuations in a frustrated magnet, *Phys. Rev. B* **68**, 104409 (2003).
 - [11] S. Wessel and M. Troyer, Supersolid Hard-Core Bosons on the Triangular Lattice, *Phys. Rev. Lett.* **95**, 127205 (2005).
 - [12] D. Heidarian and K. Damle, Persistent Supersolid Phase of Hard-Core Bosons on the Triangular Lattice, *Phys. Rev. Lett.* **95**, 127206 (2005).
 - [13] R. G. Melko, A. Paramekanti, A. A. Burkov, A. Vishwanath, D. N. Sheng, and L. Balents, Supersolid Order from Disorder: Hard-Core Bosons on the Triangular Lattice, *Phys. Rev. Lett.* **95**, 127207 (2005).
 - [14] S. V. Isakov, S. Wessel, R. G. Melko, K. Sengupta, and Y. B. Kim, Hard-Core Bosons on the Kagome Lattice: Valence-Bond Solids and Their Quantum Melting, *Phys. Rev. Lett.* **97**, 147202 (2006).
 - [15] S. V. Isakov, Y. B. Kim, and A. Paramekanti, Spin-Liquid Phase in a Spin-1/2 Quantum Magnet on the Kagome Lattice, *Phys. Rev. Lett.* **97**, 207204 (2006).
 - [16] A. Banerjee, S. V. Isakov, K. Damle, and Y. B. Kim, Unusual Liquid State of Hard-Core Bosons on the Pyrochlore Lattice, *Phys. Rev. Lett.* **100**, 047208 (2008).
 - [17] A. Sen, K. Damle, and T. Senthil, Superfluid insulator transitions of hard-core bosons on the checkerboard lattice, *Phys. Rev. B* **76**, 235107 (2007).
 - [18] K. Damle and T. Senthil, Spin Nematics and Magnetization Plateau Transition in Anisotropic Kagome Magnets, *Phys. Rev. Lett.* **97**, 067202 (2006).
 - [19] A. Sen, F. Wang, K. Damle, and R. Moessner, Triangular and Kagome Antiferromagnets with a Strong Easy-Axis Anisotropy, *Phys. Rev. Lett.* **102**, 227001 (2009).
 - [20] A. Sen, K. Damle, and A. Vishwanath, Magnetization Plateaus and Sublattice Ordering in Easy-Axis Kagome Lattice Antiferromagnets, *Phys. Rev. Lett.* **100**, 097202 (2008).
 - [21] E. Berg, M. A. Metlitski, and S. Sachdev, Sign-problem-free quantum Monte Carlo of the onset of antiferromagnetism in metals, *Science* **338**, 1606 (2012).
 - [22] Y.-P. Huang, G. Chen, and M. Hermele, Quantum Spin Ices and Topological Phases from Dipolar-Octupolar Doublets on the Pyrochlore Lattice, *Phys. Rev. Lett.* **112**, 167203 (2014).
 - [23] F. F. Assaad and H. G. Evertz, in *Proceedings of the Heraeus Summer School, Greifswald*, Springer Lecture Notes in Physics 739, edited by H. Fehske *et al.* (Springer, New York, 2008).
 - [24] C. Wu and S.-C. Zhang, Sufficient condition for absence of the sign problem in the fermionic quantum Monte Carlo algorithm, *Phys. Rev. B* **71**, 155115 (2005).
 - [25] L. Messio and F. Mila, Entropy Dependence of Correlations in One-Dimensional SU(N) Antiferromagnets, *Phys. Rev. Lett.* **109**, 205306 (2012).
 - [26] E. F. Huffman and S. Chandrasekharan, Solution to sign problems in half-filled spin-polarized electronic systems, *Phys. Rev. B* **89**, 111101 (2014).
 - [27] L. Wang, Y.-H. Liu, M. Iazzi, M. Troyer, and G. Harcos, Split Orthogonal Group: A Guiding Principle for Sign-Problem-Free Fermionic Simulations, *Phys. Rev. Lett.* **115**, 250601 (2015).

- [28] Z.-X. Li, Y.-F. Jiang, and H. Yao, Solving the fermion sign problem in quantum Monte Carlo simulations by Majorana representation, *Phys. Rev. B* **91**, 241117(R) (2015).
- [29] Z. C. Wei, C. Wu, Y. Li, S. Zhang, and T. Xiang, Majorana Positivity and the Fermion Sign Problem of Quantum Monte Carlo Simulations, *Phys. Rev. Lett.* **116**, 250601 (2016).
- [30] P. Henelius and A. W. Sandvik, Sign problem in Monte Carlo simulations of frustrated quantum spin systems, *Phys. Rev. B* **62**, 1102 (2000).
- [31] W. Bietenholz, A. Pochinsky, and U.-J. Wiese, Meron Cluster Simulation of the Theta Vacuum in the 2-d O(3) Model, *Phys. Rev. Lett.* **75**, 4524 (1995).
- [32] S. Chandrasekharan and U.-J. Wiese, Meron-Cluster Solution of Fermion Sign Problems, *Phys. Rev. Lett.* **83**, 3116 (1999).
- [33] M. Nyfeler, F.-J. Jiang, F. Kämpfer, and U.-J. Wiese, Nested Cluster Algorithm for Frustrated Quantum Antiferromagnets, *Phys. Rev. Lett.* **100**, 247206 (2008).
- [34] T. Nakamura, Vanishing of the negative-sign problem of quantum Monte Carlo simulations in one-dimensional frustrated spin systems, *Phys. Rev. B* **57**, R3197 (1998).
- [35] K. Okunishi and K. Harada, Symmetry-protected topological order and negative-sign problem for SO(N) bilinear-biquadratic chains, *Phys. Rev. B* **89**, 134422 (2014).
- [36] S. D. Geraedts and O. I. Motrunich, Model of fractionalization of Faraday lines in compact electrodynamics, *Phys. Rev. B* **90**, 214505 (2014).
- [37] S. D. Geraedts and O. I. Motrunich, Model Realization and Numerical Studies of a Three-Dimensional Bosonic Topological Insulator and Symmetry-Enriched Topological Phases, *Phys. Rev. X* **4**, 041049 (2014).
- [38] S. D. Geraedts and O. I. Motrunich, Exact realization of integer and fractional quantum Hall phases in $U(1) \times U(1)$ models in $(2+1)d$, *Ann. Phys. (Amsterdam)* **334**, 288 (2013).
- [39] M. P. Gelfand, Linked-tetrahedra spin chain: Exact ground state and excitations, *Phys. Rev. B* **43**, 8644 (1991).
- [40] A. K. Kolezhuk and H.-J. Mikeska, Finitely correlated generalized spin ladders, *Int. J. Mod. Phys. B* **12**, 2325 (1998).
- [41] A. Honecker, F. Mila, and M. Troyer, Magnetization plateaus and jumps in a class of frustrated ladders: A simple route to a complex behaviour, *Eur. Phys. J. B* **15**, 227 (2000).
- [42] W. Brenig and K. W. Becker, Magnetism of a tetrahedral cluster spin chain, *Phys. Rev. B* **64**, 214413 (2001).
- [43] E. Chattopadhyay and I. Bose, Magnetization of coupled spin clusters in ladder geometry, *Phys. Rev. B* **65**, 134425 (2002).
- [44] K. Totsuka and H.-J. Mieska, Low-lying excitations and magnetization process of coupled tetrahedral systems, *Phys. Rev. B* **66**, 054435 (2002).
- [45] K. Takano, K. Kubo, and H. Sakamoto, Ground states with cluster structures in a frustrated Heisenberg chain, *J. Phys. Condens. Matter* **8**, 6405 (1996).
- [46] H. Niggemann, G. Uimin, and J. Zittartz, Mixed Heisenberg chains: I. The ground-state problem, *J. Phys. Condens. Matter* **9**, 9031 (1997).
- [47] A. Honecker, S. Hu, R. Peters, and J. Richter, Dynamic and thermodynamic properties of the generalised diamond chain model for azurite, *J. Phys. Condens. Matter* **23**, 164211 (2011).
- [48] N. B. Ivanov and J. Richter, Competition between plaquette and dimer phases in Heisenberg chains, *Phys. Lett. A* **232**, 308 (1997).
- [49] J. Richter, N. B. Ivanov, and J. Schulenburg, The antiferromagnetic spin-1/2 chain with competing dimers and plaquettes: Numerical versus exact results, *J. Phys. Condens. Matter* **10**, 3635 (1998).
- [50] A. Koga, K. Okunishi, and N. Kawakami, First-order quantum phase transition in the orthogonal-dimer spin chain, *Phys. Rev. B* **62**, 5558 (2000).
- [51] C. Gros, R. Valentí, J. V. Alvarez, K. Hamacher, and W. Wenzel, Test of the frustrated spin-cluster model to describe the low-temperature physics of NaV_2O_5 , *Phys. Rev. B* **62**, R14617 (2000).
- [52] J. Schulenburg and J. Richter, Infinite series of magnetization plateaus in the frustrated dimer-plaquette chain, *Phys. Rev. B* **65**, 054420 (2002).
- [53] A. Honecker and W. Brenig, Magnetic properties of a spin-1/2 quadrumer chain, *Phys. Rev. B* **63**, 144416 (2001).
- [54] S. R. Manmana, J.-D. Picon, K. P. Schmidt, and F. Mila, Unconventional magnetization plateaus in a Shastry-Sutherland spin tube, *Europhys. Lett.* **94**, 67004 (2011).
- [55] A. Honecker, S. Wessel, R. Kerkdyk, T. Pruschke, F. Mila, and B. Normand, Thermodynamic properties of highly frustrated quantum spin ladders: Influence of many-particle bound states, *Phys. Rev. B* **93**, 054408 (2016).
- [56] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.117.197203> for details of the QMC simulation of H_{mixed} (including a proof of positive-weight), and results on one-dimensional systems which have been used to benchmark the method.
- [57] O. F. Syljuåsen and A. W. Sandvik, Quantum Monte Carlo with directed loops, *Phys. Rev. E* **66**, 046701 (2002).
- [58] F. Alet, S. Wessel, and M. Troyer, Generalized directed loop method for quantum Monte Carlo simulations, *Phys. Rev. E* **71**, 036706 (2005).
- [59] K. Hukushima and K. Nemoto, Exchange Monte Carlo method and application to spin glass simulations, *J. Phys. Soc. Jpn.* **65**, 1604 (1996).
- [60] J. Richter, O. Derzhko, and T. Krokhamalskii, Finite-temperature order-disorder phase transition in a frustrated bilayer quantum Heisenberg antiferromagnet in strong magnetic fields, *Phys. Rev. B* **74**, 144430 (2006).
- [61] O. Derzhko, T. Krokhamalskii, and J. Richter, Emergent Ising degrees of freedom in frustrated two-leg ladder and bilayer $s = 1/2$ Heisenberg antiferromagnets, *Phys. Rev. B* **82**, 214412 (2010).
- [62] G. Kamieniarz and H. W. J. Blöte, Universal ratio of magnetization moments in two-dimensional Ising models, *J. Phys. A* **26**, 201 (1993).
- [63] E. Müller-Hartmann, R. R. P. Singh, C. Knetter, and G. S. Uhrig, Exact Demonstration of Magnetization Plateaus and First Order Dimer-Néel Phase Transitions in a Modified

- Shastry-Sutherland Model for $\text{SrCu}_2(\text{BO}_3)_2$, *Phys. Rev. Lett.* **84**, 1808 (2000).
- [64] H. Q. Lin, J. L. Shen, and H. Y. Shik, Exactly soluble quantum spin models on a double layer: The net spin model, *Phys. Rev. B* **66**, 184402 (2002).
- [65] J. Rössler and D. Gottlieb, Analysis of a family of Heisenberg systems with simple eigenfunctions for the ground state and low lying excitations, *Phys. Rev. B* **72**, 024443 (2005).
- [66] M. Mambrini, J. Trébosch, and F. Mila, Residual entropy and spin gap in a one-dimensional frustrated antiferromagnet, *Phys. Rev. B* **59**, 13806 (1999).
- [67] O. Rojas and F. C. Alcaraz, Phase diagram of a coupled tetrahedral Heisenberg model, *Phys. Rev. B* **67**, 174401 (2003).
- [68] A. F. Albuquerque *et al.*, The ALPS project release 1.3: Open source software for strongly correlated systems, *J. Magn. Magn. Mater.* **310**, 1187 (2007).