

Influence of grassland management on carbon allocation in a semiarid temperate steppe

Liu N, Zhang Y Jun, Kan H Ming and Lu W Jie

Institute of Grassland Science, College of Animal Science and Technology, China Agricultural University, No.2, Yuan Mingyuan Xilu RD, Beijing 100193, People's Republic of China

Contact email: liunan_2293@163.com

Keywords: Carbon allocation, grazing, hay-making, fencing.

Introduction

Grazing lands in North China are often excessively grazed and widely degraded, while hay-making lands appear to be in relatively good condition due to grazing exclusion, but they are facing a continuous loss of nutrients in the harvested biomass. In semiarid grasslands, plant productivity and community composition are significantly altered by grazing and haying. Grazing mostly leads to negative effects on aboveground productivity, however root biomass seems to increase with moderate grazing (Gao *et al.* 2009; Derner *et al.* 2006), although responses can vary. Aboveground biomass removal can increase C3 grass dominance and productivity (Hofer and Bragg 1981). Grazing exclusion is a valuable mechanism of sequestering soil C (He 2008). However, grazing can change C allocation patterns and affect the amount of C entering the soil. Here we examine the potential effects of common management practices (exclusion with fencing, grazing and hay-making) on semiarid grasslands above- and below-ground C pools. The primary objective of this study was to evaluate the potential of grazing exclusion and annual last-summer haying in previous grazing lands on the storage of C in semiarid grasslands of northern China.

Methods

This study was conducted in a semiarid temperate steppe in Duolun County, Inner Mongolia, China. Long-term mean annual precipitation and temperature are approximately 383 mm and 2.1°C. The dominant plant species were perennial plants, including *Stipa krylovii*, *Artimesia frigida*, and *Potentilla acaulis*, *Cleistogenes squarrosa*.

The fencing lands were fenced from 2003, and completely excluded livestock grazing. The hay-making lands were cut at the end of the growing season every year from 2003 to the time of the study. The grazing lands were moderately grazed by cattle. We selected five 50 m transects at each management area. Five random sampling quadrats (1 × 1 m) and five root and soil cores were collected manually to 40 cm with a root sampler (80 mm diameter) and a soil sampler (50 mm diameter) respectively at 10 m intervals along each 50 m sampling transect in 2011. The plant biomass and C storage were measured. Data analyses were done with SPSS version

16.0 (SPSS Inc., Chicago, IL, USA).

Results and discussion

Hay-making lands had significantly greater aboveground biomass, especially increased *Stipa krylovii* biomass ($P < 0.05$) in comparison to the other management practices. This result was consistent with the finding of Hofer and Bragg (1981). Seven years of exclusion resulted in significantly increased litter biomass and standing dead biomass (Table 1). Similar to the biomass results, C pool in aboveground biomass and root was significantly higher in fenced and hay-making lands relative to grazing lands. However, the amount of C stored in soil decreased significantly with the introduction of fencing (Table 2). The grasslands subjected to fencing were mature in terms of aboveground productivity, including vegetation, litter, standing dead and roots. Total carbon storage in biomass and soil after 8 years of grazing exclusion decreased significantly mainly due to a loss of SOC. The relatively lower C storage in fenced lands resulted in an increase in aboveground biomass relative to grazing lands that likely caused greater competition for resources and greater consumption of SOM (He 2008). Animal traffic and machinery enhances physical breakdown and incorporation of litter into the soil, which can increase

Table 1. Dominant species biomass, aboveground total biomass, ground litter biomass, standing dead biomass and roots biomass in fenced, grazing and hay-making lands in this study.

Biomass (g/m ²)	Management types		
	Fenced lands	Grazing lands	Hay-making lands
<i>Stipa krylovii</i>	13.58±2.57a	7.86±1.73a	40.28±8.58b
<i>Artimesia frigida</i>	53.01±9.47a	62.18±10.61a	57.35±13.49a
<i>Potentilla acaulis</i>	12.17±3.93a	13.66±3.68a	12.91±6.22a
<i>Cleistogenes squarrosa</i>	5.21±1.93a	6.29±0.95a	17.63±4.18b
Aboveground	118.96±3.03b	103.79±6.84a	139.00±4.89c
Ground litter	48.97±10.08b	31.19±3.92ab	14.77±0.87a
Standing dead	20.46±5.05b	4.43±0.77a	11.82±5.37ab
Root (0-40 cm)	2039.60 ± 137.76b	1511.52 ± 125.93a	1961.11 ± 221.26ab

Table 2. Carbon distribution among different pools in fenced, grazing and hay-making lands

C pool	Management types		
	Fenced	Grazing	Hay-making
Plant (g/m ²)	52.07±2.01b	40.23±2.98a	57.44±5.28b
Litter (g/m ²)	19.37±1.13c	11.38±3.32b	2.92±0.14a
Standing dead (g/m ²)	8.06±0.36c	1.70±0.15b	2.54±0.09a
Root (kg/m ²)	0.62±0.07b	0.44±0.04a	0.60±0.05b
Soil microbial biomass (mg/kg)	544.80±125.69a	563.71±120.85a	365.00±59.91a
SOC (g/kg 0-40 cm)	13.32±1.24a	16.58±1.22ab	17.55±1.69b
SOC (kg/m ² 0-40 cm)	6.49±0.28a	7.95±0.31b	7.27±0.25ab

the rate of decomposition of litter and transfer of C and nutrients into the soil (Naeth 1991). In grazing lands, there is a higher carbon input from roots, litter and standing dead, while the fencing treatment strongly decreased this input and promoted aboveground allocation (Liu, 2012). In Addition, higher microbial biomass carbon (Table 2) further indicated higher organic matter inputs from plant litter and root exudates in the grazing soil (Bird 2002).

Conclusion

The results indicated that long-term grazing lands that are fenced to exclude grazing could be construed as proper management for increasing vegetation productivity, but not for soil C sequestration. However, grazing increased the amount of C entering the soil, and changed C allocation patterns. Annual last-summer

haymaking has potential for grassland restoration and sustaining C sequestration in semiarid grassland.

Acknowledgement

Research was supported by the earmarked fund for Modern Agro-industry Technology Research System. (CARS-35) and Beijing Key Laboratory of Grassland Science. We are grateful for the support and technical assistance from friends and colleagues at China Agriculture University.

References

- Gao YH, Schumann M, Chen H, Wu N, Luo P (2009) Impacts of grazing intensity on soil carbon and nitrogen in an alpine meadow on the eastern Tibetan Plateau. *Journal of Food and Agricultural Environment* **7**, 749-754.
- Derner J, Boutton T, Briske D (2006) Grazing and ecosystem carbon storage in the North American Great Plains. *Plant and Soil* **280**, 77-90
- Hofer EL, Bragg TB (1981) Effect of season of burning and mowing on an eastern Nebraska *Stipa-Andropogon* prairie. *American Midland Naturalist*, **150**, 13-18.
- He NP, Yu Q, Wu L, Wang YS, Han XG (2008) Carbon and nitrogen store and storage potential as affected by land-use in a *Leymus chinensis* grassland of northern China. *Soil Biology and Biochemistry* **40**, 2952-2959
- Naeth MA, Bailey AW, Pluth DJ, Chanasyk DS, Hardin RT (1991) Grazing impacts on litter and soil organic matter in mixed prairie and fescue grassland ecosystems of Alberta. *Journal of Arid Environments* **44**, 7-12.
- Liu N, Zhang YJ, Chang SJ, Kan HM, Lin LJ (2012) Impact of Grazing on Soil Carbon and Microbial Biomass in Typical Steppe and Desert Steppe of Inner Mongolia. *PLoS ONE* **7**, e36434. doi:10.1371/journal.pone.0036434
- Bird SB, Herrick JE, Wander MM, Wright SF (2002) Spatial heterogeneity of aggregate stability and soil carbon in semi-arid rangeland. *Environment Pollution* **116**, 445-455.