

# Soil organic carbon stocks in a Brazilian oxisol under different pasture systems

José C Pinto, Róberson M Pimentel, Yuri L Zinn and Fernanda HM Chizzotti

Universidade Federal de Lavras (UFLA), Lavras-MG, Brazil, 37200-000.

Contact email: [josecard@dzo.ufla.br](mailto:josecard@dzo.ufla.br)

**Keywords:** Soil organic carbon stocks, pasture systems.

## Introduction

Pastures are the main land use systems in the world and in Brazil they occupy 115 M ha. A major part of Brazil's greenhouse gas emissions are due to land use change and agriculture. Livestock production comprises >90% of methane and about 55% of CO<sub>2</sub>-equivalent emissions due to agriculture (Cerri *et al.* 2009). However, productive, well-managed pastures can improve degraded soils and increase soil organic carbon (SOC) stocks through humification of grass and root residues. In order to enhance pasture yields and SOC sequestration, nutrient availability in soils must also be improved, especially for N and P. This work aimed to assess how SOC stocks were affected by the combination of grasses and legumes, in comparison to only-grass pastures and other land uses in a clay Oxisol in southeast Brazil.

## Methods


This research was conducted in the campus of the Universidade Federal de Lavras (21°13'42" S; 44°58'13" W, 925 m.a.s.l.). Five neighboring areas were selected for sampling, comprising: (1) a native, semi-deciduous forest; (2) pure *Brachiaria decumbens* pasture (planted in 1990, under continuous grazing by bovines); (3) mixed *B. brizantha* and *Arachis pintoii* pasture; (4) mixed *B. brizantha* and *Stylosanthes guianensis*; and (5) corn under annual tillage for 25 yrs. Both pasture+legume plots were planted in 2007 after 15 yrs under conventional tillage corn, and kept under cyclic bovine grazing. The local soil is a Humic Rhodic Acrudox with clayey (>60% clay) texture and granular structure. Soil sampling for SOC and bulk density (core method) was done in March 2010 using 3 pits per treatment, at the depths of 0-5, 5-10, 10-20, 20-40, 40-60 and 60-100 cm. Soil samples were air-dried, sieved <2mm and ground to pass a 100 mesh (0.150 mm) sieve for SOC determination by dry combustion in a Vario Micro Cube (Hanau, Germany) apparatus. SOC stocks were calculated using an equivalent mass approach due to soil compaction. The experimental design was completely randomized in triplicate. Treatment means were compared by the Tukey test.

## Results and discussion

Bulk density increased significantly in pastures and corn plots in comparison to the native forest (Fig. 1a), the

highest levels were found in the areas currently or formerly cultivated with corn. Remarkably, such compaction occurred to a depth of 1 m, due to the very low densities and macroporosities in the native forest and the intensity of plowing/harrowing. The opposite occurred for SOC concentration, since the highest values were as expected found in the native forest, especially in the top 20 cm (Fig. 1b). At the 0-5 cm depth, the lowest values were found for the *Brachiaria*+*Stylosanthes* pasture, which can be ascribed to intense organic matter decomposition due to a very low C:N ratio of 7.0 (Pimentel 2012), and in the corn site.

The effect of land use change on SOC stocks occurred for all depths, although with some differences on specific comparisons (Table 1). Pure *Brachiaria*



**Figure 1.** Means of: a) bulk density ( $\text{g/cm}^3$ ) and b) soil organic carbon concentrations ( $\text{g/kg}$ ) to a depth of 100 cm. Bars show standard error of the mean. Land use systems are native forest (MN), corn (MPC), *Brachiaria*+*Arachis* pasture (PA), *Brachiaria*+*Stylosanthes* pasture (PE), and pure grass pasture (PP)

**Table 1. Soil organic carbon stocks for different standardized depths. Means followed by the same letter in the column do not differ by the Tukey test at  $P < 0.05$ .**

Land use	Soil organic C stocks (t/ha)			
	0-20 cm	0-40 cm	0-60 cm	0-100 cm
Native forest	71.8 A	128.5 A	171.8 A	239.9 A
<i>Brachiaria</i>	50.7 B	100.2 B	142.2 A	207.8 A
<i>Brachiaria</i> + <i>Arachis</i>	39.4 BC	71.3 C	100.3 B	155.9 B
<i>Brachiaria</i> + <i>Stylosanthes</i>	33.2 CD	71.8 C	102.2 B	145.9 BC
Corn	24.1 D	49.5 C	74.0 B	113.7 C

pastures significantly lost SOC when compared to the native forest for the 0-20 cm (most intensively cultivated layer) and 0-40 cm (maximal arable layer) depths, but not for the 0-60 and 0-100 depths. Since most SOC changes typically occur in the top 20 cm, such trend is primarily due to the decreasing rate of SOC change:total SOC as increasingly deeper layers are considered for the *Brachiaria* pasture. However, SOC losses for the grass-legume pastures increased from 30 t C/ha for the 0-20 cm depth to 90 t C/ha for the 0-100 cm layer in comparison to the forest, and even more for the corn. Therefore, the data showed that intense SOC losses can occur with annual cultivation even to very deep layers, contrasting with the typically superficial effect noted for Brazil as a whole (Zinn *et al.* 2005). It is possible that such heavy losses are also due to originally very high SOC stocks of

>200 t/ha for this area, which are not common in Brazilian savannic and Amazonian soils. However, mixed pastures of *Brachiaria*+*Arachis* seem to increase SOC levels in comparison to corn, which did not happen in the case of *Brachiaria*+*Stylosanthes* pastures since the organic matter generated is too labile due to high N contents.

## Conclusion

Replacing native forests with pastures where SOC stocks are originally high apparently leads to heavy SOC losses even when pastures are well managed and productive. However, pastures are more SOC-conserving than annual crops with conventional tillage, and can sequester SOC after croplands are turned into grazing lands. Nevertheless, this study also points that excessive N inputs through biological fixation in legume-grass consortia can unexpectedly accelerate organic matter decomposition and delay or preclude SOC sequestration in degraded or overexploited lands.

## References

- Cerri CC, *et al.* (2009) Brazilian greenhouse gas emissions: the importance of agriculture and livestock. *Scientia Agricola* **66**, 831-843.
- Pimentel RM (2012) Propriedades físicas, carbono e nitrogênio do solo em sistemas agropecuários. M.Sc. Dissertation, Universidade Federal de Lavras.
- Zinn YL, Lal R, Resck DWS (2005) Changes in soil organic carbon stocks under agriculture in Brazil. *Soil and Tillage Research* **84**, 28-40.