



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

22nd International Grassland Congress

Assessing Resilience of Pasture Production to Climatic Changes

Brendan R. Cullen

University of Melbourne, Australia

Richard P. Rawnsley

University of Tasmania, Australia

Richard J. Eckard

University of Melbourne, Australia

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/22/2-7/9>

The 22nd International Grassland Congress (Revitalising Grasslands to Sustain Our Communities) took place in Sydney, Australia from September 15 through September 19, 2013.

Proceedings Editors: David L. Michalk, Geoffrey D. Millar, Warwick B. Badgery, and Kim M.

Broadfoot

Publisher: New South Wales Department of Primary Industry, Kite St., Orange New South Wales, Australia

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Assessing resilience of pasture production to climatic changes

Brendan R Cullen^A, Richard P Rawnsley^B and Richard J Eckard^A

^A Melbourne School of Land and Environment, University of Melbourne, Victoria 3010 Australia.

^B Tasmanian Institute of Agriculture, University of Tasmania, PO Box 3532, Burnie, Tasmania, 7320 Australia.

Contact email: bcullen@unimelb.edu.au

Keywords: Climate change, perennial ryegrass, SGS Pasture model.

Introduction

Increasing temperatures and atmospheric carbon dioxide (CO₂) concentrations, together with changes to rainfall patterns, will influence seasonal pasture production; however climate change projections for south eastern Australia are uncertain (CSIRO and BoM 2007). Despite this, climate change impact assessments generally rely on specific climate projections, but in this study an alternative approach was developed to test the resilience of production to incremental changes in climate.

Methods

Sites and pasture systems modelled

Representative pasture types were simulated for Wagga Wagga (phalaris, subterranean clover, annual ryegrass and a native C₄ grass), Hamilton (perennial ryegrass and subterranean clover) and Elliott (perennial ryegrass and white clover) in south eastern Australia (Table 1). Annual pasture production was simulated as a cutting trial, with pasture mass cut to 1 t DM/ha on the last day of each month, using the Sustainable Grazing Systems pasture model (Johnson *et al.* 2003). In all simulations soil nutrients were non-limiting.

Climate scenarios

Twenty-five future climate scenarios were developed by scaling the historical climate (1971-2000) by increments of 0, 1, 2, 3 and 4°C (with corresponding changes to atmospheric CO₂ concentrations and relative humidity) and rainfall by +10, 0, -10, -20 and -30%, following the procedure of Cullen *et al.* (2012). These scenarios represent the range of climate change projections for southern Australia to 2070.

Data analysis

Surface charts showing annual average pasture production for the 25 climate scenarios were produced,

and on them the range of climatic changes for 2030, 2050 and 2070 (CSIRO and BoM 2007) were indicated.

Results and Discussion

With warming alone, annual production increased for the pasture sward at Wagga Wagga that consisted of a mixture of plant species with C₃ and C₄ photosynthetic pathways (Fig. 1). At the cool temperate site of Elliott warming of up to 3°C increased production, while warming up to 2°C increased production at Hamilton. At each site, production declined with lower rainfall. Generally annual pasture production was not lower than that simulated in the historical climate with the climate change projections for 2030, but it may be reduced under the 2050 and 2070 climate projections particularly if rainfall declines are >10%.

These results indicate that pasture production is more resilient to warming in cooler regions, and where there is a mix of C₃ and C₄ species. In cooler regions, warming overcomes low temperature limitations to growth, while it increases production of the C₄ species within mixed C₃/C₄ swards (Cullen *et al.* 2012). For C₃ pastures in temperate regions warming of more than 2°C will decrease annual production as increased growth rates in winter and early spring are counteracted by a shorter spring growing season.

Conclusion

Testing the sensitivity of production to incremental changes in climate can provide insights into the characteristics of regions and pasture systems that make them more or less resilient to climate change.

Acknowledgments

This research was funded by Australian Government Department of Agriculture, Fisheries and Forestry Climate Change Research Program, Dairy Australia and Meat and Livestock Australia.

Table 1 Location, soil types and average annual rainfall at Wagga Wagga, Hamilton and Elliott.

| Site | Lat./Long. | Soil type | Climatic zone | Rainfall (mm) |
|-------------|----------------|---------------------------------|----------------|---------------|
| Wagga Wagga | -35.10, 147.30 | Red chromosol/leptic tenosol | Mediterranean | 565 |
| Hamilton | -37.83, 142.06 | Brown chromosol | Temperate | 689 |
| Elliott | -41.08, 145.77 | Red mesotrophic haplic ferrosol | Cool temperate | 1220 |




Figure 1. Surfaces of average annual pasture harvested (t DM/ha) under increasing temperature (°C) and change in rainfall (%) scenarios at (a) Wagga Wagga, (b) Hamilton, and (c) Elliott. Contours show equal values across the response surfaces. Dashed ovals show the range of climate projections for 2030, 2050 and 2070

References

- Cullen BR, Eckard RJ, Rawnsley RP (2012) Resistance of pasture production to projected climate changes in south-eastern Australia. *Crop and Pasture Science* **63**, 77-86.
- CSIRO and BoM (2007) 'Climate change in Australia. Technical report 2007.' (Eds KB Pearce, PN Holper, M Hopkins, WJ Bouma, PH Whetton, KJ Hensessy, SB

Power) (CSIRO Marine and Atmospheric Research: Aspendale, Victoria) <http://www.climatechangeinaustralia.gov.au/>.

- Johnson IR, Lodge GM, White RE (2003) The Sustainable Grazing Systems Pasture Model: Description, philosophy and application to the SGS national experiment. *Australian Journal of Experimental Agriculture* **43**, 711-728.