

University of Kentucky

UKnowledge

Physics and Astronomy Faculty Publications

Physics and Astronomy

6-12-2017

Lifetime Measurements of Low-Spin Negative-Parity Levels in ^{160}Gd

S. R. Leshner

University of Wisconsin - La Crosse

C. Casarella

University of Notre Dame

L. M. Robledo

Universidad Autónoma de Madrid, Spain

Benjamin P. Crider

Michigan State University, bpscriders@gmail.com

R. Ikeyama

University of Wisconsin - La Crosse

See next page for additional authors https://uknowledge.uky.edu/physastron_facpub

 Part of the [Nuclear Commons](#)

[Right click to open a feedback form in a new tab to let us know how this document benefits you.](#)

Repository Citation

Leshner, S. R.; Casarella, C.; Robledo, L. M.; Crider, Benjamin P.; Ikeyama, R.; Marsh, I. R.; McEllistrem, Marcus T.; Peters, Erin E.; Prados-Estévez, Francisco M.; Smith, M. K.; Tully, Z. R.; Vanhoy, J. R.; and Yates, Steven W., "Lifetime Measurements of Low-Spin Negative-Parity Levels in ^{160}Gd " (2017). *Physics and Astronomy Faculty Publications*. 542.

https://uknowledge.uky.edu/physastron_facpub/542

This Article is brought to you for free and open access by the Physics and Astronomy at UKnowledge. It has been accepted for inclusion in Physics and Astronomy Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Lifetime Measurements of Low-Spin Negative-Parity Levels in ^{160}Gd

Digital Object Identifier (DOI)

<https://doi.org/10.1103/PhysRevC.95.064309>

Notes/Citation Information

Published in *Physical Review C*, v. 95, issue 6, 064309, p. 1-10.

©2017 American Physical Society

The copyright holder has granted permission for posting the article here.

Authors

S. R. Leshar, C. Casarella, L. M. Robledo, Benjamin P. Crider, R. Ikeyama, I. R. Marsh, Marcus T. McEllistrem, Erin E. Peters, Francisco M. Prados-Estévez, M. K. Smith, Z. R. Tully, J. R. Vanhoy, and Steven W. Yates

Lifetime measurements of low-spin negative-parity levels in ^{160}Gd

S. R. Leshner,^{1,2,*} C. Casarella,² A. Aprahamian,² L. M. Robledo,³ B. P. Crider,^{4,†} R. Ikeyama,¹ I. R. Marsh,¹ M. T. McEllistrem,⁴ E. E. Peters,⁵ F. M. Prados-Estévez,^{4,5} M. K. Smith,^{2,†} Z. R. Tully,¹ J. R. Vanhoy,⁶ and S. W. Yates^{4,5}

¹*Department of Physics, University of Wisconsin–La Crosse, La Crosse, Wisconsin 54601-3742, USA*

²*Department of Physics, University of Notre Dame, Notre Dame, Indiana 46556, USA*

³*Dep. Física Teórica (Módulo 15), Universidad Autónoma de Madrid, E-28049 Madrid, Spain*

⁴*Department of Physics and Astronomy, University of Kentucky, Lexington, Kentucky 40506-0055, USA*

⁵*Department of Chemistry, University of Kentucky, Lexington, Kentucky 40506-0055, USA*

⁶*Department of Physics, United States Naval Academy, Annapolis, Maryland 21402, USA*

(Received 1 February 2016; revised manuscript received 2 May 2017; published 12 June 2017)

$^{160}\text{Gd}(n,n'\gamma)$ experiments were performed with accelerator-produced monoenergetic neutrons. Excitation functions at neutron energies from 1.5 to 2.8 MeV aided in the placement of γ rays in the level scheme and angular distributions at three neutron energies resulted in the determination of 28 excited-level lifetimes or limits in ^{160}Gd , including the lifetimes of several negative-parity levels attributed to octupole vibrations.

DOI: [10.1103/PhysRevC.95.064309](https://doi.org/10.1103/PhysRevC.95.064309)

I. INTRODUCTION

The ^{160}Gd nucleus lies in the well-known rare-earth region of deformation and has an $R_{4/2}[\equiv E(4_{\text{gs}}^+)/E(2_{\text{gs}}^+)]$ ratio of 3.30, placing it close to the rotational limit. An open question in nuclear structure is the viability of vibrational excitations in deformed nuclei. The lowest-lying excited vibrational modes include quadrupole and octupole excitations. Single-phonon quadrupoles are the γ and β vibrations. γ vibrations seem to be well characterized as the first $K^\pi = 2^+$ bands and exhibit a systematic behavior across the region of deformed nuclei with typical $B(E2; 2_{\gamma}^+ \rightarrow 0_{\text{gs}}^+)$ values of a few Weisskopf units (W.u.). In contrast, the β vibrations do not exhibit these systematic characteristics because the $B(E2)$ values of transitions deexciting the first-excited $K^\pi = 0^+$ band vary greatly throughout the deformed region [1–3]. The question regarding the viability of the β vibrations in deformed nuclei remains open to debate [4–9].

Octupole excitations in spherical nuclei are identified by large $B(E3; 0_{\text{gs}}^+ \rightarrow 3^-)$ [10] values, and a compilation of $B(E3)$ values of the lowest 3^- states can be found in Ref. [11]. In deformed nuclei, the octupole mode splits into $K^\pi = 0^-, 1^-, 2^-,$ and 3^- bands. The increasing level density with excitation energy and the proximity of the pairing gap complicate the identification of vibrations in deformed nuclei. Two-phonon octupole excitations have been observed in spherical nuclei of the rare-earth region— ^{144}Sm [12], ^{146}Sm [13], ^{146}Gd [14], ^{147}Gd [15], and ^{148}Gd [16,17]—as has the combined two-phonon quadrupole-octupole vibrational mode ($2^+ \otimes 3^-$) in nearly spherical nuclei; e.g., ^{144}Sm [12], ^{146}Sm [18], and ^{148}Gd [13]. A few cases have been suggested as two-phonon double- γ vibrations ($K^\pi = 4^+$) in ^{164}Dy [19], ^{166}Er [20], ^{168}Er [21–23], and ($K^\pi = 0^+$) in ^{166}Er [24]. A

two-phonon double- β vibration has also been proposed in ^{178}Hf [25].

The low-lying structure of ^{160}Gd has been extensively studied; there are numerous positive- and negative-parity bands identified in this nucleus, including an excited $K^\pi = 2^+$ band, several excited $K^\pi = 0^+$ bands, and $K^\pi = 0^-, 1^-,$ and 2^- bands. There is, however, a paucity of level lifetimes and transition probabilities to clearly identify the collectivity or lack thereof for these excitations. Our recent paper reported on the characteristics of the $K^\pi = 0^+$ states [26] in this nucleus. This work reports mainly on the lifetime measurements of levels in low-lying $K^\pi = 0^-, 1^-,$ and 2^- bands. Using the Doppler-shift attenuation method following inelastic neutron scattering, we have determined the lifetimes of a large number of excited levels in ^{160}Gd .

II. EXPERIMENT

The low-lying states of ^{160}Gd were studied with the $(n,n'\gamma)$ reaction at the University of Kentucky Accelerator Laboratory (UKAL) with monoenergetic neutrons produced by the $^3\text{H}(p,n)^3\text{He}$ reaction. The scattering sample was 29.456 g of 98.12% enriched $^{160}\text{Gd}_2\text{O}_3$ contained in a thin-walled polyethylene cylinder 3.1 cm in height and 2.3 cm in diameter. The emitted γ rays were recorded with a HPGe detector with a relative efficiency of $\approx 50\%$. A bismuth germanate (BGO) active shield was used for Compton suppression. The HPGe detector was located at a distance of 119.3 cm from the scattering sample, and time-of-flight gating was used to suppress background radiation. Standard radioactive sources, ^{226}Ra and ^{152}Eu , were used for energy and efficiency calibration. In angular distribution measurements, a ^{60}Co source was placed near the detector as a continuous check on gain stability. At higher neutron energies, ^{24}Na was used as an additional in-beam source for accurate energy determination. Additional descriptions of the experimental setup and techniques are documented in Refs. [27–29].

The excitation function measurement provided yields of γ rays as a function of incident neutron energy (E_n), and spectra

*slesher@uwlax.edu

[†]Present address: National Superconducting Cyclotron Laboratory, Michigan State University, East Lansing, Michigan 48824, USA.




FIG. 1. Excitation functions for the 1493.40 and 1568.70 keV γ rays originating from the 1568.77 keV level. Both exhibit the same excitation-energy threshold and overall shape.

were measured for $E_n = 1.5$ to 2.8 MeV in 80 or 100 keV steps, with the detector at 90° with respect to the beam axis. Examples of the excitation functions obtained for γ rays from the same level are shown in Fig. 1. Excitation thresholds were used to place γ rays and establish levels. Angular-distribution spectra were recorded at neutron energies of 1.5, 2.0, and 2.8 MeV at ten angles over a range of 40° to 150° . To obtain the most accurate lifetimes, these neutron energies were chosen to minimize feeding to the levels of interest. The angular distributions of the γ -ray intensities, $W(\theta)$ were fit with a function of even-order Legendre polynomials,

$$W(\theta) = A_0[1 + a_2 P_2(\cos \theta_\gamma) + a_4 P_4(\cos \theta_\gamma)], \quad (1)$$

where the parameters a_2 and a_4 depend on the multiplicities and mixing amplitudes of the transitions. The experimental values of these parameters were then compared with statistical model calculations from the code CINDY [30] to determine multipole mixing ratios (δ) of the transitions. If the intensity of the γ ray was small or if the statistical uncertainty was large, the a_2 and a_4 values were compared with those in Govor *et al.* [31]. Provided the values matched within uncertainties, the values of Ref. [31] were used in relevant calculations and noted.

The angular-distribution measurements were also used to extract lifetimes of excited states shorter than about 1 ps by the Doppler-shift attenuation method (DSAM) [32]. The energies of the γ rays have an angular dependence described by

$$E_\gamma(\theta_\gamma) = E_{\gamma 0} \left[1 + \frac{v_{c.m.}}{c} F(\tau) \cos \theta_\gamma \right], \quad (2)$$

where $E_{\gamma 0}$ is the unshifted γ -ray energy, $v_{c.m.}$ is the recoil velocity of the center of mass, θ_γ is the angle of observation, and $F(\tau)$ is the attenuation factor, which depends on the stopping powers. By examining the energy of a γ ray as a function of angle, the $F(\tau)$ value can be extracted and

compared with those calculated using the Winterbon stopping power formalism [33], and the lifetime of the state, τ , can be obtained. An extensive discussion on the DSAM can be found in Ref. [32].

III. RESULTS

The levels, depopulating transitions, multipole mixing ratios, lifetimes, and resulting $B(E\lambda)$ transition probabilities are listed in Table I. Energies given for the γ rays were obtained from the angular-distribution measurements at the lowest incident neutron energy for which the γ -ray yields were observed to have good statistics. This procedure allowed for accurate energies of peaks which may evolve into complex structures at higher incident neutron energies. The branching ratios and lifetimes were also obtained at the lowest feasible neutron energies. The level energies were determined by a weighted least-squares fit of all the γ rays to and from a level.

Since the last evaluation of ^{160}Gd [34], three papers have made additions to the ^{160}Gd level structure using the $(n, n'\gamma)$ reaction [26,31,35]. This information has been considered along with the current results and is presented in Table I. Gamma rays from a level were only included if the excitation functions were consistent in shape and threshold for all γ rays. There were also previously placed γ rays that were below our experimental observation threshold and these were not included in the level scheme [31].

In the following sections, we discuss levels which are germane to the $K^\pi = 2_\gamma^+$ band and the negative-parity bands in ^{160}Gd . A partial level scheme is shown in Fig. 2.

A. $K^\pi = 2_\gamma^+$ band

The 2_γ^+ bandhead at 988.72 keV had a previously measured level lifetime of 1876(87) fs. This value is the weighted average of four Coulomb excitation experiments [36–39] as given in the latest nuclear data compilation of $A = 160$ [34]. We observed three γ rays from this level, but only the 913.43 and 988.68 keV γ rays were used in extracting the level lifetime limit of >1800 fs. This value is near the limit of our method, but is consistent with the accepted literature value. The 739.96 keV γ ray is near the energy of a background γ ray and, therefore, was not used in determining the level lifetime. In Table I, transition probabilities were calculated by using the literature lifetime and the associated branching ratios from Ref. [31] to calculate the $B(E2)$ values presented, after ensuring the consistency of the reported branching ratios with our data.

The 4_γ^+ member of the band at 1148.15 keV depopulates by 899.59 and 1072.85 keV γ rays. Govor *et al.* [31] observed a doublet at a γ -ray energy of 632.89 keV, which they placed as decays from both the 1148.14 and 1621.44 keV levels. From the excitation function shape and threshold, our work assigns this γ ray to the 1621.56 keV level.

The level at 1148.15 keV is assigned as $J^\pi = 4^+$ from polarized proton-scattering measurements, and the authors concluded a hexadecapole component for this state [40]. Our work yields a level lifetime $\tau = 1080_{-320}^{+730}$ fs, leading to $B(E2; 4_\gamma^+ \rightarrow 4_{gs}^+) = 16_{-10}^{+5}$ W.u. and $B(E2; 4_\gamma^+ \rightarrow 2_{gs}^+) = 3.8_{-1.1}^{+2.6}$ W.u., consistent with the expected quadrupole collec-

TABLE I. Energy levels (E_L) observed in the $^{160}\text{Gd}(n,n'\gamma)$ measurements. I_γ is the relative γ -ray intensity; $F(\tau)$ is the experimental attenuation factor; lifetimes τ are mean lives in fs. Multipole mixing ratios δ are obtained in fits to measured angular distributions. When two values are given, they have similar χ^2 values. If one value is in brackets, the one with the smaller χ^2 value is given first and is the δ value adopted. Literature lifetime values (τ_{lit}) are taken from Ref. [34] unless otherwise noted and are shown under the measured lifetime. Parentheses denote tentative assignments or placements. For reference, $B(E1)_{\text{m.w.u.}} = 0.190 e^2 \text{ b}$ and $B(E2)_{\text{w.u.}} = 5.16 \times 10^{-7} e^2 \text{ b}^2$.

E_L (keV)	J_i^π, K	J_f^π, K	E_γ (keV)	I_γ	$F(\tau)$	τ (fs)	Mult.	δ	$B(E1)$ ($\times 10^{-3}$ W.u.)	$B(E2)$ (W.u.)	Notes
75.24(5)	$2_{\text{gs}}^+, 0$	$0_{\text{gs}}^+, 0$	75.26	100		$\tau_{\text{lit}} = 3.9 \times 10^6$					
248.55(6)	$4_{\text{gs}}^+, 0$	$2_{\text{gs}}^+, 0$	173.34(5)	100							
515.00(8)	$6_{\text{gs}}^+, 0$	$4_{\text{gs}}^+, 0$	266.52(5)	100							
988.72(6)	$2_\gamma^+, 2$	$4_{\text{gs}}^+, 0$	739.96(7)	<2	0.008(14)	>1800	$E2$			0.96(7)	a .b.c
						$\tau_{\text{lit}} = 1876(87)$					
		$2_{\text{gs}}^+, 0$	913.43(5)	100(1)			$E2/M1$	$-0.45_{-0.05}^{+0.04}$		$1.2_{-0.18}^{+0.22}$	
		$0_{\text{gs}}^+, 0$	988.68(5)	88.7(10)			$E2$			4.1(2)	b
1057.42(6)	$3_\gamma^+, 2$	$4_{\text{gs}}^+, 0$	809.06(5)	20.6(2)	0.011(13)	>2200	$E2/M1$	0.11(3)		<0.04	b
		$2_{\text{gs}}^+, 0$	982.28(5)	100(1)			$E2/M1$	47_{-10}^{+18}		<6.5	d
1070.57(7)	$4_1^+, 4$	$6_{\text{gs}}^+, 0$	555.45(8)	1.6(1)							
		$4_{\text{gs}}^+, 0$	822.06(5)	100(2)							
		$2_{\text{gs}}^+, 0$	995.30(5)	64.2(13)							
1148.15(7)	$4_\gamma^+, 2$	$4_{\text{gs}}^+, 0$	899.59(5)	100(1)	0.037(15)	1080_{-320}^{+730}	$E2/M1$	21_{-7}^{+21}		16_{-10}^{+5}	d
		$2_{\text{gs}}^+, 0$	1072.85(5)	58.5(9)			$E2$			$3.8_{-1.1}^{+2.6}$	
1224.33(6)	$1_1^-, 0$	$2_{\text{gs}}^+, 0$	1149.12(5)	100(1)	0.676(7)	20 ± 2	$E1$		6.5(6)		
						$\tau_{\text{lit}} = 21.6(60)$					
		$0_{\text{gs}}^+, 0$	1224.38(5)	67.5(8)			$E1$		3.6(4)		
1261.24(9)	$5_\gamma^+, 2$	$6_{\text{gs}}^+, 0$	746.34(6)	19.6(9)	0.101(26)	350_{-80}^{+120}	$E2/M1$	8_{-4}^{+13}		31_{-11}^{+7}	
								[0.24(10)]		$1.7_{-1.5}^{+1.4}$	
		$4_{\text{gs}}^+, 0$	1012.65(5)	100(1)			$E2/M1$	15_{-6}^{+17}		35_{-12}^{+8}	
1289.90(7)	$3_1^-, 0$	$4_{\text{gs}}^+, 0$	1041.37(5)	54.6(7)	0.565(19)	34 ± 3	$E1$		3.0(3)		e
						$\tau_{\text{lit}} = 74(20)$					
		$2_{\text{gs}}^+, 0$	1214.79(5)	100(1)			$E1$		3.5(3)		
1351.30(6)	$1_2^-, 1$	$2_{\text{gs}}^+, 0$	1276.06(5)	100(2)	0.184(8)	180 ± 20	$E1$		0.74(8)		
		$0_{\text{gs}}^+, 0$	1351.30(5)	20.0(4)			$E1$		0.12(1)		
1376.70(8)	$2_1^-, 1$	$3_\gamma^+, 2$	319.38(6)	1.8(1)	0.035(47)	>550	$E1$		<0.18		b .f
		$2_{\text{gs}}^+, 0$	1301.46(5)	100(1)			$E1$		<0.27		
1379.71(10)	$0_2^+, 0$	$2_{\text{gs}}^+, 0$	1304.46(5)	100	0.015(14)	>1350	$E2$			<3.1	g
1427.40(12)	$5_1^-, 0$	$4_{\text{gs}}^+, 0$	1178.85(6)	100	0.514(69)	50 ± 10	$E1$		4.0(8)		
1436.34(7)	$2_2^+, 0$	$4_{\text{gs}}^+, 0$	1187.81(5)	100(1)	0.036(28)	>340	$E2$			<13	g
		$2_{\text{gs}}^+, 0$	1361.05(6)	36.4(4)			$E2/M1$	0.00(8)		<2.4	
								[$2.4_{-0.4}^{+0.6}$]		<2.0	
		$0_{\text{gs}}^+, 0$	1436.34(6)	13.5(2)			$E2$			<0.70	
1464.00(10)	$3_2^-, 1$	$2_{\text{gs}}^+, 0$	1388.75(5)	100	0.344(43)	50 ± 5	$E1$		2.5(2)		f
1498.94(10)	$4_1^-, 1$	$4_{\text{gs}}^+, 0$	1250.39(5)	100	0.053(51)	>400	$E1$		<0.41		f
1532.29(13)	$3_3^-, (3)$	$4_\gamma^+, 2$	384.14(6)	100							f
		$4_{\text{gs}}^+, 0$	1283.10(8)								i
1558.31(12)	$0_3^+, 0$	$2_{\text{gs}}^+, 0$	1483.06(6)	100	0.004(69)	>590	$E2$			<3.7	g
1561.63(10)	$4_3^+, 0$	$6_{\text{gs}}^+, 0$	1046.67(6)	100(1)	0.049(53)	>320	$E2$			<22	g
		$4_{\text{gs}}^+, 0$	1313.03(6)	74.8(3)			$E2/M1$	$0.28_{-0.12}^{+0.34}$		<0.40	

TABLE I. (*Continued.*)

E_L (keV)	J_i^π, K	J_f^π, K	E_γ (keV)	I_γ	$F(\tau)$	τ (fs)	Mult.	δ	$B(E1)$ ($\times 10^{-3}$ W.u.)	$B(E2)$ (W.u.)	Notes
1568.77(6)	$1_1^+, 1$	$1_2^-, 1$	217.51(6)	8.8(15)	0.044(28)	1000_{-400}^{+1800}	$E1$		$0.97_{-1.8}^{+0.7}$		b, f
		$2_\gamma^+, 2$	580.20(6)	89.2(16)			$E2/M1$	$0.28_{-0.18}^{+0.25}$		$5.3_{-13}^{+6.7}$	
		$2_{gs}^+, 0$	1493.40(6)	94.4(16)			$E2/M1$	$1.34_{-0.6}^{+1.6}$		$0.44_{-0.88}^{+0.23}$	
		$0_{gs}^+, 0$	1568.70(6)	100(2)			$M1$				
1586.61(12)	$2_3^+, 1$	$2_{gs}^+, 0$	1511.36(6)	100	0.044(39)	>500	$E2/M1$			<4.01	f
1599.05(6)	$2_4^+, 0$	$3_1^-, 0$	309.32(6)	8.9(4)	0.056(26)	800_{-300}^{+730}	$E1$		$0.34_{-0.31}^{+0.13}$		b, f
		$1_1^-, 0$	374.78(6)	14.8(3)			$E1$		$0.32_{-0.29}^{+0.12}$		
		$3_\gamma^+, 2$	541.53(6)	36.8(3)			$E2/M1$	$-5.57_{-5.00}^{+1.91}$ $-0.01(9)$		41_{-37}^{+16}	
		$2_{gs}^+, 0$	1523.59(6)	100(1)			$E2/M1$	$-1.04_{-2.10}^{+0.22}$		$0.004(77)$	
		$0_{gs}^+, 0$	1598.85(6)	78.7(1)			$E2$			$0.34_{-0.32}^{+0.68}$ $0.41_{-0.37}^{+0.15}$	
1621.56(9)	$2_2^-, 2$	$3_\gamma^+, 2$	564.06(6)	29.0(7)	0.167(155)	240_{-140}^{+3600}	$E1$		$1.7_{-0.1}^{+0.6}$		j
		$2_\gamma^+, 2$	632.94(6)	100(2)			$E1$		$4.2_{-0.3}^{+1.4}$		
1648.06(12)	4_5^+	$2_{gs}^+, 0$	1572.81(6)	100	0.134(58)	300_{-100}^{+260}	$E2$			$5.5_{-2.6}^{+2.8}$	f
1653.32(12)	$5_2^-, 1$	$6_{gs}^+, 0$	(1138.51(6))		0.454(78)	60_{-15}^{+20}					b, f
		$4_{gs}^+, 0$	1404.77(6)	100							
1661.75(9)		$4_{gs}^+, 0$	1413.16(7)	5.6(2)	0.050(32)	880_{-360}^{+1600}					i, f
		$2_{gs}^+, 0$	1586.52(6)	100(1)							
1691.68(7)	$3_4^-, 2$	$4_\gamma^+, 2$	543.45(6)	60.8(15)	0.170(95)	230_{-100}^{+350}	$E1$		2_{-3}^{+1}		j
		$3_\gamma^+, 2$	634.56(6)	99.5(25)			$E1$		$2.1_{-3.2}^{+0.9}$		
		$2_\gamma^+, 2$	702.84(6)	100(2)			$E1$		$1.5_{-2.3}^{+0.7}$		
		$4_{gs}^+, 0$	1442.93(8)	8.6(6)			$E1$		$0.01_{-0.02}^{+0.01}$		i
1782.67(10)	$4_2^-, 2$	$5_\gamma^+, 2$	521.53(8)	23.6(16)							
		$3_\gamma^+, 2$	725.19(6)	100(2)							
1805.13(9)	2_5^+	$4_1^+, 4$	734.50(6)	44.3(14)	0.055(96)	>300	$E2$			<76	f
		$2_\gamma^+, 2$	816.46(6)	100(2)			$E2/M1$	$-0.76_{-0.13}^{+0.10}$ $[-3.90_{-1.34}^{+0.97}]$		<37 <94	
1931.96(7)	2_6^+	$3_\gamma^+, 2$	874.51(6)	43.4(21)	0.057(39)	760_{-330}^{+1800}	$E2/M1$			$8_{-19}^{+3.5}$	b, f
		$4_{gs}^+, 0$	1683.45(7)	45.4(20)			$E2$			$0.32_{-0.75}^{+0.14}$	b
		$2_{gs}^+, 0$	1856.65(6)	100(3)			$E2/M1$	$0.92_{-0.64}^{+0.41}$ $0.50_{-0.24}^{+0.87}$		$0.20_{-0.47}^{+0.17}$ $0.08_{-0.31}^{+0.08}$	
		$0_{gs}^+, 0$	(1932.00(7))	31.5(17)			$E2$			$0.11_{-0.26}^{+0.05}$	
1966.66(10)	1_3^-	$2_{gs}^+, 0$	1891.33(6)	100(8)	0.606(48)	37_{-7}^{+8}	$E1$		$0.84_{-0.21}^{+0.28}$		
		$0_{gs}^+, 0$	1966.97(13)	57.0(70)			$E1$		$0.43_{-0.11}^{+0.16}$		
2030.90(9)	(2^+)	$4_{gs}^+, 0$	1782.14(6)	16.1(22)							f
		$2_{gs}^+, 0$	1955.93(7)	100(3)							
2060.43(11)		$0_{gs}^+, 0$	2060.42(6)	100	0.187(44)	230_{-50}^{+90}					
2109.32(7)	1^+	$3_\gamma^+, 2$	1051.87(6)	47.4(18)	0.138(34)	330_{-70}^{+120}	$E2$			$6.2(1.7)$	i, f
		$2_\gamma^+, 2$	1119.71(13)	57.9(29)			$E2/M1$			$5.6(1.5)$	i
		$2_{gs}^+, 0$	2034.26(6)	100(2)			$E2/M1$			$0.50(13)$	
		$0_{gs}^+, 0$	2109.31(6)	73.3(21)			$M1$				

TABLE I. (Continued.)

E_L (keV)	J_i^π, K	J_f^π, K	E_γ (keV)	I_γ	$F(\tau)$	τ (fs)	Mult.	δ	$B(E1)$ ($\times 10^{-3}$ W.u.)	$B(E2)$ (W.u.)	Notes
2135.76(13)		$0_{gs}^+, 0$	2135.74(7)	100	0.111(73)	420_{-180}^{+880}					^f

^aThis γ ray overlaps with an experimental background line, 739.42(5) keV from the $^{72}\text{Ge}(n, \gamma)$ reaction.

^bThis γ ray was not used in the level-lifetime determination.

^cThe branching ratios were taken from Ref. [31] and τ_{fit} was used for the transition probability calculations.

^dThe a_2 and a_4 values of this γ ray were within uncertainties of Ref. [31] and, therefore, the published δ value was used in this work.

^eLevel lifetime from Ref. [39]; see text for discussion.

^fThis level or spin assignment adopted from Ref. [31].

^gLifetime reported in Ref. [26].

^hLifetime modified from Ref. [26].

ⁱThis γ ray is not reliable for excitation function and/or lifetime determination due to its low intensity.

^jThis level or spin assignment adopted from Ref. [35].

tivity of this level and the $K^\pi = 2^+$ γ bands in the rare-earth deformed region of nuclei.

The 1261.24 keV 5_γ^+ level decays to the ground-state band, $B(E2; 5_\gamma^+ \rightarrow 4_{gs}^+) = 35_{-12}^{+8}$ W.u. and $B(E2; 5_\gamma^+ \rightarrow 6_{gs}^+) = 31_{-10}^{+7}$ W.u. The ratios of these transition probabilities are consistent, within error, with the Alaga rules.

B. $K^\pi = 0^-$ band

The 3_1^- level at 1289.90 keV deexcites by two γ rays, 1041.37 and 1214.79 keV. The 1214.79 keV γ ray is observed with an $a_2 = -0.276(33)$, which supports the assignment of an $E1$ transition. Ref. [31] assigns the 1214.79 keV γ ray as a doublet, but we see only a single γ ray in our measurements. A lifetime of $\tau = 74 \pm 20$ fs was reported from Coulomb excitation and Doppler-broadening measurements [39], but our value of $\tau = 34 \pm 3$ fs is not in agreement with this value. An advantage of the $(n, n'\gamma)$ reaction with accelerator-produced neutrons of variable energies is the ability to nonselectively

populate levels up to the incident neutron energy. By determining the lifetime of the 1289.90 keV level at $E_n = 1.5$ MeV, the feeding of the level has been greatly reduced from levels with significantly longer lifetimes. If the neutron energy is increased to 2.0 MeV, increasing feeding from higher levels, the apparent lifetime is increased to $\tau = 60 \pm 10$ fs. The method of selecting the neutron energy for population of the level gives us confidence in our measured lifetime values. The $K^\pi = 0^-$ band with $1_1^-, 3_1^-, 5_1^-$ members at 1224.33, 1289.90, and 1427.40 keV, respectively, decay by $E1$ transitions to the ground-state band. $E1$ transitions to the γ band were not observed.

C. $K^\pi = 1^-$ band

The 1351.30 keV level is the bandhead of the $K^\pi = 1^-$ band and was introduced in Ref. [41] by the placement of 1351.0 and 1275.90 keV γ rays. We observe both of these γ rays, and our spectra are not complicated by the doublet




FIG. 2. A partial level scheme of ^{160}Gd highlighting the $K^\pi = 2^+$ γ and negative-parity bands. The $B(E2)$ values in W.u. are shown in green and $B(E1)$ values in mW.u. in purple and scaled separately. All values are also listed in Table I.

at 1351.09 keV reported in Ref. [31]. We determine a level lifetime of $\tau = 180 \pm 20$ fs.

A level at 1388.64 keV was assigned by Berzin *et al.* [41] with five deexciting γ rays, where those at 1388.7 and 1313.0 keV were the most intense. In later work [31], this level was rejected on the basis of a measured angular distribution and the 1388.56 keV γ ray was assigned to a new 3^- level at 1464.00 keV. This 1464.00 keV level was assigned as a $J^\pi = 3^-$ member of the 1^- band [31]. The 1388.75 keV γ ray has an excitation threshold of 1.5 MeV which favors the placement from the 1464.00 keV level with an a_2 value in agreement with Ref. [31]. Further evidence to support the removal of the 1388.7 keV level is obtained from the γ ray at 1313.03 keV. The excitation threshold is higher than for the 1388.64 keV γ ray and is consistent with the placement at the 1561.63 keV level. Our data do not support any other decays to or from the 1388.7 or the 1464.00 keV level. The 1388.64 keV level has been removed from the level scheme.

D. $K^\pi = 2^-$ band

A $K^\pi = 2^-$ band was proposed in Ref. [35] with levels at 1621.4 keV (2^-), 1691.4 keV (3^-), 1782.5 keV (4^-), and 1884.2 keV (5^-).

In Ref. [31], a doublet at 632.82 keV was reported and the γ rays were placed from the 1148.14 keV level (4_4^+) and a 1621 keV level. The energy thresholds in our data support placement at 1621.56 keV. Unlike Ref. [31], the 632.94 keV γ ray is a singlet with $a_2 = 0.180(42)$ and $a_4 = -0.030(69)$. CINDY calculations support the placement of both the 632.94 and 564.06 keV γ rays to a $J^\pi = 2^-$ state, in agreement with Ref. [35].

The γ rays observed in this work from the 3^- level at 1691.68 keV agree with those proposed in Ref. [31]. We observe a level lifetime of $\tau = 230_{-100}^{+350}$ fs with a preferred decay to the γ band, $B(E1; 3_4^- \rightarrow 4_4^+) = 2_{-3}^{+1}$ mW.u., $B(E1; 3_4^- \rightarrow 3_4^+) = 2.1_{-3.2}^{+0.9}$ mW.u., and $B(E1; 3_4^- \rightarrow 2_4^+) = 1.5_{-2.3}^{+0.7}$ mW.u.

The 4^- level at 1782.67 keV was proposed in Ref. [35]. We confirm the placement of the 725.19 and 521.53 keV γ rays but are unable to calculate a lifetime due to low statistics.

Recent studies of quadrupole and octupole states in ^{168}Yb ($R_{4/2} = 3.27$) noted a pattern of strong decay from a $K^\pi = 2^-$ band into the γ -vibrational band when compared with the decay into the ground-state band [42]. This is consistent with the degree of K forbiddenness. Earlier work by Aprahamian [43] showed a difference of three orders of magnitude in $E1$ strength for transitions between states with the same K ($\Delta K = 0$) and for those with $\Delta K = 2$. The authors note [42] that this interesting decay pattern has also been observed in the $J^\pi = 2^-$ states belonging to the $K^\pi = 2^-$ band in the nuclei from Gd ($Z = 64$) to W ($Z = 74$). The preferred decay ^{168}Yb is to the $J^\pi = 4^-$ state, providing more evidence of a $K^\pi = 2^-$ band. The enhanced $E1$ transitions indicate some overlap with the γ band but further study of the negative-parity states, including $E3$ strengths, are needed.

E. $K^\pi = 0^+$ bands

The 0^+ states in ^{160}Gd were discussed in Ref. [26]. 0^+ states at 1379.7 and 1558.3 keV were confirmed and assigned lifetime limits of >1350 and >590 fs, respectively, and states at 1325.73 and 2236 keV were rejected as possible 0^+ states. The band structure of these levels was also explored and is included in Table I.

After evaluation of the full angular distribution data set at neutron energies of 1.5, 2.2, and 2.8 MeV, the $F(\tau)$ values were scrutinized. From a careful evaluation of the full data set, only $F(\tau)$ values with uncertainties which overlapped the other values were used in the lifetime calculation. This process led to the recalculation of the lifetime of the 1599.05 keV level. The reevaluated value of $\tau = 800_{-300}^{+730}$ fs is consistent with our formerly published limit of >300 fs [26], albeit with a large uncertainty. The γ rays of 1599.05 and 988.72 keV overlap with background lines and were not observed in this set of experiments.

F. Discussion

We have studied levels in ^{160}Gd with the $(n, n'\gamma)$ reaction. Excitation functions aided in the placement of γ rays in the level scheme and angular distributions resulted in the determination of multipole mixing ratios and level lifetimes of excited states and, hence, the depopulating $B(E2)$ and $B(E1)$ values for transitions from $K^\pi = 0^+, 2^+, 4^+, 0^-, 1^-$, and 2^- bands. In a previous paper [26], we identified excited $K^\pi = 0^+$ bandheads at 1379 and 1558 keV and the excited levels built on them. The measured lifetimes for the first- and second-excited $K^\pi = 0^+$ band members were determined as limits. Nonetheless, the limits indicate potentially enhanced collective transitions from the 2^+ and 4^+ members of the band. In this work, we have measured the level lifetimes for several members of the negative-parity bands. Figure 3 shows the revised lifetime of the 2^+ state of the $K^\pi = 0^+$ band at 1599 keV with an enhanced $B(E2)$ of 41_{-37}^{+16} W.u. connecting this state to the $K^\pi = 2^+$ band member, the 3^+ state at 1057. This kind of enhanced $B(E2)$ transition probability could be due to a collective excitation built on the $K^\pi = 2^+$ band. $B(E2)$ values to the ground-state band are 100 times weaker.

The lifetimes of the excited states in negative-parity bands are listed in Table I. A partial level scheme is shown for the depopulation of the negative-parity bands for $K^\pi = 0^-, 1^-$, and 2^- bands along with the $K^\pi = 2^+\gamma$ band. These transitions are $E1$ in nature and show $B(E1)$ values that are strongly enhanced for the $K = 0^-$ band depopulating on the order of a few mW.u. These are three orders of magnitude more enhanced than typical $E1$ values in deformed nuclei [43,44]. In comparison, the negative-parity bands with $K^\pi = 1^-$ and 2^- are one and two orders of magnitude weaker in $B(E1)$ to the ground-state band while the $K^\pi = 2^-$ band member at 1692 shows enhanced $E1$ transitions connecting this level to the $K^\pi = 2^+$ band. The Alaga values for the ratio of $B(E1; 1_{K^\pi=0^-}^- \rightarrow 2_{\text{gs}}^+)$ to $B(E1; 1_{K^\pi=0^-}^- \rightarrow 0_{\text{gs}}^+)$ is 2.0 and the experimental ratio is 1.7.

A disagreement over the assignment of the 1622 keV between Ref. [31] ($J^\pi = 4^+$) and Ref. [35] ($J^\pi = 2^-$) can be addressed. The current data support the 2^- assignment and




FIG. 3. A level scheme of ^{160}Gd highlighting the $K^\pi = 0^+$ bands. The $B(E2)$ values in W.u. are shown in green and $B(E1)$ values in mW.u. in purple and are scaled separately. All values are also listed in Table I.

experimental $B(E1; 2_{K^\pi=2^-}^- \rightarrow 3_{K^\pi=2^+}^+)$ to $B(E1; 2_{K^\pi=2^-}^- \rightarrow 2_{K^\pi=2^+}^+)$ of 0.41, in good agreement with the theoretical Alaga value of 0.36. Similarly, the $B(E1; 3_{K^\pi=0^-}^- \rightarrow 4_{\text{gs}}^+)$ to $B(E1; 3_{K^\pi=0^-}^- \rightarrow 2_{\text{gs}}^+)$ = 1.3 and the experimental $B(E1)$ ratio is 0.9, in good agreement within errors. The ratio of $B(E1; 3_{K^\pi=2^-}^- \rightarrow 4_{K^\pi=2^+}^+)$ to $B(E1; 3_{K^\pi=2^-}^- \rightarrow 3_{K^\pi=2^+}^+)$ to $B(E1; 3_{K^\pi=2^-}^- \rightarrow 2_{K^\pi=2^+}^+)$ for the 1691.68 keV level to the 4⁺, 3⁺, 2⁺ members of the $K^\pi = 2^+$ γ band, should be 1.8 : 1.4 : 1.0 from the Alaga rules and the experimental equivalent is 1.3 : 1.4 : 1.

The agreement of the Alaga rules with the measured $B(E1)$ values at 1692 as the 3⁻ member of a $K^\pi = 2^-$ band is further evidence of the $K^\pi = 2^-$ assignment. If the 1691.68 keV level belongs to a $K^\pi = 3^-$ band, then the Alaga ratio would be 0.05 : 0.35 : 1, in complete disagreement with measured values.

The collective quadrupole and octupole degrees of freedom were analyzed in ^{160}Gd by using the well-known generator coordinate method (GCM) with the Gogny D1S energy density function. Mean-field Hartree-Fock-Bogoliubov (HFB) constrained calculations were carried out by using the axially symmetric quadrupole and octupole mass moments in order to explore the shape of the potential-energy surface (PES) and to check for the existence of octupole deformed minima. An in-depth discussion of this method can be found in Refs. [44–46]. The computed PES reveals a reflection symmetric (i.e., with zero octupole deformation) for the ground state but the PES along the octupole direction is rather soft, indicating the need of a beyond-mean-field calculation using the GCM. Figure 4 shows a deep minimum potential well that is quadrupole deformed. The lowest oblate minimum does not appear until 5 MeV. The calculations show a first-excited $K = 0^-$ state appears at an excitation energy of 1.84 MeV with a $B(E1; 1_{K^\pi=0^-}^- \rightarrow 0_{\text{gs}}^+)$ value of 5.1 mW.u., as shown in Fig. 5(a). The $B(E3; 3^- \rightarrow 0_{\text{gs}}^+) = 13.50$ W.u. The second 0⁺ state is a one-phonon β vibration with an excitation energy of 3.36 MeV and the 2⁺ member decays to the ground state with

a $B(E2; 2^+ \rightarrow 0_{\text{gs}}^+) = 3.7$ W.u. and the $10^3 \rho^2 (E_0)$ transition strength of 230. The third 0⁺ state in this calculation is a two-octupole phonon at 3.94 MeV with a $B(E2; 2^+ \rightarrow 0_{\text{gs}}^+) = 0.71$ W.u. and a $10^3 \rho^2 (E_0) = 9$. These calculated values are compared with the experimental values in Fig. 5. We have not measured any $E3$ transitions in this work, but the $B(E3)$ value used in comparison was established previously. Our experimental level scheme is compressed in comparison to the GCM calculations. There is, however, very good agreement between the calculated and experimental $B(E1; 1_{K^\pi=0^-}^- \rightarrow 0_{\text{gs}}^+)$ values. The prediction is for an excited $K^\pi = 0^+$ band at 3.9 MeV as a double octupole phonon built on the $K^\pi = 0^-$ band at 1840 keV. Our work shows an excited $K^\pi = 0^+$ band at 1558 keV that is connected to the first-excited $K^\pi = 2^+$ band, as shown in Table I and Fig. 3. All indications are that this $K^\pi = 0^+$ band at 1558 keV is the double-phonon $\gamma\gamma$ vibrational band. The $K^\pi = 2^+$ γ bandhead is at 989 keV. The two-phonon $\gamma\gamma$ vibration in this case is 1.6 times the energy of the single γ -vibrational band. Negative anharmonicities similar to this have been observed in ^{232}Th [47] in the actinide region and ^{178}Hf [25] in the rare earths.




FIG. 4. Potential-energy surface as a function of the quadrupole and octupole degrees of freedom for ^{160}Gd . The dot is located at the ground-state minimum.




FIG. 5. Partial level schemes for ^{160}Gd . The $B(E2)$ values in W.u. are shown in green, $B(E1)$ values in mW.u. in purple, and $B(E3)$ values in W.u. in white outlined in black. The two level schemes are scaled separately. (a) GCM calculations using the Gogny D1S energy density functional is shown along with calculated transition probabilities. In this calculation, the first-excited $K^\pi = 0^-$ state appears at an excitation energy of 1.84 MeV and a $B(E1; 1_{K^\pi=0^-}^- \rightarrow 0_{\text{gs}}^+)$ value of 5.1 mW.u. The $B(E3; 3^- \rightarrow 0_{\text{gs}}^+) = 13.50$ W.u. The second-excited state is a one-phonon β vibration with an excitation energy of 3.36 MeV and the 2^+ member decays to the ground state with a $B(E2; 2^+ \rightarrow 0_{\text{gs}}^+) = 3.7$ W.u. The third $K^\pi = 0^+$ band is a two-phonon β vibration at 3.94 MeV, and $B(E2; 2^+ \rightarrow 0_{\text{gs}}^+) = 0.71$ W.u. The GCM calculation predicts a double-phonon excited $K^\pi = 0^+$ band at 3.9 MeV built on the $K^\pi = 0^-$ band at 1840 keV. (b) The experimental partial level scheme with experimental transition probabilities. The $B(E3; 3^- \rightarrow 0_{\text{gs}}^+)$ for the 1289 keV level is taken from Ref. [39].

IV. SUMMARY

We report the results of $(n, n'\gamma)$ measurements in the low-lying excitation-energy regime of ^{160}Gd . Transition probabilities were calculated from the level lifetimes and, where possible, multipole mixing ratios are determined. The $B(E2)$ values from the $K^\pi = 2_\gamma^+$ band indicate that this is indeed a quadrupole collective vibrational excitation. The level lifetimes of the 4^+ and 5^+ members of this band show transition rates consistent with other $K^\pi = 2_\gamma^+$ -vibrational bands in this region of deformed nuclei. Our characterization of the 1599.05 keV level with a revised multipole mixing ratio and the assignment or placement of the transitions to the γ band would make the $B(E2; 2_4^+ \rightarrow 3_\gamma^+) = 41_{-37}^{+16}$ W.u., potentially pointing to a two-phonon $\gamma\gamma$ -vibrational character for the $K^\pi = 0^+$ band at 1558.31 keV. The uncertainty on this $B(E2)$ value is large, but the observed collective strength, indicating a preferred decay to the $K^\pi = 2^+$ γ band, is difficult to ignore.

In deformed nuclei, low-lying negative-parity states are generally associated with octupole vibrations, and a simple pattern of $K^\pi = 0^-, 1^-, 2^-,$ and 3^- bands is frequently seen. In pioneering work, Neergård and Vogel [48] described the octupole states in deformed nuclei microscopically, and Barfield, Wood, and Barrett [49] treated the detailed spectra and $E3$ transition rates of nuclei in the rare-earth region within the interacting boson model. Modern approaches to the problem, such as those used in this paper, are required for a more robust prediction of transition strengths and excitation energies of the negative-parity states.

Experimentally, octupole bands are identified by enhanced $E3$ transitions between the $J^\pi = 3^-$ member of the band

and the ground state in spherical nuclei. The compilation of experimental data by Spear [50] and the update by Kibédi and Spear [11] give the excitation energies of the 3^- states and reduced electric octupole transition probabilities, $B(E3; 0_1^+ \rightarrow 3_1^-)$, for the first 3^- states of even-even nuclides. Moreover, it is frequently found that known octupole states are populated strongly in single-nucleon transfer reactions, indicating that these states have a complex character with one or more large two-quasiparticle components. While a number of $E1$ transition rates from negative-parity states are reported in this work, the correlation between these $B(E1)$ and octupole strength is less straightforward than that for the $B(E3)$. We note that, for example, the 1224.33 and 1289.90 keV levels of the $K^\pi = 0^-$ band have deexciting transitions of several mW.u. compared with the 1^- band where the $B(E1)$ are less than 1 mW.u. Structural assignments based solely on $B(E1)$ values should be made with caution [51]; therefore, we note these values, along with the previous reported $B(E3; 3^- \rightarrow 0_{\text{gs}}^+) = 11.1(7)$ W.u. [39], indicate an octupole vibration. However, no new $B(E3)$ values are available from the current work. Also of note is a preferred $\Delta K = 0$ decay from the $K^\pi = 2^-$ band to the $K^\pi = 2^+$ γ band from the 3^- state at 1691.68 keV.

Although the energies of the calculated bands in the GCM are higher than the experimental results, the agreement in the transition probabilities suggesting the population of a one-phonon β vibration and a two-phonon β vibration is striking. Further work is underway to include two-quasiparticle excitations in the GCM. New measurements of extended $B(E2)$ and $B(E0)$ values in ^{160}Gd can further clarify the situation.

ACKNOWLEDGMENTS

We thank H.E. Baber for his contributions to accelerator maintenance and operation. This material is based upon work supported by the National Science Foundation under Grants No. PHY-1606890, No. PHY-1419765, No. PHY-1205412, No. PHY-1507053, No. PHY-1068192, and No.

PHY-1305801. L.M. Robledo has been supported in part by Spanish Grants No. FPA2015-65929-P/MINECO and No. FIS2015-63770-P/MINECO. The enriched isotope used in this research was supplied by the United States Department of Energy Office of Science by the Isotope Program in the Office of Nuclear Physics.

-
- [1] A. Aprahamian, *Phys. At. Nucl.* **67**, 1750 (2004).
- [2] X. Wu, A. Aprahamian, S. M. Fischer, W. Reviol, G. Liu, and J. X. Saladin, *Phys. Rev. C* **49**, 1837 (1994).
- [3] A. Aprahamian, S. R. Leshner, C. Casarella, H. G. Börner, and M. Jentschel, *Phys. Rev. C* **95**, 024329 (2017).
- [4] P. E. Garrett, M. Kadi, C. A. McGrath, V. Sorokin, M. Li, M. Yeh, and S. W. Yates, *Phys. Lett. B* **400**, 250 (1997).
- [5] D. Bés, *Nucl. Phys.* **49**, 544 (1963).
- [6] W.-T. Chou, R. F. Casten, and P. von Brentano, *Phys. Rev. C* **45**, 9(R) (1992).
- [7] R. F. Casten and P. von Brentano, *Phys. Rev. C* **50**, 1280(R) (1994).
- [8] P. E. Garrett, *J. Phys. G* **27**, R1 (2001).
- [9] J. Smallcombe, P. J. Davies, C. J. Barton, D. G. Jenkins, L. L. Andersson, P. A. Butler, D. M. Cox, R. D. Herzberg, A. Mistry, E. Parr *et al.*, *Phys. Lett. B* **732**, 161 (2014).
- [10] P. D. Cottle, M. A. Kennedy, and K. A. Stuckey, *Phys. Rev. C* **42**, 2005 (1990).
- [11] T. Kibédi and R. H. Spear, *At. Data Nucl. Data Tables* **80**, 35 (2002).
- [12] R. A. Gatenby, J. R. Vanhoy, E. M. Baum, E. L. Johnson, S. W. Yates, T. Belgya, B. Fazekas, A. Veres, and G. Molnár, *Phys. Rev. C* **41**, 414(R) (1990).
- [13] L. Bargioni, P. G. Bizzei, A. M. Bizzeti-Dona, D. Bazzacco, S. Lunardi, P. Pavan, C. Rossi-Alvarez, G. de Angelis, G. Maron, and J. Rico, *Phys. Rev. C* **51**, 1057(R) (1995).
- [14] L. Caballero, B. Rubio, P. Kleinheinz, S. W. Yates, A. Algora, A. Dewald, A. Fitzler, A. Gadea, J. Jolie, R. Julin *et al.*, *Phys. Rev. C* **81**, 031301(R) (2010).
- [15] P. Kleinheinz, J. Styczen, M. Piiparinen, J. Blomqvist, and M. Kortelahti, *Phys. Rev. Lett.* **48**, 1457 (1982).
- [16] S. Lunardi, P. Kleinheinz, M. Piiparinen, M. Ogawa, M. Lach, and J. Blomqvist, *Phys. Rev. Lett.* **53**, 1531 (1984).
- [17] M. Piiparinen, P. Kleinheinz, J. Blomqvist, A. Virtanen, A. Atac, D. Muller, J. Nyberg, T. Ramsay, and G. Sletten, *Phys. Rev. Lett.* **70**, 150 (1993).
- [18] C. Küppersbusch, S. Pascu, P. von Brentano, D. Buscurescu, G. Căta-Danil, D. Deleanu, J. Endres, M. Elvers, D. Filipescu, C. Friesner *et al.*, *Eur. Phys. J. A* **48**, 1 (2012).
- [19] F. Corminboeuf, J. Jolie, H. Lehmann, K. Föhl, F. Hoyler, H. G. Börner, C. Doll, and P. E. Garrett, *Phys. Rev. C* **56**, 1201(R) (1997).
- [20] C. Fahlander, A. Axelsson, M. Heinebrodt, T. Härtlein, and D. Schwalm, *Phys. Lett. B* **388**, 475 (1996).
- [21] H. G. Börner, J. Jolie, S. J. Robinson, B. Krusche, R. Piepenbring, R. F. Casten, A. Aprahamian, and J. P. Draayer, *Phys. Rev. Lett.* **66**, 691 (1991).
- [22] M. Oshima, T. Morikawa, Y. Hatsukawa, S. Ichikawa, N. Shinohara, M. Matsuo, H. Kusakari, N. Kobayashi, M. Sugawara, and T. Inamura, *Phys. Rev. C* **52**, 3492 (1995).
- [23] T. Härtlein, M. Heinebrodt, D. Schwalm, and C. Fahlander, *Eur. Phys. J. A* **2**, 253 (1998).
- [24] P. E. Garrett, M. Kadi, M. Li, C. A. McGrath, V. Sorokin, M. Yeh, and S. W. Yates, *Phys. Rev. Lett.* **78**, 4545 (1997).
- [25] A. Aprahamian, R. C. de Haan, S. R. Leshner, J. Döring, A. M. Bruce, H. G. Börner, M. Jentschel, and H. Lehmann, *J. Phys. G* **25**, 685 (1999).
- [26] S. R. Leshner, C. Casarella, A. Aprahamian, B. P. Crider, R. Ikeyama, I. R. Marsh, M. T. McEllistrem, E. E. Peters, F. M. Prados-Estévez, M. K. Smith *et al.*, *Phys. Rev. C* **91**, 054317 (2015).
- [27] P. E. Garrett, N. Warr, and S. W. Yates, *J. Res. Natl. Inst. Stand. Technol.* **105**, 141 (2000).
- [28] P. E. Garrett, H. Lehmann, J. Jolie, C. A. McGrath, M. Yeh, W. Younes, and S. W. Yates, *Phys. Rev. C* **64**, 024316 (2001).
- [29] E. E. Peters, A. Chakraborty, B. P. Crider, B. H. Davis, M. K. Gnanamani, M. T. McEllistrem, F. M. Prados-Estévez, J. R. Vanhoy, and S. W. Yates, *Phys. Rev. C* **88**, 024317 (2013).
- [30] E. Shelton and V. C. Rodgers, *Comput. Phys. Commun.* **6**, 99 (1973).
- [31] L. I. Govor, A. M. Demidov, V. A. Kurkin, and I. V. Mikhailov, *Yad. Fiz.* **72**, 1799 (2009).
- [32] T. Belgya, G. Molnár, and S. W. Yates, *Nucl. Phys. A* **607**, 43 (1996).
- [33] K. B. Winterbon, *Nucl. Phys. A* **246**, 293 (1975).
- [34] C. W. Reich, *Nucl. Data Sheets* **105**, 557 (2005).
- [35] E. P. Grigoriev, *Phys. At. Nucl.* **75**, 1427 (2012).
- [36] Y. Yoshizawa, B. Elbek, B. Herskind, and M. C. Olesen, *Nucl. Phys.* **73**, 273 (1965).
- [37] C. Baktash, J. X. Saladin, J. O'Brien, I. Y. Lee, and J. E. Holder, *Phys. Rev. C* **10**, 2265 (1974).
- [38] R. Ronningen, J. Hamilton, A. V. Ramayya, L. Varnell, G. Garcia-Bermudez, J. Lange, W. Lourens, L. Riedinger, R. Robinson, P. Stelson *et al.*, *Phys. Rev. C* **15**, 1671 (1977).
- [39] F. K. McGowan and W. T. Milner, *Phys. Rev. C* **23**, 1926 (1981).
- [40] T. Ichihara, H. Sakaguchi, M. Nakamura, M. Yosoi, M. Ieiri, Y. Takeuchi, H. Togawa, T. Tsutsumi, and S. Kobayashi, *Phys. Rev. C* **36**, 1754 (1987).
- [41] Y. Y. Berzin, E. P. Grigórev, T. V. Guseva, and Yu. Ya. Tamberg, *Bull. Acad. Sci. USSR, Phys. Ser.* **53**, 76 (1989).
- [42] S. Pascu, D. Buscurescu, G. Căta-Danil, V. Derya, M. Elvers, D. Filipescu, D. G. Ghită, T. Glodariu, A. Hennig, C. Mihai *et al.*, *Phys. Rev. C* **91**, 034321 (2015).
- [43] A. Aprahamian, *Phys. Rev. C* **46**, 2093 (1992).
- [44] L. M. Robledo and G. F. Bertsch, *Phys. Rev. C* **84**, 054302 (2011).
- [45] L. M. Robledo and G. F. Bertsch, *Phys. Rev. C* **86**, 054306 (2012).

- [46] K. Nomura, R. Rodríguez-Guzmán, and L. M. Robledo, *Phys. Rev. C* **92**, 014312 (2015).
- [47] W. Korten, P. Brocking, H. Hubel, W. Pohler, U. van Severen, P. Willsau, T. Haertlein, C. Ender, P. Reiter, D. Schwalm *et al.*, *Z. Phys. A: Hadrons Nucl.* **A351**, 143 (1995).
- [48] K. Neergård and P. Vogel, *Nucl. Phys. A* **145**, 33 (1970).
- [49] A. F. Barfield, J. L. Wood, and B. R. Barrett, *Phys. Rev. C* **34**, 2001 (1986).
- [50] R. H. Spear, *At. Data Nucl. Data Tables* **42**, 55 (1989).
- [51] D. P. DiPrete, E. L. Johnson, E. M. Baum, C. A. McGrath, D. Wang, M. F. Villani, S. W. Yates, T. Belgya, B. Fazekas, and G. Molnár, *Phys. Rev. C* **52**, 2831(R) (1995).