



Assessment of Methane Production by Goats under Intermittent Grazing

Naomi C. Meister

Sao Paulo State University, Brazil

Nailson L. S. Lemos

University of Northeast Bahia State, Brazil

Fernando O. Alari

Sao Paulo State University, Brazil

Victor C. Silva

Sao Paulo State University, Brazil

William Koury Filho

Brasilcomz Company, Brazil

See next page for additional authors

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/22/2-8/14>

The XXII International Grassland Congress (Revitalising Grasslands to Sustain Our Communities) took place in Sydney, Australia from September 15 through September 19, 2013.

Proceedings Editors: David L. Michalk, Geoffrey D. Millar, Warwick B. Badgery, and Kim M.

Broadfoot

Publisher: New South Wales Department of Primary Industry, Kite St., Orange New South Wales, Australia

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Presenter Information

Naomi C. Meister, Nailson L. S. Lemos, Fernando O. Alari, Victor C. Silva, William Koury Filho, and Ana C. Ruggieri

Assessment of methane production by goats under intermittent grazing

Naomi C Meister^A, Nailson L S Lemos^B, Fernando O Alari^A, Victor C Silva^A, William Koury Filho^C and Ana C Ruggieri^A

^A Animal Science Department, University of Sao Paulo State, Brazil, www.fcav.unesp.br

^B University of Northeast Bahia State, Brazil, www.faneb.com.br

^C Brasilcomz Company, Jaboticabal, SP, Brazil

Contact email: acruggieri@fcav.unesp.br

Keywords: Leaf area index, tropical forage, greenhouse gas, sulfur hexafluoride, ruminants.

Introduction

There are 875 million goats worldwide (FAO 2012) and the majority are in developing countries where they are typically raised on pasture. Ruminants release gross energy as methane, which is a greenhouse gas. Methane mitigation strategies require prior knowledge about emission of methane by animals under different management systems. Intake, quality and type of feed are factors that influence methane emissions, therefore the objective of this study was to investigate how grazing intensity, measured by the residual leaf area index (RLAI), influences methane emission by goats under rotational grazing on Tanzania grass pasture.

Methods

The experiment was conducted at Jaboticabal, Sao Paulo, Brazil. The climate is tropical with a defined dry winter season between April and September and rainfall concentrated during the summer months (October to March). The experimental period started in November 2010, at the beginning of the rainy season. The 1.2 ha pasture area, formed 14 years ago with *Panicum maximum* Jacq cv. Tanzania grass, was divided into paddocks of 243 m² each. The criterion adopted for animal input into the paddock was 95% light interception (LI) and the cessation of grazing criterion were RLAI targets of 0.8, 1.6 and 2.4, when methane emissions and forage morphology were determined. LI and RLAI were measured using the canopy analyzer – AccuPAR Model LP – 80 PAR/LAI (Decagon Devices®). A rotational grazing system with three days of occupation was adopted. The animals remained in the paddocks for 11 hours during the day with free access to water and mineral salt. The morphological variables were evaluated within a randomized block design with six replicates. Canopy heights were determined pre- and post-grazing using a graduated ruler. The Tanzania grass morphological components of leaf, stem (stem + sheath) and dead material as well as forage dry matter (DM) were determined for a 0.5 m² area in the paddock defined using a metal quadrat. The morphological components were dried in a forced air oven at 55°C, for 72 h. Methane (CH₄) production was determined for 15 adult Anglo Nubian goats weighing average 56.8 kg that were neither

pregnant nor lactating and randomly distributed among the three RLAI treatments. Methane collection were conducted during April 2011, in two consecutive paddocks using the tracer gas sulfur hexafluoride (SF₆) technique, described by Jonhson *et al.* (1994). The concentrations of CH₄ and SF₆ were analyzed by gas chromatography (Agilent® model 6890). Feed consumption was estimated for 5 goats using two markers: LIPE® - purified lignin extracted from *Eucalyptus grandis*, which is modified and enriched hydroxyphenyl propane (Saliba e Araujo 2005) and indigestible neutral detergent fiber (NDFi). The chemical composition of Tanzania grass was determined from a sample of the grazable extract. The statistical analyses were performed by the software SAS (version 9.2, SAS Institute, Cary, NC, USA) using the PROC MIXED.

Results

LI average values were 97, 98 and 97% while RLAI were 0.8, 1.7 and 2.4, respectively, for the target treatments of 0.8; 1.6 and 2.4. The RLAI were considered close to the target for animal input and output within the experimental paddocks. The high values for total dry matter mass (TDMM) and stem DM mass (SDMM) shown in Table 1, were expected and are compatible with the tropical forage production of the genus *Panicum* (Quadros *et al.* 2002). The variables TDMM and SDMM were affected by RLAI and increased linearly ($P < 0.10$) in the pre-grazing. However, the variables, leaf DM (LDM), dead material DM (DMDM) and leaf/stem ratio (L/S) were not influenced by increasing RLAI (Table 1). The linear increase of TDMM and SDMM was possibly caused by tiller accumulation due to higher RLAI, which became old and increased in mass with time (Martha Junior *et al.* 2002). Another hypothesis is that post-grazing, the remaining leaves shaded the lower layers thus inducing stem elongation.

The linear increase of intake with increasing RLAI (Table 2) results from increased forage supply provided by the RLAI of 2.4, which was confirmed by greater leaf leftover in the pre-grazing (Table 1) and the lower number of animals in this treatment.

The 17% increase in intake between RLAI of 0.8 and 2.4 did not influence the variables related to methane production (Table 2). The methane emission value of 5.1

Table 1. Mean values (kg/ha) and corresponding contrast comparisons of morphological components of Tanzania grass as a function of residual leaf area index

Morphological Components		RLAI			Effect ¹	P-value ²
		0.8	1.6	2.4		
Pre grazing	LDMM	3,494	3,656	4,496	-	>0.10
	SDMM	5,106	5,601	6,380	L	0.035
	DMDMM	163	2,240	1,541	-	>0.10
	TDMM	10,237	11,498	12,418	L	0.010
	L/S	0.71	0.67	0.71	-	>0.10
Post Grazing	LDMM	933	1,502	1,658	L	0.021
	SDMM	3,637	3,607	3,916	-	>0.10
	DMDMM	929	861	962	-	>0.10
	TDMM	5,500	5,971	6,537	-	>0.10

¹ L (linear effect, considering $P < 0.10$); ² Probability associated with the F-test for contrasts. RLAI, residual leaf area index; LDMM, leaf dry matter mass; SDMM, stem dry matter mass; DMDMM, dead material dry matter mass; TDMM, total dry matter mass.

Table 2. Intake, average methane emission per metabolic weight, dry matter intake and energy loss in goats as a function of RLAI of Tanzania grasslands.

	RLAI			Effect ¹	P-value ²
	0.8	1.6	2.4		
Dry matter intake(g/day)	505 ± 15	514 ± 15	591 ± 15	L	<0.0001
CH ₄ /metabolic weight (g/body weight ^{0.75})	0.91 ± 0.05	0.88 ± 0.05	0.84 ± 0.05	-	>0.10
CH ₄ /MS intake (g/kg)	35.7 ± 2.7	37.3 ± 2.7	31.8 ± 2.7	-	>0.10
Loss of gross energy (%)	12 ± 0.9	12 ± 0.9	10 ± 0.9	-	>0.10

¹ L (linear effect, considering $P < 0.10$); ² Probability associated with F-test, for contrast; CH₄ = methane; RLAI = residual leaf area index; DM = dry matter; Metabolic weight = body weight^{0.75}; Loss of gross energy as methane was calculated using forage average energy of the two paddocks where methane was measured.

kg/year for a 40 kg animal in this study agrees with the IPCC (2006) of 5.0 kg/year. The similar methane emission for all three treatments can be explained by the similar chemical composition found for all three treatments, which displayed average 14% crude protein, 70 % NDF, 29% acid detergent fiber, 9% ash and 16.5 MJ/kg DM gross energy (GE). If the emission measurements had been made at different times during the rainy season, methane emission by goats grazing on Tanzania grass pasture may have differed among treatments.

Conclusion

The RLAI changed the features of Tanzania grass, without changing methane production. The methane emitted by goats under rotational grazing on Tanzania grassland was 0.88 g/kg metabolic weight (body weight^{0.75}) or 5.1 kg/year for a 40 kg animal.

Acknowledgments

The authors gratefully acknowledge the funding from FAPESP.

References

- Johnson KA, Huyler MT, Westberg HH, Lamb BK, Zimmerman P (1994) *Environmental Science and Technology* **28**, 359-362; DOI 10.1021/es00051a025.
- FAO – Food and Agriculture Organization. Visualizado em: <http://faostat.fao.org/site/573>
- Saliba EOS, Araújo VL (2005) Teleconferencia sobre o uso de indicadores em nutrição animal. UFMG, 45p.
- Quadros DG, Rodrigues LRA, Favoretto V, Malheiros EB, Herling VR, Ramos AKB (2002) Componentes da produção de forragem em pastagens dos capins Tanzânia e Mombaça adubadas com quatro doses de NPK. R. Bras. Zootec **31**, p.1333-1342, (suplemento)
- Martha Júnior GB, Corsi M, Barioni LG, Vilela, L (2004) Intensidade de desfolha e produção de forragem do capim-tanzânia irrigado na primavera e no verão. Pesquisa Agropecuária Brasileira., *Brasília* **39**, p.927-936, set. IPCC: Revised
- IPCC (2006) Guidelines for national greenhouse gas inventories: Reference manual. - Emissions from livestock and manure management. Cambridge: University Press.