



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

22nd International Grassland Congress

Carbon and Nitrogen Stocks in Integrated Crop Livestock

Valdinei T. Paulino

Instituto de Zootecnia, Brazil

Erika M. L. C. Teixeira

Instituto de Zootecnia, Brazil

Marcos Siqueira Neto

Universidade de São Paulo, Brazil

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/22/2-11/11>

The 22nd International Grassland Congress (Revitalising Grasslands to Sustain Our Communities) took place in Sydney, Australia from September 15 through September 19, 2013.

Proceedings Editors: David L. Michalk, Geoffrey D. Millar, Warwick B. Badgery, and Kim M.

Broadfoot

Publisher: New South Wales Department of Primary Industry, Kite St., Orange New South Wales, Australia

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Carbon and nitrogen stocks in integrated crop livestock

Valdinei T. Paulino^A, Erika M. L. C. Teixeira^A and Marcos Siqueira Neto^B

^A Instituto de Zootecnia, Brazil, www.iz.sp.gov.br

^B Universidade de São Paulo, Centro de Energia Nuclear na Agricultura, Piracicaba, São Paulo State, Brazil

Contact email: paulino@iz.sp.gov.br

Keywords: Conventional tillage, integrated crop-livestock system, *Urochloa brizantha*, *Urochloa ruziziensis*.

Introduction

Soil organic matter (SOM) is recognized as an important characteristic of sustainable agricultural systems in the tropics and temperate regions. It is considered an integrated characteristic of physical, chemical and biological soil conditions and has been used as a soil quality index in different land uses and management practices (Lal 2011). Environmental, social, and financial imbalances have occurred, requiring producers to become more efficient. A system of intercropping grain crops with forages can provide enough forage in the dry season to maintain nutritional quality of livestock and even promote weight gain (Castro Filho *et al.* 1991). An integrated crop-livestock system (ICLS) that includes a crop phase and a pasture phase in rotation has strong potential for sustainable beef and grain crop production (Salton *et al.* 2011). Perennial pastures can contribute to soil C sequestration. Several studies have shown significant soil organic C accumulation with well-managed grazing of pastures (Maia *et al.* 2009; Salton *et al.* 2011). However, there are few studies that compare the efficiency of different grasses to maintain productivity and promote an increase in soil organic C and stocks. We hypothesized that an ICLS with maize cultivated under no-tillage in rotation with grazed pasture (*Urochloa* P. Beauv. spp. (syn. *Brachiaria* (Trin.) Griseb. spp.) would result in accumulation of soil organic C and N compared with continuously grazed, degraded pasture or with conventionally tilled monoculture production of maize. Our objective was to evaluate the stocks of soil organic C and N under alternative land uses from the prevailing degraded pasture condition in Sao Paulo state of Brazil.

Methods

The study was located at the Institute of Animal Science (22°46'28.8"S; 47°17'38.2"W) in the municipality of Nova Odessa, São Paulo state (Brazil). The soil, a typical Acrudox, presented amounts of clay, silt and sand of 550, 160 and 290 g/kg, respectively. A continuously grazed pasture area cultivated with *Urochloa brizantha* cv. Marandu was selected as the base condition for this study. The pasture had low productivity and high degradation from the original condition. In August 2003 initial management upon conversion of a portion of the land area to ICLS consisted of surface application of 1.2 Mg/ha of lime. In November 2004, the first planting of maize (*Zea mays* L.) was made with no-tillage in a mulch

of grassy residues, using a chisel plow with cutting discs and winged tips. Maize was fertilized with 350 kg/ha of 8-26-16 (NPK) applied in the groove planting and 360 kg/ha of 5-20-20 (NPK) as top dressing with incorporation into soil mechanical incorporation to the soil with a trencher at 5 cm depth.

Five land use systems were compared: (1) integrated crop-livestock with maize and *Urochloa brizantha* cv. Marandu (ICL- Ub- Marandu); (2) integrated crop-livestock with maize and *U. ruziziensis* (ICL- Ruz); (3) integrated crop-livestock with maize and *U. brizantha* cv. Piata (ICL- Ub- Piata); (4) long-term (25-yr-old) pasture with *U. brizantha* (Cont - Pas); and (5) conventional-tillage cropping with– maize only (CT-maize). The experimental design was a completely randomized block with five replications; experimental units were paddocks of 0.8 ha. For the present purposes, soil evaluations were performed from sampling made in May 2010, after seven years of the experiment establishment. Soil carbon and nitrogen content and stocks were evaluated at depths of 0-10, 10-20, 20-30 and 30-40 cm. Total C and N concentrations were determined by dry combustion using an autoanalyzer LECO CN-2000 model. The Soil C and N stocks were calculated based on equivalent soil masses (Ellert and Bettany 1996), taking the soil mass of the native forest as a reference value (woodland Atlantic Forest). Analysis of variance was used to detect significant differences among treatments. Means were statistically separated using the Student Newman-Keuls test ($P < 0.05$). Analyses were performed using SAS (2010).

Results and Discussion

Soil organic C and N concentrations were greatest near the surface. In the surface layer (0-10 cm), C concentration ranged between 14.9 and 16.8 g/kg for ICL-Ub-Piata and ICL-Ub-Marandu, respectively; while N concentration ranged between 1.2 and 1.4 g/kg for CT-maize and ICL-Ub-Marandu, respectively. Soil organic C and N concentrations were typical, *i.e.* higher concentrations at the surface and lower concentrations with increasing depth. With depth, soil organic C and N concentrations became more similar among treatments due to the decreasing influence of surface deposition of organic matter inputs. Surface soil organic C accumulation can occur due to deposition of leaves, stems and roots. Highest soil organic C stocks were obtained under ICL-Ub-Marandu and ICL-Ruz followed




Figure 1. Soil organic C (a) and N (b) stocks in different depths (0-10, 10-20, 20-30 and 30-40 cm) under long-term pasture (R1 – cont pas), conventionally tilled maize silage (CT-maize), integrated crop-livestock systems with maize and *Urochloa brizantha* cv. Marandu (ICL Mar), with *U. ruziziensis* (ICL Ruz), and with *U. brizantha* cv. Piata (ICL Pia). Means (n = 5) with different letters at a depth 0–40 cm differ significantly at $P < 0.05$.

by ICL-Ub-Piata, long-term pasture, and CT maize (Fig. 1a). Similarly, soil N stock was greater in the ICL-Ub-Marandu and ICL-Ruz than other systems (Fig. 1b). No differences occurred among ICL-Ub-Piata and the two references areas.

Annual plowing and harrowing of maize leads to organic matter mineralization and, consequently, lowest soil organic C stock. Productive pastures can maintain soil organic C stock. The initial condition of soil physical and chemical limitations for cultivation resulted in low organic carbon contents. The amounts of N added and / or present in the soil were deficient and limited the addition of C in systems constituted mainly by grasses. This is showed as the difference in soil organic C and N between ICL and CT- maize was not so large in this soil. The C:N ratio of SOM was similar among depths sampled and values were similar to those recognized for stable SOM in the tropics. This gives a hint that the high C input from plant residues were being offset by additional N from somewhere in the system. Additional N could be due to fertilizer input in ICL (although there is high N demand by maize that is subsequently exported in grain harvest), or due to biological N fixation by diazotrophic communities present in the rhizosphere of grasses or as endophytic mutualists (Boddey *et al.* 2001). Paul and Clark (1989) attributed soil organic C accumulation under conservation tillage systems to two main factors: (1) physical protection of organic compounds against microbial decomposition, favored by the C occlusion in soil aggregates; and (2) chemical protection of organic compounds through their interaction with minerals and cations hindering their decomposition.

Conclusion

Integrated crop-livestock systems have a great

potential to increase SOM, represented in this study in the form of C and N stock accumulation. It was the combination of crops and pastures that improved soil organic C and N accumulation compared with continuation of long-term continuously grazed pasture or monoculture cultivation of maize alone. We found larger accumulation rates of soil organic C and N with *U. brizantha* cv. Marandu and *U. ruziziensis* than with *U. brizantha* cv. Piata and with the reference conditions of long-term pasture or conventional tillage cultivation of maize.

References

- Boddey RM, Polidoro JC, Resende AS, *et al.* (2001) Use of the ^{15}N natural abundance technique for the quantification of the contribution of N_2 fixation to sugar cane and other grasses. *Australian Journal of Plant Physiology* **28**, 889–895.
- Castro Filho C, Henklain JC, Vieira MJ, *et al.* (1991) Tillage methods and soil and water conservation in southern Brazil. *Soil Tillage Research* **20**, 271–283.
- Ellert BH, Bettany JR (1996) Calculation of organic matter and nutrients stored in soils under contrasting management regimes. *Canadian Journal of Soil Science* **75**, 529–538.
- Lal R (2011) Sequestering carbon in soils of agro-ecosystems. *Food Policy* **36**, 33–39.
- Maia SMF, Ogle SM, Cerri CEP, *et al.* (2009) Effect of grassland management on soil carbon sequestration in Rondônia and Mato Grosso states, Brazil. *Geoderma, Amsterdam* **149**, 84–91.
- Paul EA, Clark FE (1989) 'Soil microbiology and biochemistry.' (Academic Press, San Diego)
- Salton JC, Mielniczuk J, Bayer C, *et al.* (2011) Teor e dinâmica do carbono no solo em sistemas de integração lavoura-pecuária. *Pesquisa Agropecuária Brasileira*, **46**, 1349–1356.