



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

XXII International Grassland Congress

---

## Effect of Organic Nitrogen Levels on N Fixation in Pea-Barley Mixture

H. W. Lee

*Korea National Open University, South Korea*

J. D. Kim

*Cheonan Yonam College, South Korea*

C. H. Kim

*Kongju National University, South Korea*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/22/2-11/15>

The XXII International Grassland Congress (Revitalising Grasslands to Sustain Our Communities) took place in Sydney, Australia from September 15 through September 19, 2013.

Proceedings Editors: David L. Michalk, Geoffrey D. Millar, Warwick B. Badgery, and Kim M.

Broadfoot

Publisher: New South Wales Department of Primary Industry, Kite St., Orange New South Wales, Australia

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

# Effect of organic nitrogen levels on N fixation in pea-barley mixture

HW Lee<sup>A</sup>, JD Kim<sup>B</sup> and CH Kim<sup>C</sup>

<sup>A</sup> Department of Agricultural Science, Korea National Open University, Seoul 110-791, Republic of Korea

<sup>B</sup> Cheonan Yonam College, Sunghwan, Cheonan-Si 330-709, Republic of Korea

<sup>C</sup> Department of Plant Resources, Kongju National University, Chongnam Ye-san Daehak-ro 54, Republic of Korea

Contact email: [hyowon@knou.ac.kr](mailto:hyowon@knou.ac.kr)

**Keywords:** Organic farming, N fixation.

## Introduction

One limitation in organic farming is short supply of forage. Organic fertilizer including cattle manure is overproduced in Korea but farmers are reluctant to use these kinds of resources because of heavy weight and handling cost. Legumes are of crucial importance for the N supply of grassland farming. They convert atmospheric dinitrogen (N<sub>2</sub>) to plant available N through symbiosis. Information on N<sub>2</sub> fixation and transfer is lacking in Asian countries with a lot of results coming from EU and America. The objective of this report was to estimate potential N fixation and determine related benefits from intercropping pea with barley at different organic fertilizer levels. We also wished to measure the efficacy of N<sub>2</sub> fixation by <sup>15</sup>N dilution and the difference method using barley as reference crop.

## Materials and Methods

Pure stands of pea (*Pisum sativum* L. cv. Chungmi) and barley (*Hordeum vulgare* L. cv. Albori) and a mixture of the two were sown on 15<sup>th</sup> of March in 2009. Seed rate was 40 kg/ha of barley and 70 kg/ha of pea. The experiments were arranged in a complete block design with 3 replications. Treatments were three levels organic fertilizer N, 40, 80 and 120 kg/ha. P and K were applied in a chemical fertilizer. When the crops were at the early stage of plant growth, a subplot of 600cm<sup>2</sup> was fertilized with a solution of <sup>15</sup>NH<sub>4</sub>SO<sub>4</sub> at 99.7 atom% excess at a rate to corresponding to 7 kg/ha. Crops were harvested at maturity from each plot and separated into pea and barley. Total N content and <sup>15</sup>N values of the sample were analyzed using a continuous flow stable isotope ratio mass spectrometer. Nitrogen fixation of pea was determined with a procedure described by Peoples *et al.* (1989) and Chalk (1996).

## Results and Discussion

Table 1 shows the estimate of N fixation and transfer by <sup>15</sup>N dilution. The percent of total N derived from atmosphere (%Ndfa) at maturity ranged from 0.89 to 4.45, indicating a very low value. There were significant differences between treatments with 40kg organic N lower than other N levels. Sakai *et al.* (2011) reported intercropping with lupin and cowpea transferred 18 and 17% of N, and transfer N from pea to oat was 10.7 to 10.3 kg/ha. The greatest fixation was from the 80 kg organic N plots. Using the applied difference method, the transfer rate and transfer amount were highest in the 120 kg N

**Table 1. Estimate of N fixation and transfer to barley at different levels of organic N fertilization by <sup>15</sup>N dilution method.**

Treatments (N/ha)	N fixation rate (%)	Total N (kg/ha)	Fixed amount (kg/ha)	Transfer amount (kg/ha)
Organic N 40 kg	0.89a	192	1.71	0
Organic N 80kg	4.43b	233	10.32	0
Organic N 120kg	4.65b	195	9.08	0

<sup>15</sup>N-dilution method. Pnon-leg(<=atm) = 1 - (Enon-leg(m)/Enon-leg(p))  
Pleg(<=atm) = 1 - (Eleg(m)/Enon-leg(p))

**Table 2. Estimate of N transfer from pea to barley at different levels of organic N fertilization by difference method**

Treatment (N/ha)	N transfer rate (%)	Total N (kg/ha)	Transfer amount (kg/ha)
Organic N 40 kg	23.1a	192	44.1
Organic N 80 kg	23.2a	232	56.9
Organic N 120 kg	47.5b	195	91.7

The N-difference method. Nleg(=>non-leg) = Nnon-leg(m) - Nnon-leg(p)-R(1, Pnon-leg(<=leg) = Nleg(=>non-leg)/Nnon-leg(m) = 1 - (Nnon-leg(p)-R/Nnon-leg(m))

treatment (Table 2). N transfer rate was from 23.1 to 47.5% and the amount ranged from 44.1 to 91.7 kg/ha. Haby *et al.* (2006), reported %Ndfa was 42 and the transfer N yield was less than 18kg/ha/yr. Anderson *et al.* (2004) reported that the greatest relative amount of N<sub>2</sub> fixed at maturity was measured for the pea grown in association with rape under condition of low N fertilization. Because the <sup>15</sup>N dilution method is based on concentration (of <sup>15</sup>N) in the N uptake and not on the total N amount as in the difference method, there was difference between two methods in terms of N fixation and transfer amount. (Geijersstam *et al.* 2006).

## Conclusion

Our findings indicate that there was large difference between <sup>15</sup>N dilution and difference method for determination of N fixation from atmosphere. N transfer to barley was 0.89 to 4.65% in <sup>15</sup>N dilution and 23.1 to 47.5% in difference method when organic N fertilization levels were different. We also confirm that estimates of N<sub>2</sub> fixation were varied with measurement method, maturity, soil, plant and so on.

## References

- Andersen MK, Hauggaard-Nielsen H, Ambus P, Jensen ES (2004) Biomass production, symbiotic nitrogen fixation and inorganic N use in dual and tri-component annual intercrops. *Plant and Soil* **266**, 273-287.
- Chalk P M (1996) Nitrogen transfer from legumes to cereals in intercropping. Proceedings of International Workshop on Dynamic of Roots and Nitrogen in Cropping systems of the Semi-Arid Tropics, November 21-25, 1996, Andhra Pradesh, India, pp: 351-374
- Geijersstam L, Martensson A (2006) Nitrogen fixation and residual effects of field pea intercropped with oats. *Acta Agricultura Scandinavica Section B-Soil and Plant Science* **56**, 186-196.
- Haby VA, Stout SA, Hons FM, Leonard AT (2006) Nitrogen fixation and transfer in a mixed stand of alfalfa and Bermuda grass. *Agronomy Journal* **98**, 890-898
- Peoples MB, Faizah AW, Rerkasem B, Herridge DF (Eds.) (1988) Methods for evaluating nitrogen fixation by nodulated legumes in the field. ACIAR Monograph No. 11, vii + 76 p. Canberra, Australia.
- Sakai R, et al., 2011. N transfer from green manure to lettuce in an intercropping cultivation system. *Acfa Scientiarum. Agronomy Maringa* **33**, 679-686.