



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

23rd International Grassland Congress

---

## Mineral Content of Course Cereals Roughages as well as Their Requirement in Dairy Animals

Asim Kumar Misra

*Indian Grassland and Fodder Research Institute, India*

K. K. Singh

*Indian Grassland and Fodder Research Institute, India*

M. M. Das

*Indian Grassland and Fodder Research Institute, India*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/23/2-1-2/18>

The 23rd International Grassland Congress (Sustainable use of Grassland Resources for Forage Production, Biodiversity and Environmental Protection) took place in New Delhi, India from November 20 through November 24, 2015.

Proceedings Editors: M. M. Roy, D. R. Malaviya, V. K. Yadav, Tejveer Singh, R. P. Sah, D. Vijay, and A. Radhakrishna

Published by Range Management Society of India

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## Mineral content of coarse cereals roughages as well as their requirement in dairy animals

Asim Kumar Misra\*, K. K. Singh, M. M. Das

ICAR-Indian Grassland and Fodder Research institute, Jhansi-284003, India

\*Corresponding author e-mail: asimkmisra@gmail.com

**Keywords:** Coarse cereal *stovers*, Dairy animals, Mineral contents, Mineral requirement

### Introduction

The necessity of mineral for growth, reproduction and production as well as normal physiological functions of animal body is well documented. In most of the tropical countries coarse cereal *stovers* are second major feed resource after wheat and paddy straw used in the livestock feeding. The concentration of macro and trace minerals in crop residues depends on various factors including species (Singh *et al.*, 2005). Further, limited information is available on coarse cereals *stovers* mineral content. Therefore, the present study was under taken to see the macro and micro mineral content of some coarse cereal *stovers* in relation to animal requirement.

### Materials and Methods

Samples of maize (*Zea mays*), sorghum (*Sorghum bicolor*) and Bajra (*Pennisetum glaucum*) *stover* were collected from major growing states *viz.*, Andhra Pradesh, Bihar, Gujrat, Haryana, Karnataka, Maharashtra, Uttar Pradesh, Rajasthan and Uttarakhand. After drying to a constant weight samples were ground and required quantity (1-1.25g) were digested in tri-acid mixture (HNO<sub>3</sub>:H<sub>2</sub>SO<sub>4</sub>:HClO<sub>4</sub> in the ratio of 15:2:4) and extractable aliquot were subjected to estimation of Calcium (Ca), Magnesium (Mg), Copper (Cu), Zinc (Zn), Manganese (Mn), Cobalt (Co) and Iron (Fe) using Atomic Absorption Spectroscopy model Varian AA240. Phosphorus (P) was analyzed using ammonium molybdate (AOAC, 1980). The data were analyzed statistically (Snedecor and Cochran, 1967).

### Results and Discussion

The calcium content was sufficient (0.45 to 0.64%) to meet the requirements of dairy animals in all the three course roughages. However, phosphorus (0.06-0.13%) and magnesium (0.05- 0.14%) were deficient invariably in all the *stover* samples of maize, sorghum and bajra as compared to the required level of 0.22 and 0.20% (McDowell, 1985), respectively (Table 1). There was a wide variation in Cu (6.49 to 8.25 ppm) and Zn (17.25 to 22.01 ppm) contents of maize, sorghum and bajra *stover* samples. Similar findings have also been reported by Garg *et al.* (2003). However, the concentration of iron in course roughage samples was many folds higher than its required levels of 50 ppm. Cobalt content was adequate in all samples while manganese was low in most of the coarse cereal roughage samples.

**Table 1:** Macro and micro mineral status of coarse cereal roughages

| Minerals        | Maize <i>stover</i> (n=30) | Sorghum <i>stover</i> (n=38) | Bajra <i>stover</i> (n=29) | SEM   |
|-----------------|----------------------------|------------------------------|----------------------------|-------|
| Calcium (%)     | 0.45                       | 0.64                         | 0.49                       | 0.02  |
| Phosphorus (%)  | 0.07                       | 0.06                         | 0.13                       | 0.01  |
| Magnesium (%)   | 0.14                       | 0.05                         | 0.10                       | 0.01  |
| Copper (ppm)    | 8.25                       | 6.66                         | 6.49                       | 0.37  |
| Zinc (ppm)      | 17.83                      | 17.25                        | 22.01                      | 0.65  |
| Manganese (ppm) | 46.66                      | 46.80                        | 40.88                      | 3.30  |
| Cobalt (ppm)    | 0.58                       | 0.12                         | 0.68                       | 0.01  |
| Iron (ppm)      | 370.40                     | 418.29                       | 160.79                     | 38.20 |

### Conclusion

From the results it is evident that only Ca and Co content were adequate in meeting the dietary need whereas P, Mg, Cu and Zn were deficient thus necessitate dietary supplementation and higher levels of Fe also require suitable counteractive mineral supplementation strategies in *stover* based feeding systems.

### References

- A.O.A.C. 1980. *Official Methods of Analysis*. 15<sup>th</sup> ed. Association of Official Analytical Chemists, USDA, Washington, DC.
- Garg, M. R., S. P. Arora, B. M. Bhanderi and P. L. Serasia. 2003. Trace mineral status of feeds and fodder in Dahod and Panchmahal districts of Gujrat. *Anim. Nutr. Feed Technol.* 3:27-30.
- McDowell, L. R. 1985. *Nutrient Requirement of Grazing Ruminants in Warm Climate*, 2<sup>nd</sup> ed. Academic Press Inc. California, USA.
- Singh, K. K., S. K. Nag, M. R. Garg and A. B. Mojumdar. 2005. Mineral status of some commonly used feeds and forages *vis-a-vis* their requirements. *Anim. Nutr. Feed Technol.* 5:171-177.
- Snedecor, G.W. and W. G. Cochran. 1967. *Statistical Methods*. 6<sup>th</sup> ed. Oxford and IBH Publishing Co. New Delhi.

### **Acknowledgement**

Authors are thankful to Director, Indian Grassland and Fodder Research Institute, Jhansi, for providing necessary facilities in carrying out the experimentation.