



Herbaceous Perennial Legumes: New Options for Pasture in Dryland Livestock Systems in Sub-Humid Mediterranean Chile

Carlos Ovalle
INIA, Chile

Soledad Espinoza
INIA, Chile

Viviana Barahona
INIA, Chile

Macarena Gerding
Universidad de Concepción, Chile

Alan Humpries
South Australian Research and Development Institute, Australia

See next page for additional authors

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/23/2-3-2/1>

The 23rd International Grassland Congress (Sustainable use of Grassland Resources for Forage Production, Biodiversity and Environmental Protection) took place in New Delhi, India from November 20 through November 24, 2015.

Proceedings Editors: M. M. Roy, D. R. Malaviya, V. K. Yadav, Tejveer Singh, R. P. Sah, D. Vijay, and A. Radhakrishna

Published by Range Management Society of India

Presenter Information

Carlos Ovalle, Soledad Espinoza, Viviana Barahona, Macarena Gerding, Alan Humpries, and Alejandro del Pozo

Herbaceous perennial legumes: new options for pasture in dryland livestock systems in sub-humid Mediterranean Chile

Carlos Ovalle^{1*}, Soledad Espinoza², Viviana Barahona³, Macarena Gerding⁴, Alan Humphries⁵ and Alejandro del Pozo⁶

¹Instituto de Investigaciones Agropecuarias (INIA), Olmué, Chile

²INIA, Cauquenes, Chile

³INIA, Cauquenes, INIA

⁴Universidad de Concepción, Chillán, Chile

⁵Sustainable Systems SARDI, Adelaide, Australia

⁶Universidad de Talca, Talca, Chile

*Corresponding author e-mail: covalle@inia.cl

Keywords: Dryland alfalfa, Mediterranean pastures, Perennial legumes

Introduction

In a climate with winter-dominant rainfall and long dry summers, like in Central Chile, the introduction and use of annual legumes and grasses, to improve productivity of livestock farming systems and to the rehabilitation of degraded agroecosystems have been the main strategies in the last twenty years. Unfortunately, the growth rate of annual legumes is low during autumn and winter, and also plant senescence starts by the end of October due to low soil water availability. As a consequence, the distribution of biomass production is mainly concentrated in spring when temperatures are moderate and soil water is available. In other Mediterranean-type climates, particularly in Western and Southern Australia, research on perennial legumes has been a priority to increase the availability, quality and timing of forage production, and to tackle environmental problems such as salinization and erosion (Humphries *et al.*, 2008). In this paper we present the first results of a new research line oriented to the introduction and assessment of perennial legumes with depth rooting systems in order to improve pasture productivity by extending the production period. The objectives were to introduce and evaluate the productivity and persistence of perennial forage legumes to summer drought in rainfed Mediterranean environments of Chile.

Materials and Methods

The field experiment was conducted at the Experimental Center of Cauquenes-INIA (35° 58' S, 72° 17' W; 140 m.a.s.l.), located in the interior dryland, of the sub-humid Mediterranean zone. The soil was a granitic classified as Ultic Palexeralfs. Soil pH (1:5 in water, 0–20 cm) was 7.0, the organic matter content was 1.6%. Long-term average annual precipitation is 695 mm, with a six-month (November to April) dry season.

Field experiments: Twenty genotypes from six perennial legume species were established in May 2012: nine cv of *Medicago sativa*, two *Hedysarum coronarium*, two *Bituminaria bituminosa*, one *Cullen australasicum*, three *Lotus tenuis*, and two *Lotus corniculatus* (Table 1). The experiment was evaluated till December 2014. Seedlings were grown in a glasshouse from seeds inoculated and lime pelleted with the appropriate strain of root nodule bacteria; *M. sativa* cultivars were inoculated with strain WSM2141, *Lotus* spp. with WSM166 and *Hedysarum coronarium* was inoculated with WSM1592. Plant survival (%) was assessed at the start of the second growing season (autumn) in order to determine the percentage of plants that survived the summer which had five months free of precipitation, by counting live plants in 3 m rows. Plant growth (height in cm) was evaluated at the end of the second winter (August) to measure growth and winter dormancy. Biomass was evaluated at the end of the growing season (December) in the first year of the experiment, and in September, November and January in the second and third seasons, by cutting the whole plot above 5 cm. Samples were oven-dried with forced air ventilation at 70°C until reaching a constant weight for dry matter (DM) determination.

Results and Discussion

Plant survival and winter growth:

A number of species, genus *Medicago*, *Psoralea* and *Cullen* experienced high survival rates (over 70%). On the contrary, cultivars of *Lotus tenuis*, *Lotus corniculatus*, and *Hedysarum coronarium*, did not persist properly. The low persistence of the latter two species could be due to failures in the persistence of rhizobia and therefore in the lack of nodulation (Fig. 1)

<i>Lotus tenuis</i>	Maitén		11	1,176	304	406	0	710	--	--	--	--
	Ecotipo 4		11	630	248	100	0	348	--	--	--	--
<i>Lotus corniculatus</i>	San Gabriel		19	1,317	942	584	1,644	3,170	1,710	--	--	1,710
	Quimey		19	1,184	1,131	691	1,088	2,909	1,209	--	--	1,209
5% LSD				396	618	991	964	2,003	651	1,674	962	238
					0.54	0.82	0.63	0.88	0.39	0.31	0.37	0.39
Correlation with known winter activity class (lucerne only, r value)												

Winter production: In 2013, the Australian-origin cvs did not differ statistically in winter production (August evaluation), and their production ranged 1,629 - 2,092 kg DM ha⁻¹ (Table 1); the Californian cvs WL326HQ and WL456HQ had lower winter growth compared to the Australian ones. However, in the third year (02 September 2014) no statistical differences were detected among lucerne cvs (Table 1). The production obtained at the end of winter is a resource of high value for a period in which the annual pasture virtually does not produce.

Summer production: SARDI Ten was significantly more productive (3,275 kg DM ha⁻¹) than the other lucerne cvs, which did not differ in summer production. Cultivars San Gabriel and Quimey of *Lotus corniculatus* produced between 1,088 and 1,644 kg DM ha⁻¹ (Table 1).

Conclusion

Several cv of alfalfa presented a high rate of survival of plants when they were establish under conditions of severe summer water stress prevailing in the interior dryland of the Mediterranean Sub-humid zone of Chile. On the contrary, the genotypes of the genus *Lotus*, *Hedysarum*, *Psoralea* and *Cullen*, did not persist in this environment.

Among alfalfa cvs, large differences in total production were not detected. A large group of cvs exceeded 8000 KgDMha⁻¹. Among the most productive cvs: Genesis, Sardi ten, Sardi seven, WL 903, Venus, and Aquarius were highlighted.

Cultivars with low or no winter dormancy were most productive and appropriate for rainfed Mediterranean conditions of Chile.

Enhanced distribution of forage production could be obtained with low winter dormancy alfalfa cvs. Indeed, the most productive allow obtaining 2 ton MS in late winter, which is none achieved by any of the annual legumes options, currently used in the production systems. Finally, the production of late spring early summer, would prolong the period of animal grazing for about two months, compared to that achieved with annual legumes.

References

Humphries A.W., X. G. Zhang, K. S. McDonald, R.A. Latta and G. C. Auricht. 2008. Persistence of diverse lucerne (*Medicago sativa* spp.) germplasm under farmer management across a range of soil types in southern Australia. *Australian Journal of Agricultural Research* 59:139-148.

Acknowledgement

This study was carried out thanks to the financial support of the Fondo Nacional de Desarrollo Científico and Tecnológico, Project FONDECYT N°1128829.