



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

XXIII International Grassland Congress

Effects of Abscisic Acid on Alleviation of Aluminum Toxicity in Alfalfa

Yuan An
Shanghai Jiao Tong University, China

Shengyin Wang
Shanghai Jiaotong University, China

Peng Zhou
Shanghai Jiaotong University, China

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/23/2-3-1/2>

The XXIII International Grassland Congress (Sustainable use of Grassland Resources for Forage Production, Biodiversity and Environmental Protection) took place in New Delhi, India from November 20 through November 24, 2015.

Proceedings Editors: M. M. Roy, D. R. Malaviya, V. K. Yadav, Tejveer Singh, R. P. Sah, D. Vijay, and A. Radhakrishna

Published by Range Management Society of India

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Effects of abscisic acid on alleviation of aluminum toxicity in alfalfa

Yuan, An*, Shengyin Wang, Peng Zhou

Shanghai Jiaotong University, Shanghai, P. R. China

*Corresponding author e-mail: anyuan@sjtu.edu.cn

Keywords: Aluminium toxicity, Callose, IAA, PIN, Root tip

Introduction

On a global scale, acidic soils cover an estimated 37.8 million km² of the earth's surface, and up to 50% of the world's potentially arable soils are acidic. Aluminum (Al) is highly abundant in acid soil conditions. At low (<5.5) pH, a toxic form of aluminum, Al³⁺, is solubilized from aluminosilicate clay minerals into soil solutions. High concentrations of Al³⁺ in soil solutions caused seriously damage to plants by inhibiting cell elongation and division in roots, leading to swollen root apices with poor or no root-hair development, and decreasing plant growth (Kochian *et al.*, 2005). Al toxicity represents one of the most important constraints for agricultural production in areas with acidic soils. Plant hormones are involved in plant adaptation to environmental stresses (Zhu *et al.*, 2013). However, limited information is available for hormones effects on plant tolerance to Al³⁺ toxicity. Alfalfa (*Medicago sativa* L.) is an important legume used as a forage crop worldwide. Al toxicity is a major factor limiting alfalfa production in soils with low pH. The purpose of this study was to investigate Al³⁺-induced abscisic acid (IAA) transport, distribution and their relation to Al³⁺-inhibition of root growth in alfalfa by exogenous foliar application of IAA on plants with or without apical buds.

Materials and Methods

Alfalfa seedlings with or without apical buds were grown in pH 4.5 nutrient solution containing Al at 0 (-Al) or 100 μM L⁻¹ AlCl₃·6H₂O (Al), and were foliar sprayed with water or IAA during 10-d treatment. The measurements of root length, root weight, above-ground biomass, root viability reflected by triphenyltetrazolium chloride (TTC) reduction, leaf membrane lipid peroxidation reflected by malondialdehyde (MDA) content, callose content, IAA concentration and expression of three genes related to IAA transport were taken.

Results and Discussion

Aluminum stress caused a significant reduction in plant growth, and significant increases in leaf membrane lipid peroxidation and callose production in root tips; Aluminum stress also induced a significant increase of IAA concentrations in root base, but a significant reduction in apical buds, root tips and shoot base, with most pronounced reduction in root tips when apical bud was removed. The variation of IAA distribution in Al³⁺-stressed plants would influence its function on regulating plant development and growth, while the decreased IAA concentrations in apical buds and root tips may directly restrict the growth of shoots and roots, and disturb physiological process in Al³⁺-stressed alfalfa. Exogenous application of IAA markedly alleviated the Al³⁺-induced inhibition of plant growth by increasing IAA concentrations in root tips, whereas the increased degree of IAA accumulations was significantly lower in the treatments with Al³⁺ than that without Al³⁺ and in plants with or without apical buds. In addition, Al³⁺ stress up-regulated *AUX1* and *PIN2*, but had no significant effects on *PIN1* gene. These results indicate that Al³⁺-induced reduction of plant growth could be associated with the inhibition of IAA synthesis in apical buds and IAA transport from shoot to root.

Data are means ±SE of three replicates. Bars with different letters indicate significant difference at P < 0.05 (Least significant difference test).




Fig. 1: Root length (A) and root fresh weight (FW) (B) of alfalfa seedlings grown for 1, 3 and 10 days on ½-strength Hoagland’s medium (pH 4.5) with treatments of without aluminum (-Al), aluminum stress (Al) (100 μM AlCl₃), aluminum with additional IAA (Al+IAA) (100 μM AlCl₃+IAA) and -Al with additional IAA (-Al+IAA).

Conclusion

Our results demonstrated that IAA played an important role in alfalfa responses to Al³⁺ stress. IAA synthesis and transport in alfalfa was sensitive to Al³⁺ stress, as manifested by quick changes in the distribution of IAA in apical buds, shoot base, root base and root tip of alfalfa in response to Al³⁺ stress. The significant reduction of IAA concentrations in apical buds and root tips could be the fundamental factor attributing to restricted plant growth by Al³⁺ stress. The reduction in IAA concentration in root tips was partly due to disturbance of the auxin signal transduction pathways by inhibiting IAA transport from shoots to roots, and promoting IAA transport from root tip to root base. IAA transport from shoots to roots could play key roles in plant tolerance to high Al³⁺ stress.

References

- Kochian, L.V., M. A. Pineros and O. A. Hoekenga. 2005. The physiology, genetics and molecular biology of plant aluminum resistance and toxicity. *Plant Soil* 274:175-195.
- Zhu, X.F., Z. W. Wang, F. Dong, G. J. Lei, Y. Z. Shi, G. X. Li and S. J. Zheng. 2013. Exogenous auxin alleviates cadmium toxicity in *Arabidopsis thaliana* by stimulating synthesis of hemicellulose 1 and increasing the cadmium fixation capacity of root cell walls. *Journal of Hazardous Materials* 263: 398-403.