



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

XXIII International Grassland Congress

---

## Effect of Planting Methods and Forage Crop Combinations on Fodder Productivity through Moisture Conservation

Kauthale Vitthal Keshav  
*BAIF-Development Research Foundation, India*

P. S. Takawale  
*BAIF-Development Research Foundation, India*

S. D. Patil  
*BAIF-Development Research Foundation, India*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/23/2-4-1/10>

The XXIII International Grassland Congress (Sustainable use of Grassland Resources for Forage Production, Biodiversity and Environmental Protection) took place in New Delhi, India from November 20 through November 24, 2015.

Proceedings Editors: M. M. Roy, D. R. Malaviya, V. K. Yadav, Tejveer Singh, R. P. Sah, D. Vijay, and A. Radhakrishna

Published by Range Management Society of India

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## **Effect of planting methods and forage crop combinations on fodder productivity through moisture conservation**

**Kauthale Vitthal Keshav\*, P. S. Takawale, S. D. Patil**

BAIF Development Research Foundation, Pune, India

\*Corresponding author e-mail: vkkauthale@gmail.com

**Keywords:** Benefit cost ratio, Forage yield, Grasses legume combinations, Moisture conservation, Planting methods

### **Introduction**

Out of 140.02 million ha cultivated area of the country, 76.76 million ha (54.82 %) is rainfed having soils with poor fertility and numerous physiochemical constraints. The productivity of animals is very low because of shortage of green fodder during most of the years. The poor feed quality and dry season feed shortage are the serious limitations for livestock production in rainfed areas & farmers maintain a large herd of animals to compensate for the low productivity of the livestock, which adds to the pressure on land and fodder resources (Pathak, 2005).

Western Maharashtra region suffers from very low rainfall with uncertainty and ill distribution. Occurrence of drought noticed once in three years. The water scarcity and delayed monsoon are major problems of crop production. The efforts needed to increase production through suitable agro forestry system fit to the land, climate and resources. Incorporation of trees and bushes particularly of fodder values in agricultural production system through a systematic manner will help in providing sufficient fodder to the farmers without affecting arable crop yields (Deb Roy, 1994).

*Cenchrus ciliaris* and *Dichanthium annulatum* are potential fast growing range grasses and having good regeneration capacity and can withstand moisture stress for fairly long time. *Desmanthus*, *Siratro* and *Stylosanthes* are legume fodder species which give nutritious fodder and could be grown under rainfed situations. Grasses and legume contributed to binding and soil stabilization due to their extensive root system. The overall impact of the soil conservation measures and pastoral systems resulted in increased infiltration rate, reduced soil loss and increased water stable aggregates. Therefore high productive, more palatable, perennial and persistent legumes like *Stylosanthes* and *Desmanthus* are thought to be the best suitable to overcome protein deficiency. Keeping the above points in view, the present study was undertaken to develop appropriate fodder production technology under rainfed conditions through moisture conservation.

### **Materials and Methods**

A field experiment was conducted at Central Research Station of BAIF Development Research Foundation, Urulikanchan, Pune during 2009-12 to study the effect of planting methods and forage crop combinations on fodder productivity through moisture conservation. The experiment was conducted in light gravel soil having initial soil status of 0.35 dS m<sup>-1</sup> EC, 7.36 pH, 0.30 % organic carbon, 128, 34 and 265 Kg ha<sup>-1</sup> NPK respectively. There were four combinations of grasses with legume viz. *Cenchrus ciliaris* + *Desmanthus virgatus*, *Cenchrus ciliaris* + *Stylosanthes seabrana*, *Dichanthium annulatum* + *Desmanthus virgatus*, *Dichanthium annulatum* + *Stylosanthes seabrana* in 1:1 proportion planted with two moisture conservation techniques viz. ridges and furrows & flat bed. Therefore, eight treatment combinations replicated three times in a factorial randomized block design (FRBD).

The grasses were established by planting seedlings at a spacing of 90 cm X 45 cm & legumes sown in between the two rows of grass, i.e. 1:1 proportion of 45 cm spacing maintained in between two rows. Soil fertility status (pH, EC, OC, N, P and K), microbial population count was determined using the methods described in (AOAC, 1995) and soil moisture data were recorded. The grasses and legumes were cut at 45-50 days interval. The growth and yield parameters were recorded at every cut and samples were analyzed in laboratory by using standard analytical methods. The pooled four years data was statistically analyzed.

### **Results and Discussion**

The pooled data on green fodder, dry matter and crude protein yields as influenced by different planting methods and forage crop combinations is given in Table 1.

**Table 1:** Effect of planting methods and forage crop combinations on green fodder, dry matter & crude protein yields

Treat. no.	Treatments details	Yield (ha <sup>-1</sup> )		
		Green fodder	Dry matter	Crude protein
T <sub>1</sub>	Ridges and furrows, <i>Cenchrus ciliaris</i> + <i>Desmanthus virgatus</i>	763.11	165.54	18.92
T <sub>2</sub>	Ridges and furrows, <i>Cenchrus ciliaris</i> + <i>Stylosanthes seabrana</i>	664.78	139.51	16.18
T <sub>3</sub>	Ridges and furrows, <i>Dichanthium annulatum</i> + <i>Desmanthus virgatus</i>	456.83	101.54	11.82
T <sub>4</sub>	Ridges and furrows, <i>Dichanthium annulatum</i> + <i>Stylosanthes seabrana</i>	423.19	92.86	11.05
T <sub>5</sub>	Flat bed, <i>Cenchrus ciliaris</i> + <i>Desmanthus virgatus</i>	724.34	156.75	17.75
T <sub>6</sub>	Flat bed, <i>Cenchrus ciliaris</i> + <i>Stylosanthes seabrana</i>	649.53	141.41	16.66
T <sub>7</sub>	Flat bed, <i>Dichanthium annulatum</i> + <i>Desmanthus virgatus</i>	413.31	89.67	10.62
T <sub>8</sub>	Flat bed, <i>Dichanthium annulatum</i> + <i>Stylosanthes seabrana</i>	360.60	82.24	10.01
SE (m) <sub>±</sub>		5.66	1.14	0.16
CD at 5%		15.68	3.16	0.44
CV %		19.26	18.14	17.41

Based on the pooled data for four years (2009-12), it was revealed that significantly increase in the green fodder, dry matter and crude protein yields (763.11, 165.54 and 18.92 ha<sup>-1</sup> per year respectively) in treatment combination of planting *Cenchrus ciliaris* with *Desmanthus virgatus* in 1:1 proportion on ridges and furrows over all the other planting methods & moisture conservation technique. Patel *et al.*, (2007) reported that Marvel grass may be fertilized with 60 kg ha<sup>-1</sup> and harvested once at the time of maturity to obtain higher dry fodder yield. Treatment combination of planting *Cenchrus ciliaris* with *Desmanthus virgatus* in 1:1 proportion on ridges and furrows showed significantly higher net monetary return of Rs. 55529 ha<sup>-1</sup> per year, benefit cost ratio of 2.72 and maize fodder equivalent yield of 308 qha<sup>-1</sup>. The experiment conducted by Kumar and Faruqui (2009) recorded maximum dry matter yield of 4.55 tonnes ha<sup>-1</sup> in Aonla + *Cenchrus ciliaris* + *Stylosanthes hamata* over the 3.92 tonnes ha<sup>-1</sup> in pure *Cenchrus ciliaris* + *Stylosanthes hamata* pasture system when applied with highest doses of nitrogen (60 kg ha<sup>-1</sup>). The pooled data on economics of the crops as influenced by different planting methods and forage crop combinations are given in Table 2.

**Table 2:** Effect of planting methods and forage crop combinations on economics of the crops

Treat. no.	Treatments details	Green fodder yield (qha <sup>-1</sup> )	Gross income (Rs.)	Cost of cultivation (Rs. ha <sup>-1</sup> )	Net monetary returns (Rs. ha <sup>-1</sup> )	Maize fodder equivalent yield (q ha <sup>-1</sup> )	Benefit : cost ratio
T <sub>1</sub>	Ridges and furrows, <i>Cenchrus ciliaris</i> + <i>Desmanthus virgatus</i>	763.11	87757	32228	55529.21	308.50	2.72
T <sub>2</sub>	Ridges and furrows, <i>Cenchrus ciliaris</i> + <i>Stylosanthes seabrana</i>	664.78	76449	31646	44803.50	248.91	2.42
T <sub>3</sub>	Ridges and furrows, <i>Dichanthium annulatum</i> + <i>Desmanthus virgatus</i>	456.83	52535	30548	21987.07	122.15	1.72
T <sub>4</sub>	Ridges and furrows, <i>Dichanthium annulatum</i> + <i>Stylosanthes seabrana</i>	423.19	48666	30000	18666.46	103.70	1.62
T <sub>5</sub>	Flat bed, <i>Cenchrus ciliaris</i> + <i>Desmanthus virgatus</i>	724.34	83299	32228	51071.43	283.73	2.58
T <sub>6</sub>	Flat bed, <i>Cenchrus ciliaris</i> + <i>Stylosanthes seabrana</i>	649.53	74696	31646	43050.20	239.17	2.36
T <sub>7</sub>	Flat bed, <i>Dichanthium annulatum</i> + <i>Desmanthus virgatus</i>	413.31	47531	30548	16982.72	94.35	1.56
T <sub>8</sub>	Flat bed, <i>Dichanthium annulatum</i> + <i>Stylosanthes seabrana</i>	360.60	41469	30000	11468.71	63.72	1.38
SE (m) <sub>±</sub>					651.07	3.62	0.02
CD at 5%					1803.46	10.03	0.05

## **Conclusion**

Based on the pooled data for four years (2009-12), it was concluded that the green fodder, dry matter & crude protein yields, net monetary returns, benefit cost ratio and maize fodder equivalent yield were significantly higher in treatment combination of planting *Cenchrus ciliaris* with *Desmanthus virgatus* in 1:1 proportion on ridges and furrows. The different planting methods + grasses & legume combinations also recorded higher level of organic carbon and available N, P, K in soil as compared initial status after completion of experiment under rainfed condition of Western Maharashtra. This technology is recommended for small & marginal farmers for cut & carries method of green forage under rainfed condition.

## **References**

- AOAC, 1995. *Official Methods of Analysis*, 16<sup>th</sup> Ed., Association of Analytical Chemists, Washington, USA.
- Deb Roy, R. 1994. *Principles and practices of Agroforestry for sustainable agriculture*, Proceedings of summer institute on advances in Agroforestry and its role, NRCAF, Jhansi, Pp: 61-74.
- Kumar, S. and S. A. Faruqui, 2009. Production potential and economic viability of food-forage based cropping system under irrigated conditions. *Indian J. Agron.* 54(1):36-41.
- Patel, P. C., L. N. Baraiya, D. V. Patel and R. H. Patel. 2007. Effect of fertilizer and cutting management on dry fodder yield of Marvel grass (*Dichanthium annulatum*). *Range Mgmt. & Agroforestry* 28(2): 414-415.
- Pathak, P. S. 2005. *Importance of feed resources for livestock production improvement in India*, IGFRI, Jhansi, Book chapter XVIII.

## **Acknowledgement**

The author is grateful to Director, Indian Grassland and Fodder Research Institute, Jhansi for providing necessary facilities and thanks to Project Coordinator, All India Coordinated Research Project on Forage Crops & Utilization for encouragement and valuable discussions.