



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

23rd International Grassland Congress

---

## Preliminary Trial to Establish Artificial Grassland in Tibet, China

Guowen Cui

*Northeast Agricultural University, China*

Bing Li

*Northeast Agricultural University, China*

Xiu-jie Yin

*Northeast Agricultural University, China*

Ming-jun Wang

*Northeast Agricultural University, China*

Wen-hua He

*Northeast Agricultural University, China*

*See next page for additional authors*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

---

Cui, Guowen; Li, Bing; Yin, Xiu-jie; Wang, Ming-jun; He, Wen-hua; and Liu, Sheng-yong, "Preliminary Trial to Establish Artificial Grassland in Tibet, China" (2020). *International Grassland Congress Proceedings*. 29. <https://uknowledge.uky.edu/igc/23/2-6-1/29>

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

---

**The 23rd International Grassland Congress (Sustainable use of Grassland Resources for Forage Production, Biodiversity and Environmental Protection) took place in New Delhi, India from November 20 through November 24, 2015.**

Proceedings Editors: M. M. Roy, D. R. Malaviya, V. K. Yadav, Tejveer Singh, R. P. Sah, D. Vijay, and A. Radhakrishna

Published by Range Management Society of India

**Presenter Information**

Guowen Cui, Bing Li, Xiu-jie Yin, Ming-jun Wang, Wen-hua He, and Sheng-yong Liu

## **Preliminary trial to establish artificial grassland in Tibet, China**

**Cui Guowen<sup>\*</sup>, Li Bing, Yin Xiu-jie, wang Ming-jun, He Wen-hua, Liu Sheng-yong**

Northeast Agricultural University, Harbin, China

<sup>\*</sup>Corresponding author e-mail : cgw603@163.com

**Keywords:** Artificial grassland, Mixture sowing, Sequential cropping; Tibet sorting cultivars

### **Introduction**

Tibet Autonomous Region with an average altitude of more than 4,000 m above sea level, altogether cover an area of 1.22 million km<sup>2</sup>, equivalent to one-eight of the total area of China. High mountains along with forests and rivers in Tibet consist of an important ecological barrier for protecting parts of inland ecosystems. Tibet is also one of the five major pastoral areas in China. The grassland in Tibet covers 0.83 million km<sup>2</sup>, but the grass is sparse and low, and with low yield in this area. There are nearly 2.8 million people in this region, while the area of arable land is less than 2.533 thousand km<sup>2</sup>. Nearly 40.5 million tons of hay are needed for about 45 million sheep units in Tibet (Yu *et al.*, 2010), while the actual amount is less than 20 million tons (including 15 million tons per year from natural grassland, and 2~3 million tons per year from the straw of naked barley). It exacerbating the degradation of natural grassland for the contradiction between livestock and grass. Research shows that the most effective measure to maintain local economic growth, coordinated development of ecological security and curb the further degradation of the grassland in Tibet is by planting high yielding and high quality forage crops, changing the traditional way of free grazing and implementing captive feeding gradually.

### **Materials and Methods**

The author participated in research and extension of cultivated artificial grassland in Tibet from 2005 to 2012. The preliminary results are as follow:

### **Results and Discussion**

**Variety selection:** The forage species suitable for planting in most areas at altitude below 4000 m in Tibet are oat (*Avena sativa* L.), common vetch (*Vicia sativa* L.), alfalfa (*Medicago sativa* L.), Forage-type triticale (*Triticosecale Wittmack*), forage-type rape (*Brassica napus*), sainfoin (*Onobrychis viciaefolia* Scop.), *Elymus dahuricus* Turcz. and *Elymus sibiricus* L.; while above 4000 m areas they are oat, forage-type Triticale and *Elymus nutans* Turcz.

**Sequential cropping:** The total hay yield can be up to 12.42 t·ha<sup>-1</sup> in the sequential cropping plot, and 43% higher than single cropped plot in the south central and other areas at altitude below 4000 m.

**The effect of soil repression after sowing on productivity of forage crops:** The evaporation of soil moisture in Tibet is numerous. The seedling emergence is often low due to insufficient soil compactness caused by the lack of rolling after sowing. With the “traditional plowing + roller compaction”, seedling emergence rate of oats, forage-type triticale and naked barley improved 18%, 8%, 30%, respectively, resulting in seed saving 117 kg·ha<sup>-1</sup>, 46 kg·ha<sup>-1</sup> and 244 kg·ha<sup>-1</sup>, respectively (Guo-Wen Cui *et al.*, 2014).

**Mixed cropping of oats and common vetch:** The highest crude protein yield which was obtained from the mixture group (seeding rate of oat was 210 kg·ha<sup>-1</sup> and common vetch was 75 kg·ha<sup>-1</sup>) was 1513.06 kg·ha<sup>-1</sup>. This was and 415%, 17% higher than that of single cropped oat and single cropped common vetch, respectively (Li Ji-kai, *et al.*, 2011).

**Seed production:** In order to reduce the cost of seed transport, and accelerate the planting of artificial grassland, seed production of oat base has to be established in Tibet. This will accelerate the spread and planting of artificial grassland.

### **References**

Guo-Wen Cui, Hong-Ying Li, Tao Sun, *et al.*, 2014. An experimental study of variety screening, sequential cropping, compaction and mixed cropping techniques for the cultivation of annual forage crops in agro-pastoral area of Tibet, China. *International Journal of Agriculture and Biology* 16: 97-103.

Li Ji-kai, Sun Tao, Wang Zha, *et al.*, 2011. Effects on mixture sowing ratio on the yield and quality of both Vetch and Oat in Tibet. *Acta Agrestia Sinica* 19(5): 830-833.

Yu L., L. Zhou, W. Liu and H. K. Zhou. 2010. Using remote sensing and GIS technologies to estimate grass yield and livestock carrying capacity of alpine grasslands in golog prefecture, China. *Pedosphere* 20: 342–351.