



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

23rd International Grassland Congress

---

## Biochemical Evaluation of Forage Sorghum for Stress Tolerance

Manoj Kumar Srivastava

*Indian Grassland and Fodder Research Institute, India*

Chandan Kumar Gupta

*Indian Grassland and Fodder Research Institute, India*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/23/4-1-2/11>

The 23rd International Grassland Congress (Sustainable use of Grassland Resources for Forage Production, Biodiversity and Environmental Protection) took place in New Delhi, India from November 20 through November 24, 2015.

Proceedings Editors: M. M. Roy, D. R. Malaviya, V. K. Yadav, Tejveer Singh, R. P. Sah, D. Vijay, and A. Radhakrishna

Published by Range Management Society of India

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## Biochemical evaluation of forage sorghum for stress tolerance

Manoj Kumar Srivastava\*, Chandan Kumar Gupta

ICAR-Indian Grassland and Fodder Research Institute, Jhansi-284003, India

\*Corresponding author e-mail: [srimanoj@hotmail.com](mailto:srimanoj@hotmail.com)

**Keywords:** Forage sorghum, Proline, Stress

### Introduction

Abiotic stresses are the major limiting factors in plant growth. With continuous increment in scarcity of water and increased salinization of soil and water, it is estimated that more than 50% of all arable lands will be affected by salinity and drought by 2050. Plants are often exposed to both, soil and atmospheric water deficit during their life cycle. The frequency and intensity of droughts is increasing as a result of global climate change. Understanding how plants respond to water stress is thus crucial for the estimation of impacts of climate change on crop productivity and ecosystem functioning. The knowledge about the mechanism adapted by plant to respond to drought, salt and co-occurring stresses can play an important role in stabilizing crop performance under drought and saline conditions. Sorghum (*Sorghum bicolor* L.) is an important crop in many parts of the world. It is utilized as food, fodder and several industrial purposes. In general, sorghum is known to be more tolerant to any stresses including heat, drought, salinity and flooding.

### Materials and Methods

Nine selected lines with two checks were sown in glass house and field for the evaluation of drought tolerance at morphological and biochemical levels. Fresh weight, dry weight, water content, and levels of various photosynthetic pigments were measured in leaves. Proline was measured by the acid-ninhydrin method of Bates *et al.* (1973). 0.5 g leaf tissue was homogenized in 5.0 ml sulphosalicylic acid (3%). The homogenate was centrifuged at 11,000 x g for 15 min at 4 °C. The clear supernatant was used for estimation of proline.

### Results and Discussion

Water content and specific leaf weight of sorghum leaves in control and water stressed conditions indicated that some of the sorghum lines have higher water retention capacity than others (Tables 1 and 2). These lines may be better performing under drought conditions. Line IG-03-285 was early flowering line identified both in green house and field conditions. This line may be used as early maturing short duration crop. This may also be utilized to develop short duration hybrid lines. Proline content in these lines was estimated to evaluate their capacity to tolerate the stress. The data is presented in Table 3. In general, the SGS lines have more proline as compared to non SGS lines.

Proline accumulation has been correlated with improved plant performance under stress. It was suggested that proline might confer a protective role by inducing biosynthesis of stress protective proteins. Exogenous application of proline induces the expression of ubiquitin, antioxidative enzymes and dehydrins. Glycine betaine is an osmoprotectant which accumulates under osmotic stress. Choline dehydrogenase catalyzes the oxidation of choline to glycine betaine *via* betaine aldehyde as intermediate. When flowering time was measured, it was found that flowering was delayed in all stay green lines. Furthermore, the stay green lines having delayed flowering showed a better performance and resistance to drought. Flowering was delayed in some SGS lines, which may be good candidates to be used as fodder in drought prone areas.

**Table 1:** Leaf morphological parameters of different lines of sorghum

| S. No. | Line      | Leaf length (cm) | Leaf width (cm) | Plant height (cm) |
|--------|-----------|------------------|-----------------|-------------------|
| 1      | Control 1 | 40.00 ± 8.89     | 1.67 ± 0.35     | 56.33± 1.53       |
| 2      | Control 2 | 51.00 ± 2.65     | 1.57 ± 0.25     | 55.00 ± 1.00      |
| 3      | IG-03-285 | 42.67 ± 8.39     | 1.23 ± 0.06     | 56.33 ± 1.53      |
| 4      | EC-507882 | 45.67 ± 5.13     | 2.23 ± 0.31     | 66.00 ± 2.00      |
| 5      | SGS-96    | 54.67 ± 7.57     | 2.67 ± 0.15     | 72.00 ± 2.00      |
| 6      | SGS-154   | 62.33 ± 4.04     | 2.67 ± 0.21     | 69.33 ± 3.06      |
| 7      | SGS-89    | 49.67 ± 7.09     | 2.53 ± 0.55     | 66.33 ± 1.53      |
| 8      | SGS-168   | 48.67 ± 1.53     | 1.90 ± 0.20     | 60.67 ± 0.58      |

|    |         |              |             |               |
|----|---------|--------------|-------------|---------------|
| 9  | SGS-33  | 42.33 ± 0.58 | 2.93 ± 0.06 | 51.00 ± 1.00  |
| 10 | SGS-45  | 76.67 ± 4.16 | 2.43 ± 0.38 | 91.67 ± 1.53  |
| 11 | SGS-118 | 73.67 ± 1.53 | 3.23 ± 0.47 | 114.33 ± 4.04 |

**Table 2:** Dry matter and water percentage in leaves of different lines of sorghum

| S. No. | Line      | Dry Matter % | Water %      |
|--------|-----------|--------------|--------------|
| 1      | Control 1 | 29.58 ± 2.09 | 70.42 ± 2.09 |
| 2      | Control 2 | 46.85 ± 7.94 | 53.15 ± 7.94 |
| 3      | IG-03-285 | 29.66 ± 6.23 | 70.34 ± 6.23 |
| 4      | EC-507882 | 28.55 ± 3.77 | 71.45 ± 3.77 |
| 5      | SGS-96    | 29.93 ± 2.29 | 70.07 ± 2.29 |
| 6      | SGS-154   | 25.48 ± 2.10 | 74.52 ± 2.10 |
| 7      | SGS-89    | 30.72 ± 1.33 | 69.28 ± 1.33 |
| 8      | SGS-168   | 32.82 ± 2.80 | 67.18 ± 2.80 |
| 9      | SGS-33    | 26.55 ± 0.79 | 73.45 ± 0.79 |
| 10     | SGS-45    | 18.68 ± 2.98 | 81.32 ± 2.98 |
| 11     | SGS-118   | 29.55 ± 1.34 | 70.45 ± 1.34 |

**Table 3:** Proline content in leaves of different lines of sorghum

| S. No. | Line      | Proline (µmoles/g Fr. Wt.) |
|--------|-----------|----------------------------|
| 1      | Control 1 | 0.226                      |
| 2      | Control 2 | 0.488                      |
| 3      | IG-03-285 | 0.448                      |
| 4      | EC-507882 | 0.988                      |
| 5      | SGS-96    | 0.485                      |
| 6      | SGS-154   | 1.330                      |
| 7      | SGS-89    | 0.572                      |
| 8      | SGS-168   | 0.514                      |
| 9      | SGS-33    | 0.157                      |
| 10     | SGS-45    | 1.618                      |
| 11     | SGS-118   | 0.496                      |

### Conclusion

The three SGS lines, SGS 33, SGS 45 and SGS 118 showed delayed flowering and higher growth among all the lines tested. These are being used to develop as stay green drought tolerant sorghum using breeding techniques.

### References

Bates L. S., R. P. Waldram and I. D. Tease. 1973. Rapid determination of free proline for water stress studies. *Plant Soil* 39: 205-208.

### Acknowledgement

The authors acknowledge the support and encouragement of Head-CI Division, Head-ST Division and Director IGFRI for present work.